

# Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ *Deutsch*

## ICD-10 C18.1: Appendix cancer

### Incidence and Mortality

Year of diagnosis	1998-2020
Patients	988
Diseases	989
Creation date	12/20/2021
Database export	12/20/2021
Population	4.95 m




Munich Cancer Registry  
Cancer Registry Bavaria - Upper Bavaria Regional Center  
at Klinikum Grosshadern/IBE  
Marchioninstr. 15  
Munich, 81377  
Germany

<https://www.tumorregister-muenchen.de/en>

[https://www.tumorregister-muenchen.de/en/facts/base/bC181\\_E-ICD-10-C18.1-Appendix-cancer-incidence-and-mortality.pdf](https://www.tumorregister-muenchen.de/en/facts/base/bC181_E-ICD-10-C18.1-Appendix-cancer-incidence-and-mortality.pdf)

### Index of figures and tables

Fig./Tbl.		Page
1	Annual cases, mult. malignancies, follow-up / yr	4
2	Incidence by year of diagnosis	7
3	Age distribution parameters by year of diagnosis	8
4	Age distribution by 5-year age group and sex	11
5	Age-specific incidence, proportion malignancies	12
6	Age distribution and age-specific incidence (chart)	13
6a	Age-specific incidence internationally (chart)	14
7	Standardized incidence ratio of further malignancies	15
8a	Map of cancer incidence (BRD-S) by county (chart)	17
8b	Standardized incidence ratio (SIR) by county (chart)	18
9a	Pts incident cohorts and mortality / yr	19
9b	Incidence and mortality by year of diagnosis	20
9c	Cancer-related deaths, death certification available / yr	21
10	Medians of age at death / yr	22
11	Mortality by year of death	24
12	Distribution of age at death	26
13	Age-specific mortality	27
14	Further malignancies in deaths	28
15	Age-specific mortality (first primaries)	30
16	Age-specific mortality (single primaries)	31
17	Age distribution and age-specific mortality (chart)	32
18a	Map of cancer mortality (BRD-S) by county (chart)	33
18b	Standardized mortality ratio (SMR) by county (chart)	34

**Global Statements about the statistics on the Internet –  
Baseline Statistics** (grey button ) , **Survival** (red button )

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut<sup>#</sup>, with a total of 4.69 million inhabitants, account for the frequency of cancer diseases<sup>##</sup> and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases<sup>###</sup> are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to [tumor@ibe.med.uni-muenchen.de](mailto:tumor@ibe.med.uni-muenchen.de).

Munich Cancer Registry, December 2021

- <sup>#</sup> Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.65 million to 4.10 in 2002, and to 4.69 million in 2007).
- <sup>##</sup> Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- <sup>###</sup> DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate.

### ICD-10 codes (ICD-10 2015) used for specifying cancer site

Code	Description
C18.1	Malignant neoplasm: Appendix

## INCIDENCE

Table 1

Cases with invasive cancer by year of diagnosis, proportions of further malignancies, deaths, and active follow-up (ALL PATIENTS)

Year of diagnosis	All cases n	Prop. at least 1 further malign. prior + synchron. %	Prop. at least 1 further malign. after %	Prop. deaths %	Prop. actively followed %
1998	12	8.3	8.5	58.3	100.0
1999	15	14.8	8.1	40.0	93.3
2000	17	13.6	7.9	35.3	94.1
2001	13	15.8	7.7	76.9	100.0
2002	35	17.4	7.7	51.4	97.1 #
2003	22	20.2	7.2	54.5	95.5
2004	20	20.1	7.1	45.0	100.0
2005	31	20.6	7.0	77.4	93.5
2006	25	20.5	6.5	52.0	88.0
2007	35	19.1	6.1	51.4	88.6 #
2008	32	19.1	5.6	50.0	100.0
2009	39	18.2	5.5	53.8	100.0
2010	44	17.4	5.2	50.0	95.5
2011	54	16.2	5.3	33.3	96.3
2012	77	15.1	4.4	26.0	94.8
2013	67	14.9	4.8	28.4	97.0
2014	74	15.5	5.0	29.7	97.3
2015	68	15.4	3.8	20.6	89.7
2016	66	16.1	3.3	21.2	98.5
2017	79	16.1	3.0	15.2	98.7
2018	66	16.8	1.9	25.8	100.0
2019	57	17.1	1.1	8.8	100.0
2020	41	17.5	0.0	4.9	100.0 ##
1998-2020	989	17.5	8.5	32.9	96.6

989 cases diagnosed 1998-2020 are related to a total of 988 patients. Currently, in 259 (26.2 %) of these 988 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 191 / 47 / 21 (19.3 % / 4.8 % / 2.1 %) patients exist having 2 / 3 / 4+ malignancies.

# The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

## Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retrieved from the respective headings.

How to interpret:

In 2018, a subgroup of 66 cases has been diagnosed, of which 16.8 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 1.9 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

Table 1a

Cases with invasive cancer by year of diagnosis, proportions of further malignancies, deaths, and active follow-up (MALES)

Year of diagnosis	Males n	Males %	Prop. at least 1 further malign. prior + synchron. %	Prop. at least 1 further malign. after %	Prop. deaths %	Prop. actively followed %
1998	7	58.3	14.3	10.8	42.9	100.0
1999	8	53.3	20.0	9.8	25.0	100.0
2000	7	41.2	22.7	9.8	57.1	100.0
2001	8	61.5	20.0	9.7	87.5	100.0
2002	15	42.9	15.6	9.6	40.0	100.0 #
2003	10	45.5	14.5	8.7	40.0	90.0
2004	9	45.0	12.5	8.4	44.4	100.0
2005	12	38.7	14.5	8.6	75.0	100.0
2006	13	52.0	15.7	8.3	61.5	100.0
2007	22	62.9	15.3	7.5	63.6	95.5 #
2008	16	50.0	16.5	7.4	43.8	100.0
2009	18	46.2	15.9	7.5	72.2	100.0
2010	22	50.0	16.2	7.2	45.5	90.9
2011	27	50.0	16.5	7.4	29.6	96.3
2012	32	41.6	15.9	6.9	34.4	93.8
2013	36	53.7	16.4	7.9	36.1	100.0
2014	38	51.4	17.7	8.2	36.8	97.4
2015	32	47.1	17.2	6.9	18.8	87.5
2016	32	48.5	18.1	6.3	25.0	100.0
2017	26	32.9	18.5	6.3	19.2	100.0
2018	26	39.4	19.2	2.8	26.9	100.0
2019	28	49.1	20.0	2.2	10.7	100.0
2020	20	48.8	20.5	0.0		100.0 ##
1998-2020	464	46.9	20.5	10.8	35.8	97.4

464 cases diagnosed 1998-2020 are related to a total of 463 patients. Currently, in 142 (30.7 %) of these 463 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 99 / 28 / 15 (21.4 % / 6.0 % / 3.2 %) patients exist having 2 / 3 / 4+ malignancies.

# The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

## Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retrieved from the respective headings.

How to interpret:

In 2018, a subgroup of 26 cases has been diagnosed, of which 19.2 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 2.8 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

Table 1b

Cases with invasive cancer by year of diagnosis, proportions of further malignancies, deaths, and active follow-up (FEMALES)

Year of diagnosis	Females n	Females %	Prop. at least 1 further malign. prior + synchron. %	Prop. at least 1 further malign. after %	Prop. deaths %	Prop. actively followed %
1998	5	41.7	0.0	6.5	80.0	100.0
1999	7	46.7	8.3	6.5	57.1	85.7
2000	10	58.8	4.5	6.2	20.0	90.0
2001	5	38.5	11.1	5.9	60.0	100.0
2002	20	57.1	19.1	6.0	60.0	95.0 #
2003	12	54.5	25.4	5.8	66.7	100.0
2004	11	55.0	27.1	6.0	45.5	100.0
2005	19	61.3	25.8	5.7	78.9	89.5
2006	12	48.0	24.8	5.0	41.7	75.0
2007	13	37.1	22.8	4.9	30.8	76.9 #
2008	16	50.0	21.5	4.0	56.3	100.0
2009	21	53.8	20.5	3.9	38.1	100.0
2010	22	50.0	18.5	3.6	54.5	100.0
2011	27	50.0	16.0	3.5	37.0	96.3
2012	45	58.4	14.3	2.2	20.0	95.6
2013	31	46.3	13.4	2.2	19.4	93.5
2014	36	48.6	13.5	2.5	22.2	97.2
2015	36	52.9	13.8	1.5	22.2	91.7
2016	34	51.5	14.1	1.2	17.6	97.1
2017	53	67.1	14.0	0.7	13.2	98.1
2018	40	60.6	14.7	1.1	25.0	100.0
2019	29	50.9	14.5	0.0	6.9	100.0
2020	21	51.2	14.9	0.0	9.5	100.0 ##
1998–2020	525	53.1	14.9	6.5	30.3	95.8

525 cases diagnosed 1998-2020 are related to a total of 525 patients. Currently, in 117 (22.3 %) of these 525 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 92 / 19 / 6 (17.5 % / 3.6 % / 1.1 %) patients exist having 2 / 3 / 4+ malignancies.

# The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

## Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retrieved from the respective headings.

How to interpret:

In 2018, a subgroup of 40 cases has been diagnosed, of which 14.7 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 1.1 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

Table 2

Incidence measures by year of diagnosis  
(with respect to registry area expansion from 2.65 to 4.10 m as of 2002,  
and from 4.10 to 4.94 m as of 2007, respectively)

Year of diagnosis	Males n	Females n	Males Inc. raw	Fem. Inc. raw	Males Inc. WS	Fem. Inc. WS	Males Inc. ES	Fem. Inc. ES	Males Inc. BRD-S	Fem. Inc. BRD-S
1998	7	5	0.6	0.4	0.4	0.3	0.5	0.4	0.5	0.3
1999	8	7	0.7	0.6	0.6	0.5	0.6	0.6	0.9	0.6
2000	7	10	0.6	0.8	0.4	0.6	0.5	0.7	0.6	0.9
2001	8	5	0.7	0.4	0.4	0.2	0.6	0.3	0.6	0.3
2002	15	20	0.8	1.0	0.5	0.7	0.7	0.9	0.8	1.0
2003	10	12	0.5	0.6	0.3	0.3	0.5	0.4	0.6	0.5
2004	9	11	0.5	0.6	0.3	0.4	0.4	0.5	0.5	0.5
2005	12	19	0.6	1.0	0.4	0.6	0.5	0.8	0.6	0.9
2006	13	12	0.7	0.6	0.4	0.4	0.6	0.5	0.7	0.5
2007	22	13	1.0	0.6	0.7	0.5	0.8	0.5	1.1	0.5
2008	16	16	0.7	0.7	0.5	0.5	0.6	0.5	0.7	0.6
2009	18	21	0.8	0.9	0.5	0.7	0.7	0.8	0.8	0.9
2010	22	22	1.0	0.9	0.6	0.7	0.8	0.8	0.9	0.8
2011	27	27	1.2	1.2	0.8	0.8	1.0	1.0	1.1	1.1
2012	32	45	1.4	1.9	1.0	1.9	1.2	1.9	1.3	2.0
2013	36	31	1.6	1.3	0.9	1.1	1.2	1.2	1.5	1.4
2014	38	36	1.6	1.5	1.2	1.1	1.4	1.2	1.6	1.4
2015	32	36	1.3	1.5	0.9	1.2	1.2	1.3	1.3	1.4
2016	32	34	1.3	1.4	0.9	1.1	1.1	1.2	1.3	1.4
2017	26	53	1.1	2.2	0.8	1.9	0.9	2.0	1.1	2.2
2018	26	40	1.1	1.6	0.6	1.1	0.8	1.3	0.9	1.5
2019	28	29	1.2	1.2	0.7	0.8	0.9	1.0	1.1	1.1
2020	20	21	0.8	0.8	0.6	0.6	0.7	0.7	0.8	0.8
1998-2020	464	525	1.0	1.1	0.7	0.8	0.8	0.9	1.0	1.1

The computation of the incidence measures includes all cancers, irrespective of first or subsequent malignancy.

Table 3

Age distribution parameters by year of diagnosis (ALL PATIENTS)

Year of diagnosis	Cases n	Std.		Median						
		Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	12	54.8	20.9	13.2	87.9	32.1	43.3	55.8	63.8	86.0
1999	15	46.2	18.7	24.9	80.5	26.2	27.4	43.0	59.8	75.1
2000	17	50.6	13.7	24.7	81.6	32.7	41.6	49.5	60.1	64.1
2001	13	60.9	17.1	34.3	88.5	35.2	54.0	62.6	68.9	84.4
2002	35	56.4	19.1	17.7	90.9	29.9	37.0	60.3	72.8	78.6
2003	22	59.8	17.9	23.5	88.5	32.4	53.6	59.7	77.1	79.4
2004	20	51.9	21.3	13.8	81.0	22.4	33.8	57.8	71.9	75.9
2005	31	63.5	14.5	16.1	89.9	46.9	56.7	67.2	71.8	76.7
2006	25	61.2	11.9	40.8	81.2	44.8	50.6	63.7	70.9	76.0
2007	35	52.6	22.0	13.4	84.4	19.1	38.6	60.0	70.7	78.7
2008	32	54.3	19.7	18.9	86.2	27.5	41.7	52.5	72.0	80.6
2009	39	58.7	21.3	12.4	91.8	24.1	49.6	59.1	76.4	84.7
2010	44	59.6	19.6	14.9	94.1	31.8	46.3	64.6	70.8	81.9
2011	54	56.4	20.9	15.5	88.8	23.3	41.0	57.7	73.0	83.8
2012	77	48.8	21.7	9.7	89.9	18.5	28.7	49.0	66.4	77.3
2013	67	53.5	19.7	15.7	83.5	23.6	35.1	56.5	72.0	76.6
2014	74	54.0	19.8	15.8	88.6	24.4	39.3	54.5	71.4	78.1
2015	68	51.6	19.0	11.4	82.6	24.7	40.3	52.7	67.7	76.3
2016	66	53.2	20.2	13.8	93.8	21.2	42.6	53.8	64.7	79.4
2017	79	50.2	20.6	9.4	84.4	20.4	29.9	54.8	68.8	79.0
2018	66	58.8	18.0	14.3	90.4	31.6	51.1	60.2	74.1	78.1
2019	57	58.5	19.9	17.7	93.2	29.1	47.1	62.5	73.8	84.8
2020	41	55.1	20.1	18.5	87.8	23.9	39.6	60.3	68.4	79.8
1998-2020	989	54.8	19.8	9.4	94.1	24.6	40.5	57.0	70.5	79.2



Table 3a

Age distribution parameters by year of diagnosis (MALES)

Year of diagnosis	Cases n	Std.		Median						
		Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	7	52.6	12.2	32.1	64.1	32.1	40.0	56.0	63.6	64.1
1999	8	45.5	22.5	24.9	80.5	24.9	26.8	36.1	66.3	80.5
2000	7	50.1	9.9	37.4	64.0	37.4	39.0	49.5	59.2	64.0
2001	8	59.2	16.7	34.3	88.5	34.3	48.0	60.0	67.5	88.5
2002	15	56.9	14.9	27.1	79.8	35.0	53.4	59.7	65.9	74.9
2003	10	58.0	13.1	32.4	78.0	40.6	53.6	57.4	60.0	77.6
2004	9	53.4	19.8	27.8	76.4	27.8	32.9	58.2	74.3	76.4
2005	12	64.7	11.9	34.1	77.0	53.3	59.5	69.5	71.5	74.8
2006	13	62.0	11.2	44.8	78.3	49.4	51.7	64.2	70.9	76.0
2007	22	56.4	21.3	15.8	84.4	24.8	39.2	62.5	70.7	81.3
2008	16	51.8	16.7	19.3	80.6	27.5	38.6	55.1	63.4	71.9
2009	18	62.1	17.5	12.4	84.7	40.8	53.2	63.8	75.4	83.8
2010	22	63.0	15.0	27.9	86.5	43.5	57.4	64.9	70.8	81.4
2011	27	57.6	17.7	15.5	85.6	34.7	42.6	62.9	68.7	82.1
2012	32	54.8	17.2	9.7	79.8	28.8	45.0	57.7	67.6	75.3
2013	36	60.7	16.7	19.4	83.4	35.6	48.1	64.4	74.6	79.3
2014	38	55.1	19.3	20.3	88.6	24.1	39.3	57.4	69.1	82.2
2015	32	51.4	16.3	24.7	82.6	29.0	40.9	51.1	63.9	72.3
2016	32	54.1	18.2	15.8	90.7	32.0	44.8	53.0	67.8	78.9
2017	26	55.3	22.7	12.9	84.4	21.0	34.1	59.1	72.4	79.4
2018	26	61.8	17.8	14.3	90.4	40.8	54.1	61.9	75.9	77.1
2019	28	60.6	19.2	22.6	91.4	29.1	48.2	63.5	77.3	86.5
2020	20	55.8	20.4	18.5	87.8	22.7	40.2	60.0	71.0	80.4
1998-2020	464	57.0	17.8	9.7	91.4	29.3	45.3	58.6	70.8	78.6

Table 3b

Age distribution parameters by year of diagnosis (FEMALES)

Year of diagnosis	Cases n	Std.		Median						
		Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	5	57.9	30.9	13.2	87.9	13.2	46.7	55.5	86.0	87.9
1999	7	47.0	14.8	26.9	64.7	26.9	28.4	52.7	59.8	64.7
2000	10	51.0	16.4	24.7	81.6	28.7	41.6	49.3	60.8	72.8
2001	5	63.5	19.4	35.2	84.4	35.2	56.7	62.6	78.5	84.4
2002	20	56.1	22.2	17.7	90.9	26.3	35.0	62.6	74.0	83.3
2003	12	61.3	21.6	23.5	88.5	30.6	46.3	64.0	79.1	84.0
2004	11	50.6	23.4	13.8	81.0	17.0	34.6	57.4	70.4	73.4
2005	19	62.8	16.2	16.1	89.9	45.7	54.2	63.8	72.2	83.1
2006	12	60.3	13.0	40.8	81.2	43.1	49.0	62.3	70.7	72.3
2007	13	46.2	22.4	13.4	76.0	17.8	27.1	42.9	68.5	74.4
2008	16	56.7	22.5	18.9	86.2	22.8	43.9	49.2	77.1	85.5
2009	21	55.8	24.1	15.9	91.8	24.1	35.1	58.0	76.5	86.7
2010	22	56.3	23.2	14.9	94.1	26.0	35.1	64.2	70.7	85.4
2011	27	55.3	23.9	16.5	88.8	18.7	33.0	57.0	79.8	87.0
2012	45	44.6	23.7	13.7	89.9	16.4	21.9	40.8	62.4	78.1
2013	31	45.1	19.7	15.7	83.5	22.7	28.6	39.3	62.8	72.7
2014	36	52.8	20.5	15.8	86.7	24.4	35.4	50.7	72.2	76.6
2015	36	51.8	21.3	11.4	81.3	18.9	38.6	55.6	70.1	78.6
2016	34	52.3	22.1	13.8	93.8	20.3	34.5	54.6	64.7	79.4
2017	53	47.7	19.3	9.4	81.3	20.4	29.9	51.3	59.3	71.9
2018	40	56.8	18.1	19.3	88.0	30.2	46.9	60.2	71.8	78.7
2019	29	56.4	20.7	17.7	93.2	27.8	45.8	62.5	68.2	84.8
2020	21	54.5	20.4	20.7	87.4	26.2	38.0	60.3	67.4	79.8
1998–2020	525	52.8	21.2	9.4	94.1	21.2	35.1	54.8	69.7	79.8

Table 4

Age distribution by 5-year age group and sex for period 2007–2020


Age at diagnosis Years	Cases			Males			Females		
	n	%	Cum.%	n	%	Cum.%	n	%	Cum.%
0–4									
5–9	2	0.3	0.3	1	0.3	0.3	1	0.2	0.2
10–14	9	1.1	1.4	3	0.8	1.1	6	1.4	1.7
15–19	40	5.0	6.4	8	2.1	3.2	32	7.5	9.2
20–24	43	5.4	11.8	16	4.3	7.5	27	6.4	15.6
25–29	44	5.5	17.3	15	4.0	11.5	29	6.8	22.4
30–34	30	3.8	21.0	11	2.9	14.4	19	4.5	26.9
35–39	35	4.4	25.4	16	4.3	18.7	19	4.5	31.4
40–44	44	5.5	30.9	23	6.1	24.8	21	5.0	36.3
45–49	67	8.4	39.3	28	7.5	32.3	39	9.2	45.5
50–54	63	7.9	47.2	34	9.1	41.3	29	6.8	52.4
55–59	74	9.3	56.4	41	10.9	52.3	33	7.8	60.1
60–64	72	9.0	65.5	34	9.1	61.3	38	9.0	69.1
65–69	70	8.8	74.2	37	9.9	71.2	33	7.8	76.9
70–74	61	7.6	81.9	37	9.9	81.1	24	5.7	82.5
75–79	75	9.4	91.2	38	10.1	91.2	37	8.7	91.3
80–84	38	4.8	96.0	24	6.4	97.6	14	3.3	94.6
85+	32	4.0	100.0	9	2.4	100.0	23	5.4	100.0
All ages	799	100.0		375	100.0		424	100.0	

Table 5


Age-specific incidence and proportion of all cancers for period 2007-2020

Age at diagnosis Years	Males n	Females n	Males Age- spec. incid.	Females Age- spec. incid.	Males Prop.all cancers n=153686 %	Females Prop.all cancers n=155051 %
0- 4						
5- 9	1	1	0.1	0.1	0.9	1.0
10-14	3	6	0.2	0.4	2.2	4.7
15-19	8	32	0.5	2.0	2.5	12.1
20-24	16	27	0.8	1.4	2.5	5.2
25-29	15	29	0.7	1.3	1.6	2.4
30-34	11	19	0.5	0.8	0.8	0.9
35-39	16	19	0.7	0.8	0.9	0.5
40-44	23	21	0.9	0.9	0.8	0.3
45-49	28	39	1.0	1.5	0.6	0.4
50-54	34	29	1.3	1.2	0.4	0.2
55-59	41	33	1.9	1.5	0.3	0.2
60-64	34	38	1.9	2.0	0.2	0.2
65-69	36	33	2.2	1.8	0.1	0.2
70-74	37	24	2.5	1.4	0.1	0.1
75-79	38	37	3.1	2.5	0.2	0.2
80-84	24	14	3.3	1.3	0.2	0.1
85+	9	23	1.9	2.2	0.1	0.1
All ages	374	424			0.2	0.3
Incidence						
Raw			1.1	1.3		
WS			0.8	1.0		
ES			1.0	1.1		
BRD-S			1.1	1.2		

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).



**Figure 6.** Age distribution (males: mean=57.0 yrs, median=58.8 yrs; females: mean=51.7 yrs, median=52.1 yrs) and age-specific incidence.



**Figure 6a.** Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).

Reference:  
 Surveillance, Epidemiology, and End Results (SEER) Program SEER\*Stat Database: Incidence - SEER 21 Regs Research Data, released April 2021, based on the November 2020 submission. <http://www.seer.cancer.gov>.

Table 7a

Standardized incidence ratio (SIR, with 95% confidence limits),  
excess absolute risk (EAR) and DCO rate of further malignancies  
for period 1998–2020

## MALES

Diagnosis	Observed n	Expected n	SIR	CI 95%	CI 95%	EAR	DCO %
C07–C08 Salivary gland	1	0.0	22.7	0.6	126.4	5.6	
C12–C13 Hypopharynx	1	0.1	8.5	0.2	47.4	5.2	
C15 Oesophagus	5	0.4	12.2	4.0	28.6 #	27.0	
C17 Small intestine	5	0.1	40.0	13.0	93.3 #	28.6	
C18 Colon	19	1.7	11.0	6.6	17.2 #	101.4	
C19–C20 Rectum	7	1.0	7.0	2.8	14.4 #	35.2	
C25 Pancreas	1	0.7	1.4	0.0	7.6	1.6	
C33–C34 Lung	3	2.2	1.4	0.3	4.0	4.6	
C40–C41 Bone	1	0.0	54.9	1.4	305.7 #	5.8	
C43 Malign. melanoma	1	0.9	1.1	0.0	5.9	0.4	
C61 Prostate	19	5.1	3.8	2.3	5.9 #	81.8	
C64 Kidney	4	0.7	6.0	1.6	15.3 #	19.6	
C66 Ureter	1	0.0	20.2	0.5	112.7	5.6	
C67 Bladder	1	0.8	1.2	0.0	6.7	1.0	100.0
C70–C72 CNS cancer	1	0.3	4.0	0.1	22.0	4.4	
C74–C80 Cancer others	1	0.0	28.8	0.7	160.2	5.7	
C82–C85 NHL	3	0.8	3.8	0.8	11.1	13.0	
C91–C96 Leukaemia	1	0.3	3.6	0.1	19.8	4.2	
Not observed	0	3.6	0.0	0.0	1.0	-21.1	
All further malignancies	75	18.9	4.0	3.1	5.0 #	329.3	1.3
Patients		458					
Median age at next malignancy (years)		72.2					
Person-years		1704					
Mean observation time (years)		3.7					
Median observation time (years)		2.1					

# The occurrence of further specified malignancy is statistically significant.

Table 7b

Standardized incidence ratio (SIR, with 95% confidence limits),  
excess absolute risk (EAR) and DCO rate of further malignancies  
for period 1998–2020


## FEMALES

Diagnosis	Observed n	Expected n	SIR	CI 95%	CI 95%	EAR	DCO %
C16 Stomach	3	0.4	8.5	1.8	24.9 #	15.2	33.3
C17 Small intestine	3	0.1	46.7	9.6	136.6 #	16.8	
C18 Colon	9	1.0	8.7	4.0	16.6 #	45.7	
C19–C20 Rectum	3	0.4	7.1	1.5	20.6 #	14.8	
C25 Pancreas	2	0.5	4.0	0.5	14.5	8.6	
C33–C34 Lung	5	0.9	5.8	1.9	13.5 #	23.7	
C43 Malign. melanoma	2	0.5	4.0	0.5	14.3	8.6	
C50 Breast	7	3.8	1.8	0.7	3.8	18.1	14.3
C56 Ovary	6	0.5	13.1	4.8	28.5 #	31.8	
C64 Kidney	3	0.2	12.2	2.5	35.5 #	15.8	
C67 Bladder	2	0.2	9.7	1.2	35.1 #	10.3	
C81 Hodgkin lymphoma	1	0.0	27.5	0.7	153.4	5.5	
C82–C85 NHL	1	0.4	2.4	0.1	13.1	3.3	
C91–C96 Leukaemia	1	0.2	6.0	0.2	33.4	4.8	
Not observed	0	2.7	0.0	0.0	1.4	-15.3	
All further malignancies	48	11.8	4.1	3.0	5.4 #	207.9	4.2
Patients		519					
Median age at next malignancy (years)		66.4					
Person-years		1742					
Mean observation time (years)		3.4					
Median observation time (years)		1.8					


# The occurrence of further specified malignancy is statistically significant.



Average incidence (Germany 1987 standard population) 2007 - 2020: Males




Average incidence (Germany 1987 standard population) 2007 - 2020: Females




**Figure 8a.** Map of cancer incidence (german standard population) by county averaged for period 2007 to 2020. According to their individual incidence rates, the counties are displayed in different red and blue hues, being the fine white color attributed to the population mean (males 1.1/100,000 WS N=374, females 1.2/100,000 WS N=424).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,727 female residents (averaged) in the period from 2007 to 2020 a total of 8 women were identified with newly diagnosed appendix cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.9/100,000 (german standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.3 and 2.1/100,000.

Standardized incidence ratio (SIR) 2007 - 2020: Males



Standardized incidence ratio (SIR) 2007 - 2020: Females



**Figure 8b.** Map of standardized incidence ratio (SIR) by county averaged for period 2007 to 2020. According to their individual SIR values, the counties are displayed in different red and blue hues, being the fine white color attributed to the population overall of 1.0 (males N=374, females N=424).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,153 female residents (averaged) in the period from 2007 to 2020 a total of 8 women were identified with newly diagnosed appendix cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.68. Though, the value of this parameter may vary with an underlying probability of 99% between 0.22 and 1.58, and is therefore not statistically striking.

## MORTALITY

Table 9a

Annual cohorts: Incident cancers, follow-up status,  
and deaths among the annual cohorts

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002,  
and from 4.10 to 4.94 m as of 2007, respectively)

Year of diagnosis	Incident cases n	Prop. actively followed %	Deaths n	Prop. deaths %	Prop. deaths with death certific. %
1998	12	100.0	7	58.3	100.0
1999	15	93.3	6	40.0	83.3
2000	17	94.1	6	35.3	83.3
2001	13	100.0	10	76.9	100.0
2002	35	97.1	18	51.4	100.0
2003	22	95.5	12	54.5	100.0
2004	20	100.0	9	45.0	100.0
2005	31	93.5	24	77.4	95.8
2006	25	88.0	13	52.0	92.3
2007	35	88.6	18	51.4	88.9
2008	32	100.0	16	50.0	87.5
2009	39	100.0	21	53.8	95.2
2010	44	95.5	22	50.0	100.0
2011	54	96.3	18	33.3	83.3
2012	77	94.8	20	26.0	85.0
2013	67	97.0	19	28.4	89.5
2014	74	97.3	22	29.7	86.4
2015	68	89.7	14	20.6	85.7
2016	66	98.5	14	21.2	92.9
2017	79	98.7	12	15.2	66.7
2018	66	100.0	17	25.8	64.7
2019	57	100.0	5	8.8	100.0
2020	41	100.0	2	4.9	100.0
1998-2020	989	96.6	325	32.9	89.8

Table 9b

Annual cohorts of incident cancers and deaths,  
and cases deceased within the same year of being diagnosed with cancer

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002,  
and from 4.10 to 4.94 m as of 2007, respectively)

Year of diagnosis/ death	Incident cases n	Deaths n	Deaths in same year n	Prop. deaths in same year %
1998	12	5	2	16.7
1999	15	5	2	13.3
2000	17			
2001	13	5	2	15.4
2002	35	5	2	5.7
2003	22	9	2	9.1
2004	20	7		
2005	31	11	5	16.1
2006	25	10	1	4.0
2007	35	4	1	2.9
2008	32	15	1	3.1
2009	39	14	4	10.3
2010	44	24	4	9.1
2011	54	15	2	3.7
2012	77	18	6	7.8
2013	67	21	2	3.0
2014	74	22	3	4.1
2015	68	26	1	1.5
2016	66	23	1	1.5
2017	79	22	1	1.3
2018	66	28	2	3.0
2019	57	27	1	1.8
2020	41	24		
1998–2020	989	340	45	4.6

Table 9c

Annual cohorts of deaths, and proportion of cancer-related and non-cancer-related deaths

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.94 m as of 2007, respectively)

Year of death	Deaths n	Prop. cancer- related %	Prop. non-cancer- related %	Prop. cancer recorded on death certificate %
1998	5	80.0	20.0	80.0
1999	5	100.0		100.0
2000				
2001	5	100.0		100.0
2002	5	60.0	40.0	60.0
2003	9	66.7	33.3	88.9
2004	7	85.7	14.3	85.7
2005	11	72.7	27.3	80.0
2006	10	80.0	20.0	88.9
2007	4	100.0		100.0
2008	15	86.7	13.3	78.6
2009	14	78.6	21.4	100.0
2010	24	70.8	29.2	79.2
2011	15	100.0		93.3
2012	18	72.2	27.8	82.4
2013	21	85.7	14.3	85.7
2014	22	68.2	31.8	72.7
2015	26	69.2	30.8	70.8
2016	23	60.9	39.1	59.1
2017	22	86.4	13.6	76.2
2018	28	53.6	46.4	57.9
2019	27	51.9	48.1	72.7
2020	24	50.0	50.0	73.9
1998–2020	340	71.5	28.5	78.0

Table 10a

Medians of age at death according to the grouping in Table 9  
MALES

Year of death	Deaths n	Age at death (all causes) Years	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate) Years
1998	1	56.4	56.4		56.4
1999	2	71.5	71.5		71.5
2000					
2001	4	64.3	64.3		64.3
2002	2	71.0	71.0		63.2
2003	2	60.1	60.1		60.1
2004	4	69.6	69.6		82.0
2005	4	75.1	74.9	79.9	74.9
2006	5	76.4	58.5	76.8	58.5
2007					
2008	10	65.7	65.4	81.2	65.1
2009	5	78.9	78.9		78.9
2010	7	78.1	75.9	78.1	75.9
2011	8	67.7	67.7		67.8
2012	8	79.2	79.8	71.4	79.8
2013	13	71.9	66.8	71.9	66.8
2014	11	70.8	72.9	67.2	72.9
2015	10	67.1	67.1		66.3
2016	15	71.1	71.2	71.1	71.2
2017	11	81.0	78.9	87.3	81.0
2018	19	67.3	64.0	72.0	64.0
2019	11	68.5	57.6	71.5	54.5
2020	19	76.1	75.0	81.3	78.3
1998–2020	171	72.4	71.0	76.4	73.5

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 10b

Medians of age at death according to the grouping in Table 9  
FEMALES

Year of death	Deaths n	Age at death (all causes) Years	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate) Years
1998	4	85.9	85.8	88.1	85.8
1999	3	72.9	72.9		76.9
2000					
2001	1	49.6	49.6		49.6
2002	3	91.0	95.7	76.9	79.2
2003	7	74.8	76.0	74.8	76.0
2004	3	79.7	79.9	79.7	79.7
2005	7	70.4	70.4	70.7	70.4
2006	5	66.8	66.8		66.8
2007	4	68.3	68.3		68.3
2008	5	72.2	72.2		73.8
2009	9	81.7	80.1	84.7	81.7
2010	17	87.4	80.8	89.4	87.4
2011	7	68.4	68.4		68.4
2012	10	79.1	71.8	83.8	78.1
2013	8	67.6	67.6		67.6
2014	11	77.4	49.8	79.2	65.2
2015	16	75.7	72.6	85.5	72.6
2016	8	77.2	77.2	67.8	77.7
2017	11	78.8	78.8		68.1
2018	9	83.7	83.7	80.8	85.6
2019	16	79.3	64.4	83.6	64.4
2020	5	81.9	73.6	92.2	73.6
1998–2020	169	77.7	74.8	83.6	75.8

By 2018, Bavarians' life expectancy at birth is estimated at 79.3 years for boys and 83.8 years for girls.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 11a

Mortality measures (cancer-related death) and mortality-incidence-index  
by year of death

MALES

Year of death	Deaths n	Mort. raw	MI-Index raw	Mort. WS	MI-Index WS	Mort. ES	MI-Index ES	Mort. BRD-S	MI-Index BRD-S
1998	1	0.1	0.14	0.0	0.11	0.1	0.13	0.1	0.13
1999	2	0.2	0.25	0.1	0.18	0.2	0.24	0.3	0.31
2000									
2001	4	0.3	0.50	0.2	0.50	0.3	0.50	0.3	0.52
2002	2	0.1	0.13	0.1	0.10	0.1	0.12	0.1	0.16
2003	2	0.1	0.20	0.1	0.17	0.1	0.17	0.1	0.20
2004	4	0.2	0.44	0.1	0.44	0.2	0.46	0.3	0.50
2005	3	0.2	0.25	0.1	0.24	0.1	0.25	0.2	0.30
2006	3	0.2	0.23	0.1	0.24	0.1	0.26	0.2	0.26
2007									
2008	8	0.4	0.50	0.2	0.42	0.3	0.44	0.3	0.49
2009	5	0.2	0.28	0.1	0.15	0.1	0.21	0.3	0.32
2010	6	0.3	0.27	0.1	0.18	0.2	0.22	0.2	0.26
2011	8	0.4	0.30	0.2	0.24	0.3	0.26	0.3	0.27
2012	7	0.3	0.22	0.1	0.13	0.2	0.17	0.3	0.24
2013	10	0.4	0.28	0.2	0.24	0.3	0.27	0.4	0.27
2014	8	0.3	0.21	0.2	0.14	0.2	0.17	0.3	0.18
2015	10	0.4	0.31	0.2	0.23	0.3	0.28	0.4	0.30
2016	8	0.3	0.25	0.2	0.18	0.2	0.21	0.3	0.24
2017	8	0.3	0.31	0.1	0.16	0.2	0.23	0.3	0.27
2018	10	0.4	0.38	0.3	0.41	0.3	0.40	0.4	0.42
2019	7	0.3	0.26	0.2	0.23	0.2	0.26	0.3	0.25
2020	8	0.3	0.40	0.1	0.24	0.2	0.30	0.3	0.35
1998-2020	124	0.3	0.27	0.1	0.20	0.2	0.24	0.3	0.27



Table 11b

Mortality measures (cancer-related death) and mortality-incidence-index  
by year of death

FEMALES

Year of death	Deaths n	Mort. raw	MI-Index raw	Mort. WS	MI-Index WS	Mort. ES	MI-Index ES	Mort. BRD-S	MI-Index BRD-S
1998	3	0.3	0.60	0.0	0.15	0.1	0.28	0.2	0.47
1999	3	0.3	0.43	0.1	0.25	0.2	0.33	0.2	0.43
2000									
2001	1	0.1	0.20	0.1	0.40	0.1	0.33	0.1	0.29
2002	1	0.1	0.05	0.0	0.01	0.0	0.02	0.0	0.02
2003	4	0.2	0.33	0.1	0.23	0.1	0.27	0.2	0.30
2004	2	0.1	0.18	0.0	0.08	0.1	0.11	0.1	0.12
2005	5	0.3	0.26	0.1	0.15	0.1	0.18	0.2	0.19
2006	5	0.2	0.42	0.1	0.38	0.2	0.39	0.2	0.39
2007	4	0.2	0.31	0.1	0.17	0.1	0.23	0.2	0.28
2008	5	0.2	0.31	0.1	0.21	0.1	0.25	0.2	0.27
2009	6	0.3	0.29	0.1	0.09	0.1	0.15	0.2	0.23
2010	11	0.5	0.50	0.1	0.22	0.2	0.30	0.3	0.39
2011	7	0.3	0.26	0.1	0.15	0.2	0.18	0.2	0.21
2012	6	0.3	0.13	0.1	0.06	0.2	0.09	0.2	0.11
2013	8	0.3	0.26	0.2	0.14	0.2	0.19	0.3	0.21
2014	7	0.3	0.19	0.1	0.14	0.2	0.15	0.2	0.16
2015	8	0.3	0.22	0.1	0.13	0.2	0.17	0.3	0.19
2016	6	0.2	0.18	0.1	0.07	0.1	0.10	0.2	0.13
2017	11	0.4	0.21	0.2	0.09	0.3	0.12	0.3	0.16
2018	5	0.2	0.13	0.0	0.04	0.1	0.06	0.1	0.08
2019	7	0.3	0.24	0.2	0.18	0.2	0.21	0.2	0.22
2020	4	0.2	0.19	0.1	0.10	0.1	0.13	0.1	0.14
1998-2020	119	0.2	0.23	0.1	0.12	0.1	0.15	0.2	0.18

Table 12

Age distribution of age at death (cancer-related) for period 2007-2020  
(incl. multiple malignancies)

Age at death Years	Cases			Males			Females		
	n	%	Cum.%	n	%	Cum.%	n	%	Cum.%
0-4									
5-9									
10-14									
15-19									
20-24									
25-29	1	0.5	0.5	1	1.0	1.0			0.0
30-34	1	0.5	1.0	1	1.0	1.9			0.0
35-39	4	2.0	3.0	1	1.0	2.9	3	3.2	3.2
40-44	5	2.5	5.6	4	3.9	6.8	1	1.1	4.2
45-49	8	4.0	9.6	3	2.9	9.7	5	5.3	9.5
50-54	8	4.0	13.6	6	5.8	15.5	2	2.1	11.6
55-59	14	7.1	20.7	10	9.7	25.2	4	4.2	15.8
60-64	20	10.1	30.8	9	8.7	34.0	11	11.6	27.4
65-69	27	13.6	44.4	14	13.6	47.6	13	13.7	41.1
70-74	22	11.1	55.6	15	14.6	62.1	7	7.4	48.4
75-79	39	19.7	75.3	19	18.4	80.6	20	21.1	69.5
80-84	27	13.6	88.9	12	11.7	92.2	15	15.8	85.3
85+	22	11.1	100.0	8	7.8	100.0	14	14.7	100.0
All ages	198	100.0		103	100.0		95	100.0	

Table 13

Age-specific mortality (cancer-related) and proportion of all cancers  
for period 2007–2020  
(incl. multiple malignancies)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	Males MI-index	Females Age- spec. mortal.	Females MI-index	Males Prop.all cancers %	Females Prop.all cancers %
0- 4								
5- 9								
10-14								
15-19								
20-24								
25-29	1		0.0	0.07			1.1	
30-34	1		0.0	0.09			0.7	
35-39	1	3	0.0	0.06	0.1	0.16	0.4	0.7
40-44	4	1	0.2	0.17	0.0	0.05	0.7	0.1
45-49	3	5	0.1	0.11	0.2	0.13	0.2	0.3
50-54	6	2	0.2	0.18	0.1	0.07	0.2	0.1
55-59	10	4	0.5	0.24	0.2	0.12	0.2	0.1
60-64	9	11	0.5	0.26	0.6	0.29	0.1	0.2
65-69	14	13	0.9	0.39	0.7	0.39	0.2	0.2
70-74	15	7	1.0	0.41	0.4	0.29	0.1	0.1
75-79	19	20	1.6	0.50	1.3	0.54	0.2	0.2
80-84	12	15	1.7	0.50	1.4	1.07	0.1	0.2
85+	8	14	1.7	0.89	1.3	0.61	0.1	0.1
All ages	103	95					0.1	0.2
Mortality								
Raw			0.3	0.28	0.3	0.22		
WS			0.2	0.20	0.1	0.11		
ES			0.2	0.24	0.2	0.15		
BRD-S			0.3	0.26	0.2	0.18		
PYLL-70								
per 100,000			2.2		1.5			
ES			1.9		1.3			
AYLL-70			12.7		11.2			

Table 14a

Further malignancies in deaths in period 1998-2020  
MALES

Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C12-C13 Hypopharynx	1	1.4					1	100.0
C15 Oesophagus	3	4.3					3	100.0
C16 Stomach	1	1.4					1	100.0
C17 Small intestine	3	4.3	1	33.3	1	33.3	1	33.3
C18 Colon	9	13.0			7	77.8	2	22.2
C19-C20 Rectum	9	13.0	5	55.6	3	33.3	1	11.1
C25 Pancreas	3	4.3	1	33.3			2	66.7
C33-C34 Lung	3	4.3					3	100.0
C43 Malign. melanoma	1	1.4					1	100.0
C44 Skin others	3	4.3	3	100.0				
C48 Peritoneal	1	1.4	1	100.0				
C61 Prostate	16	23.2	7	43.8	3	18.8	6	37.5
C64 Kidney	2	2.9	1	50.0			1	50.0
C65 Renal pelvis	2	2.9					2	100.0
C67 Bladder	5	7.2	4	80.0			1	20.0
C69 Eye melanoma	1	1.4	1	100.0				
C70-C72 CNS cancer	1	1.4					1	100.0
C73 Thyroid	1	1.4	1	100.0				
C76-C79 CUP	1	1.4					1	100.0
C82-C85 NHL	1	1.4			1	100.0		
C91-C96 Leukaemia	2	2.9	1	50.0			1	50.0
All further malignancies	69	100.0	26	37.7	15	21.7	28	40.6

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a further malignancy.

Table 14b

Further malignancies in deaths in period 1998-2020  
FEMALES

Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C03-C06 Oral cavity	1	1.6					1	100.0
C16 Stomach	3	4.7	1	33.3			2	66.7
C17 Small intestine	1	1.6			1	100.0		
C18 Colon	11	17.2			6	54.5	5	45.5
C19-C20 Rectum	5	7.8	2	40.0	2	40.0	1	20.0
C25 Pancreas	3	4.7			1	33.3	2	66.7
C33-C34 Lung	4	6.3	1	25.0			3	75.0
C44 Skin others	2	3.1	1	50.0			1	50.0
C46,C49 Soft tissue	2	3.1	2	100.0				
C48 Peritoneal	1	1.6	1	100.0				
C50 Breast	11	17.2	9	81.8			2	18.2
C51 Vulva	1	1.6	1	100.0				
C53 Cervix uteri	2	3.1	1	50.0	1	50.0		
C54 Corpus uteri	2	3.1			1	50.0	1	50.0
C55,C57 Fem. genitals un	1	1.6	1	100.0				
C56 Ovary	8	12.5	3	37.5	4	50.0	1	12.5
C64 Kidney	4	6.3	2	50.0	1	25.0	1	25.0
C67 Bladder	2	3.1	1	50.0			1	50.0
All further malignancies	64	100.0	26	40.6	17	26.6	21	32.8

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a further malignancy.

Table 15

Age-specific mortality (cancer-related) and proportion of all cancers  
for period 2007-2020  
(First primaries only \*)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	Males MI-index	Females Age- spec. mortal.	Females MI-index	Males Prop.all cancers %	Females Prop.all cancers %
0- 4								
5- 9								
10-14								
15-19								
20-24								
25-29	1		0.0	0.07			1.2	
30-34	1		0.0	0.09			0.7	
35-39	1	1	0.0	0.06	0.0	0.07	0.4	0.3
40-44	4	1	0.2	0.17	0.0	0.06	0.7	0.1
45-49	3	4	0.1	0.14	0.2	0.12	0.2	0.3
50-54	6	1	0.2	0.20	0.0	0.05	0.3	0.0
55-59	8	3	0.4	0.25	0.1	0.11	0.2	0.1
60-64	4	6	0.2	0.15	0.3	0.20	0.1	0.1
65-69	11	11	0.7	0.50	0.6	0.42	0.1	0.2
70-74	11	6	0.7	0.55	0.3	0.29	0.1	0.1
75-79	13	16	1.1	0.57	1.1	0.62	0.1	0.2
80-84	7	8	1.0	0.54	0.8	0.73	0.1	0.1
85+	4	11	0.9	1.00	1.1	0.58	0.1	0.1
All ages	74	68					0.1	0.1
Mortality								
Raw			0.2	0.26	0.2	0.19		
WS			0.1	0.18	0.1	0.09		
ES			0.2	0.22	0.1	0.12		
BRD-S			0.2	0.24	0.2	0.14		
PYLL-70								
per 100,000			1.9		1.0			
ES			1.7		0.8			
AYLL-70			14.2		10.3			


\* See corresponding tables with multiple malignancies.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers  
for period 2007-2020  
(**Single primaries only** \*)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	Males MI-index	Females Age- spec. mortal.	Females MI-index	Males Prop.all cancers %	Females Prop.all cancers %
0- 4								
5- 9								
10-14								
15-19								
20-24								
25-29	1		0.0	0.07			1.2	
30-34	1		0.0	0.09			0.7	
35-39	1	1	0.0	0.06	0.0	0.07	0.4	0.3
40-44	4	1	0.2	0.17	0.0	0.06	0.7	0.1
45-49	3	4	0.1	0.14	0.2	0.12	0.2	0.3
50-54	6	1	0.2	0.20	0.0	0.05	0.3	0.0
55-59	8	3	0.4	0.28	0.1	0.12	0.2	0.1
60-64	3	4	0.2	0.13	0.2	0.15	0.1	0.1
65-69	9	8	0.6	0.45	0.4	0.35	0.1	0.1
70-74	10	4	0.7	0.63	0.2	0.20	0.1	0.1
75-79	9	13	0.7	0.41	0.9	0.57	0.1	0.2
80-84	5	6	0.7	0.50	0.6	0.60	0.1	0.1
85+	2	8	0.4	0.50	0.8	0.47	0.0	0.1
All ages	62	53					0.1	0.1
Mortality								
Raw			0.2	0.23	0.2	0.15		
WS			0.1	0.17	0.1	0.07		
ES			0.1	0.20	0.1	0.10		
BRD-S			0.2	0.22	0.1	0.12		
PYLL-70								
per 100,000			1.9		0.9			
ES			1.6		0.7			
AYLL-70			15.0		11.6			

\* See corresponding tables with multiple malignancies.




**Figure 17.** Distribution of age at death (bars; males: mean=63.7 yrs, median=65.4 yrs; females: mean=68.1 yrs, median=70.7 yrs) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).


The difference between age at diagnosis (Table 3) and age at appendix cancer-related death (see Table 10) should be considered.



Average mortality (Germany 1987 standard population) 2007 - 2020: Males




Average mortality (Germany 1987 standard population) 2007 - 2020: Females




**Figure 18a.** Map of cancer mortality (german standard population) by county averaged for period 2007 to 2020. According to their individual mortality rates, the counties are displayed in different red and blue hues, being the fine white color attributed to the population mean (males 0.3/100,000 WS N=103, females 0.2/100,000 WS N=95).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,727 female residents (averaged) in the period from 2007 to 2020 a total of 3 women died from appendix cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.3/100,000 (german standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.1/100,000.

Standardized mortality ratio (SMR) 2007 - 2020: Males



Standardized mortality ratio (SMR) 2007 - 2020: Females



**Figure 18b.** Map of standardized mortality ratio (SMR) by county averaged for period 2007 to 2020. According to their individual SMR values, the counties are displayed in different red and blue hues, being the fine white color attributed to the population overall of 1.0 (males N=103, females N=95).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,153 female residents (averaged) in the period from 2007 to 2020 a total of 3 women died from appendix cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 1.16. Though, the value of this parameter may vary with an underlying probability of 99% between 0.13 and 4.26, and is therefore not statistically striking.

### Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases, where applicable. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

#### 1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

#### 2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

#### 3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head&neck and lung).

The ratio of mortality and incidence (mortality-to-incidence ratio, **MIR, MI-Index**) is a statistical index that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MIR. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

**Shortcuts**

MCR	Munich Cancer Registry (Tumorregister München)
GEKID	Association of Population-based Cancer Registries in Germany (Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)
SEER	Surveillance, Epidemiology, and End Results (USA)
DCO	Death certificate only
BRD-S	German (FRG) standard population
ES	European standard population (old)
WS	World standard population
SIR	Standardized incidence ratio
CI	Confidence interval
EAR	Excess absolute risk = excess cancer cases (O - E) per 10,000 person-years
PYLL-70	Potential years of life lost prior to age 70 given a person dies before that age
AYLL-70	Average years of life lost prior to age 70 given a person dies before that age
SMR	Standardized mortality ratio
MI-index	Ratio of mortality to incidence, MIR
FRG	Federal Republic of Germany

**Recommended Citation**

Munich Cancer Registry. ICD-10 C18.1: Appendix cancer - Incidence and Mortality [Internet]. 2021 [updated 2021 Dec 20; cited 2022 Feb 1]. Available from: [https://www.tumorregister-muenchen.de/en/facts/base/bC181\\_E-ICD-10-C18.1-Appendix-cancer-incidence-and-mortality.pdf](https://www.tumorregister-muenchen.de/en/facts/base/bC181_E-ICD-10-C18.1-Appendix-cancer-incidence-and-mortality.pdf)

**Copyright**

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

**Disclaimer**

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.