

Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ *Deutsch*

ICD-10 C24: Biliary tract cancer

Incidence and Mortality

Year of diagnosis	1998-2019
Patients	3,271
Diseases	3,271
Creation date	01/25/2021
Database export	01/07/2021
Population	4.92 m





Munich Cancer Registry
Cancer Registry Bavaria - Upper Bavaria Regional Center
at Klinikum Grosshadern/IBE
Marchioninstr. 15
Munich, 81377
Germany

<https://www.tumorregister-muenchen.de/en>

https://www.tumorregister-muenchen.de/en/facts/base/bC24__E-ICD-10-C24-Biliary-tract-cancer-incidence-and-mortality.pdf

Index of figures and tables

Fig./Tbl.		Page
1	Annual cases, DCO, mult. malignancies, follow-up / yr	4
2	Incidence by year of diagnosis	7
3	Age distribution parameters by year of diagnosis	8
4	Age distribution by 5-year age group and sex	11
5	Age-specific incidence, DCO rate, proportion malignancies	12
6	Age distribution and age-specific incidence (chart)	13
6a	Age-specific incidence internationally (chart)	14
7	Standardized incidence ratio of further malignancies	15
8a	Map of cancer incidence (BRD-S) by county (chart)	17
8b	Standardized incidence ratio (SIR) by county (chart)	18
9a	Pts incident cohorts and mortality / yr	19
9b	Incidence and mortality by year of diagnosis	20
9c	Cancer-related deaths, death certification available / yr	21
10	Medians of age at death / yr	22
11	Mortality by year of death	24
12	Distribution of age at death	26
13	Age-specific mortality	27
14	Further malignancies in deaths	28
15	Age-specific mortality (first primaries)	30
16	Age-specific mortality (single primaries)	31
17	Age distribution and age-specific mortality (chart)	32
18a	Map of cancer mortality (BRD-S) by county (chart)	33
18b	Standardized mortality ratio (SMR) by county (chart)	34

**Global Statements about the statistics on the Internet –
Baseline Statistics** (grey button ) , **Survival** (red button )

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.69 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases^{###} are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, January 2021

[#] Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.65 million to 4.10 in 2002, and to 4.69 million in 2007).

^{##} Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.

^{###} DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate.

ICD-10 codes (ICD-10 2016) used for specifying cancer site

Code	Description
C24.-	Malignant neoplasm of other and unspecified parts of biliary tract
C24.0	Extrahepatic bile duct
C24.1	Ampulla of Vater
C24.8	Overlapping lesion of biliary tract
C24.9	Biliary tract, unspecified

INCIDENCE

Table 1

Cases with invasive cancer by year of diagnosis, proportions of DCO, further malignancies, deaths, and active follow-up (ALL PATIENTS) (incl. DCO)

Year of diagnosis	All cases n	DCO cases n	Prop. DCO %	Prop. at least 1 further malign. prior + synchron. %	Prop. at least 1 further malign. after %	Prop. deaths %	Prop. actively followed %
1998	69	15	21.7	5.8	3.6	91.3	100.0
1999	74	15	20.3	9.8	3.6	94.6	100.0
2000	59	17	28.8	9.9	3.5	98.3	100.0
2001	80	29	36.3	10.3	3.5	95.0	98.8
2002	180	79	43.9	12.3	3.6	97.8	100.0 #
2003	164	54	32.9	11.7	3.5	95.7	99.4
2004	154	48	31.2	11.3	3.4	90.3	97.4
2005	152	46	30.3	11.4	3.5	89.5	99.3
2006	163	40	24.5	12.4	3.6	93.9	98.8
2007	188	42	22.3	12.6	3.7	91.0	98.4 #
2008	188	48	25.5	13.5	3.4	91.5	98.9
2009	175	24	13.7	13.3	3.3	89.7	100.0
2010	152	27	17.8	13.7	3.1	84.9	100.0
2011	175	28	16.0	14.4	3.0	90.3	98.9
2012	162	23	14.2	14.8	3.2	87.7	98.8
2013	150	33	22.0	15.3	2.9	81.3	98.7
2014	179	33	18.4	15.4	2.6	86.0	99.4
2015	157	29	18.5	15.9	2.7	80.9	99.4
2016	183	37	20.2	16.1	2.7	77.0	99.5
2017	198	17	8.6	16.7	2.2	71.2	100.0
2018	131	5	3.8	17.1	1.9	52.7	100.0
2019	138			17.3	1.5	42.8	80.4 ##
1998-2019	3271	689	21.1	17.3	3.6	84.7	98.5

3,271 cases diagnosed 1998-2019 are related to a total of 3,271 patients. Currently, in 698 (21.3 %) of these 3,271 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 564 / 104 / 30 (17.2 % / 3.2 % / 0.9 %) patients exist having 2 / 3 / 4+ malignancies.

The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retrieved from the respective headings.

How to interpret:

In 2017, a subgroup of 198 cases has been diagnosed, of which 16.7 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 2.2 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

Table 1a

Cases with invasive cancer by year of diagnosis, proportions of DCO, further malignancies, deaths, and active follow-up (MALES) (incl. DCO)

Year of diagnosis	Males n	Males %	DCO cases n	Prop. DCO %	Prop. at least 1 further malign. prior + synchron. %	Prop. at least 1 further malign. after %	Prop. deaths %	Prop. actively followed %
1998	28	40.6	5	17.9	10.7	3.6	82.1	100.0
1999	39	52.7	8	20.5	11.9	3.4	89.7	100.0
2000	32	54.2	5	15.6	12.1	3.3	100.0	100.0
2001	33	41.3	6	18.2	12.1	3.3	97.0	100.0
2002	84	46.7	30	35.7	13.4	3.4	96.4	100.0 #
2003	83	50.6	25	30.1	12.7	3.2	96.4	100.0
2004	73	47.4	19	26.0	12.1	3.2	90.4	97.3
2005	74	48.7	17	23.0	12.6	3.1	89.2	100.0
2006	75	46.0	17	22.7	13.8	3.1	94.7	98.7
2007	88	46.8	15	17.0	14.0	3.2	87.5	98.9 #
2008	103	54.8	18	17.5	14.6	3.0	91.3	100.0
2009	99	56.6	11	11.1	13.8	2.7	90.9	100.0
2010	84	55.3	8	9.5	14.2	2.9	81.0	100.0
2011	92	52.6	8	8.7	15.3	3.0	84.8	97.8
2012	97	59.9	11	11.3	15.7	3.4	86.6	99.0
2013	81	54.0	11	13.6	16.1	3.3	80.2	100.0
2014	93	52.0	15	16.1	16.0	3.2	80.6	98.9
2015	86	54.8	14	16.3	16.7	3.4	82.6	98.8
2016	109	59.6	18	16.5	16.7	3.6	75.2	100.0
2017	120	60.6	5	4.2	17.8	3.2	68.3	100.0
2018	83	63.4	2	2.4	18.5	2.4	49.4	100.0
2019	90	65.2			18.6	2.3	48.9	84.4 ##
1998-2019	1746	53.4	268	15.3	18.6	3.6	82.3	98.7

1,746 cases diagnosed 1998-2019 are related to a total of 1,746 patients. Currently, in 396 (22.7 %) of these 1,746 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 315 / 64 / 17 (18.0 % / 3.7 % / 1.0 %) patients exist having 2 / 3 / 4+ malignancies.

The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retrieved from the respective headings.

How to interpret:

In 2017, a subgroup of 120 cases has been diagnosed, of which 17.8 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 3.2 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

Table 1b

Cases with invasive cancer by year of diagnosis, proportions of DCO, further malignancies, deaths, and active follow-up (FEMALES) (incl. DCO)

Year of diagnosis	Females n	Females %	DCO cases n	Prop. DCO %	Prop. at least 1 further malign. prior + synchron. %	Prop. at least 1 further malign. after %	Prop. deaths %	Prop. actively followed %
1998	41	59.4	10	24.4	2.4	3.7	97.6	100.0
1999	35	47.3	7	20.0	7.9	3.8	100.0	100.0
2000	27	45.8	12	44.4	7.8	3.8	96.3	100.0
2001	47	58.8	23	48.9	8.7	3.8	93.6	97.9
2002	96	53.3	49	51.0	11.4	3.9	99.0	100.0 #
2003	81	49.4	29	35.8	10.7	3.9	95.1	98.8
2004	81	52.6	29	35.8	10.5	3.7	90.1	97.5
2005	78	51.3	29	37.2	10.3	3.9	89.7	98.7
2006	88	54.0	23	26.1	11.1	4.1	93.2	98.9
2007	100	53.2	27	27.0	11.4	4.4	94.0	98.0 #
2008	85	45.2	30	35.3	12.4	4.0	91.8	97.6
2009	76	43.4	13	17.1	12.8	4.1	88.2	100.0
2010	68	44.7	19	27.9	13.3	3.4	89.7	100.0
2011	83	47.4	20	24.1	13.6	3.1	96.4	100.0
2012	65	40.1	12	18.5	14.0	3.0	89.2	98.5
2013	69	46.0	22	31.9	14.5	2.3	82.6	97.1
2014	86	48.0	18	20.9	14.8	1.7	91.9	100.0
2015	71	45.2	15	21.1	15.1	1.6	78.9	100.0
2016	74	40.4	19	25.7	15.5	1.2	79.7	98.6
2017	78	39.4	12	15.4	15.5	0.6	75.6	100.0
2018	48	36.6	3	6.3	15.6	1.1	58.3	100.0
2019	48	34.8			15.8	0.0	31.3	72.9 ##
1998-2019	1525	46.6	421	27.6	15.8	3.7	87.4	98.2

1,525 cases diagnosed 1998-2019 are related to a total of 1,525 patients. Currently, in 302 (19.8 %) of these 1,525 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 249 / 40 / 13 (16.3 % / 2.6 % / 0.9 %) patients exist having 2 / 3 / 4+ malignancies.

The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retrieved from the respective headings.

How to interpret:

In 2017, a subgroup of 78 cases has been diagnosed, of which 15.5 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 0.6 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

Table 2

Incidence measures by year of diagnosis including DCO cases
(with respect to registry area expansion from 2.65 to 4.10 m as of 2002,
and from 4.10 to 4.92 m as of 2007, respectively)

Year of diagnosis	Males n	Females n	Males Inc. raw	Fem. Inc. raw	Males Inc. WS	Fem. Inc. WS	Males Inc. ES	Fem. Inc. ES	Males Inc. BRD-S	Fem. Inc. BRD-S
1998	28	41	2.5	3.5	1.4	1.6	2.3	2.4	3.4	2.9
1999	39	35	3.5	2.9	2.0	1.2	3.1	1.9	3.8	2.6
2000	32	27	2.8	2.2	1.6	0.7	2.5	1.2	3.5	1.8
2001	33	47	2.8	3.9	1.6	1.6	2.5	2.5	3.5	3.3
2002	84	96	4.5	4.9	2.5	1.6	3.8	2.7	4.9	3.8
2003	83	81	4.4	4.1	2.4	1.4	3.7	2.2	4.9	3.1
2004	73	81	3.9	4.1	2.1	1.5	3.2	2.3	4.2	3.1
2005	74	78	3.9	3.9	2.1	1.6	3.1	2.4	3.9	3.2
2006	75	88	3.9	4.4	2.0	1.5	3.0	2.3	4.0	3.2
2007	88	100	4.0	4.3	2.2	1.5	3.2	2.4	4.0	3.3
2008	103	85	4.6	3.7	2.2	1.4	3.4	2.1	4.7	2.8
2009	99	76	4.4	3.3	2.1	1.2	3.3	1.8	4.4	2.5
2010	84	68	3.7	2.9	1.8	1.1	2.8	1.6	3.7	2.2
2011	92	83	4.1	3.6	1.8	1.2	2.9	1.9	4.0	2.6
2012	97	65	4.3	2.8	2.0	1.0	3.0	1.5	3.9	2.0
2013	81	69	3.5	2.9	1.5	1.1	2.4	1.6	3.3	2.1
2014	93	86	4.0	3.6	1.8	1.1	2.7	1.8	3.6	2.6
2015	86	71	3.6	2.9	1.5	0.9	2.3	1.5	3.3	2.1
2016	109	74	4.5	3.0	2.0	1.0	3.1	1.5	4.1	2.1
2017	120	78	5.0	3.2	2.0	1.0	3.2	1.6	4.3	2.3
2018	83	48	3.4	1.9	1.6	0.7	2.3	1.0	3.1	1.4
2019	90	48	3.7	1.9	1.4	0.7	2.3	1.1	3.2	1.4
1998-2019	1746	1525	4.0	3.3	1.9	1.2	2.9	1.8	3.9	2.5

The computation of the incidence measures includes all cancers, irrespective of first or subsequent malignancy.

Table 3

Age distribution parameters by year of diagnosis (ALL PATIENTS)
(incl. DCO)

Year of diagnosis	Cases n	Std.		Min.	Max.	10%	25%	Median		
		Mean	dev.					50%	75%	90%
1998	69	71.4	13.1	28.5	99.5	54.3	62.3	72.7	79.5	89.3
1999	74	70.3	11.8	35.3	96.5	56.5	63.4	71.0	77.5	83.7
2000	59	74.4	10.5	49.5	94.0	60.0	66.4	75.3	81.8	86.8
2001	80	72.9	11.9	40.2	99.1	57.1	64.4	74.5	81.1	88.2
2002	180	74.0	12.2	34.1	94.5	56.8	65.4	75.7	82.7	89.0
2003	164	74.3	10.4	48.5	96.8	60.2	66.3	74.7	82.2	87.6
2004	154	73.0	12.3	34.0	100	55.6	65.5	73.7	82.6	88.0
2005	152	71.5	11.5	44.5	98.0	57.0	63.3	71.4	80.1	85.7
2006	163	73.6	12.8	36.7	99.2	56.9	65.3	74.1	83.5	87.1
2007	188	71.9	12.0	40.0	97.1	55.3	64.0	71.9	81.0	87.2
2008	188	73.3	11.5	32.9	99.3	58.5	66.8	73.2	82.4	86.5
2009	175	72.0	12.1	26.5	93.7	54.4	65.8	73.3	81.0	86.1
2010	152	72.2	11.2	45.3	93.8	55.9	64.4	73.8	79.5	87.0
2011	175	74.0	11.2	36.1	100	58.6	66.5	75.3	82.8	87.5
2012	162	72.4	10.8	29.3	93.9	57.9	64.8	73.6	79.6	86.7
2013	150	73.6	10.2	48.6	96.2	58.3	66.9	74.1	80.6	87.7
2014	179	73.2	12.0	30.3	97.2	54.9	66.9	75.5	81.4	87.1
2015	157	74.2	11.7	34.2	98.4	58.2	69.0	76.2	81.3	86.8
2016	183	73.8	11.1	41.1	96.4	58.9	67.1	75.4	81.1	86.9
2017	198	73.1	11.1	41.2	98.5	55.8	66.4	76.2	80.3	84.7
2018	131	71.5	10.6	39.1	94.0	56.1	65.3	73.8	79.0	82.3
2019	138	72.8	10.4	39.2	89.0	57.6	68.1	75.4	80.3	83.7
1998-2019	3271	73.0	11.5	26.5	100	57.0	65.8	74.4	81.3	86.8

Table 3a

Age distribution parameters by year of diagnosis (MALES)
(incl. DCO)

Year of diagnosis	Cases n	Std.		Median						
		Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	28	71.0	10.9	51.0	91.0	55.5	61.1	73.7	79.2	81.8
1999	39	66.7	13.2	35.3	89.8	49.9	58.2	68.1	76.9	85.6
2000	32	70.6	11.2	49.5	94.0	56.4	62.5	72.1	79.8	82.2
2001	33	72.7	10.6	52.5	89.8	58.4	66.1	72.8	80.5	88.0
2002	84	70.5	10.9	44.7	93.5	57.2	62.3	69.7	78.9	86.0
2003	83	72.0	9.7	52.5	95.2	58.9	63.9	72.0	79.7	84.8
2004	73	70.3	11.6	34.0	91.5	54.1	64.5	71.8	77.3	84.0
2005	74	69.2	10.2	47.0	98.0	57.0	63.3	69.1	75.8	82.1
2006	75	69.6	12.3	36.7	92.8	51.8	62.2	68.8	78.3	83.8
2007	88	67.7	11.3	40.0	93.1	53.6	61.4	66.4	76.2	84.2
2008	103	72.1	10.6	37.0	89.5	58.5	65.7	72.3	81.0	84.6
2009	99	70.6	10.9	43.3	91.0	52.8	64.7	71.3	79.3	82.3
2010	84	70.8	10.2	45.3	93.7	56.7	63.7	72.1	78.1	84.2
2011	92	71.9	10.9	38.9	92.1	56.9	65.4	74.3	79.5	84.3
2012	97	70.9	9.7	49.8	93.5	56.5	64.3	72.5	76.2	83.2
2013	81	72.9	9.3	48.6	93.8	60.7	66.9	74.1	79.6	83.0
2014	93	70.9	12.6	30.3	97.2	51.8	64.8	74.0	79.4	84.4
2015	86	73.3	11.1	34.2	95.1	58.2	68.5	75.6	79.4	84.8
2016	109	72.0	10.6	41.1	95.5	56.7	66.4	73.1	79.8	83.2
2017	120	72.4	11.2	43.5	93.5	53.7	64.8	75.2	80.2	84.8
2018	83	70.1	10.5	39.1	94.0	55.1	64.7	71.2	77.8	80.2
2019	90	73.2	8.4	51.9	87.4	59.7	68.1	75.2	78.6	83.2
1998-2019	1746	71.1	10.8	30.3	98.0	56.4	64.1	72.5	79.1	84.0

Table 3b

Age distribution parameters by year of diagnosis (FEMALES)
(incl. DCO)

Year of diagnosis	Cases n	Std.		Min.	Max.	Median				
		Mean	dev.			10%	25%	50%	75%	90%
1998	41	71.7	14.6	28.5	99.5	54.3	62.6	71.6	83.7	89.9
1999	35	74.3	8.6	56.3	96.5	65.5	68.8	73.2	79.0	83.7
2000	27	78.9	7.4	64.9	92.4	66.4	74.0	78.9	82.4	90.2
2001	47	73.0	12.8	40.2	99.1	54.7	63.4	75.7	81.6	88.5
2002	96	77.0	12.6	34.1	94.5	56.8	72.7	80.5	84.9	90.0
2003	81	76.6	10.6	48.5	96.8	62.5	67.8	77.8	84.1	89.0
2004	81	75.5	12.4	44.2	100	58.0	67.7	77.0	84.2	90.0
2005	78	73.7	12.2	44.5	98.0	55.5	62.9	74.2	82.5	90.5
2006	88	76.9	12.2	43.2	99.2	59.3	71.2	78.9	85.7	91.3
2007	100	75.6	11.4	45.3	97.1	59.5	69.2	77.1	83.6	88.6
2008	85	74.7	12.4	32.9	99.3	59.1	68.6	76.9	83.5	88.4
2009	76	73.7	13.3	26.5	93.7	58.5	67.5	76.3	83.6	88.2
2010	68	74.0	12.2	45.3	93.8	55.3	68.1	75.4	83.7	88.3
2011	83	76.2	11.1	36.1	100	62.4	69.4	76.2	84.7	89.0
2012	65	74.6	12.1	29.3	93.9	60.4	67.8	74.7	83.8	90.3
2013	69	74.4	11.2	50.9	96.2	56.0	68.6	74.2	83.0	88.4
2014	86	75.8	10.9	44.5	91.6	56.5	70.6	78.3	82.9	88.7
2015	71	75.3	12.3	35.4	98.4	58.7	70.1	77.6	83.6	88.9
2016	74	76.5	11.2	46.3	96.4	61.6	70.0	78.1	85.9	89.7
2017	78	74.2	10.9	41.2	98.5	57.2	68.2	76.8	80.6	84.1
2018	48	73.9	10.5	49.0	92.3	56.8	68.1	77.0	80.5	84.0
2019	48	72.0	13.4	39.2	89.0	46.2	67.6	76.6	82.4	83.7
1998-2019	1525	75.1	11.8	26.5	100	58.6	68.3	76.7	83.4	88.9

Table 4

Age distribution by 5-year age group and sex for period 2007–2019
(incl. DCO)

Age at diagnosis Years	Cases			Males			Females		
	n	%	Cum.%	n	%	Cum.%	n	%	Cum.%
0–4									
5–9									
10–14									
15–19									
20–24									
25–29	2	0.1	0.1		0.0	0.0	2	0.2	0.2
30–34	4	0.2	0.3	3	0.2	0.2	1	0.1	0.3
35–39	8	0.4	0.6	4	0.3	0.6	4	0.4	0.7
40–44	17	0.8	1.4	9	0.7	1.3	8	0.8	1.6
45–49	43	2.0	3.4	23	1.9	3.2	20	2.1	3.7
50–54	93	4.3	7.7	60	4.9	8.1	33	3.5	7.2
55–59	134	6.2	13.8	91	7.4	15.5	43	4.5	11.7
60–64	185	8.5	22.3	122	10.0	25.5	63	6.6	18.3
65–69	281	12.9	35.2	175	14.3	39.8	106	11.1	29.4
70–74	357	16.4	51.7	221	18.0	57.8	136	14.3	43.7
75–79	448	20.6	72.2	252	20.6	78.4	196	20.6	64.4
80–84	344	15.8	88.1	177	14.4	92.8	167	17.6	81.9
85+	260	11.9	100.0	88	7.2	100.0	172	18.1	100.0
All ages	2176	100.0		1225	100.0		951	100.0	

Table 5

Age-specific incidence, DCO rate and proportion of all cancers for period 2007–2019

Age at diagnosis Years	Males n	Females n	Males Age- spec. incid.	Females Age- spec. incid.	Males DCO rate n=136 %	Females DCO rate n=210 %	Males Prop.all cancers n=143063 %	Females Prop.all cancers n=144724 %
0- 4								
5- 9								
10-14								
15-19								
20-24								
25-29		2		0.1				0.2
30-34	3	1	0.1	0.0			0.3	0.1
35-39	4	4	0.2	0.2		25.0	0.2	0.1
40-44	9	8	0.4	0.4			0.3	0.1
45-49	23	20	0.9	0.8			0.5	0.2
50-54	60	33	2.6	1.4	5.0	3.0	0.8	0.3
55-59	91	43	4.7	2.2	2.2	4.7	0.8	0.3
60-64	122	63	7.5	3.6	5.7	11.1	0.7	0.4
65-69	175	106	11.5	6.3	5.1	6.6	0.8	0.6
70-74	221	136	15.8	8.5	9.5	5.1	0.9	0.7
75-79	252	196	22.8	14.2	9.1	15.8	1.1	1.1
80-84	177	167	27.0	17.2	18.6	31.1	1.3	1.2
85+	88	172	20.6	17.8	43.2	59.3	0.9	1.1
All ages	1225	951			11.1	22.1	0.9	0.7
Incidence								
Raw			4.1	3.1				
WS			1.8	1.1				
ES			2.8	1.7				
BRD-S			3.8	2.3				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

ICD-10 C24: Malignant neoplasm of other and unspecified parts of biliary tract

Age distribution and age-specific incidence 2007 - 2019 (Males: 1225, Females: 951)




Figure 6. Age distribution (males: mean=71.5 yrs, median=73.4 yrs; females: mean=74.8 yrs, median=76.5 yrs) and age-specific incidence.

CD-10 C24: Malignant neoplasm of other and unspecified parts of biliary tract

Age-specific incidence rates: international comparison




Figure 6a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).

Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2019, based on the November 2018 submission. <http://www.seer.cancer.gov>.

Table 7a

Standardized incidence ratio (SIR, with 95% confidence limits),
excess absolute risk (EAR) and DCO rate of further malignancies
for period 1998–2019

MALES

Diagnosis	Observed n	Expected n	SIR	CI 95%	CI 95%	EAR	DCO %
C09–C10 Oropharynx	1	0.4	2.6	0.1	14.4	2.6	
C15 Oesophagus	2	0.8	2.6	0.3	9.3	5.2	
C16 Stomach	8	1.6	5.0	2.2	9.8 #	27.1	12.5
C17 Small intestine	9	0.2	38.3	17.5	72.6 #	37.0	
C18 Colon	11	3.9	2.8	1.4	5.0 #	30.0	18.2
C19–C20 Rectum	1	2.1	0.5	0.0	2.6	-4.7	
C22 Liver	4	1.2	3.4	0.9	8.8	12.0	
C25 Pancreas	8	1.6	5.1	2.2	10.1 #	27.2	25.0
C33–C34 Lung	12	4.7	2.5	1.3	4.5 #	30.8	
C38,C45 Mesothelioma	1	0.3	3.5	0.1	19.7	3.0	100.0
C43 Malign. melanoma	5	1.7	2.9	0.9	6.7	13.8	
C50 Breast	1	0.1	9.1	0.2	50.8	3.8	
C61 Prostate	13	11.4	1.1	0.6	2.0	6.8	15.4
C64 Kidney	1	1.4	0.7	0.0	4.1	-1.6	
C65 Renal pelvis	1	0.2	5.6	0.1	31.3	3.5	
C67 Bladder	5	1.9	2.7	0.9	6.2	13.2	
C82–C85 NHL	2	1.7	1.2	0.1	4.3	1.4	
Not observed	0	5.2	0.0	0.0	0.7 #	-22.1	
All further malignancies	85	40.3	2.1	1.7	2.6 #	188.9	9.4
Patients		1521					
Median age at next malignancy (years)		73.1					
Person-years		2366					
Mean observation time (years)		1.6					
Median observation time (years)		0.7					

The occurrence of further specified malignancy is statistically significant.

Table 7b


Standardized incidence ratio (SIR, with 95% confidence limits),
excess absolute risk (EAR) and DCO rate of further malignancies
for period 1998–2019

FEMALES

Diagnosis	Observed n	Expected n	SIR	CI 95%	CI 95%	EAR	DCO %
C16 Stomach	4	0.8	4.7	1.3	12.1 #	16.7	25.0
C18 Colon	9	2.4	3.8	1.7	7.2 #	35.0	11.1
C19–C20 Rectum	3	1.0	3.1	0.6	9.0	10.7	33.3
C23–C24 Bile	2	0.4	5.5	0.7	19.9	8.7	
C25 Pancreas	8	1.2	7.0	3.0	13.7 #	36.3	12.5
C33–C34 Lung	10	1.8	5.6	2.7	10.3 #	43.5	
C43 Malign. melanoma	1	0.8	1.2	0.0	6.6	0.8	
C46,C49 Soft tissue	1	0.1	7.5	0.2	41.9	4.6	
C48 Peritoneal	1	0.1	10.4	0.3	57.8	4.8	
C50 Breast	4	6.9	0.6	0.2	1.5	-15.4	
C53 Cervix uteri	2	0.3	7.5	0.9	27.2	9.2	50.0
C54 Corpus uteri	1	1.3	0.8	0.0	4.2	-1.7	100.0
C56 Ovary	5	1.0	5.2	1.7	12.1 #	21.4	20.0
C64 Kidney	4	0.6	6.9	1.9	17.6 #	18.1	25.0
C69 Eye carcinoma	1	0.0	185.3	4.7	1032 #	5.3	
C74–C80 Cancer others	1	0.1	11.3	0.3	62.8	4.8	100.0
C76–C79 CUP	2	0.4	4.5	0.5	16.4	8.3	
C82–C85 NHL	4	0.9	4.2	1.1	10.8 #	16.2	50.0
C91–C96 Leukaemia	3	0.4	8.6	1.8	25.0 #	14.0	
Not observed	0	3.2	0.0	0.0	1.1	-17.0	
All further malignancies	66	23.7	2.8	2.2	3.5 #	224.2	16.7
Patients		1199					
Median age at next malignancy (years)		76.5					
Person-years		1888					
Mean observation time (years)		1.6					
Median observation time (years)		0.7					

The occurrence of further specified malignancy is statistically significant.

Average incidence (Germany 1987 standard population) 2007 - 2019: Males



Average incidence (Germany 1987 standard population) 2007 - 2019: Females





Figure 8a. Map of cancer incidence (german standard population, incl. DCO cases) by county averaged for period 2007 to 2019. According to their individual incidence rates, the counties are displayed in different red and blue hues, being the fine white color attributed to the population mean (males 3.8/100,000 WS N=1,225, females 2.3/100,000 WS N=951).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,462 female residents (averaged) in the period from 2007 to 2019 a total of 23 women were identified with newly diagnosed biliary tract cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 2.1/100,000 (german standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 1.1 and 3.5/100,000.

Standardized incidence ratio (SIR) 2007 - 2019: Males



Standardized incidence ratio (SIR) 2007 - 2019: Females




Figure 8b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2019. According to their individual SIR values, the counties are displayed in different red and blue hues, being the fine white color attributed to the population overall of 1.0 (males N=1,225, females N=951).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,153 female residents (averaged) in the period from 2007 to 2019 a total of 23 women were identified with newly diagnosed biliary tract cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.90. Though, the value of this parameter may vary with an underlying probability of 99% between 0.49 and 1.50, and is therefore not statistically striking.

MORTALITY

Table 9a

Annual cohorts: Incident cancers, follow-up status, proportion of DCO, deaths among the annual cohorts and proportion of available death certificates (with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.92 m as of 2007, respectively)

Year of diagnosis	Incident cases n	Prop. actively followed %	Prop. DCO %	Deaths n	Prop. deaths %	Prop. deaths with death certific. %
1998	69	100.0	21.7	63	91.3	92.1
1999	74	100.0	20.3	70	94.6	98.6
2000	59	100.0	28.8	58	98.3	91.4
2001	80	98.8	36.3	76	95.0	97.4
2002	180	100.0	43.9	176	97.8	99.4
2003	164	99.4	32.9	157	95.7	96.8
2004	154	97.4	31.2	139	90.3	98.6
2005	152	99.3	30.3	136	89.5	99.3
2006	163	98.8	24.5	153	93.9	100.0
2007	188	98.4	22.3	171	91.0	98.8
2008	188	98.9	25.5	172	91.5	97.7
2009	175	100.0	13.7	157	89.7	96.8
2010	152	100.0	17.8	129	84.9	98.4
2011	175	98.9	16.0	158	90.3	98.1
2012	162	98.8	14.2	142	87.7	95.8
2013	150	98.7	22.0	122	81.3	99.2
2014	179	99.4	18.4	154	86.0	96.8
2015	157	99.4	18.5	127	80.9	92.9
2016	183	99.5	20.2	141	77.0	90.1
2017	198	100.0	8.6	141	71.2	75.2
2018	131	100.0	3.8	69	52.7	47.8
2019	138	80.4		59	42.8	78.0
1998-2019	3271	98.5	21.1	2770	84.7	94.3

Table 9b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased within the same year of being diagnosed with cancer (incl. DCO)

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.92 m as of 2007, respectively)

Year of diagnosis/ death	Incident cases n	Deaths n	Prop. deaths with death certific. %	Deaths in same year n	Prop. deaths in same year %
1998	69	56	91.1	32	46.4
1999	74	63	95.2	33	44.6
2000	59	65	95.4	29	49.2
2001	80	69	97.1	34	42.5
2002	180	142	99.3	116	64.4
2003	164	109	97.2	79	48.2
2004	154	105	99.0	73	47.4
2005	152	109	99.1	63	41.4
2006	163	140	97.9	86	52.8
2007	188	118	98.3	71	37.8
2008	188	143	97.9	89	47.3
2009	175	150	98.7	69	39.4
2010	152	136	100.0	47	30.9
2011	175	146	99.3	66	37.7
2012	162	158	97.5	70	43.2
2013	150	123	98.4	53	35.3
2014	179	133	99.2	71	39.7
2015	157	148	100.0	62	39.5
2016	183	153	99.3	83	45.4
2017	198	150	98.0	74	37.4
2018	131	110	26.4	34	26.0
2019	138	85	50.6	28	20.3
1998–2019	3271	2611	93.7	1362	41.6

Table 9c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates
(incl. DCO)

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002,
and from 4.10 to 4.92 m as of 2007, respectively)

Year of death	Deaths n	Prop. cancer- related %	Prop. non-cancer- related %	Prop. cancer recorded on death certificate %
1998	56	85.7	14.3	94.1
1999	63	81.0	19.0	95.0
2000	65	92.3	7.7	98.4
2001	69	89.9	10.1	97.0
2002	142	90.1	9.9	94.3
2003	109	90.8	9.2	97.2
2004	105	89.5	10.5	96.2
2005	109	93.6	6.4	98.1
2006	140	90.0	10.0	93.4
2007	118	92.4	7.6	97.4
2008	143	94.4	5.6	97.1
2009	150	90.0	10.0	94.6
2010	136	89.0	11.0	94.9
2011	146	91.1	8.9	95.2
2012	158	89.2	10.8	94.8
2013	123	90.2	9.8	95.0
2014	133	90.2	9.8	94.7
2015	148	90.5	9.5	95.3
2016	153	89.5	10.5	94.1
2017	150	92.7	7.3	95.2
2018	110	60.0	40.0	89.7
2019	85	63.5	36.5	90.7
1998–2019	2611	88.3	11.7	95.3

Table 10a

Medians of age at death according to the grouping in Table 9
MALES

Year of death	Deaths n	Age at death (all causes) Years	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate) Years
1998	25	75.6	73.6	77.4	75.6
1999	28	70.8	67.5	75.3	68.5
2000	29	75.5	75.5	65.8	76.3
2001	35	69.8	67.5	75.2	67.8
2002	66	71.9	72.0	71.6	71.7
2003	53	72.3	71.1	74.3	73.0
2004	46	73.5	73.5	73.7	73.8
2005	70	70.8	70.4	73.9	70.7
2006	65	70.9	70.9	72.0	70.9
2007	55	68.2	67.9	73.8	68.9
2008	62	70.8	70.8	70.5	70.7
2009	89	71.2	70.7	74.7	71.2
2010	67	73.1	71.6	82.2	72.0
2011	85	75.3	75.3	72.5	75.0
2012	87	73.8	72.8	76.9	73.6
2013	75	74.6	73.8	80.0	74.0
2014	75	74.9	74.8	79.3	74.8
2015	69	74.9	74.7	79.7	74.8
2016	68	76.0	76.1	74.7	75.9
2017	89	78.1	78.1	76.3	78.5
2018	70	76.6	74.3	79.9	79.5
2019	54	77.2	76.7	78.0	76.5
1998-2019	1362	74.1	73.5	76.3	73.8

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 10b

Medians of age at death according to the grouping in Table 9
FEMALES

Year of death	Deaths n	Age at death (all causes) Years	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate) Years
1998	31	76.2	74.2	97.1	76.2
1999	35	76.3	76.8	72.0	76.7
2000	36	75.8	75.5	90.2	76.2
2001	34	78.6	78.3	91.9	78.9
2002	76	80.3	79.6	82.2	80.3
2003	56	79.6	78.8	85.0	79.5
2004	59	78.6	78.8	78.0	78.6
2005	39	77.1	77.4	64.9	77.1
2006	75	78.8	78.3	82.4	77.8
2007	63	77.2	76.9	83.4	77.2
2008	81	78.1	77.9	88.4	78.1
2009	61	79.2	77.9	84.1	77.9
2010	69	79.1	78.6	83.7	79.4
2011	61	77.7	77.6	87.5	77.7
2012	71	76.4	76.0	81.9	76.1
2013	48	78.8	77.8	81.2	78.8
2014	58	77.7	78.1	64.6	78.1
2015	79	79.5	79.2	85.2	79.1
2016	85	79.5	79.0	82.5	79.5
2017	61	77.2	77.1	80.3	77.1
2018	40	77.6	77.1	78.2	76.4
2019	31	79.0	78.2	82.9	77.4
1998-2019	1249	78.1	77.9	82.3	78.0

By 2018, Bavarians' life expectancy at birth is estimated at 79.3 years for boys and 83.8 years for girls.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 11a

Mortality measures (cancer-related death) and mortality-incidence-index by year of death
 MALES

Year of death	Deaths n	Mort. raw	MI-Index raw	Mort. WS	MI-Index WS	Mort. ES	MI-Index ES	Mort. BRD-S	MI-Index BRD-S
1998	19	1.7	0.68	1.0	0.73	1.6	0.70	2.5	0.74
1999	19	1.7	0.49	1.0	0.48	1.5	0.48	1.8	0.47
2000	27	2.4	0.84	1.3	0.78	2.2	0.86	3.0	0.88
2001	29	2.5	0.88	1.4	0.92	2.2	0.86	2.8	0.79
2002	57	3.1	0.68	1.7	0.66	2.6	0.68	3.5	0.71
2003	47	2.5	0.57	1.3	0.56	2.1	0.57	2.8	0.57
2004	42	2.2	0.58	1.1	0.55	1.8	0.57	2.5	0.59
2005	64	3.4	0.86	1.8	0.86	2.6	0.86	3.4	0.88
2006	57	3.0	0.76	1.5	0.74	2.3	0.76	3.1	0.78
2007	50	2.3	0.57	1.2	0.53	1.7	0.53	2.1	0.54
2008	58	2.6	0.56	1.3	0.59	2.0	0.59	2.7	0.58
2009	79	3.5	0.80	1.8	0.83	2.7	0.82	3.5	0.80
2010	60	2.7	0.71	1.3	0.70	2.0	0.71	2.6	0.70
2011	77	3.4	0.84	1.4	0.78	2.4	0.81	3.3	0.83
2012	76	3.3	0.78	1.5	0.77	2.4	0.78	3.2	0.81
2013	67	2.9	0.83	1.2	0.81	1.9	0.83	2.6	0.81
2014	65	2.8	0.70	1.2	0.66	1.9	0.68	2.6	0.71
2015	61	2.6	0.71	1.1	0.74	1.7	0.73	2.3	0.72
2016	59	2.5	0.54	0.9	0.46	1.5	0.49	2.2	0.53
2017	85	3.5	0.71	1.3	0.65	2.2	0.68	3.1	0.71
2018	37	1.5	0.45	0.6	0.41	1.0	0.42	1.3	0.43
2019	32	1.3	0.36	0.5	0.32	0.8	0.34	1.1	0.35
1998-2019	1167	2.6	0.67	1.2	0.65	1.9	0.66	2.6	0.67

Table 11b

Mortality measures (cancer-related death) and mortality-incidence-index
by year of death
FEMALES

Year of death	Deaths n	Mort. raw	MI-Index raw	Mort. WS	MI-Index WS	Mort. ES	MI-Index ES	Mort. BRD-S	MI-Index BRD-S
1998	29	2.5	0.71	1.1	0.66	1.6	0.68	2.0	0.70
1999	32	2.7	0.91	1.0	0.86	1.6	0.86	2.3	0.87
2000	33	2.7	1.22	1.1	1.55	1.7	1.41	2.3	1.24
2001	33	2.7	0.70	0.9	0.58	1.5	0.63	2.3	0.69
2002	71	3.6	0.74	1.3	0.78	2.0	0.75	2.8	0.74
2003	52	2.6	0.64	1.0	0.67	1.5	0.67	2.0	0.65
2004	52	2.6	0.64	0.9	0.59	1.4	0.62	2.0	0.65
2005	38	1.9	0.49	0.7	0.43	1.1	0.44	1.5	0.48
2006	69	3.4	0.78	1.2	0.80	1.9	0.79	2.5	0.79
2007	59	2.6	0.59	0.9	0.56	1.4	0.57	1.9	0.58
2008	77	3.3	0.91	1.1	0.82	1.8	0.85	2.5	0.90
2009	56	2.4	0.74	0.7	0.60	1.2	0.67	1.9	0.76
2010	61	2.6	0.90	0.9	0.79	1.4	0.83	1.9	0.84
2011	56	2.4	0.67	0.9	0.71	1.3	0.68	1.8	0.68
2012	65	2.8	1.00	1.0	0.93	1.5	0.97	2.0	0.97
2013	44	1.8	0.64	0.6	0.58	0.9	0.58	1.3	0.62
2014	55	2.3	0.64	0.7	0.62	1.1	0.64	1.6	0.63
2015	73	3.0	1.03	0.8	0.88	1.4	0.92	2.1	0.99
2016	78	3.2	1.05	0.9	0.92	1.5	0.97	2.2	1.04
2017	54	2.2	0.69	0.7	0.73	1.2	0.71	1.6	0.69
2018	29	1.2	0.60	0.3	0.52	0.6	0.55	0.9	0.60
2019	22	0.9	0.46	0.3	0.36	0.4	0.40	0.6	0.42
1998-2019	1138	2.5	0.75	0.8	0.70	1.3	0.72	1.8	0.74

Table 12

Age distribution of age at death (cancer-related) for period 2007-2019
(incl. multiple malignancies)

Age at death Years	Cases			Males			Females		
	n	%	Cum.%	n	%	Cum.%	n	%	Cum.%
0-4									
5-9									
10-14									
15-19									
20-24									
25-29	2	0.1	0.1			0.0	2	0.3	0.3
30-34	0	0.0	0.1			0.0			0.3
35-39	2	0.1	0.3	1	0.1	0.1	1	0.1	0.4
40-44	6	0.4	0.7	5	0.6	0.7	1	0.1	0.5
45-49	21	1.4	2.0	9	1.1	1.9	12	1.6	2.2
50-54	57	3.7	5.7	38	4.7	6.6	19	2.6	4.8
55-59	82	5.3	11.1	51	6.3	12.9	31	4.3	9.1
60-64	118	7.7	18.8	79	9.8	22.7	39	5.3	14.4
65-69	168	10.9	29.7	106	13.2	35.9	62	8.5	22.9
70-74	250	16.3	46.0	140	17.4	53.2	110	15.1	38.0
75-79	333	21.7	67.7	173	21.5	74.7	160	21.9	59.9
80-84	260	16.9	84.6	119	14.8	89.5	141	19.3	79.3
85+	236	15.4	100.0	85	10.5	100.0	151	20.7	100.0
All ages	1535	100.0		806	100.0		729	100.0	

Table 13

Age-specific mortality (cancer-related) and proportion of all cancers
for period 2007–2019
(incl. multiple malignancies)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	MI-index	Females Age- spec. mortal.	MI-index	Males Prop.all cancers %	Females Prop.all cancers %
0- 4								
5- 9								
10-14								
15-19								
20-24								
25-29		2			0.1	1.00		2.2
30-34								
35-39	1	1	0.0	0.25	0.0	0.25	0.4	0.3
40-44	5	1	0.2	0.56	0.0	0.13	0.9	0.1
45-49	9	12	0.4	0.39	0.5	0.60	0.7	0.8
50-54	38	19	1.6	0.63	0.8	0.58	1.5	0.8
55-59	51	31	2.6	0.56	1.6	0.72	1.2	0.9
60-64	79	39	4.8	0.65	2.2	0.62	1.3	0.8
65-69	106	62	7.0	0.61	3.7	0.58	1.2	1.0
70-74	140	110	10.0	0.63	6.8	0.81	1.3	1.3
75-79	173	160	15.6	0.69	11.6	0.82	1.5	1.8
80-84	119	141	18.1	0.67	14.5	0.84	1.3	1.7
85+	85	151	19.9	0.97	15.6	0.88	1.0	1.4
All ages	806	729					1.3	1.3
Mortality								
Raw			2.7	0.66	2.3	0.77		
WS			1.2	0.64	0.7	0.70		
ES			1.8	0.65	1.2	0.72		
BRD-S			2.5	0.66	1.7	0.75		
PYLL-70								
per 100,000			9.5		6.0			
ES			8.1		5.0			
AYLL-70			8.8		9.5			

Table 14a

Further malignancies in deaths in period 1998-2019
MALES

Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C00 Lip	1	0.4	1	100.0				
C03-C06 Oral cavity	2	0.7	1	50.0			1	50.0
C09-C10 Oropharynx	1	0.4	1	100.0				
C12-C13 Hypopharynx	2	0.7	1	50.0			1	50.0
C15 Oesophagus	2	0.7			1	50.0	1	50.0
C16 Stomach	11	3.9	5	45.5	3	27.3	3	27.3
C17 Small intestine	6	2.1	1	16.7	4	66.7	1	16.7
C18 Colon	26	9.2	17	65.4	6	23.1	3	11.5
C19-C20 Rectum	17	6.0	16	94.1	1	5.9		
C22 Liver	4	1.4	1	25.0	3	75.0		
C23-C24 Bile	5	1.8					5	100.0
C25 Pancreas	9	3.2	1	11.1	3	33.3	5	55.6
C26 GI cancer	1	0.4			1	100.0		
C30-C31 Sinuses	2	0.7	2	100.0				
C32 Larynx	6	2.1	6	100.0				
C33-C34 Lung	13	4.6	4	30.8	3	23.1	6	46.2
C38,C45 Mesothelioma	1	0.4					1	100.0
C43 Malign. melanoma	14	5.0	10	71.4	1	7.1	3	21.4
C44 Skin others	28	9.9	19	67.9	2	7.1	7	25.0
C46,C49 Soft tissue	2	0.7	2	100.0				
C48 Peritoneal	1	0.4			1	100.0		
C60 Penis	1	0.4	1	100.0				
C61 Prostate	77	27.3	64	83.1	2	2.6	11	14.3
C62 Testis	5	1.8	5	100.0				
C64 Kidney	10	3.5	9	90.0	1	10.0		
C65 Renal pelvis	1	0.4	1	100.0				
C67 Bladder	12	4.3	7	58.3	3	25.0	2	16.7
C69 Eye melanoma	2	0.7	1	50.0			1	50.0
C70-C72 CNS cancer	1	0.4	1	100.0				
C73 Thyroid	1	0.4	1	100.0				
C76-C79 CUP	6	2.1	4	66.7	2	33.3		
C81 Hodgkin lymphoma	2	0.7	2	100.0				
C82-C85 NHL	8	2.8	6	75.0	1	12.5	1	12.5
C90 Mult. myeloma	1	0.4	1	100.0				
C91-C96 Leukaemia	1	0.4	1	100.0				
All further malignancies	282	100.0	192	68.1	38	13.5	52	18.4

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a further malignancy.

Table 14b

Further malignancies in deaths in period 1998-2019
FEMALES

Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C03-C06 Oral cavity	3	1.2	3	100.0				
C07-C08 Salivary gland	1	0.4	1	100.0				
C15 Oesophagus	1	0.4			1	100.0		
C16 Stomach	6	2.4	1	16.7			5	83.3
C17 Small intestine	1	0.4	1	100.0				
C18 Colon	30	11.9	20	66.7	3	10.0	7	23.3
C19-C20 Rectum	10	4.0	8	80.0			2	20.0
C22 Liver	1	0.4					1	100.0
C23-C24 Bile	3	1.2			2	66.7	1	33.3
C25 Pancreas	12	4.7	3	25.0	5	41.7	4	33.3
C33-C34 Lung	13	5.1	1	7.7	2	15.4	10	76.9
C37 Thymus	1	0.4			1	100.0		
C43 Malign. melanoma	10	4.0	9	90.0			1	10.0
C44 Skin others	13	5.1	7	53.8	2	15.4	4	30.8
C46,C49 Soft tissue	2	0.8	2	100.0				
C50 Breast	69	27.3	66	95.7	2	2.9	1	1.4
C51 Vulva	3	1.2	3	100.0				
C53 Cervix uteri	7	2.8	6	85.7			1	14.3
C54 Corpus uteri	20	7.9	19	95.0			1	5.0
C56 Ovary	15	5.9	8	53.3	2	13.3	5	33.3
C64 Kidney	8	3.2	5	62.5	2	25.0	1	12.5
C67 Bladder	3	1.2	3	100.0				
C70-C72 CNS cancer	1	0.4	1	100.0				
C73 Thyroid	2	0.8	2	100.0				
C74-C80 Cancer others	1	0.4			1	100.0		
C76-C79 CUP	4	1.6	1	25.0	1	25.0	2	50.0
C82-C85 NHL	10	4.0	8	80.0	1	10.0	1	10.0
C91-C96 Leukaemia	3	1.2					3	100.0
All further malignancies	253	100.0	178	70.4	25	9.9	50	19.8

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a further malignancy.

Table 15

Age-specific mortality (cancer-related) and proportion of all cancers
for period 2007-2019
(First primaries only *)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	Males MI-index	Females Age- spec. mortal.	Females MI-index	Males Prop.all cancers %	Females Prop.all cancers %
0- 4								
5- 9								
10-14								
15-19								
20-24								
25-29		2			0.1	1.00		2.3
30-34								
35-39	1	1	0.0	0.33	0.0	0.33	0.4	0.3
40-44	5	1	0.2	0.71	0.0	0.13	0.9	0.1
45-49	9	12	0.4	0.45	0.5	0.60	0.7	0.9
50-54	33	17	1.4	0.63	0.7	0.59	1.5	0.8
55-59	47	28	2.4	0.57	1.4	0.72	1.3	0.9
60-64	64	32	3.9	0.62	1.8	0.59	1.3	0.8
65-69	91	46	6.0	0.62	2.7	0.55	1.3	0.9
70-74	114	90	8.1	0.67	5.6	0.89	1.3	1.4
75-79	127	121	11.5	0.73	8.8	0.80	1.5	1.8
80-84	83	105	12.6	0.67	10.8	0.80	1.2	1.6
85+	65	127	15.2	1.00	13.2	0.90	1.1	1.5
All ages	639	582					1.3	1.3
Mortality								
Raw			2.1	0.67	1.9	0.76		
WS			1.0	0.65	0.6	0.69		
ES			1.5	0.66	1.0	0.72		
BRD-S			2.0	0.67	1.3	0.74		
PYLL-70								
per 100,000			8.5		5.4			
ES			7.2		4.5			
AYLL-70			9.0		10.2			

* See corresponding tables with multiple malignancies.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers
for period 2007-2019
(**Single primaries only** *)

Age at death Years	Males n	Females n	Males Age- spec. mortal. MI-index	Females Age- spec. mortal. MI-index	Males Prop.all cancers %	Females Prop.all cancers %		
0- 4								
5- 9								
10-14								
15-19								
20-24								
25-29		2		0.1	1.00	2.4		
30-34								
35-39	1	1	0.0	0.33	0.0	0.33	0.4	0.3
40-44	5	1	0.2	0.71	0.0	0.13	1.0	0.1
45-49	9	12	0.4	0.45	0.5	0.60	0.7	0.9
50-54	33	16	1.4	0.63	0.7	0.62	1.5	0.8
55-59	46	25	2.4	0.58	1.3	0.68	1.3	0.9
60-64	62	32	3.8	0.63	1.8	0.62	1.3	0.9
65-69	88	45	5.8	0.62	2.7	0.56	1.3	0.9
70-74	110	86	7.8	0.66	5.4	0.90	1.3	1.4
75-79	123	114	11.1	0.72	8.3	0.81	1.5	1.7
80-84	83	100	12.6	0.69	10.3	0.78	1.3	1.6
85+	61	122	14.3	0.98	12.6	0.87	1.1	1.5
All ages	621	556					1.3	1.3
Mortality								
Raw			2.1	0.67	1.8	0.76		
WS			0.9	0.65	0.6	0.70		
ES			1.4	0.66	0.9	0.72		
BRD-S			1.9	0.67	1.3	0.74		
PYLL-70								
per 100,000			8.3		5.2			
ES			7.1		4.3			
AYLL-70			9.1		10.1			

* See corresponding tables with multiple malignancies.

ICD-10 C24: Malignant neoplasm of other and unspecified parts of biliary tract

Age distribution and age-specific mortality 2007 - 2019 (Males: 806, Females: 729)





Figure 17. Distribution of age at death (bars; males: mean=71.3 yrs, median=73.1 yrs; females: mean=74.9 yrs, median=76.4 yrs) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at biliary tract cancer-related death (see Table 10) should be considered.

Average mortality (Germany 1987 standard population) 2007 - 2019: Males



Average mortality (Germany 1987 standard population) 2007 - 2019: Females





Figure 18a. Map of cancer mortality (german standard population) by county averaged for period 2007 to 2019. According to their individual mortality rates, the counties are displayed in different red and blue hues, being the fine white color attributed to the population mean (males 2.5/100,000 WS N=806, females 1.7/100,000 WS N=729).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,462 female residents (averaged) in the period from 2007 to 2019 a total of 16 women died from biliary tract cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 1.4/100,000 (german standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.7 and 2.7/100,000.

Standardized mortality ratio (SMR) 2007 - 2019: Males



Standardized mortality ratio (SMR) 2007 - 2019: Females




Figure 18b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2019. According to their individual SMR values, the counties are displayed in different red and blue hues, being the fine white color attributed to the population overall of 1.0 (males N=806, females N=729).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,153 female residents (averaged) in the period from 2007 to 2019 a total of 16 women died from biliary tract cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.82. Though, the value of this parameter may vary with an underlying probability of 99% between 0.39 and 1.51, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases, where applicable. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head&neck and lung).

The ratio of mortality and incidence (mortality-to-incidence ratio, **MIR, MI-Index**) is a statistical index that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MIR. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

MCR	Munich Cancer Registry (Tumorregister München)
GEKID	Association of Population-based Cancer Registries in Germany (Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)
SEER	Surveillance, Epidemiology, and End Results (USA)
DCO	Death certificate only
BRD-S	German (FRG) standard population
ES	European standard population (old)
WS	World standard population
SIR	Standardized incidence ratio
CI	Confidence interval
EAR	Excess absolute risk = excess cancer cases (O - E) per 10,000 person-years
PYLL-70	Potential years of life lost prior to age 70 given a person dies before that age
AYLL-70	Average years of life lost prior to age 70 given a person dies before that age
SMR	Standardized mortality ratio
MI-index	Ratio of mortality to incidence, MIR
FRG	Federal Republic of Germany

Recommended Citation

Munich Cancer Registry. ICD-10 C24: Biliary tract cancer - Incidence and Mortality [Internet]. 2021 [updated 2021 Jan 25; cited 2021 Mar 1]. Available from: https://www.tumorregister-muenchen.de/en/facts/base/bC24__E-ICD-10-C24-Biliary-tract-cancer-incidence-and-mortality.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.