Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ Deutsch

ICD-10 C38: Heart, mediastinum and pleura

Incidence and Mortality

Year of diagnosis	1998-2020
Patients	386
Diseases	386
Creation date	12/21/2021
Database export	12/20/2021
Population	4.95 m



Munich Cancer Registry
Cancer Registry Bavaria - Upper Bavaria Regional Center
at Klinikum Grosshadern/IBE
Marchioninistr. 15
Munich, 81377
Germany

https://www.tumorregister-muenchen.de/en

https://www.tumorregister-muenchen.de/en/facts/base/bC38__E-ICD-10-C38-Heart-mediastinum-and-pleura-incidence-and-mortality.pdf

Index of figures and tables

Fig./Tb	d.	Page
1	Annual cases, DCO, mult. malignancies, follow-up / yr	4
2	Incidence by year of diagnosis	7
3	Age distribution parameters by year of diagnosis	8
4	Age distribution by 5-year age group and sex	11
5	Age-specific incidence, DCO rate, proportion malignancies	12
6	Age distribution and age-specific incidence (chart)	13
6a	Age-specific incidence internationally (chart)	14
7	Standardized incidence ratio of further malignancies	15
8a	Map of cancer incidence (BRD-S) by county (chart)	17
8b	Standardized incidence ratio (SIR) by county (chart)	18
9a	Pts incident cohorts and mortality / yr	19
9b	Incidence and mortality by year of diagnosis	20
9с	Cancer-related deaths, death certification available / yr	21
10	Medians of age at death / yr	22
11	Mortality by year of death	24
12	Distribution of age at death	26
13	Age-specific mortality	27
14	Further malignancies in deaths	28
15	Age-specific mortality (first primaries)	30
16	Age-specific mortality (single primaries)	31
17	Age distribution and age-specific mortality (chart)	32
18a	Map of cancer mortality (BRD-S) by county (chart)	33
18b	Standardized mortality ratio (SMR) by county (chart)	34

Global Statements about the statistics on the Internet – Baseline Statistics (grey button ——), Survival (red button ——)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.69 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases### are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, December 2021

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.65 million to 4.10 in 2002, and to 4.69 million in 2007).
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate.

ICD-10 codes (ICD-10 2016) used for specifying cancer site

Code	Description
C38	Malignant neoplasm of heart, mediastinum and pleura
C38.0	Heart
C38.1	Anterior mediastinum
C38.2	Posterior mediastinum
C38.3	Mediastinum, part unspecified
C38.4	Pleura
C38.8	Overlapping lesion of heart, mediastinum and pleura

INCIDENCE

Table 1

Cases with invasive cancer by year of diagnosis, proportions of DCO, further malignancies, deaths, and active follow-up (ALL PATIENTS) (incl. DCO)

				Prop.			
				at least	Prop.		
				1 further	at least		
				malign.	1 further		Prop.
	All	DCO	Prop.	prior +	malign.	Prop.	actively
Year of	cases	cases	DCO	synchron.	after	deaths	followed
diagnosis	n	n	용	%	%	용	용
1998	8	4	50.0	12.5	3.4	100.0	100.0
1999	11	5	45.5	10.5	3.5	100.0	100.0
2000	14	6	42.9	6.1	3.3	85.7	100.0
2001	10			14.0	3.4	100.0	100.0
2002	27	2	7.4	12.9	3.3	77.8	96.3 #
2003	23	/4	17.4	10.8	3.5	95.7	100.0
2004	18	2	11.1	11.7	3.8	100.0	100.0
2005	18	1	5.6	10.9	3.7	94.4	100.0
2006	24	\ 4	16.7	13.7	3.6	95.8	100.0
2007	21	3	14.3	13.8	3.5	90.5	95.2 #
2008	17			15.7	3.8	82.4	100.0
2009	20	1	5.0	15.6	3.6	80.0	100.0
2010	16	2	12.5	15.9	3.4	87.5	100.0
2011	17			16.0	3.8	88.2	100.0
2012	18	1	5.6	16.4	3.5	77.8	100.0
2013	26	3	11.5	16.0	3.3	92.3	100.0
2014	22	1	4.5	15.8	2.1	90.9	100.0
2015	20	7	35.0	17.3	1.3	95.0	100.0
2016	18	2	11.1	17.2	1.8	77.8	100.0
2017	13			17.2	2.7	76.9	100.0
2018	13			17.9	4.0	69.2	100.0
2019	6			17.6	0.0	33.3	100.0
2020	6			17.6	0.0	33.3	100.0 ##
1998-2020	386	48	12.4	17.6	3.4	86.5	99.5

386 cases diagnosed 1998-2020 are related to a total of 386 patients. Currently, in 81 (21.0 %) of these 386 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 58 / 19 / 4 (15.0 % / 4.9 % / 1.0 %) patients exist having 2 / 3 / 4+ malignancies.

How to interpret:

In 2018, a subgroup of 13 cases has been diagnosed, of which 17.9 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 4.0 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

[#] The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

^{##} Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retreived from the respective headings.

Table 1a

Cases with invasive cancer by year of diagnosis, proportions of DCO, further malignancies, deaths, and active follow-up (MALES) (incl. DCO)

			DCO	Prop.	Prop. at least 1 further malign. prior +	Prop. at least 1 further malign.	Prop.	Prop. actively
Year of	Males	Males	cases	DCO	synchron.	after	deaths	followed
diagnosis	n	୪	n	૾ૢ	90	90	00	ଚ୍ଚ
1998	4	50.0	_ 2	50.0	25.0	3.9	100.0	100.0
1999	6	54.5	3	50.0	20.0	3.9	100.0	100.0
2000	8	57.1	4	50.0	11.1	4.0	100.0	100.0
2001	7	70.0	-	30.0	16.0	4.2	100.0	100.0
2002	16	59.3	2	12.5	12.2	3.8	68.8	93.8 #
2003	16	69.6	4	25.0	8.8	4.2	93.8	100.0
2004	13	72.2	1	7.7	11.4	4.5	100.0	100.0
2005	11	61.1	1	9.1	9.9	4.3	100.0	100.0
2006	12	50.0	2	16.7	12.9	4.6	91.7	100.0
2007	12	57.1	1	8.3	12.4	4.3	91.7	91.7 #
2008	8	47.1			13.3	4.7	75.0	100.0
2009	14	70.0			12.6	4.1	71.4	100.0
2010	9	56.3	1	11.1	13.2	3.7	77.8	100.0
2011	11	64.7			12.9	4.1	90.9	100.0
2012	13	72.2	1	7.7	13.8	3.4	84.6	100.0
2013	20	76.9	3	15.0	13.9	2.7	90.0	100.0
2014	12	54.5	1	8.3	13.5	1.9	91.7	100.0
2015	14	70.0	6	42.9	15.0	2.4	100.0	100.0
2016	7	38.9	1	14.3	15.0	3.6	85.7	100.0
2017	6	46.2			15.1	4.8	66.7	100.0
2018	9	69.2			16.2	6.3	66.7	100.0
2019	4	66.7			15.9	0.0	50.0	100.0
2020	3	50.0			16.2	0.0	33.3	100.0 ##
1998-2020	235	60.9	33	14.0	16.2	3.9	86.4	99.1

235 cases diagnosed 1998-2020 are related to a total of 235 patients. Currently, in 45 (19.1 %) of these 235 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 31 / 12 / 2 (13.2 % / 5.1 % / 0.9 %) patients exist having 2 / 3 / 4+ malignancies.

- # The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.
- ## Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retreived from the respective headings.

How to interpret:

In 2018, a subgroup of 9 cases has been diagnosed, of which 16.2 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 6.3 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

Table 1b

Cases with invasive cancer by year of diagnosis, proportions of DCO, further malignancies, deaths, and active follow-up (FEMALES) (incl. DCO)

					Prop.			
					at least	Prop.		
					1 further	at least		
					malign.	1 further		Prop.
			DCO	Prop.	prior +	malign.	Prop.	actively
Year of	Females	Females	cases	DCO	synchron.	after	deaths	followed
diagnosis	n	용	n	용	용	90	%	િ
1998	4	50.0	2	50.0	0.0	2.7	100.0	100.0
1999	5	45.5	2	40.0	0.0	2.8	100.0	100.0
2000	6	42.9	2	33.3	0.0	2.2	66.7	100.0
2001	3	30.0			11.1	2.3	100.0	100.0
2002	11	40.7			13.8	2.3	90.9	100.0 #
2003	7	30.4			13.9	2.5	100.0	100.0
2004	5	27.8	1	20.0	12.2	2.7	100.0	100.0
2005	7	38.9			12.5	2.8	85.7	100.0
2006	12 /	50.0	2	16.7	15.0	2.0	100.0	100.0
2007	9	42.9	2	22.2	15.9	2.2	88.9	100.0 #
2008	9	52.9			19.2	2.5	88.9	100.0
2009	6	30.0	1	16.7	20.2	2.7	100.0	100.0
2010	7\	43.8	1	14.3	19.8	3.0	100.0	100.0
2011	6	35.3			20.6	3.3	83.3	100.0
2012	5	27.8			20.6	3.7	60.0	100.0
2013	6	23.1			19.4	4.1	100.0	100.0
2014	10	45.5			19.5	2.3	90.0	100.0
2015	6	30.0	1	16.7	21.0	0.0	83.3	100.0
2016	11	61.1	1	9.1	20.7	0.0	72.7	100.0
2017	7	53.8			20.4	0.0	85.7	100.0
2018	4	30.8			20.5	0.0	75.0	100.0
2019	2	33.3			20.3	0.0		100.0
2020	3	50.0			19.9	0.0	33.3	100.0 ##
1998-2020	151	39.1	15	9.9	19.9	2.7	86.8	100.0

151 cases diagnosed 1998-2020 are related to a total of 151 patients. Currently, in 36 (23.8 %) of these 151 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 27 / 7 / 2 (17.9 % / 4.6 % / 1.3 %) patients exist having 2 / 3 / 4+ malignancies.

- # The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.
- ## Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retreived from the respective headings.

How to interpret:

In 2018, a subgroup of 4 cases has been diagnosed, of which 20.5 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 0.0 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

Table 2

Incidence measures by year of diagnosis including DCO cases (with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.94 m as of 2007, respectively)

			M-1	F	Malaa	П	Malaa		Malaa	E
V	Malaa	E-m-l	Males	-/		Fem.	Males		Males	
Year of		Females	Inc.							
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BKD-2
1000	4	4	0.4	0 2	0 0	0 1	0 2	0 0	0 4	0 0
1998	4	4	0.4	0.3	0.2	0.1	0.3	0.2	0.4	0.3
1999	6	5	0.5	0.4	0.3	0.3	0.5	0.3	0.8	0.4
2000	8	6	0.7	0.5	0.4	0.2	0.6	0.3	0.8	0.4
2001	7	3	0.6	0.2	0.3	0.2	0.5	0.2	0.7	0.2
2002	16	11 🛴	0.9	0.6	0.6	0.3	0.8	0.4	1.0	0.5
2003	16	7	0.9	0.4	0.6	0.2	0.8	0.2	0.9	0.3
2004	13	5	0.7	0.3	0.4	0.1	0.6	0.2	0.7	0.2
2005	11	7	0.6	0.4	0.4	0.2	0.5	0.2	0.6	0.3
2006	12	12	0.6	0.6	0.3	0.3	0.5	0.4	0.6	0.5
2007	12	9	0.5	0.4	0.4	0.2	0.5	0.2	0.6	0.3
2008	8	9	0.4	0.4	0.2	0.2	0.3	0.3	0.3	0.3
2009	14	6	0.6	0.3	0.5	0.1	0.6	0.2	0.6	0.2
2010	9	7/	0.4	0.3	0.3	0.2	0.4	0.2	0.4	0.3
2011	11/	6	0.5	0.3	0.2	0.1	0.3	0.2	0.5	0.2
2012	13	5	0.6	0.2	0.3	0.2	0.4	0.2	0.6	0.2
2013	20	6	0.9	0.3	0.5	0.1	0.6	0.2	0.8	0.2
2014	12	10	0.5	0.4	0.3	0.2	0.4	0.3	0.5	0.3
2015	14	6	0.6	0.2	0.2	0.1	0.4	0.2	0.5	0.2
2016	7	11	0.3	0.4	0.1	0.2	0.2	0.3	0.3	0.4
2017	6	7	0.2	0.3	0.2	0.1	0.2	0.2	0.2	0.2
2018	9	4	0.4	0.2	0.2	0.1	0.3	0.1	0.3	0.1
2019	4	2	0.2	0.1	0.1	0.1	0.1	0.1		0.1
2020	3	3	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
2020	J	S	V	· · ·	··-	V • I	J, 1	Ŭ • ·	· · ·	V • ±
1998-2020	235	151	0.5	0.3	0.3	0.2	0.4	0.2	0.5	0.3
7					, , ,					

The computation of the incidence measures includes all cancers, irrespective of first or subsequent malignancy.

Table 3 $\label{eq:Age_age} \mbox{Age distribution parameters by year of diagnosis (ALL PATIENTS) } \mbox{(incl. DCO)}$

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	8	65.8	13,4	51.1	83.9	51.1	56.6	58.7	80.3	83.9
1999	11	63.3	15.8	38.6	86.7	47.2	51.7	60.8	80.3	81.7
2000	14	71.8	16.0	36.7	95.3	57.1	61.0	73.3	78.1	93.7
2001	10	64.8	10.7	44.6	78.7	47.7	62.7	65.5	69.7	78.5
2002	27	63.3	19.5	14.1	89.7	40.7	53.8	64.3	80.3	83.6
2003	23	67.5	18.8	0.3	92.4	53.9	61.1	67.9	80.3	85.8
2004	18	65.6	13.2	35.7	83.0	48.2	53.2	67.1	78.3	82.0
2005	18	64.9	13.4	33.4	86.5	48.5	55.0	66.7	77.5	78.7
2006	24	70.3	11.8	46.5	92.2	51.8	63.9	72.4	77.8	84.1
2007	21	63.9	18.3	22.0	84.2	39.9	54.6	68.1	77.9	82.6
2008	17	68.5	12.7	43.3	93.9	47.0	65.9	67.2	77.6	81.6
2009	20	57.2	15.4	25.6	89.7	36.4	49.8	56.3	65.3	77.8
2010	16	61.0	15.2	29.7	93.2	34.7	56.6	61.8	69.6	74.8
2011	17	65.3	15.3	28.1	82.5	37.7	62.6	70.4	74.4	81.0
2012	18	63.1	19.1	18.5	90.7	27.0	53.1	66.5	78.6	83.7
2013	26	68.0	13.2	22.1	86.7	50.7	62.6	69.8	75.2	80.6
2014	22	71.7	14.1	24.3	93.6	61.4	64.9	75.0	78.7	87.6
2015	20	73.4	14.3	43.5	90.4	51.7	61.2	77.8	85.3	87.9
2016	18	67.2	21.1	18.3	84.9	27.9	57.8	78.5	81.5	83.4
2017	13 \	68.4	20.7	25.8	95.7	39.5	59.1	73.9	81.2	87.5
2018	13	66.6	12.7	39.1	82.1	49.7	62.5	65.6	76.9	78.5
2019	6	53.0	25.7	31.3	98.7	31.3	33.7	43.9	66.5	98.7
2020	6	62.7	13.9	37.8	78.2	37.8	60.7	63.6	72.5	78.2
1998-2020	386	66.1	16.1	0.3	98.7	44.7	57.3	67.8	77.9	83.6

Table 3a

Age distribution parameters by year of diagnosis (MALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	4	61.1	11,0	51.1	76.8	51.1	54.1	58.2	68.1	76.8
1999	6	68.5	12.6	53.3	81.7	53.3	57.0	69.5	80.3	81.7
2000	8	73.0	10.8	59.3	93.7	59.3	65.0	73.3	77.1	93.7
2001	7	69.0	6.9	62.7	78.7	62.7	62.8	65.6	78.3	78.7
2002	16	62.5	21.5	14.1	89.7	17.7	55.2	64.1	80.4	83.6
2003	16	65.9	21.2	0.3	92.4	50.0	59.0	67.6	78.2	85.8
2004	13	63.3	14.1	35.7	82.0	48.2	52.8	66.8	74.5	79.2
2005	11	60.3	14.4	33.4	78.7	48.5	52.3	57.6	77.6	77.6
2006	12	69.4	8.6	46.5	78.2	63.0	66.6	71.4	75.3	77.4
2007	12	60.9	17.4	22.2	84.2	39.9	54.5	61.5	73.7	77.9
2008	8	68.0	14.5	47.0	93.9	47.0	60.6	66.7	74.4	93.9
2009	14	54.3	16.6	25.6	89.7	29.1	44.7	53.8	63.4	72.2
2010	9	58.6	18.8	29.7	93.2	29.7	55.9	59.1	65.1	93.2
2011	11	65.5	18.4	28.1	82.5	37.7	52.5	71.0	80.6	81.0
2012	13 /	67.5	16.2	27.0	90.7	52.3	62.0	67.9	78.6	83.3
2013	20	66.5	14.2	22.1	85.4	49.8	60.9	69.6	75.4	79.7
2014	12	71.0	17.3	24.3	93.6	64.2	64.9	74.7	78.8	87.6
2015	14	78.4	12.1	50.8	90.4	54.0	77.0	81.7	86.3	89.4
2016	7	64.8	24.4	27.9	84.9	27.9	32.4	79.2	83.4	84.9
2017	6	59.3	26.8	25.8	95.7	25.8	39.5	58.6	77.4	95.7
2018	9	65.1	14.5	39.1	82.1	39.1	56.5	65.6	76.9	82.1
2019	4	61.2	28.4	33.7	98.7	33.7	39.8	56.3	82.6	98.7
2020	3	68.4	8.6	62.3	78.2	62.3	62.3	64.9	78.2	78.2
1998-2020	235	65.5	16.8	0.3	98.7	44.1	56.5	67.3	77.6	83.4

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	4	70.5	15.5	56.2	83.9	56.2	57.1	70.9	83.9	83.9
1999	5	57.0	18.4	38.6	86.7	38.6	47.2	51.7	60.8	86.7
2000	6	70.2	22.2	36.7	95.3	36.7	57.1	70.6	91.1	95.3
2001	3	55.1	13.1	44.6	69.7	44.6	44.6	50.9	69.7	69.7
2002	11	64.4	17.3	40.7	85.4	40.9	41.5	71.4	80.3	81.0
2003	7	71.1	12.3	53.9	87.6	53.9	61.1	70.9	80.9	87.6
2004	5	71.7	9.0	63.8	83.0	63.8	64.3	67.9	79.6	83.0
2005	7	72.0	7.9	63.9	86.5	63.9	66.5	69.0	77.5	86.5
2006	12	71.3	14.6	47.4	92.2	51.8	58.4	75.1	82.8	85.5
2007	9	67.9	19.8	22.0	83.2	22.0	65.4	74.6	81.5	83.2
2008	9	68.9	11.9	43.3	79.8	43.3	66.9	71.3	77.6	79.8
2009	6	64.1	10.5	56.1	83.5	56.1	56.1	60.4	68.0	83.5
2010	7	64.2	9.4	46.8	74.8	46.8	61.0	62.9	73.8	74.8
2011	6	65.0	8.6	50.5	74.4	50.5	62.6	64.8	73.1	74.4
2012	5 /	51.5	23.2	18.5	83.7	18.5	48.5	53.1	53.9	83.7
2013	6	72.7	8.1	63.5	86.7	63.5	68.1	71.2	75.2	86.7
2014	10	72.6	10.0	57.4	89.6	59.4	62.6	75.4	78.7	84.7
2015	6	61.7	12.9	43.5	77.8	43.5	52.6	61.2	73.7	77.8
2016	11	68.7	19.8	18.3	82.7	50.0	57.8	77.8	81.5	81.7
2017	7 \	76.3	9.9	59.1	87.5	59.1	71.2	74.5	86.8	87.5
2018	4	70.1	7.8	62.5	77.5	62.5	63.4	70.2	76.8	77.5
2019	2	36.5	7.3	31.3	41.7	31.3	31.3	36.5	41.7	41.7
2020	3	57.0	17.6	37.8	72.5	37.8	37.8	60.7	72.5	72.5
1998-2020	151	67.1	15.0	18.3	95.3	47.2	58.5	68.1	79.2	83.7

Age at								
diagnosis	Cases		Males			Females		
Years	n	% Cum.%	'n	%	Cum.%	n	%	Cum.%
0 - 4								
5-9								
10-14								
15-19	2	0.9 0.9			0.0	2	2.2	2.2
20-24	4	1.7 2.6	3	2.1	2.1/	1	1.1	3.3
25-29	7	3.0 5.6	7	4.9	7.0			3.3
30-34	4	1.7 7.3	3	2.1	9.2	1	1.1	4.4
35-39	5	2.1 9.4	4	2.8	12.0	1	1.1	5.5
40 - 44	6	2.6 12.0	3	2.1	14.1	3	3.3	8.8
45-49	7	3.0 15.0	5	3.5	17.6	2	2.2	11.0
50-54	15	6.4 21.5	9	6.3	23.9	6	6.6	17.6
55-59	20	8.6 30.0	12	8.5	32.4	8	8.8	26.4
60-64	26	11.2 41.2	13	9.2	41.5	13	14.3	40.7
65-69	28	12.0 53.2	18	12.7	54.2	10	11.0	51.6
70-74	32	13.7 67.0	18	12.7	66.9	14	15.4	67.0
75-79	34	14.6 81.5	18	12.7	79.6	16	17.6	84.6
80-84	24	10.3 91.8	14	9.9	89.4	10	11.0	95.6
85+	19	8.2 100.0	15	10.6	100.0	4	4.4	100.0
All ages	233	100.0	142	100.0		91	100.0	
J -								

Table 5 $\label{eq:Age-specific} \mbox{Age-specific incidence, DCO rate and proportion of all cancers} \\ \mbox{for period 2007-2020}$

							Males	Females
			Males	Females	Males	Females		Prop.all
Age at			Age-	Age-	DCO rate	DCO rate	cancers	cancers
diagnosis	Males	Females	spec.	spec.	n=14	n=6	n=153686	n=155051
Years	n	n	incid.	incid.	%	%	%	%
0 - 4								
5- 9								
10-14								
15-19		2		0.1				0.8
20-24	3	1	0.1	0.1			0.5	0.2
25-29	7		0.3				0.7	
30-34	3	1	0.1	0.0			0.2	0.0
35-39	4	1	0.2	0.0			0.2	0.0
40 - 44	3	3	0.1	0.1			0.1	0.0
45-49	5	2	0.2	0.1			0.1	0.0
50-54	9	6	0.4	0.2			0.1	0.0
55-59	12	8	0.6	0.4			0.1	0.1
60-64	13	13 /	0.7	0.7		15.4	0.1	0.1
65-69	18	10	1.1	0.6	5.6		0.1	0.1
70-74	18	14	1.2	0.8	11.1		0.1	0.1
75-79	18	16	1.5	1.1	11.1	6.3	0.1	0.1
80-84	14	10	1.9	0.9	14.3	30.0	0.1	0.1
85+	15	4	3.2	0.4	46.7		0.1	0.0
All ages	142	91			9.9	6.6	0.1	0.1
Incidence								
Raw			0.4	0.3				
WS			0.2	0.1				
ES			0.3	0.2				
BRD-S			0.4	0.2				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

ICD-10 C38: Malignant neoplasm of heart, mediastinum and pleura Age distribution and age-specific incidence 2007 - 2020 (Males: 142, Females: 91)




Figure 6. Age distribution (males: mean=65.4 yrs, median=67.2 yrs; females: mean=66.6 yrs, median=68.1 yrs) and age-specific incidence.



ICD-10 C38: Malignant neoplasm of heart, mediastinum and pleura Age-specific incidence rates: international comparison




Figure 6a. Age-specific incidence in MCR registry areas compared to Germany (FRG, RKI estimates) and SEER (Surveillance, Epidemiology, and End Results, USA).



Reference:

Estimated age-specific patient population of Germany, latest update: 16 March 2021. German Centre for Cancer Registry Data, Robert Koch Institute (RKI), based on data of the population based cancer registries. http://www.krebsdaten.de. Last access: 08/17/2021 Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 21 Regs Research Data, released April 2021, based on the November 2020 submission. http://www.seer.cancer.gov.

Table 7a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of further malignancies for period 1998-2020

MALES

	Observed	Expected		CI	CI		DCO
Diagnosis	/ n /	n	SIR	95%	95%	EAR	%
C18 Colon	2	0.3	6.4	0.8	23.0	55.8	100.0
C32 Larynx	/ 1/	0.0	29.5	0.7	164.1	32.0	100.0
C38,C45 Mesothelioma	/ 1/	0.0	47.4	1.2	264.0 #	32.4	
C61 Prostate	1	0.9	1.1	0.0	6.3	3.9	
C62 Testis	1	0.0	46.1	1.2	256.6 #	32.4	
C82-C85 NHL	1	0.1	7.2	0.2	40.1	28.5	100.0
C90 Mult. myeloma	1	0.0	24.2	0.6	134.6	31.7	
Not observed	0	1.8	0.0	0.0	2.0	-60.9	
All further malignancies	8	3.3	2.4	1.0	4.8 #	155.7	50.0
Patients		207					
Median age at next maligna	ncy (years	s) 71.3					
Person-years		302					
Mean observation time (yea	rs)	1.5					
Median observation time (y	ears)	0.5					

The occurrence of further specified malignancy is statistically significant.

Table 7b


Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of further malignancies for period 1998-2020

FEMALES

		Observed	Expected		CI	CI		DCO
Diagnos	is	/ n /	n	SIR	95%	95%	EAR	%
C33-C34	Lung	/ 1/	0.2	6.3	0.2	34.9	42.2	
C43	Malign. melanoma	/ 1/	0.1	12.7	0.3	70.9	46.3	
C50	Breast	4	0.7	6.1	1.7	15.6	# 167.9	50.0
C81	Hodgkin lymphoma	4	0.0	268.7	6.8	1497	# 50.0	
Not obs	erved	0	1.1	0.0	0.0	3.3	-56.9	
All fur	ther malignancies	7	2.0	3.4	1.4	7.1	# 249.5	28.6
Patients			139					
Median age	at next malignan	cy (years)	80.8					
Person-year	rs		199					
Mean obser	vation time (year	s)	1.4					
Median obs	ervation time (ye	ars)	0.7					

The occurrence of further specified malignancy is statistically significant.

Average incidence (Germany 1987 standard population) 2007 - 2020: Males



werage incidence (Germany 1987 standard population) 2007 - 2020: Females





Figure 8a. Map of cancer incidence (german standard population, incl. DCO cases) by county averaged for period 2007 to 2020. According to their individual incidence rates, the counties are displayed in different red and blue hues, being the fine white color attributed to the population mean (males 0.4/100,000 WS N=142, females 0.2/100,000 WS N=91).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,727 female residents (averaged) in the period from 2007 to 2020 a total of 3 women were identified with newly diagnosed heart, mediastinum and pleura. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.3/100,000 (german standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.0/100,000.

Standardized incidence ratio (SIR) 2007 - 2020: Males



Standardized incidence ratio (SIR) 2007 - 2020: Females




Figure 8b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2020. According to their individual SIR values, the counties are displayed in different red and blue hues, being the fine white color attributed to the population overall of 1.0 (males N=142, females N=91).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,153 female residents (averaged) in the period from 2007 to 2020 a total of 3 women were identified with newly diagnosed heart, mediastinum and pleura. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 1.20. Though, the value of this parameter may vary with an underlying probability of 99% between 0.13 and 4.38, and is therefore not statistically striking.

MORTALITY

Table 9a

Annual cohorts: Incident cancers, follow-up status, proportion of DCO, deaths among the annual cohorts and proportion of available death certificates (with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.94 m as of 2007, respectively)

						Prop.
		Prop.				deaths
	Incident	actively	Prop.		Prop.	with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	90	00	n	%	90
1998	8	100.0	50.0	8	100.0	100.0
1999	11	100.0	45.5	11	100.0	90.9
2000	14	100.0	42.9	12	85.7	100.0
2001	10	100.0		10	100.0	90.0
2002	27	96.3	7.4	21/	77.8	100.0
2003	23	100.0	17.4	22	95.7	100.0
2004	18	100.0	11.1	18	100.0	94.4
2005	18	100.0	5.6	17	94.4	94.1
2006	24	100.0	16.7	23	95.8	100.0
2007	21	95.2	14.3	19	90.5	100.0
2008	17	100.0		14	82.4	100.0
2009	20	100.0	5.0	16	80.0	100.0
2010	16	100.0	12.5	14	87.5	100.0
2011	17	100.0		15	88.2	93.3
2012	18	100.0	5.6	14	77.8	92.9
2013	26	100.0	11.5	24	92.3	100.0
2014	22	100.0	4.5	20	90.9	90.0
2015	20	100.0	35.0	19	95.0	100.0
2016	18	100.0	11.1	14	77.8	100.0
2017	13	100.0		10	76.9	80.0
2018	13	100.0		9	69.2	66.7
2019	6	100.0		2	33.3	100.0
2020	6	100.0		2	33.3	100.0
1998-2020	386	99.5	12.4	334	86.5	96.1

Table 9b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased within the same year of being diagnosed with cancer (incl. DCO)

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.94 m as of 2007, respectively)

			Prop.		
			deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n /	n	ଚ୍ଚ	n	ଚ୍ଚ
1998	8	7	100.0	6	75.0
1999	11	5	100.0	4	36.4
2000	14	13	92.3	8	57.1
2001	10	13	100.0	5	50.0
2002	27	18	100.0	9	33.3
2003	23	24	100.0	11	47.8
2004	18	17	100.0	6	33.3
2005	18	16	93.8	9	50.0
2006	24	19	89.5	12	50.0
2007	21 /	23	100.0	14	66.7
2008	17	16	100.0	5	29.4
2009	20	13	100.0	5	25.0
2010	16	15	100.0	7	43.8
2011	17\	17	100.0	7	41.2
2012	18	11	100.0	6	33.3
2013	26	22	95.5	13	50.0
2014	22	11	100.0	7	31.8
2015	20	21	100.0	13	65.0
2016	18	19	94.7	9	50.0
2017	13	18	100.0	7 /	53.8
2018	13	10	60.0	5	38.5
2019	6	5	20.0		
2020	6	10	100.0	2	33.3
1998-2020	386	343	95.9	170	44.0

Table 9c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.94 m as of 2007, respectively)

				Prop.
				cancer
		Prop.	Prop.	recorded
		cancer-	non-cancer-	on death
Year of	Deaths	related	related	certificate
death	n/	90	ୃଚ	%
1998	7	71.4	28.6	100.0
1999	5	60.0	40.0	100.0
2000	13	92.3	7.7	100.0
2001	13	92.3	7.7	100.0
2002	18	94.4	5.6	100.0
2003	24	91.7	8.3	95.8
2004	17	88.2	11.8	94.1
2005	16	87.5	12.5	93.3
2006	/ 19	84.2	15.8	94.1
2007	23	100.0		95.7
2008	16	100.0		93.8
2009	13	92.3	7.7	100.0
2010	\ 15	100.0		100.0
2011	17	94.1	5.9	94.1
2012	11	81.8	18.2	81.8
2013	22	90.9	9.1	90.5
2014	11	81.8	18.2	90.9
2015	21	90.5	9.5	90.5
2016	19	89.5	10.5	88.9
2017	18	88.9	11.1	94.4
2018	10	70.0	30.0	83.3
2019	5	80.0	20.0	100.0
2020	10	70.0	30.0	80.0
1998-2020	343	89.2	10.8	93.9

 $\begin{tabular}{ll} Table 10a \\ \hline \begin{tabular}{ll} Medians of age at death according to the grouping in Table 9 \\ \hline \begin{tabular}{ll} MALES \end{tabular}$

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(non-cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
acacii	11	rears	icais	rearb	Icaib
1998	5	59.8	57.6	76.4	59.8
1999	3	80.3	81.0	76.9	80.3
2000	8	66.7	66.7		66.7
2001	8	67.8	67.8		67.8
2002	9	79.0	77.6	79.0	79.0
2003	16	66.2	66.2	71.2	65.2
2004	12	69.8	69.3	78.3	69.8
2005	9	66.1	62.2	78.9	67.3
2006	13	67.7	67.7	67.7	67.7
2007	11/	71.5	71.5		71.5
2008	1.4	66.5	66.5		66.1
2009	8	56.7	49.5	77.6	56.7
2010	6	62.6	62.6		62.6
2011	10	65.6	65.3	86.0	65.3
2012	8	78.6	80.8	73.4	80.8
2013	16	72.8	71.9	77.7	72.6
2014	6	78.4	81.6	64.4	78.4
2015	14	79.9	78.0	90.4	78.0
2016	11	76.7	77.1	74.0	76.7
2017	8	70.6	74.1	65.2	73.1
2018	7	65.7	64.7	76.8	64.7
2019	2	61.2	61.2		
2020	6	67.9	67.5	99.4	67.5
1998-2020	210	69.9	69.1	76.9	69.5

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Age at death death death death death (according to death) Year of death n n Years Years Years Years Years 1998						
Year of death Deaths causes) causes) related) related) <td></td> <td></td> <td>- /</td> <td>/ -</td> <td>٦ - ١</td> <td>death</td>			- /	/ -	٦ - ١	death
Year of death Deaths of death causes) related) related) related) related) certificate) 1998 2 70.1 70.1 70.1 1999 2 61.7 84.8 38.6 61.7 2000 5 57.1 57.4 57.1 55.4 2001 5 62.6 61.7 88.8 62.6 2002 9 71.5 71.5 71.5 2003 8 70.4 70.4 70.4 2004 5 64.7 59.9 83.1 59.9 2005 7 72.9 71.2 80.6 71.2 2006 6 70.9 70.9 70.9 2007 12 78.0 78.0 76.2 2008 2 72.4 72.4 72.4 2009 5 72.2 72.2 72.2 2010 9 62.6 62.6 62.6 2011 7 74.1 74.						_
death n Years Years Years Years 1998 2 70.1 70.1 70.1 1999 2 61.7 84.8 38.6 61.7 2000 5 57.1 57.4 57.1 55.4 2001 5 62.6 61.7 88.8 62.6 2002 9 71.5 71.5 71.5 71.5 2003 8 70.4 70.4 70.4 70.4 70.4 2004 5 64.7 59.9 83.1 59.9 9 2005 7 72.9 71.2 80.6 71.2 70.9 2005 7 72.9 71.2 80.6 71.2 70.9 2006 6 70.9				/	. /	
1998 2 70.1 70.1 1999 2 61.7 84.8 38.6 61.7 2000 5 57.1 57.4 57.1 55.4 2001 5 62.6 61.7 88.8 62.6 2002 9 71.5 71.5 71.5 2003 8 70.4 70.4 70.4 2004 5 64.7 59.9 83.1 59.9 2005 7 72.9 71.2 80.6 71.2 2006 6 70.9 70.9 70.9 2007 12 78.0 78.0 76.2 2008 2 72.4 72.4 72.4 2009 5 72.2 72.2 72.2 2010 9 62.6 62.6 62.6 2011 7 74.1 74.1 74.1 2012 3 63.5 63.5 63.5 2013 6 74.8 74.8 74.8 2014 5 78.9 73.9 9			/	/		
1999 2 61.7 84.8 38.6 61.7 2000 5 57.1 57.4 57.1 55.4 2001 5 62.6 61.7 88.8 62.6 2002 9 71.5 71.5 71.5 2003 8 70.4 70.4 70.4 2004 5 64.7 59.9 83.1 59.9 2005 7 72.9 71.2 80.6 71.2 2006 6 70.9 70.9 70.9 2007 12 78.0 78.0 76.2 2008 2 72.4 72.4 72.4 2009 5 72.2 72.2 72.2 2010 9 62.6 62.6 62.6 2011 7 74.1 74.1 74.1 2012 3 63.5 63.5 63.5 2013 6 74.8 74.8 74.8 2014 5 78.9 73.9 90.5 73.9 2015 7 7	death	n	Years	Years	Years	Years
1999 2 61.7 84.8 38.6 61.7 2000 5 57.1 57.4 57.1 55.4 2001 5 62.6 61.7 88.8 62.6 2002 9 71.5 71.5 71.5 2003 8 70.4 70.4 70.4 2004 5 64.7 59.9 83.1 59.9 2005 7 72.9 71.2 80.6 71.2 2006 6 70.9 70.9 70.9 2007 12 78.0 78.0 76.2 2008 2 72.4 72.4 72.4 2009 5 72.2 72.2 72.2 2010 9 62.6 62.6 62.6 2011 7 74.1 74.1 74.1 2012 3 63.5 63.5 63.5 2013 6 74.8 74.8 74.8 2014 5 78.9 73.9 90.5 73.9 2015 7 7	1000	0	70.1	70.1		70.1
2000 5 57.1 57.4 57.1 55.4 2001 5 62.6 61.7 88.8 62.6 2002 9 71.5 71.5 71.5 2003 8 70.4 70.4 70.4 2004 5 64.7 59.9 83.1 59.9 2005 7 72.9 71.2 80.6 71.2 2006 6 70.9 70.9 70.9 2007 12 78.0 78.0 76.2 2008 2 72.4 72.4 72.4 2009 5 72.2 72.2 72.2 2010 9 62.6 62.6 62.6 2011 7 74.1 74.1 74.1 2012 3 63.5 63.5 63.5 2013 6 74.8 74.8 74.8 2014 5 78.9 73.9 90.5 73.9 2015 7 73.4 70.3 73.4 70.3 2017 10					20/6	
2001 5 62.6 61.7 88.8 62.6 2002 9 71.5 71.5 71.5 2003 8 70.4 70.4 70.4 2004 5 64.7 59.9 83.1 59.9 2005 7 72.9 71.2 80.6 71.2 2006 6 70.9 70.9 70.9 70.9 2007 12 78.0 78.0 76.2 2008 2 72.4 72.4 72.4 2009 5 72.2 72.2 72.2 2010 9 62.6 62.6 62.6 2011 7 74.1 74.1 74.1 74.1 2012 3 63.5 63.5 63.5 2013 6 74.8 74.8 74.8 2014 5 78.9 73.9 90.5 73.9 2015 7 73.4 70.3 73.4 70.3 2016 8 80.6 79.5 90.0 79.5 <t< td=""><td></td><td></td><td></td><td></td><td></td><td></td></t<>						
2002 9 71.5 71.5 71.5 2003 8 70.4 70.4 70.4 2004 5 64.7 59.9 83.1 59.9 2005 7 72.9 71.2 80.6 71.2 2006 6 70.9 70.9 70.9 2007 12 78.0 78.0 76.2 2008 2 72.4 72.4 72.4 2009 5 72.2 72.2 72.2 2010 9 62.6 62.6 62.6 2011 7 74.1 74.1 74.1 2012 3 63.5 63.5 2013 6 74.8 74.8 74.8 2014 5 78.9 73.9 90.5 73.9 2015 7 73.4 70.3 73.4 70.3 2016 8 80.6 79.5 90.0 79.5 2017 10 75.9 75.9 75.9 75.9 2018 3 65.3						
2003 8 70.4 70.4 70.4 2004 5 64.7 59.9 83.1 59.9 2005 7 72.9 71.2 80.6 71.2 2006 6 70.9 70.9 70.9 2007 12 78.0 78.0 76.2 2008 2 72.4 72.4 72.4 2009 5 72.2 72.2 72.2 2010 9 62.6 62.6 62.6 2011 7 74.1 74.1 74.1 2012 3 63.5 63.5 63.5 2013 6 74.8 74.8 74.8 2014 5 78.9 73.9 90.5 73.9 2015 7 73.4 70.3 73.4 70.3 2016 8 80.6 79.5 90.0 79.5 2017 10 75.9 75.9 75.9 2018 3 65.3 65.1 78.0 65.1 2019 3					88.8	
2004 5 64.7 59.9 83.1 59.9 2005 7 72.9 71.2 80.6 71.2 2006 6 70.9 70.9 70.9 2007 12 78.0 78.0 76.2 2008 2 72.4 72.4 72.4 2009 5 72.2 72.2 72.2 2010 9 62.6 62.6 62.6 2011 7 74.1 74.1 74.1 2012 3 63.5 63.5 63.5 2013 6 74.8 74.8 74.8 2014 5 78.9 73.9 90.5 73.9 2015 7 73.4 70.3 73.4 70.3 2016 8 80.6 79.5 90.0 79.5 2017 10 75.9 75.9 75.9 2018 3 65.3 65.1 78.0 65.1 2019 3 75.9 75.3 79.1 74.6						
2005 7 72.9 71.2 80.6 71.2 2006 6 70.9 70.9 70.9 2007 12 78.0 78.0 76.2 2008 2 72.4 72.4 72.4 2009 5 72.2 72.2 72.2 2010 9 62.6 62.6 62.6 2011 7 74.1 74.1 74.1 2012 3 63.5 63.5 63.5 2013 6 74.8 74.8 74.8 2014 5 78.9 73.9 90.5 73.9 2015 7 73.4 70.3 73.4 70.3 2016 8 80.6 79.5 90.0 79.5 2017 10 75.9 75.9 75.9 2018 3 65.3 65.1 78.0 65.1 2019 3 75.9 75.3 79.1 74.6	2003		70.4	70.4		70.4
2006 6 70.9 70.9 70.9 2007 12 78.0 78.0 76.2 2008 2 72.4 72.4 72.4 2009 5 72.2 72.2 72.2 2010 9 62.6 62.6 62.6 2011 7 74.1 74.1 74.1 2012 3 63.5 63.5 63.5 2013 6 74.8 74.8 74.8 2014 5 78.9 73.9 90.5 73.9 2015 7 73.4 70.3 73.4 70.3 2016 8 80.6 79.5 90.0 79.5 2017 10 75.9 75.9 75.9 2018 3 65.3 65.1 78.0 65.1 2019 3 75.9 75.3 79.1 74.6	2004		64.7	59.9	83.1	59.9
2007 12 78.0 78.0 76.2 2008 2 72.4 72.4 72.4 2009 5 72.2 72.2 72.2 2010 9 62.6 62.6 62.6 2011 7 74.1 74.1 74.1 2012 3 63.5 63.5 63.5 2013 6 74.8 74.8 74.8 2014 5 78.9 73.9 90.5 73.9 2015 7 73.4 70.3 73.4 70.3 2016 8 80.6 79.5 90.0 79.5 2017 10 75.9 75.9 75.9 2018 3 65.3 65.1 78.0 65.1 2019 3 75.9 75.3 79.1 74.6	2005	7	72.9	71.2	80.6	71.2
2008 2 72.4 72.4 72.4 2009 5 72.2 72.2 72.2 2010 9 62.6 62.6 62.6 2011 7 74.1 74.1 74.1 2012 3 63.5 63.5 63.5 2013 6 74.8 74.8 74.8 2014 5 78.9 73.9 90.5 73.9 2015 7 73.4 70.3 73.4 70.3 2016 8 80.6 79.5 90.0 79.5 2017 10 75.9 75.9 75.9 2018 3 65.3 65.1 78.0 65.1 2019 3 75.9 75.3 79.1 74.6	2006	6	70.9	70.9		70.9
2009 5 72.2 72.2 72.2 2010 9 62.6 62.6 62.6 2011 7 74.1 74.1 74.1 2012 3 63.5 63.5 63.5 2013 6 74.8 74.8 74.8 2014 5 78.9 73.9 90.5 73.9 2015 7 73.4 70.3 73.4 70.3 2016 8 80.6 79.5 90.0 79.5 2017 10 75.9 75.9 75.9 2018 3 65.3 65.1 78.0 65.1 2019 3 75.9 75.3 79.1 74.6	2007	12	78.0	78.0		76.2
2010 9 62.6 62.6 62.6 2011 7 74.1 74.1 74.1 2012 3 63.5 63.5 63.5 2013 6 74.8 74.8 74.8 2014 5 78.9 73.9 90.5 73.9 2015 7 73.4 70.3 73.4 70.3 2016 8 80.6 79.5 90.0 79.5 2017 10 75.9 75.9 75.9 2018 3 65.3 65.1 78.0 65.1 2019 3 75.9 75.3 79.1 74.6	2008	2	72.4	72.4		72.4
2010 9 62.6 62.6 62.6 2011 7 74.1 74.1 74.1 2012 3 63.5 63.5 63.5 2013 6 74.8 74.8 74.8 2014 5 78.9 73.9 90.5 73.9 2015 7 73.4 70.3 73.4 70.3 2016 8 80.6 79.5 90.0 79.5 2017 10 75.9 75.9 75.9 2018 3 65.3 65.1 78.0 65.1 2019 3 75.9 75.3 79.1 74.6	2009	/5	72.2	72.2		72.2
2011 7 74.1 74.1 74.1 2012 3 63.5 63.5 63.5 2013 6 74.8 74.8 74.8 2014 5 78.9 73.9 90.5 73.9 2015 7 73.4 70.3 73.4 70.3 2016 8 80.6 79.5 90.0 79.5 2017 10 75.9 75.9 75.9 2018 3 65.3 65.1 78.0 65.1 2019 3 75.9 75.3 79.1 74.6	2010	9	62.6	62.6		
2012 3 63.5 63.5 2013 6 74.8 74.8 2014 5 78.9 73.9 90.5 2015 7 73.4 70.3 73.4 70.3 2016 8 80.6 79.5 90.0 79.5 2017 10 75.9 75.9 75.9 2018 3 65.3 65.1 78.0 65.1 2019 3 75.9 75.3 79.1 74.6	2011		74.1	74.1		74.1
2013 6 74.8 73.9 73.9 73.9 70.3 70.3 70.3 70.3 79.5 79.5 79.5 79.5 79.5 79.5 75.9 75.9 75.9 75.9 75.9 79.1 74.6 74.6 74.8 <t< td=""><td>2012</td><td></td><td>63.5</td><td>63.5</td><td></td><td></td></t<>	2012		63.5	63.5		
2014 5 78.9 73.9 90.5 73.9 2015 7 73.4 70.3 73.4 70.3 2016 8 80.6 79.5 90.0 79.5 2017 10 75.9 75.9 75.9 2018 3 65.3 65.1 78.0 65.1 2019 3 75.9 75.3 79.1 74.6	2013					
2015 7 73.4 70.3 73.4 70.3 2016 8 80.6 79.5 90.0 79.5 2017 10 75.9 75.9 75.9 2018 3 65.3 65.1 78.0 65.1 2019 3 75.9 75.3 79.1 74.6	2014		78.9		90.5	
2016 8 80.6 79.5 90.0 79.5 2017 10 75.9 75.9 75.9 2018 3 65.3 65.1 78.0 65.1 2019 3 75.9 75.3 79.1 74.6	2015	7	73.4	70.3		
2017 10 75.9 75.9 75.9 2018 3 65.3 65.1 78.0 65.1 2019 3 75.9 75.3 79.1 74.6	2016		80.6	79.5	90.0	
2018 3 65.3 65.1 78.0 65.1 2019 3 75.9 75.3 79.1 74.6						
2019 3 75.9 75.3 79.1 74.6	2018				78.0	
					1.7	
10.0						
	2020	-	, 5 • 5	73.3	, 1.0	, ,
1998-2020 133 74.0 72.9 79.9 72.4	1998-2020	133	74.0	72.9	79.9	72.4

By 2018, Bavarians' life expectancy at birth is estimated at 79.3 years for boys and 83.8 years for girls.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 11a $\begin{tabular}{ll} Mortality measures (cancer-related death) and mortality-incidence-index \\ by year of death \\ MALES \end{tabular}$

Year of	Deaths	Mort	MI-Index	Mort 1	MT-Tndev	Mort	MT-Index	Mort	MT-Index
death	n	raw	raw	WS	WS	ES ES	ES ES	BRD-S	BRD-S
deach	11	law	law	WS	WS	120	EO	מ-מאם	DKD-3
1998	3	0.3	0.75	0.2	0.79	0.2	0.73	0.2	0.59
1999	2						0.73		
		0.2	0.33	0.1	0.30	0.2		0.4	0.51
2000	8	0.7	1.00	0.4	1.03	0.6	1.01	0.8	0.91
2001	8	0.7		0.4	1.21	0.6	1.25	0.8	1.20
2002	8	0.4		0.2	0.34	0.3	0.42	0.5	0.52
2003	14	0.7	0.88	0.4	0.75	0.6	0.79	0.8	0.80
2004	11	0.6	0.85	0.4	0.90	0.5	0.89	0.6	0.89
2005	8	0.4	0.73	0.2	0.63	0.3	0.62	0.4	0.65
2006	10	0.5	0.83	0.3	0.96	0.4	0.93	0.5	0.85
2007	11	0.5	0.92	0.3	0.72	0.4	0.78	0.5	0.83
2008	14	0.6	1.75	0.4	2.00	0.5	1.94	0.6	1.96
2009	7	0.3	0.50	0.2	0.44	0.3	0.45	0.3	0.47
2010	6	0.3		0.2	0.55	0.2	0.63	0.2	0.62
2011	9	0.4	0.82	0.2	0.90	0.3	0.92	0.4	0.77
2012	6	0.3		0.1	0.32	0.2	0.38	0.3	0.48
2013	14	0.6	0.70	0.3	0.64	0.4	0.69	0.5	0.67
2013	5	0.2		0.1	0.31	0.1	0.37	0.2	0.40
2014	13	0.5	0.93	0.3	1.25	0.4	1.04	0.5	0.99
2015	10	0.3		0.3	1.30	0.3	1.38	0.3	1.35
	6	0.4						0.4	0.91
2017				0.1	0.64	0.2	0.82		
2018	5	0.2		0.1	0.64	0.2	0.61	0.2	0.56
2019	2	0.1		0.1	0.48	0.1	0.50	0.1	0.47
2020	5	0.2	1.67	0.1	1.75	0.2	1,77	0.2	1.62
1998-2020	185	0.4	0.79	0.2	0.74	0.3	0.77	0.4	0.78

Table 11b $\label{lem:mortality} \mbox{Mortality measures (cancer-related death) and mortality-incidence-index } \mbox{by year of death} \mbox{FEMALES}$

_	_		_ /				_		_
Year of	Deaths	Mort.	MI-Index						
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	2	0.2	0.50	0.1	0.50	0.1	0.50	0.1	0.50
1999	1	0.1	0.20	0.0	0.07	0.0	0.11	0.1	0.24
2000	4	0.3	0,67	0.2	0.98	0.3	0.88	0.3	0.81
2001	4	0.3	1.33	0.2	0.95	0.2	1.01	0.2	1.10
2002	9	0.5	0.82	0.2	0.75	0.3/	0.78	0.4	0.83
2003	8	0.4	1.14	0.2	1.14	0.2	1.09	0.3	1.11
2004	4	0.2	0.80	0.1	1.11	0.2	1.06	0.2	0.88
2005	6	0.3	0.86	0.1	0.85	0.2	0.85	0.2	0.81
2006	6	0.3	0.50	0.1	0.47	0.2	0.47	0.2	0.48
2007	12	0.5	1.33	0.2	1.00	0.3	1.17	0.4	1.18
2008	2	0.1	0.22	0.0	0.20	0.1	0.20	0.1	0.22
2009	5	0.2	0.83	0.1	0.64	0.1	0.66	0.2	0.77
2010	9	0.4	1.29	0.2	1.29	0.3	1.35	0.3	1.35
2011	7	0.3	1.17	0.1	0.81	0.2	0.89	0.2	1.10
2012	3	0.1	0.60	0.1	0.50	0.1	0.60	0.1	0.57
2013	6	0.3	1.00	0.1	0.87	0.1	0.91	0.2	0.99
2014	4	0.2	0.40	0.1	0.40	0.1	0.41	0.1	0.37
2015	6	0.2	1.00	0.1	0.81	0.2	0.83	0.2	0.97
2016	7	0.3	0.64	0.1	0.31	0.1	0.44	0.2	0.53
2017	10	0.4	1.43	0.1	1.37	0.2	1.35	0.3	1.46
2018	2	0.1	0.50	0.1	0.71	0.1	0.65	0.1	0.53
2019	2	0.1	1.00	0.0	0.35	0.0	0.50	0.1	0.78
2020	2	0.1	0.67	0.0	0.32	0.0	0.41	0.1	0.56
1998-2020	121	0.3	0.80	0.1	0.70	0.2	0.74	0.2	0.77

Table 12

Age distribution of age at death (cancer-related) for period 2007-2020 (incl. multiple malignancies)

Age at									
death	Cases			Males			Females		
Years	n	용	Cum.%	/n	용	Cum.%	n	용	Cum.%
0 - 4									
5-9									
10-14									
15-19	1	0.5	0.5	1	0.9	0.9			0.0
20-24	2	1.1	1.6	2	1.8	2.7/			0.0
25-29	0	0.0	1.6			2.7			0.0
30-34	1	0.5	2.1	1	0.9	3.5			0.0
35-39	1	0.5	2.6	1	0.9	4.4			0.0
40 - 44	3	1.6	4.2	2	1.8	6.2	1	1.3	1.3
45-49	8	4.2	8.4	5	4.4	10.6	3	3.9	5.2
50-54	11	5.8	14.2	8	7.1	17.7	3	3.9	9.1
55-59	15	7.9	22.1	10	8.8	26.5	5	6.5	15.6
60-64	19	10.0	32.1	9	8.0	34.5	10	13.0	28.6
65-69	25	13.2	45.3	17	15.0	49.6	8	10.4	39.0
70-74	29	15.3	60.5	16	14.2	63.7	13	16.9	55.8
75-79	32	16.8	77.4	16	14.2	77.9	16	20.8	76.6
80-84	21	11.1	88.4	10	8.8	86.7	11	14.3	90.9
85+	22	11.6	100.0	15	13.3	100.0	7	9.1	100.0
All ages	190	100.0		113	100.0		77	100.0	
_									

Table 13

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2020 (incl. multiple malignancies)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males	Females	spec.		spec.		cancers	cancers
Years	n	n		MI-index		MI-index	90	%
0- 4								
5- 9								
10-14								
15-19	1		0.1	1.00			2.1	
20-24	2		0.1	0.67			2.7	
25-29								
30-34	1		0.0	0.33			0.7	
35-39	1		0.0	0.25			0.4	
40-44	2	1	0.1		0.0	0.33	0.3	0.1
45-49	5	3	0.2	1.00	0.1	1.50	0.4	0.2
50-54	8	3	0.3		0.1	0.50	0.3	0.1
55-59	10 /	5	0.5	0.83	0.2	0.63	0.2	0.1
60-64	9	10	0.5	0.69	0.5	0.77	0.1	0.2
65-69	17	8	1.0	0.94	0.4	0.80	0.2	0.1
70-74	16	13	1.1	0.89	0.8	0.93	0.1	0.1
75-79	16	16	1.3	0.89	1.1	1.00	0.1	0.2
80-84	10	11	1.4	0.71	1.0	1.10	0.1	0.1
85+	15	7	3.2	1.00	0.7	1.75	0.2	0.1
031	10	/ //	3.2	1.00	0.7	1.75	0.2	0.1
All ages	113	77					0.2	0.1
HII ages	115	, ,					0.2	0.1
Mortality								
Raw			0.3	0.80	0.2	0.85		
WS			0.2		0.2	0.68		
ES			0.3	0.74	0.1	0.74		
BRD-S			0.3	0.78	0.1	0.74		
BKD-2			0.3	0.76	0.2	0.79		
PYLL-70								
			2.7		1.1			
per 100,000								
ES			2.4		0.9			
AYLL-70			13.6		10.2			

					Syn-	Syn-		
					chron	chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n	%↓	n	-%	n	← %	n	←%
C15 Oesophagus	/ 1	2.5	1	100.0				
C18 Colon	5	12.5	2	40.0	1	20.0	2	40.0
C19-C20 Rectum	/ 1 /	2.5			1	100.0		
C25 Pancreas	2	5.0	1	50.0			1	50.0
C33-C34 Lung	2	5.0			/ 1	50.0	1	50.0
C38,C45 Mesothelioma	2	5.0			1	50.0	1	50.0
C43 Malign. melanoma	2	5.0	2	100.0				
C44 Skin others	3	7.5	2	66.7			1	33.3
C60 Penis	1	2.5	1	100.0				
C61 Prostate	12	30.0	11	91.7	1	8.3		
C64 Kidney	2	5.0	2	100.0				
C67 Bladder	2	5.0	2	100.0				
C70-C72 CNS cancer	1	2.5	1	100.0				
C73 Thyroid	1	2.5	1	100.0				
C82-C85 NHL	3	7.5	2	66.7			1	33.3
All further malignancies	40	100.0	28	70.0	5	12.5	7	17.5

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a further malignancy.

	Total	Total	Pre	Pre	Syn- chron ±30d	Syn- chron ±30d	Post	Post
Diagnosis	n	용↓	n	← %	n	← %	n	← %
C03-C06 Oral cavity C12-C13 Hypopharynx	1 1	2.4	1 1	100.0				
C18 Colon	3 /	7.3	3	100.0				
C19-C20 Rectum	/ 1	2.4	1	100.0				
C33-C34 Lung	2	4.9			/ 1	50.0	1	50.0
C43 Malign. melanoma	3	7.3	2	66.7			1	33.3
C46,C49 Soft tissue	1	2.4	1	100.0				
C50 Breast	20	48.8	15	75.0	2	10.0	3	15.0
C54 Corpus uteri	1	2.4	1	100.0				
C56 Ovary	1	2.4	1	100.0				
C64 Kidney	1	2.4	1	100.0				
C65 Renal pelvis	1	2.4	1	100.0				
C70-C72 CNS cancer	1	2.4	1	100.0				
C81 Hodgkin lymphoma	1	2.4			1	100.0		
C82-C85 NHL	2	4.9	2	100.0				
C91-C96 Leukaemia	1	2.4	1	100.0				
7.11	4.1	100.0	2.0	70.0	4	0 0	F	10.0
All further malignancies	41	100.0	32	78.0	4	9.8	5	12.2

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a further malignancy.



Table 15

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2020 (First primaries only *)

			Males		Females		Males	Females
Age at			Age-		Age-		_	Prop.all
death		Females	spec.		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0- 4								
5- 9								
10-14								
15-19	1		0.1	1.00			2.2	
20-24	2		0.1	0.67			3.0	
25-29								
30-34	1		0.0	0.33			0.7	
35-39	1		0.0	0.33			0.4	
40-44	1		0.0	0.50			0.2	
45-49	5	3	0.2	1.25	0.1	1.50	0.4	0.2
50-54	7	2	0.3	0.78	0.1	0.40	0.3	0.1
55-59	8	3	0.4	0.89	0.1	0.60	0.2	0.1
60-64	9 /	8	0.5	0.82	0.4	0.67	0.2	0.2
65-69	16	7	1.0	0.94	0.4	0.78	0.2	0.1
70-74	14	7	0.9		0.4	1.00	0.2	0.1
75-79	11	13	0.9		0.9	1.08	0.1	0.2
80-84	8	6	1.1		0.6	0.86	0.1	0.1
85+	11	6	2.4		0.6	2.00	0.2	0.1
001		\	2 • 1	1.00	0.0	2.00	0.2	0.1
All ages	95	55					0.2	0.1
nii ages	33	93					/ 0.2	0.1
Mortality								
Raw			0.3	0.83	0.2	0.80		
WS			0.3		0.1	0.61		
ES			0.2	0.80	0.1	0.68		
			0.2					
BRD-S			0.3	0.81	0.1	0.73		
PYLL-70								
			2.4		0.8			
per 100,000								
ES			2.2		0.6			
AYLL-70			13.5		9.5			

^{*} See corresponding tables with multiple malignancies.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2020

(Single primaries only *)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males F	emales	spec.		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0- 4								
5- 9								
10-14								
15-19	1		0.1	1.00			2.2	
20-24	2		0.1	0.67			3.0	
25-29								
30-34	1		0.0	0.33			0.7	
35-39	1		0.0	0.33			0.4	
40-44	1		0.0	0.50			0.2	
45-49	5	3	0.2	1.25	0.1	1.50	0.4	0.2
50-54	7 /	2	0.3	0.78	0.1	0.40	0.3	0.1
55-59	8	3/	0.4	0.89	0.1	0.60	0.2	0.1
60-64	8 /	8	0.5	0.73	0.4	0.73	0.2	0.2
65-69	15	6	0.9	0.94	0.3	0.67	0.2	0.1
70-74	13	7	0.9	1.00	0.4	1.00	0.1	0.1
75-79	11	13	0.9		0.9	1.08	0.1	0.2
80-84	7 \	6	1.0	0.70	0.6	0.86	0.1	0.1
85+	11	5	2.4		0.5	1.67	0.2	0.1
All ages	91	53					0.2	0.1
- 3		1.						
Mortality								
Raw			0.3	0.82	0.2	0.78		
WS			0.1		0.1	0.60		
ES			0.2	0.79	0.1	0.67		
BRD-S			0.3	0.81	0.1	0.72		
DIE 5			0.3	0.01	0.1	0.72		
PYLL-70								
per 100,000			2.4		0.8			
ES ES			2.2		0.6			
AYLL-70			13.8		9.8			
77777 / 0			13.0		7.0			

^{*} See corresponding tables with multiple malignancies.

ICD-10 C38: Malignant neoplasm of heart, mediastinum and pleura Age distribution and age-specific mortality 2007 - 2020 (Males: 113, Females: 77)





Figure 17. Distribution of age at death (bars; males: mean=66.5 yrs, median=68.1 yrs; females: mean=70.1 yrs, median=73.1 yrs) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at heart, mediastinum and pleura-related death (see Table 10) should be considered.



werage mortality (Germany 1987 standard population) 2007 - 2020: Males



Average mortality (Germany 1987 standard population) 2007 - 2020: Females





Figure 18a. Map of cancer mortality (german standard population) by county averaged for period 2007 to 2020. According to their individual mortality rates, the counties are displayed in different red and blue hues, being the fine white color attributed to the population mean (males 0.3/100,000 WS N=113, females 0.2/100,000 WS N=77).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,727 female residents (averaged) in the period from 2007 to 2020 a total of 3 women died from heart, mediastinum and pleura. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.3/100,000 (german standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.0/100,000.

Standardized mortality ratio (SMR) 2007 - 2020: Males



Standardized mortality ratio (SMR) 2007 - 2020: Females




Figure 18b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2020. According to their individual SMR values, the counties are displayed in different red and blue hues, being the fine white color attributed to the population overall of 1.0 (males N=113, females N=77).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,153 female residents (averaged) in the period from 2007 to 2020 a total of 3 women died from heart, mediastinum and pleura. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 1.43. Though, the value of this parameter may vary with an underlying probability of 99% between 0.16 and 5.23, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases, where applicable. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head&neck and lung).

The ratio of mortality and incidence (mortality-to-incidence ratio, **MIR**, **MI-Index**) is a statistical index that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MIR. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

MCR Munich Cancer Registry (Tumorregister München)

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

SEER Surveillance, Epidemiology, and End Results (USA)

DCO Death certificate only

BRD-S German (FRG) standard population ES European standard population (old)

WS World standard population

SIR Standardized incidence ratio

CI Confidence interval EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

SMR Standardized mortality ratio

MI-index Ratio of mortality to incidence, MIR

FRG Federal Republic of Germany

Recommended Citation

Munich Cancer Registry. ICD-10 C38: Heart, mediastinum and pleura - Incidence and Mortality [Internet]. 2021 [updated 2021 Dec 21; cited 2022 Feb 1]. Available from: https://www.tumorregister-muenchen.de/en/facts/base/bC38_E-ICD-10-C38-Heart-mediastinum-and-pleura-incidence-and-mortality.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.