

Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ *Deutsch*

ICD-10 C50: Breast cancer (men)

Incidence and Mortality

Year of diagnosis	1998-2019
Patients	597
Diseases	604
Creation date	01/25/2021
Database export	01/07/2021
Population (males)	2.43 m



Munich Cancer Registry
Cancer Registry Bavaria - Upper Bavaria Regional Center
at Klinikum Grosshadern/IBE
Marchioninstr. 15
Munich, 81377
Germany

<https://www.tumorregister-muenchen.de/en>

https://www.tumorregister-muenchen.de/en/facts/base/bC50m_E-ICD-10-C50-Breast-cancer-men-incidence-and-mortality.pdf

Index of figures and tables

Fig./Tbl.		Page
1	Annual cases, DCO, mult. malignancies, follow-up / yr	4
2	Incidence by year of diagnosis	5
3	Age distribution parameters by year of diagnosis	6
4	Age distribution by 5-year age group	7
5	Age-specific incidence, DCO rate, proportion malignancies	8
6	Age distribution and age-specific incidence (chart)	9
6a	Age-specific incidence internationally (chart)	10
7	Standardized incidence ratio of further malignancies	11
8a	Map of cancer incidence (BRD-S) by county (chart)	12
8b	Standardized incidence ratio (SIR) by county (chart)	13
9a	Pts incident cohorts and mortality / yr	14
9b	Incidence and mortality by year of diagnosis	15
9c	Cancer-related deaths, death certification available / yr	16
10	Medians of age at death / yr	17
11	Mortality by year of death	18
12	Distribution of age at death	19
13	Age-specific mortality	20
14	Further malignancies in deaths	21
15	Age-specific mortality (first primaries)	22
16	Age-specific mortality (single primaries)	23
17	Age distribution and age-specific mortality (chart)	24
18a	Map of cancer mortality (BRD-S) by county (chart)	25
18b	Standardized mortality ratio (SMR) by county (chart)	26

**Global Statements about the statistics on the Internet –
Baseline Statistics** (grey button ) , **Survival** (red button )

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.69 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases^{###} are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, January 2021

- [#] Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.65 million to 4.10 in 2002, and to 4.69 million in 2007).
- ^{##} Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- ^{###} DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate.

ICD-10 codes (ICD-10 2015) used for specifying cancer site

Code	Description
C50.-	Malignant neoplasm of breast
C50.0	Nipple and areola
C50.1	Central portion of breast
C50.2	Upper-inner quadrant of breast
C50.3	Lower-inner quadrant of breast
C50.4	Upper-outer quadrant of breast
C50.5	Lower-outer quadrant of breast
C50.6	Axillary tail of breast
C50.8	Overlapping lesion of breast
C50.9	Breast, unspecified

Sex: Male

INCIDENCE

Table 1

Cases with invasive cancer by year of diagnosis, proportions of DCO, further malignancies, deaths, and active follow-up (incl. DCO)

Year of diagnosis	All cases n	DCO cases n	Prop. DCO %	Prop. at least 1 further malign. prior + synchron. %	Prop. at least 1 further malign. after %	Prop. deaths %	Prop. actively followed %
1998	8	2	25.0	12.5	15.7	75.0	87.5
1999	12	2	16.7	20.0	15.4	83.3	100.0
2000	14	4	28.6	20.6	15.0	78.6	92.9
2001	13			17.0	14.8	46.2	84.6
2002	20	3	15.0	17.9	14.6	90.0	95.0 #
2003	35	1	2.9	15.7	14.4	74.3	100.0
2004	27	3	11.1	17.1	13.2	66.7	100.0
2005	24			15.7	12.8	54.2	91.7
2006	25	2	8.0	14.6	12.4	76.0	100.0
2007	41	2	4.9	16.0	11.1	61.0	95.1 #
2008	26	5	19.2	16.7	10.4	69.2	100.0
2009	25	1	4.0	16.3	9.2	40.0	100.0
2010	29			15.7	9.3	51.7	100.0
2011	33	2	6.1	15.4	8.1	51.5	100.0
2012	35	2	5.7	16.3	8.3	37.1	100.0
2013	45	2	4.4	18.4	7.0	53.3	97.8
2014	34			18.8	4.3	29.4	94.1
2015	32	1	3.1	19.2	4.5	28.1	93.8
2016	38	2	5.3	19.8	3.3	23.7	97.4
2017	39	2	5.1	20.9	2.3	28.2	100.0
2018	25			21.0	0.0	16.0	100.0
2019	24			21.2	0.0	4.2	54.2 ##
1998-2019	604	36	6.0	21.2	15.7	48.5	95.7

604 cases diagnosed 1998-2019 are related to a total of 597 patients. Currently, in 206 (34.5 %) of these 597 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 153 / 39 / 14 (25.6 % / 6.5 % / 2.3 %) patients exist having 2 / 3 / 4+ malignancies.

The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retrieved from the respective headings.

How to interpret:

In 2017, a subgroup of 39 cases has been diagnosed, of which 20.9 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 2.3 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

Table 2

Incidence measures by year of diagnosis including DCO cases
(with respect to registry area expansion from 2.65 to 4.10 m as of 2002,
and from 4.10 to 4.92 m as of 2007, respectively)

Year of diagnosis	Cases n	Incidence raw	Incidence WS	Incidence ES	Incidence BRD-S
1998	8	0.7	0.5	0.6	0.9
1999	12	1.1	0.6	1.0	1.2
2000	14	1.2	0.7	1.1	1.5
2001	13	1.1	0.7	1.0	1.2
2002	20	1.1	0.6	0.9	1.2
2003	35	1.9	1.0	1.5	2.0
2004	27	1.4	0.8	1.2	1.5
2005	24	1.3	0.7	1.0	1.3
2006	25	1.3	0.7	1.0	1.3
2007	41	1.9	0.9	1.4	1.9
2008	26	1.2	0.6	0.9	1.2
2009	25	1.1	0.6	0.9	1.1
2010	29	1.3	0.6	0.9	1.2
2011	33	1.5	0.7	1.1	1.4
2012	35	1.5	0.8	1.1	1.5
2013	45	2.0	0.9	1.4	1.7
2014	34	1.5	0.7	1.0	1.3
2015	32	1.3	0.7	1.1	1.2
2016	38	1.6	0.8	1.1	1.4
2017	39	1.6	0.8	1.2	1.4
2018	25	1.0	0.5	0.7	0.9
2019	24	1.0	0.5	0.7	0.9
1998-2019	604	1.4	0.7	1.0	1.3

The computation of the incidence measures includes all cancers, irrespective of first or subsequent malignancy.

Table 3

Age distribution parameters by year of diagnosis
(incl. DCO)

Year of diagnosis	Cases n	Std.		Min.	Max.	Median				
		Mean	dev.			10%	25%	50%	75%	90%
1998	8	62.7	12.3	47.9	82.8	47.9	53.8	60.9	70.5	82.8
1999	12	68.8	10.3	52.8	85.2	55.5	62.0	66.1	78.6	79.6
2000	14	66.7	14.9	41.2	87.8	44.4	56.6	68.2	79.7	84.7
2001	13	63.4	9.9	48.9	84.7	50.4	58.4	62.6	67.0	77.8
2002	20	66.1	12.1	34.5	82.5	49.4	60.7	67.5	75.7	79.8
2003	35	66.5	13.5	30.3	89.6	46.2	59.6	68.3	76.0	82.5
2004	27	68.2	10.7	45.7	89.6	55.1	62.7	70.3	74.3	85.4
2005	24	70.1	10.1	44.3	90.9	55.2	66.7	71.7	75.8	81.1
2006	25	66.2	12.6	45.7	86.1	47.4	58.1	66.8	74.0	84.7
2007	41	69.3	11.6	41.3	96.1	58.2	60.7	71.7	76.4	80.6
2008	26	68.8	9.2	44.9	83.8	60.3	62.6	66.1	77.1	80.6
2009	25	69.0	11.7	46.0	89.5	54.4	61.4	70.4	77.3	85.9
2010	29	69.6	11.8	44.0	91.0	47.7	64.1	70.5	75.2	87.6
2011	33	71.0	11.4	48.6	90.6	55.3	61.2	72.1	80.8	84.0
2012	35	67.5	11.0	46.8	89.3	49.9	59.8	68.3	77.1	79.0
2013	45	70.4	12.9	45.1	90.4	50.5	64.7	72.2	78.0	88.6
2014	34	69.1	10.8	41.7	83.6	54.2	59.4	70.9	78.0	81.4
2015	32	66.8	10.0	50.2	92.0	55.3	59.9	66.8	73.1	77.8
2016	38	70.7	12.3	33.1	97.2	56.0	64.5	69.1	78.1	86.3
2017	39	69.6	13.5	41.8	96.5	53.7	60.2	67.5	79.4	94.2
2018	25	69.5	12.3	44.0	84.1	50.7	64.1	71.6	80.7	83.1
2019	24	70.7	12.5	36.6	90.4	61.1	64.2	68.5	79.9	85.5
1998-2019	604	68.7	11.8	30.3	97.2	52.8	61.1	69.3	77.1	83.2

Table 4

Age distribution by 5-year age group for period 2007-2019
(incl. DCO)

Age at diagnosis Years	Cases		Cum.%
	n	%	
0-4			
5-9			
10-14			
15-19			
20-24			
25-29			
30-34	1	0.2	0.2
35-39	1	0.2	0.5
40-44	6	1.4	1.9
45-49	19	4.5	6.3
50-54	23	5.4	11.7
55-59	31	7.3	19.0
60-64	65	15.3	34.3
65-69	68	16.0	50.2
70-74	63	14.8	65.0
75-79	71	16.7	81.7
80-84	45	10.6	92.3
85+	33	7.7	100.0
All ages	426	100.0	

Table 5

Age-specific incidence, DCO rate and proportion of all cancers for period 2007-2019

Age at diagnosis Years	Cases n	Age-spec. incidence	DCO rate n=19 %	Prop. all cancers n=143063 %
0- 4		0.0		
5- 9		0.0		
10-14		0.0		
15-19		0.0		
20-24		0.0		
25-29		0.0		
30-34	1	0.0		0.1
35-39	1	0.0		0.1
40-44	6	0.3	16.7	0.2
45-49	19	0.8		0.4
50-54	23	1.0		0.3
55-59	30	1.5		0.3
60-64	65	4.0	3.1	0.4
65-69	68	4.5	1.5	0.3
70-74	62	4.4		0.2
75-79	71	6.4	5.6	0.3
80-84	44	6.7	11.4	0.3
85+	32	7.5	18.8	0.3
All ages	422		4.5	0.3
Incidence				
Raw		1.4		
WS		0.7		
ES		1.0		
BRD-S		1.3		

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

ICD-10 C50: Malignant neoplasm of breast (men)

Age distribution and age-specific incidence 2007 - 2019 (n=422)




Figure 6. Age distribution (mean=69.4 yrs, median=69.9 yrs) and age-specific incidence.




Figure 6a. Age-specific incidence in MCR registry areas compared to Germany (FRG, GEKID extrapolation) and SEER (Surveillance, Epidemiology, and End Results, USA).

Reference:

Extrapolated age-specific patient population of Germany, data status middle of 2010. Association of Population-based Cancer Registries in Germany (GEKID e.V.). Berlin, 2014. <http://www.gekid.de>. Last access: 02/11/2015
 Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2019, based on the November 2018 submission. <http://www.seer.cancer.gov>.

Table 7

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of further malignancies for period 1998–2019

Diagnosis	Observed	Expected	SIR	CI		EAR	DCO %
	n	n		95%	95%		
C09–C10 Oropharynx	3	0.4	8.0	1.6	23.3 #	11.3	
C15 Oesophagus	3	0.8	3.9	0.8	11.3	9.6	
C16 Stomach	7	1.6	4.3	1.7	8.8 #	23.1	14.3
C18 Colon	7	4.0	1.7	0.7	3.6	12.9	
C19–C20 Rectum	3	2.1	1.4	0.3	4.2	3.8	
C22 Liver	2	1.2	1.7	0.2	6.1	3.5	
C23–C24 Bile	2	0.4	4.6	0.6	16.5	6.7	
C25 Pancreas	8	1.6	4.9	2.1	9.6 #	27.5	12.5
C30–C31 Sinuses	1	0.1	14.1	0.4	78.3	4.0	
C32 Larynx	1	0.4	2.6	0.1	14.5	2.7	
C33–C34 Lung	9	4.8	1.9	0.9	3.6	18.3	22.2
C43 Malign. melanoma	3	1.8	1.7	0.3	4.9	5.2	33.3
C50 Breast	7	0.1	60.7	24.4	125.0 #	29.7	
C61 Prostate	29	11.3	2.6	1.7	3.7 #	76.5	13.8
C64 Kidney	4	1.4	2.9	0.8	7.4	11.3	
C66 Ureter	1	0.1	8.9	0.2	49.6	3.8	
C67 Bladder	4	2.0	2.0	0.5	5.2	8.7	
C81 Hodgkin lymphoma	1	0.1	11.2	0.3	62.5	3.9	
C82–C85 NHL	3	1.8	1.7	0.4	5.0	5.4	
C91–C96 Leukaemia	1	0.6	1.5	0.0	8.6	1.5	100.0
Not observed	0	4.3	0.0	0.0	0.9 #	-18.6	
All further malignancies	99	40.9	2.4	2.0	2.9 #	250.9	10.1
Patients		564					
Median age at next malignancy (years)		72.5					
Person-years		2316					
Mean observation time (years)		4.1					
Median observation time (years)		2.8					

The occurrence of further specified malignancy is statistically significant.

Average incidence (Germany 1987 standard population) 2007 - 2019




Figure 8a. Map of cancer incidence (german standard population, incl. DCO cases) by county averaged for period 2007 to 2019. According to their individual incidence rates, the counties are displayed in different red and blue hues, being the fine white color attributed to the population mean (1.3/100,000 WS N=422).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 66,374 male residents (averaged) in the period from 2007 to 2019 a total of 10 men were identified with newly diagnosed breast cancer (men). Therefore, the mean incidence rate for this cancer type in this area can be calculated at 1.0/100,000 (german standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.4 and 2.3/100,000.

Standardized incidence ratio (SIR) 2007 - 2019




Figure 8b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2019. According to their individual SIR values, the counties are displayed in different red and blue hues, being the fine white color attributed to the population overall of 1.0 (N=422).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 65,991 male residents (averaged) in the period from 2007 to 2019 a total of 10 men were identified with newly diagnosed breast cancer (men). Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.83. Though, the value of this parameter may vary with an underlying probability of 99% between 0.31 and 1.77, and is therefore not statistically striking.

MORTALITY

Table 9a

Annual cohorts: Incident cancers, follow-up status, proportion of DCO, deaths among the annual cohorts and proportion of available death certificates (with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.92 m as of 2007, respectively)

Year of diagnosis	Incident cases n	Prop. actively followed %	Prop. DCO %	Deaths n	Prop. deaths %	Prop. deaths with death certific. %
1998	8	87.5	25.0	6	75.0	83.3
1999	12	100.0	16.7	10	83.3	100.0
2000	14	92.9	28.6	11	78.6	90.9
2001	13	84.6		6	46.2	83.3
2002	20	95.0	15.0	18	90.0	94.4
2003	35	100.0	2.9	26	74.3	92.3
2004	27	100.0	11.1	18	66.7	100.0
2005	24	91.7		13	54.2	100.0
2006	25	100.0	8.0	19	76.0	78.9
2007	41	95.1	4.9	25	61.0	92.0
2008	26	100.0	19.2	18	69.2	100.0
2009	25	100.0	4.0	10	40.0	70.0
2010	29	100.0		15	51.7	86.7
2011	33	100.0	6.1	17	51.5	100.0
2012	35	100.0	5.7	13	37.1	92.3
2013	45	97.8	4.4	24	53.3	95.8
2014	34	94.1		10	29.4	70.0
2015	32	93.8	3.1	9	28.1	88.9
2016	38	97.4	5.3	9	23.7	88.9
2017	39	100.0	5.1	11	28.2	45.5
2018	25	100.0		4	16.0	100.0
2019	24	54.2		1	4.2	100.0
1998-2019	604	95.7	6.0	293	48.5	89.8

Table 9b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased within the same year of being diagnosed with cancer (incl. DCO)

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.92 m as of 2007, respectively)

Year of diagnosis/ death	Incident cases n	Deaths n	Prop. deaths with death certific. %	Prop. deaths in same year	
				n	%
1998	8	4	100.0	1	12.5
1999	12	3	100.0	1	8.3
2000	14	7	100.0	5	35.7
2001	13	12	83.3		
2002	20	9	88.9	4	20.0
2003	35	10	100.0	5	14.3
2004	27	12	100.0	2	7.4
2005	24	10	100.0	1	4.2
2006	25	12	100.0	2	8.0
2007	41	14	100.0	5	12.2
2008	26	13	100.0	5	19.2
2009	25	16	100.0	2	8.0
2010	29	13	100.0		
2011	33	21	100.0	2	6.1
2012	35	21	95.2	2	5.7
2013	45	24	100.0	7	15.6
2014	34	16	100.0	1	2.9
2015	32	20	95.0	2	6.3
2016	38	24	100.0	3	7.9
2017	39	21	100.0	3	7.7
2018	25	23	30.4		
2019	24	23	60.9		
1998–2019	604	328	90.9	53	8.8

Table 9c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates
(incl. DCO)

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002,
and from 4.10 to 4.92 m as of 2007, respectively)

Year of death	Deaths n	Prop. cancer- related %	Prop. non-cancer- related %	Prop. cancer recorded on death certificate %
1998	4	75.0	25.0	75.0
1999	3	66.7	33.3	66.7
2000	7	85.7	14.3	85.7
2001	12	83.3	16.7	90.0
2002	9	66.7	33.3	75.0
2003	10	70.0	30.0	90.0
2004	12	66.7	33.3	100.0
2005	10	70.0	30.0	90.0
2006	12	83.3	16.7	91.7
2007	14	50.0	50.0	71.4
2008	13	38.5	61.5	61.5
2009	16	81.3	18.8	93.8
2010	13	61.5	38.5	61.5
2011	21	90.5	9.5	85.7
2012	21	76.2	23.8	70.0
2013	24	37.5	62.5	62.5
2014	16	81.3	18.8	87.5
2015	20	70.0	30.0	63.2
2016	24	66.7	33.3	70.8
2017	21	61.9	38.1	76.2
2018	23	43.5	56.5	57.1
2019	23	43.5	56.5	64.3
1998–2019	328	64.6	35.4	76.2

Table 10

Medians of age at death according to the grouping in Table 9

Year of death	Deaths n	Age at death (all causes) Years	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate) Years
1998	4	81.1	82.8	79.4	82.8
1999	3	68.0	67.3	79.7	67.3
2000	7	79.1	78.0	87.8	78.2
2001	12	71.9	69.1	84.5	70.2
2002	9	75.9	76.1	69.3	76.1
2003	10	71.9	70.2	88.4	71.4
2004	12	79.4	73.4	86.4	79.4
2005	10	80.8	79.5	84.5	82.1
2006	12	70.6	70.6	72.9	68.0
2007	14	78.4	69.1	87.6	76.4
2008	13	82.0	70.8	86.1	78.3
2009	16	70.7	72.1	66.9	69.3
2010	13	71.4	73.0	71.4	73.0
2011	21	74.1	74.1	84.6	74.6
2012	21	79.2	75.7	80.8	77.7
2013	24	80.3	76.9	80.9	82.8
2014	16	78.7	78.2	81.8	75.8
2015	20	78.2	76.2	78.9	76.2
2016	24	80.8	77.4	84.4	75.6
2017	21	76.3	75.1	79.1	77.6
2018	23	83.6	81.1	83.6	73.7
2019	23	81.1	74.3	81.8	79.4
1998–2019	328	78.4	75.3	81.8	76.1

By 2018, Bavarians' life expectancy at birth is estimated at 79.3 years for boys and 83.8 years for girls.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 11

Mortality measures (cancer-related death) and mortality-incidence-index by year of death

Year of death	Deaths n	Mort. raw	MI-Index raw	Mort. WS	MI-Index WS	Mort. ES	MI-Index ES	Mort. BRD-S	MI-Index BRD-S
1998	3	0.3	0.38	0.2	0.34	0.3	0.42	0.4	0.48
1999	2	0.2	0.17	0.1	0.20	0.2	0.18	0.2	0.15
2000	6	0.5	0.43	0.3	0.38	0.5	0.43	0.8	0.54
2001	10	0.9	0.77	0.5	0.74	0.8	0.79	1.0	0.89
2002	6	0.3	0.30	0.2	0.27	0.3	0.32	0.4	0.35
2003	7	0.4	0.20	0.2	0.21	0.3	0.21	0.4	0.22
2004	8	0.4	0.30	0.2	0.25	0.3	0.27	0.5	0.31
2005	7	0.4	0.29	0.2	0.27	0.3	0.31	0.5	0.35
2006	10	0.5	0.40	0.3	0.38	0.4	0.39	0.5	0.40
2007	7	0.3	0.17	0.2	0.17	0.2	0.18	0.3	0.16
2008	5	0.2	0.19	0.1	0.16	0.2	0.18	0.2	0.20
2009	13	0.6	0.52	0.3	0.50	0.5	0.52	0.6	0.55
2010	8	0.4	0.28	0.2	0.25	0.2	0.26	0.4	0.29
2011	19	0.8	0.59	0.4	0.59	0.6	0.60	0.8	0.58
2012	16	0.7	0.46	0.3	0.35	0.4	0.38	0.7	0.46
2013	9	0.4	0.21	0.1	0.16	0.2	0.19	0.4	0.22
2014	13	0.6	0.38	0.2	0.32	0.4	0.36	0.5	0.35
2015	14	0.6	0.44	0.2	0.34	0.4	0.38	0.5	0.44
2016	16	0.7	0.42	0.3	0.33	0.4	0.37	0.6	0.42
2017	13	0.5	0.33	0.2	0.28	0.3	0.29	0.5	0.33
2018	10	0.4	0.42	0.2	0.35	0.3	0.38	0.4	0.41
2019	10	0.4	0.42	0.2	0.33	0.3	0.37	0.4	0.40
1998-2019	212	0.5	0.35	0.2	0.31	0.3	0.33	0.5	0.36

Table 12

Age distribution of age at death (cancer-related) for period 2007-2019
(incl. multiple malignancies)

Age at death Years	Cases		Cum.%
	n	%	
0-4			
5-9			
10-14			
15-19			
20-24			
25-29			
30-34			
35-39			
40-44	2	1.3	1.3
45-49	3	2.0	3.3
50-54	4	2.6	5.9
55-59	7	4.6	10.5
60-64	10	6.5	17.0
65-69	23	15.0	32.0
70-74	22	14.4	46.4
75-79	28	18.3	64.7
80-84	27	17.6	82.4
85+	27	17.6	100.0
All ages	153	100.0	

Table 13

Age-specific mortality (cancer-related) and proportion of all cancers
for period 2007-2019
(incl. multiple malignancies)

Age at death Years	Cases n	Age-spec. mortality	MI-index	Prop. all cancers %
0- 4		0.0		
5- 9		0.0		
10-14		0.0		
15-19		0.0		
20-24		0.0		
25-29		0.0		
30-34		0.0		
35-39		0.0		
40-44	2	0.1	0.33	0.3
45-49	3	0.1	0.16	0.2
50-54	4	0.2	0.17	0.2
55-59	7	0.4	0.23	0.2
60-64	10	0.6	0.15	0.2
65-69	23	1.5	0.34	0.3
70-74	22	1.6	0.35	0.2
75-79	28	2.5	0.39	0.2
80-84	27	4.1	0.61	0.3
85+	27	6.3	0.84	0.3
All ages	153			0.2
Mortality				
Raw		0.5	0.36	
WS		0.2	0.31	
ES		0.3	0.33	
BRD-S		0.5	0.36	
PYLL-70				
per 100,000		1.6		
ES		1.3		
AYLL-70		8.4		

Table 14

Further malignancies in deaths in period 1998-2019

Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C09-C10 Oropharynx	4	3.2	1	25.0			3	75.0
C12-C13 Hypopharynx	1	0.8					1	100.0
C15 Oesophagus	4	3.2	3	75.0			1	25.0
C16 Stomach	8	6.5	3	37.5			5	62.5
C18 Colon	11	8.9	7	63.6			4	36.4
C19-C20 Rectum	4	3.2	4	100.0				
C22 Liver	3	2.4	2	66.7			1	33.3
C25 Pancreas	7	5.6					7	100.0
C30-C31 Sinuses	1	0.8	1	100.0				
C32 Larynx	2	1.6	2	100.0				
C33-C34 Lung	16	12.9	2	12.5	3	18.8	11	68.8
C43 Malign. melanoma	2	1.6	2	100.0				
C44 Skin others	10	8.1	4	40.0	3	30.0	3	30.0
C46,C49 Soft tissue	1	0.8					1	100.0
C50 Breast	8	6.5			3	37.5	5	62.5
C61 Prostate	26	21.0	14	53.8	3	11.5	9	34.6
C64 Kidney	2	1.6					2	100.0
C67 Bladder	3	2.4	1	33.3			2	66.7
C76-C79 CUP	1	0.8	1	100.0				
C81 Hodgkin lymphoma	2	1.6	1	50.0			1	50.0
C82-C85 NHL	6	4.8	5	83.3			1	16.7
C91-C96 Leukaemia	2	1.6					2	100.0
All further malignancies	124	100.0	53	42.7	12	9.7	59	47.6

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a further malignancy.

Table 15

Age-specific mortality (cancer-related) and proportion of all cancers
for period 2007-2019
(**First primaries only ***)

Age at death Years	Cases n	Age-spec. mortality	MI-index	Prop. all cancers %
0- 4		0.0		
5- 9		0.0		
10-14		0.0		
15-19		0.0		
20-24		0.0		
25-29		0.0		
30-34		0.0		
35-39		0.0		
40-44	1	0.0	0.20	0.2
45-49	2	0.1	0.13	0.2
50-54	3	0.1	0.13	0.1
55-59	6	0.3	0.27	0.2
60-64	8	0.5	0.16	0.2
65-69	20	1.3	0.40	0.3
70-74	12	0.9	0.28	0.1
75-79	20	1.8	0.38	0.2
80-84	19	2.9	0.66	0.3
85+	19	4.5	0.86	0.3
All ages	110			0.2
Mortality				
Raw		0.4	0.35	
WS		0.2	0.30	
ES		0.2	0.32	
BRD-S		0.3	0.35	
PYLL-70				
per 100,000		1.2		
ES		1.0		
AYLL-70		7.8		

* See corresponding tables with multiple malignancies.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers
for period 2007-2019
(**Single primaries only ***)

Age at death Years	Cases n	Age-spec. mortality	MI-index	Prop. all cancers %
0- 4		0.0		
5- 9		0.0		
10-14		0.0		
15-19		0.0		
20-24		0.0		
25-29		0.0		
30-34		0.0		
35-39		0.0		
40-44	1	0.0	0.20	0.2
45-49	2	0.1	0.13	0.2
50-54	3	0.1	0.14	0.1
55-59	6	0.3	0.30	0.2
60-64	6	0.4	0.13	0.1
65-69	16	1.1	0.36	0.2
70-74	10	0.7	0.26	0.1
75-79	14	1.3	0.29	0.2
80-84	7	1.1	0.28	0.1
85+	14	3.3	0.78	0.3
All ages	79			0.2
Mortality				
Raw		0.3	0.28	
WS		0.1	0.25	
ES		0.2	0.27	
BRD-S		0.2	0.27	
PYLL-70				
per 100,000		1.1		
ES		0.9		
AYLL-70		8.4		

* See corresponding tables with multiple malignancies.

ICD-10 C50: Malignant neoplasm of breast (men)

Age distribution and age-specific mortality 2007 - 2019 (n=153)




Figure 17. Distribution of age at death (bars; mean=68.9 yrs, median=68.5 yrs) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at breast cancer (men)-related death (see Table 10) should be considered.

Average mortality (Germany 1987 standard population) 2007 - 2019




Figure 18a. Map of cancer mortality (german standard population) by county averaged for period 2007 to 2019. According to their individual mortality rates, the counties are displayed in different red and blue hues, being the fine white color attributed to the population mean (0.5/100,000 WS N=153).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 66,374 male residents (averaged) in the period from 2007 to 2019 a total of 1 men died from breast cancer (men). Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.1/100,000 (german standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 0.9/100,000.

Standardized mortality ratio (SMR) 2007 - 2019




Figure 18b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2019. According to their individual SMR values, the counties are displayed in different red and blue hues, being the fine white color attributed to the population overall of 1.0 (N=153).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 65,991 male residents (averaged) in the period from 2007 to 2019 a total of 1 men died from breast cancer (men). Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.23. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 1.71, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases, where applicable. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head&neck and lung).

The ratio of mortality and incidence (mortality-to-incidence ratio, **MIR, MI-Index**) is a statistical index that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MIR. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

MCR	Munich Cancer Registry (Tumorregister München)
GEKID	Association of Population-based Cancer Registries in Germany (Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)
SEER	Surveillance, Epidemiology, and End Results (USA)
DCO	Death certificate only
BRD-S	German (FRG) standard population
ES	European standard population (old)
WS	World standard population
SIR	Standardized incidence ratio
CI	Confidence interval
EAR	Excess absolute risk = excess cancer cases (O - E) per 10,000 person-years
PYLL-70	Potential years of life lost prior to age 70 given a person dies before that age
AYLL-70	Average years of life lost prior to age 70 given a person dies before that age
SMR	Standardized mortality ratio
MI-index	Ratio of mortality to incidence, MIR
FRG	Federal Republic of Germany

Recommended Citation

Munich Cancer Registry. ICD-10 C50: Breast cancer (men) - Incidence and Mortality [Internet]. 2021 [updated 2021 Jan 25; cited 2021 Mar 1]. Available from: https://www.tumorregister-muenchen.de/en/facts/base/bC50m_E-ICD-10-C50-Breast-cancer-men-incidence-and-mortality.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.