

# Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ *Deutsch*

## ICD-10 C57: Female genitale cancer NOS

### Incidence and Mortality

Year of diagnosis	1998-2019
Patients	891
Diseases	891
Creation date	01/25/2021
Database export	01/07/2021
Population (females)	2.48 m



Munich Cancer Registry  
Cancer Registry Bavaria - Upper Bavaria Regional Center  
at Klinikum Grosshadern/IBE  
Marchioninstr. 15  
Munich, 81377  
Germany

<https://www.tumorregister-muenchen.de/en>

[https://www.tumorregister-muenchen.de/en/facts/base/bC57\\_\\_E-ICD-10-C57-Female-genitale-cancer-NOS-incidence-and-mortality.pdf](https://www.tumorregister-muenchen.de/en/facts/base/bC57__E-ICD-10-C57-Female-genitale-cancer-NOS-incidence-and-mortality.pdf)

### Index of figures and tables

Fig./Tbl.		Page
1	Annual cases, DCO, mult. malignancies, follow-up / yr	4
2	Incidence by year of diagnosis	5
3	Age distribution parameters by year of diagnosis	6
4	Age distribution by 5-year age group	7
5	Age-specific incidence, DCO rate, proportion malignancies	8
6	Age distribution and age-specific incidence (chart)	9
6a	Age-specific incidence internationally (chart)	10
7	Standardized incidence ratio of further malignancies	11
8a	Map of cancer incidence (BRD-S) by county (chart)	12
8b	Standardized incidence ratio (SIR) by county (chart)	13
9a	Pts incident cohorts and mortality / yr	14
9b	Incidence and mortality by year of diagnosis	15
9c	Cancer-related deaths, death certification available / yr	16
10	Medians of age at death / yr	17
11	Mortality by year of death	18
12	Distribution of age at death	19
13	Age-specific mortality	20
14	Further malignancies in deaths	21
15	Age-specific mortality (first primaries)	22
16	Age-specific mortality (single primaries)	23
17	Age distribution and age-specific mortality (chart)	24
18a	Map of cancer mortality (BRD-S) by county (chart)	25
18b	Standardized mortality ratio (SMR) by county (chart)	26

**Global Statements about the statistics on the Internet –  
Baseline Statistics** (grey button ) , **Survival** (red button )

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut<sup>#</sup>, with a total of 4.69 million inhabitants, account for the frequency of cancer diseases<sup>##</sup> and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases<sup>###</sup> are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to [tumor@ibe.med.uni-muenchen.de](mailto:tumor@ibe.med.uni-muenchen.de).

Munich Cancer Registry, January 2021

<sup>#</sup> Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.65 million to 4.10 in 2002, and to 4.69 million in 2007).

<sup>##</sup> Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.

<sup>###</sup> DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate.

### ICD-10 codes (ICD-10 2015) used for specifying cancer site

Code	Description
C57.-	Malignant neoplasm of other and unspecified female genital organs
C57.0	Fallopian tube
C57.1	Broad ligament
C57.2	Round ligament
C57.3	Parametrium
C57.4	Uterine adnexa, unspecified
C57.7	Other specified female genital organs
C57.8	Overlapping lesion of female genital organs
C57.9	Female genital organ, unspecified

## INCIDENCE

Table 1

Cases with invasive cancer by year of diagnosis, proportions of DCO, further malignancies, deaths, and active follow-up (incl. DCO)

Year of diagnosis	All cases n	DCO cases n	Prop. DCO %	Prop. at least 1 further malign. prior + synchron. %	Prop. at least 1 further malign. after %	Prop. deaths %	Prop. actively followed %
1998	21	5	23.8	9.5	6.5	66.7	95.2
1999	21	6	28.6	16.7	6.3	100.0	100.0
2000	27	7	25.9	17.4	6.1	92.6	100.0
2001	17	3	17.6	17.4	5.8	76.5	94.1
2002	55	29	52.7	18.4	5.8	85.5	98.2 #
2003	28	3	10.7	17.2	5.5	71.4	96.4
2004	35	8	22.9	17.6	4.9	85.7	100.0
2005	32	10	31.3	16.1	4.7	62.5	96.9
2006	38	8	21.1	16.8	4.7	71.1	94.7
2007	44	10	22.7	16.7	4.5	79.5	100.0 #
2008	34	4	11.8	18.2	4.5	79.4	94.1
2009	36	9	25.0	17.8	4.6	75.0	100.0
2010	53	7	13.2	17.9	4.1	60.4	100.0
2011	63	6	9.5	17.9	3.5	55.6	98.4
2012	53	6	11.3	18.0	2.1	62.3	96.2
2013	46	5	10.9	17.9	1.9	50.0	93.5
2014	58	8	13.8	18.2	1.4	70.7	100.0
2015	55	6	10.9	17.9	0.4	47.3	92.7
2016	45	9	20.0	17.7	0.6	62.2	100.0
2017	45	7	15.6	18.2	0.8	53.3	100.0
2018	46			18.1	1.2	19.6	100.0
2019	39			18.1	0.0	10.3	76.9 ##
1998–2019	891	156	17.5	18.1	6.5	63.0	96.9

891 cases diagnosed 1998-2019 are related to a total of 891 patients. Currently, in 223 (25.0 %) of these 891 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 162 / 52 / 9 (18.2 % / 5.8 % / 1.0 %) patients exist having 2 / 3 / 4+ malignancies.

# The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

## Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retrieved from the respective headings.

How to interpret:

In 2017, a subgroup of 45 cases has been diagnosed, of which 18.2 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 0.8 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

Table 2

Incidence measures by year of diagnosis including DCO cases  
(with respect to registry area expansion from 2.65 to 4.10 m as of 2002,  
and from 4.10 to 4.92 m as of 2007, respectively)

Year of diagnosis	Cases n	Incidence raw	Incidence WS	Incidence ES	Incidence BRD-S
1998	21	1.8	0.8	1.2	1.3
1999	21	1.8	0.7	1.0	1.4
2000	27	2.2	0.9	1.4	1.9
2001	17	1.4	0.6	0.9	1.2
2002	55	2.8	1.1	1.6	2.1
2003	28	1.4	0.6	0.9	1.2
2004	35	1.8	0.7	1.0	1.3
2005	32	1.6	0.8	1.1	1.4
2006	38	1.9	0.9	1.2	1.4
2007	44	1.9	0.9	1.2	1.6
2008	34	1.5	0.6	0.9	1.2
2009	36	1.5	0.6	0.9	1.2
2010	53	2.3	1.0	1.5	1.9
2011	63	2.7	1.2	1.8	2.2
2012	53	2.2	0.9	1.4	1.7
2013	46	1.9	0.9	1.3	1.6
2014	58	2.4	1.0	1.4	1.8
2015	55	2.3	1.1	1.6	1.8
2016	45	1.8	0.7	1.1	1.4
2017	45	1.8	0.7	1.0	1.3
2018	46	1.9	0.8	1.2	1.5
2019	39	1.6	0.7	1.1	1.3
1998-2019	891	1.9	0.8	1.2	1.5

The computation of the incidence measures includes all cancers, irrespective of first or subsequent malignancy.

Table 3

Age distribution parameters by year of diagnosis  
(incl. DCO)

Year of diagnosis	Cases n	Std.		Min.	Max.	10%	25%	Median		
		Mean	dev.					50%	75%	90%
1998	21	70.5	16.9	41.9	97.0	52.1	55.9	65.4	85.7	89.7
1999	21	75.5	11.9	57.7	91.1	60.0	63.3	79.1	85.4	88.6
2000	27	74.1	10.5	53.7	98.0	61.2	65.1	73.7	81.6	86.5
2001	17	73.1	10.6	54.5	91.2	58.2	63.4	72.7	81.5	86.5
2002	55	76.5	13.0	45.2	99.4	60.3	64.4	77.0	88.7	91.5
2003	28	71.1	13.3	42.2	98.6	56.8	62.3	69.1	79.9	88.3
2004	35	76.4	12.0	50.0	93.9	64.5	67.4	73.8	89.2	92.0
2005	32	69.9	19.0	1.7	103	46.9	61.9	74.7	79.9	90.0
2006	38	71.1	14.2	44.4	95.4	50.3	60.9	69.8	85.3	90.9
2007	44	71.5	14.4	18.3	100	54.8	63.5	71.9	80.6	86.8
2008	34	72.4	12.5	47.6	92.5	55.1	61.8	74.7	82.4	89.0
2009	36	73.0	12.0	46.5	96.2	58.5	65.6	73.0	81.8	86.9
2010	53	68.8	12.1	41.9	93.9	52.3	59.7	69.8	76.8	83.4
2011	63	70.4	10.6	44.3	95.0	57.5	63.0	69.7	78.8	84.3
2012	53	71.3	13.4	42.7	97.1	53.5	60.2	72.1	82.0	88.6
2013	46	69.0	12.0	47.1	105	53.3	62.4	65.8	78.5	85.3
2014	58	71.9	14.0	42.4	100	52.0	60.1	73.2	84.7	88.9
2015	55	66.4	12.7	37.3	92.9	51.3	56.7	65.9	76.6	82.8
2016	45	73.1	11.4	43.8	93.6	57.1	66.7	73.1	80.8	87.2
2017	45	73.6	12.1	45.5	95.0	58.9	65.5	75.4	83.7	87.7
2018	46	67.5	11.1	45.0	87.9	52.1	56.5	72.3	76.1	80.3
2019	39	67.3	11.5	47.0	92.2	54.3	58.0	65.9	75.7	84.5
1998-2019	891	71.3	13.0	1.7	105	54.5	62.1	71.7	81.0	88.0

Table 4

Age distribution by 5-year age group for period 2007-2019  
(incl. DCO)

Age at diagnosis Years	Cases		Cum.%
	n	%	
0-4			
5-9			
10-14			
15-19	1	0.2	0.2
20-24	0	0.0	0.2
25-29	0	0.0	0.2
30-34	0	0.0	0.2
35-39	2	0.3	0.5
40-44	8	1.3	1.8
45-49	17	2.8	4.5
50-54	49	7.9	12.5
55-59	59	9.6	22.0
60-64	69	11.2	33.2
65-69	88	14.3	47.5
70-74	85	13.8	61.3
75-79	95	15.4	76.7
80-84	64	10.4	87.0
85+	80	13.0	100.0
All ages	617	100.0	

Table 5

Age-specific incidence, DCO rate and proportion of all cancers for period 2007-2019

Age at diagnosis Years	Cases n	Age-spec. incidence	DCO rate n=77 %	Prop. all cancers n=144724 %
0- 4		0.0		
5- 9		0.0		
10-14		0.0		
15-19	1	0.1		0.4
20-24		0.0		
25-29		0.0		
30-34		0.0		
35-39	2	0.1		0.1
40-44	8	0.4		0.1
45-49	17	0.7	11.8	0.2
50-54	49	2.1		0.4
55-59	59	3.0	1.7	0.5
60-64	69	3.9		0.5
65-69	88	5.2	2.3	0.5
70-74	85	5.3	4.7	0.5
75-79	95	6.9	8.4	0.5
80-84	64	6.6	17.2	0.5
85+	80	8.3	61.3	0.5
All ages	617		12.5	0.4
Incidence				
Raw		2.0		
WS		0.9		
ES		1.3		
BRD-S		1.6		

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).



ICD-10 C57: Malignant neoplasm of other and unspecified female genital organs

Age distribution and age-specific incidence 2007 - 2019 (n=617)





Figure 6. Age distribution (mean=70.4 yrs, median=70.7 yrs) and age-specific incidence.



**Figure 6a.** Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).

Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER\*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2019, based on the November 2018 submission. <http://www.seer.cancer.gov>.


Table 7

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of further malignancies for period 1998–2019

Diagnosis	Observed n	Expected n	SIR	CI 95%	CI 95%	EAR	DCO %
C15 Oesophagus	1	0.2	5.5	0.1	30.6	3.5	
C16 Stomach	2	0.9	2.2	0.3	8.0	4.7	
C17 Small intestine	2	0.1	13.4	1.6	48.6 #	7.9	50.0
C18 Colon	7	2.6	2.7	1.1	5.6 #	18.8	14.3
C19–C20 Rectum	4	1.1	3.8	1.0	9.6 #	12.5	
C23–C24 Bile	1	0.4	2.6	0.1	14.6	2.6	100.0
C33–C34 Lung	8	2.1	3.8	1.6	7.4 #	25.1	
C43 Malign. melanoma	1	1.0	1.0	0.0	5.6	0.0	
C48 Peritoneal	1	0.1	8.7	0.2	48.3	3.8	
C50 Breast	22	8.3	2.6	1.7	4.0 #	58.5	9.1
C51 Vulva	2	0.3	7.0	0.8	25.3	7.3	
C53 Cervix uteri	2	0.3	6.2	0.8	22.5	7.2	50.0
C54 Corpus uteri	16	1.6	10.3	5.9	16.7 #	61.7	6.3
C56 Ovary	7	1.1	6.4	2.6	13.1 #	25.2	
C65 Renal pelvis	1	0.1	11.2	0.3	62.5	3.9	100.0
C67 Bladder	2	0.5	3.8	0.5	13.8	6.3	
C68 Urinary org.	1	0.0	105.7	2.7	588.7 #	4.2	100.0
C70–C72 CNS cancer	1	0.3	2.9	0.1	16.0	2.8	
C73 Thyroid	1	0.4	2.4	0.1	13.6	2.5	
C76–C79 CUP	3	0.5	6.2	1.3	18.0 #	10.7	
C82–C85 NHL	2	1.1	1.9	0.2	6.8	4.0	
C91–C96 Leukaemia	1	0.4	2.5	0.1	14.1	2.6	
Not observed	0	3.8	0.0	0.0	1.0 #	-16.4	
All further malignancies	88	27.2	3.2	2.6	4.0 #	259.5	10.2
Patients		745					
Median age at next malignancy (years)		72.7					
Person-years		2341					
Mean observation time (years)		3.1					
Median observation time (years)		1.7					

# The occurrence of further specified malignancy is statistically significant.


Average incidence (Germany 1987 standard population) 2007 - 2019



**Figure 8a.** Map of cancer incidence (german standard population, incl. DCO cases) by county averaged for period 2007 to 2019. According to their individual incidence rates, the counties are displayed in different red and blue hues, being the fine white color attributed to the population mean (1.6/100,000 WS N=617).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,462 female residents (averaged) in the period from 2007 to 2019 a total of 14 women were identified with newly diagnosed female genitale cancer NOS. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 1.3/100,000 (german standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.6 and 2.5/100,000.

Standardized incidence ratio (SIR) 2007 - 2019



**Figure 8b.** Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2019. According to their individual SIR values, the counties are displayed in different red and blue hues, being the fine white color attributed to the population overall of 1.0 (N=617).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,153 female residents (averaged) in the period from 2007 to 2019 a total of 14 women were identified with newly diagnosed female genitale cancer NOS. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.83. Though, the value of this parameter may vary with an underlying probability of 99% between 0.37 and 1.59, and is therefore not statistically striking.

## MORTALITY

Table 9a

Annual cohorts: Incident cancers, follow-up status, proportion of DCO, deaths among the annual cohorts and proportion of available death certificates (with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.92 m as of 2007, respectively)

Year of diagnosis	Incident cases n	Prop. actively followed %	Prop. DCO %	Deaths n	Prop. deaths %	Prop. deaths with death certific. %
1998	21	95.2	23.8	14	66.7	100.0
1999	21	100.0	28.6	21	100.0	100.0
2000	27	100.0	25.9	25	92.6	100.0
2001	17	94.1	17.6	13	76.5	100.0
2002	55	98.2	52.7	47	85.5	95.7
2003	28	96.4	10.7	20	71.4	100.0
2004	35	100.0	22.9	30	85.7	100.0
2005	32	96.9	31.3	20	62.5	90.0
2006	38	94.7	21.1	27	71.1	92.6
2007	44	100.0	22.7	35	79.5	100.0
2008	34	94.1	11.8	27	79.4	96.3
2009	36	100.0	25.0	27	75.0	92.6
2010	53	100.0	13.2	32	60.4	96.9
2011	63	98.4	9.5	35	55.6	100.0
2012	53	96.2	11.3	33	62.3	97.0
2013	46	93.5	10.9	23	50.0	100.0
2014	58	100.0	13.8	41	70.7	90.2
2015	55	92.7	10.9	26	47.3	100.0
2016	45	100.0	20.0	28	62.2	85.7
2017	45	100.0	15.6	24	53.3	75.0
2018	46	100.0		9	19.6	66.7
2019	39	76.9		4	10.3	75.0
1998-2019	891	96.9	17.5	561	63.0	94.8

Table 9b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased within the same year of being diagnosed with cancer (incl. DCO)

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.92 m as of 2007, respectively)

Year of diagnosis/ death	Incident cases n	Deaths n	Prop. deaths with death certific. %	Deaths in same year n	Prop. deaths in same year %
1998	21	16	100.0	7	33.3
1999	21	26	88.5	10	47.6
2000	27	15	93.3	8	29.6
2001	17	21	100.0	4	23.5
2002	55	41	95.1	36	65.5
2003	28	19	94.7	4	14.3
2004	35	28	100.0	12	34.3
2005	32	21	95.2	12	37.5
2006	38	29	96.6	12	31.6
2007	44	28	96.4	16	36.4
2008	34	23	100.0	8	23.5
2009	36	19	100.0	6	16.7
2010	53	25	100.0	9	17.0
2011	63	29	100.0	8	12.7
2012	53	31	96.8	11	20.8
2013	46	39	94.9	10	21.7
2014	58	40	100.0	14	24.1
2015	55	42	97.6	10	18.2
2016	45	51	98.0	15	33.3
2017	45	38	97.4	11	24.4
2018	46	32	40.6	1	2.2
2019	39	16	56.3	2	5.1
1998–2019	891	629	93.3	226	25.4

Table 9c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates  
(incl. DCO)

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002,  
and from 4.10 to 4.92 m as of 2007, respectively)

Year of death	Deaths n	Prop. cancer- related %	Prop. non-cancer- related %	Prop. cancer recorded on death certificate %
1998	16	68.8	31.3	100.0
1999	26	73.1	26.9	95.7
2000	15	66.7	33.3	85.7
2001	21	81.0	19.0	90.5
2002	41	73.2	26.8	94.9
2003	19	73.7	26.3	88.9
2004	28	75.0	25.0	92.9
2005	21	76.2	23.8	80.0
2006	29	75.9	24.1	85.7
2007	28	82.1	17.9	96.3
2008	23	87.0	13.0	91.3
2009	19	89.5	10.5	89.5
2010	25	80.0	20.0	96.0
2011	29	69.0	31.0	79.3
2012	31	90.3	9.7	93.3
2013	39	79.5	20.5	86.5
2014	40	80.0	20.0	87.5
2015	42	83.3	16.7	87.8
2016	51	82.4	17.6	88.0
2017	38	81.6	18.4	89.2
2018	32	75.0	25.0	100.0
2019	16	56.3	43.8	77.8
1998–2019	629	78.2	21.8	89.8



Table 10

Medians of age at death according to the grouping in Table 9

Year of death	Deaths n	Age at death (all causes) Years	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate) Years
1998	16	86.2	84.8	87.0	86.2
1999	26	83.6	80.9	86.4	84.4
2000	15	83.7	81.8	86.1	81.8
2001	21	78.0	76.0	84.7	76.0
2002	41	82.0	81.1	87.1	81.5
2003	19	75.1	70.4	84.9	73.6
2004	28	79.9	78.7	81.1	79.9
2005	21	80.3	78.8	86.2	78.8
2006	29	77.1	73.4	84.1	75.4
2007	28	79.2	78.9	80.6	79.2
2008	23	75.8	73.6	77.5	71.4
2009	19	76.3	75.9	88.6	75.9
2010	25	75.8	73.9	89.1	75.0
2011	29	79.2	76.9	81.3	79.2
2012	31	75.0	74.9	89.9	74.9
2013	39	79.4	77.6	86.7	78.4
2014	40	82.9	78.2	87.4	83.6
2015	42	77.5	76.1	89.3	76.2
2016	51	76.1	74.6	89.5	74.6
2017	38	77.0	73.6	82.6	73.8
2018	32	79.7	80.2	75.4	80.5
2019	16	77.8	78.9	76.5	78.4
1998-2019	629	78.9	77.0	85.4	77.8

By 2018, Bavarians' life expectancy at birth is estimated at 79.3 years for boys and 83.8 years for girls.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 11

Mortality measures (cancer-related death) and mortality-incidence-index by year of death

Year of death	Deaths n	Mort. raw	MI-Index raw	Mort. WS	MI-Index WS	Mort. ES	MI-Index ES	Mort. BRD-S	MI-Index BRD-S
1998	11	0.9	0.52	0.3	0.33	0.4	0.38	0.7	0.49
1999	19	1.6	0.90	0.6	0.90	1.0	0.93	1.3	0.98
2000	10	0.8	0.37	0.3	0.33	0.5	0.33	0.6	0.32
2001	17	1.4	1.00	0.5	0.91	0.8	0.94	1.1	0.97
2002	30	1.5	0.55	0.5	0.46	0.8	0.49	1.1	0.55
2003	14	0.7	0.50	0.3	0.48	0.4	0.47	0.5	0.46
2004	21	1.1	0.60	0.4	0.56	0.6	0.59	0.8	0.63
2005	16	0.8	0.50	0.3	0.33	0.4	0.40	0.6	0.45
2006	22	1.1	0.58	0.5	0.62	0.7	0.57	0.9	0.61
2007	23	1.0	0.52	0.3	0.36	0.5	0.41	0.7	0.45
2008	20	0.9	0.59	0.3	0.53	0.5	0.54	0.6	0.55
2009	17	0.7	0.47	0.3	0.45	0.4	0.45	0.6	0.49
2010	20	0.9	0.38	0.3	0.31	0.5	0.32	0.6	0.33
2011	20	0.9	0.32	0.3	0.24	0.4	0.25	0.6	0.28
2012	28	1.2	0.53	0.4	0.48	0.7	0.49	0.9	0.53
2013	31	1.3	0.67	0.4	0.43	0.6	0.49	0.9	0.59
2014	32	1.3	0.55	0.4	0.42	0.7	0.46	0.9	0.52
2015	35	1.4	0.64	0.5	0.41	0.7	0.47	1.0	0.55
2016	42	1.7	0.93	0.7	0.92	1.0	0.92	1.3	0.94
2017	31	1.3	0.69	0.5	0.71	0.7	0.70	0.9	0.68
2018	24	1.0	0.52	0.3	0.33	0.4	0.36	0.7	0.45
2019	9	0.4	0.23	0.1	0.09	0.1	0.12	0.2	0.18
1998-2019	492	1.1	0.55	0.4	0.45	0.6	0.48	0.8	0.52

Table 12

Age distribution of age at death (cancer-related) for period 2007-2019  
(incl. multiple malignancies)

Age at death Years	Cases		Cum.%
	n	%	
0-4			
5-9			
10-14			
15-19			
20-24			
25-29			
30-34			
35-39			
40-44			
45-49	6	1.8	1.8
50-54	8	2.4	4.2
55-59	15	4.5	8.7
60-64	19	5.7	14.5
65-69	47	14.2	28.6
70-74	56	16.9	45.5
75-79	57	17.2	62.7
80-84	60	18.1	80.7
85+	64	19.3	100.0
All ages	332	100.0	

Table 13

Age-specific mortality (cancer-related) and proportion of all cancers  
for period 2007-2019  
(incl. multiple malignancies)

Age at death Years	Cases n	Age-spec. mortality	MI-index	Prop. all cancers %
0- 4		0.0		
5- 9		0.0		
10-14		0.0		
15-19		0.0		
20-24		0.0		
25-29		0.0		
30-34		0.0		
35-39		0.0		
40-44		0.0		
45-49	6	0.2	0.35	0.4
50-54	8	0.3	0.16	0.3
55-59	15	0.8	0.25	0.4
60-64	19	1.1	0.28	0.4
65-69	47	2.8	0.53	0.7
70-74	56	3.5	0.66	0.7
75-79	57	4.1	0.60	0.6
80-84	60	6.2	0.94	0.7
85+	64	6.6	0.80	0.6
All ages	332			0.6
Mortality				
Raw		1.1	0.54	
WS		0.4	0.42	
ES		0.6	0.45	
BRD-S		0.8	0.50	
PYLL-70				
per 100,000		2.8		
ES		2.2		
AYLL-70		7.6		

Table 14

Further malignancies in deaths in period 1998-2019

Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C09-C10 Oropharynx	1	0.6	1	100.0				
C15 Oesophagus	1	0.6					1	100.0
C16 Stomach	2	1.2			1	50.0	1	50.0
C17 Small intestine	2	1.2	1	50.0			1	50.0
C18 Colon	17	10.0	6	35.3	3	17.6	8	47.1
C19-C20 Rectum	9	5.3	4	44.4	1	11.1	4	44.4
C23-C24 Bile	1	0.6					1	100.0
C25 Pancreas	2	1.2					2	100.0
C33-C34 Lung	13	7.6	2	15.4	1	7.7	10	76.9
C43 Malign. melanoma	3	1.8	2	66.7			1	33.3
C44 Skin others	9	5.3	4	44.4			5	55.6
C46,C49 Soft tissue	2	1.2					2	100.0
C48 Peritoneal	1	0.6			1	100.0		
C50 Breast	42	24.7	29	69.0	4	9.5	9	21.4
C51 Vulva	1	0.6	1	100.0				
C52 Vagina	3	1.8	1	33.3	1	33.3	1	33.3
C53 Cervix uteri	5	2.9	2	40.0	1	20.0	2	40.0
C54 Corpus uteri	19	11.2	3	15.8	14	73.7	2	10.5
C56 Ovary	7	4.1	4	57.1	3	42.9		
C64 Kidney	3	1.8	1	33.3			2	66.7
C65 Renal pelvis	4	2.4	2	50.0			2	50.0
C67 Bladder	3	1.8	1	33.3	1	33.3	1	33.3
C68 Urinary org.	1	0.6					1	100.0
C70-C72 CNS cancer	1	0.6					1	100.0
C73 Thyroid	2	1.2	2	100.0				
C76-C79 CUP	7	4.1	1	14.3	2	28.6	4	57.1
C81 Hodgkin lymphoma	2	1.2	2	100.0				
C82-C85 NHL	6	3.5	1	16.7			5	83.3
C91-C96 Leukaemia	1	0.6			1	100.0		
All further malignancies	170	100.0	70	41.2	34	20.0	66	38.8

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a further malignancy.

Table 15

Age-specific mortality (cancer-related) and proportion of all cancers  
for period 2007-2019  
(**First primaries only \***)

Age at death Years	Cases n	Age-spec. mortality	MI-index	Prop. all cancers %
0- 4		0.0		
5- 9		0.0		
10-14		0.0		
15-19		0.0		
20-24		0.0		
25-29		0.0		
30-34		0.0		
35-39		0.0		
40-44		0.0		
45-49	5	0.2	0.29	0.4
50-54	6	0.3	0.16	0.3
55-59	12	0.6	0.24	0.4
60-64	17	1.0	0.32	0.5
65-69	37	2.2	0.55	0.7
70-74	42	2.6	0.65	0.7
75-79	49	3.6	0.66	0.7
80-84	50	5.1	0.98	0.8
85+	48	5.0	0.75	0.6
All ages	266			0.6
Mortality				
Raw		0.9	0.55	
WS		0.3	0.43	
ES		0.5	0.46	
BRD-S		0.6	0.51	
PYLL-70				
per 100,000		2.2		
ES		1.8		
AYLL-70		7.6		

\* See corresponding tables with multiple malignancies.


Table 16

Age-specific mortality (cancer-related) and proportion of all cancers  
for period 2007-2019  
(**Single primaries only \***)

Age at death Years	Cases n	Age-spec. mortality	MI-index	Prop. all cancers %
0- 4		0.0		
5- 9		0.0		
10-14		0.0		
15-19		0.0		
20-24		0.0		
25-29		0.0		
30-34		0.0		
35-39		0.0		
40-44		0.0		
45-49	5	0.2	0.29	0.4
50-54	6	0.3	0.17	0.3
55-59	12	0.6	0.24	0.4
60-64	16	0.9	0.33	0.4
65-69	35	2.1	0.58	0.7
70-74	40	2.5	0.63	0.7
75-79	39	2.8	0.57	0.6
80-84	43	4.4	0.84	0.7
85+	41	4.2	0.66	0.5
All ages	237			0.5
Mortality				
Raw		0.8	0.51	
WS		0.3	0.41	
ES		0.4	0.44	
BRD-S		0.6	0.48	
PYLL-70				
per 100,000		2.2		
ES		1.8		
AYLL-70		7.8		

\* See corresponding tables with multiple malignancies.

ICD-10 C57: Malignant neoplasm of other and unspecified female genital organs  
Age distribution and age-specific mortality 2007 - 2019 (n=332)




**Figure 17.** Distribution of age at death (bars; n=mean=71.2 yrs, median=72.0 yrs) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at female genitale cancer NOS-related death (see Table 10) should be considered.




## Average mortality (Germany 1987 standard population) 2007 - 2019



**Figure 18a.** Map of cancer mortality (german standard population) by county averaged for period 2007 to 2019. According to their individual mortality rates, the counties are displayed in different red and blue hues, being the fine white color attributed to the population mean (0.8/100,000 WS N=332).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,462 female residents (averaged) in the period from 2007 to 2019 a total of 5 women died from female genitale cancer NOS. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.4/100,000 (german standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.1 and 1.3/100,000.

Standardized mortality ratio (SMR) 2007 - 2019



**Figure 18b.** Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2019. According to their individual SMR values, the counties are displayed in different red and blue hues, being the fine white color attributed to the population overall of 1.0 (N=332).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,153 female residents (averaged) in the period from 2007 to 2019 a total of 5 women died from female genitale cancer NOS. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.56. Though, the value of this parameter may vary with an underlying probability of 99% between 0.12 and 1.59, and is therefore not statistically striking.

### Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases, where applicable. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

#### 1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

#### 2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

#### 3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head&neck and lung).

The ratio of mortality and incidence (mortality-to-incidence ratio, **MIR, MI-Index**) is a statistical index that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MIR. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

**Shortcuts**

MCR	Munich Cancer Registry (Tumorregister München)
GEKID	Association of Population-based Cancer Registries in Germany (Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)
SEER	Surveillance, Epidemiology, and End Results (USA)
DCO	Death certificate only
BRD-S	German (FRG) standard population
ES	European standard population (old)
WS	World standard population
SIR	Standardized incidence ratio
CI	Confidence interval
EAR	Excess absolute risk = excess cancer cases (O - E) per 10,000 person-years
PYLL-70	Potential years of life lost prior to age 70 given a person dies before that age
AYLL-70	Average years of life lost prior to age 70 given a person dies before that age
SMR	Standardized mortality ratio
MI-index	Ratio of mortality to incidence, MIR
FRG	Federal Republic of Germany

**Recommended Citation**

Munich Cancer Registry. ICD-10 C57: Female genitale cancer NOS - Incidence and Mortality [Internet]. 2021 [updated 2021 Jan 25; cited 2021 Mar 1]. Available from: [https://www.tumorregister-muenchen.de/en/facts/base/bC57\\_\\_E-ICD-10-C57-Female-genitale-cancer-NOS-incidence-and-mortality.pdf](https://www.tumorregister-muenchen.de/en/facts/base/bC57__E-ICD-10-C57-Female-genitale-cancer-NOS-incidence-and-mortality.pdf)

**Copyright**

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

**Disclaimer**

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.