Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ Deutsch

ICD-10 C66: Ureteral cancer

Incidence and Mortality

Year of diagnosis	1998-2020
Patients	821
Diseases	832
Creation date	12/21/2021
Database export	12/20/2021
Population	4.95 m



Munich Cancer Registry
Cancer Registry Bavaria - Upper Bavaria Regional Center
at Klinikum Grosshadern/IBE
Marchioninistr. 15
Munich, 81377
Germany

https://www.tumorregister-muenchen.de/en

https://www.tumorregister-muenchen.de/en/facts/base/bC66___E-ICD-10-C66-Ureteral-cancer-incidence-and-mortality.pdf

Index of figures and tables

Fig./Tbl	l.	Page
1	Annual cases, DCO, mult. malignancies, follow-up / yr	4
2	Incidence by year of diagnosis	7
3	Age distribution parameters by year of diagnosis	8
4	Age distribution by 5-year age group and sex	11
5	Age-specific incidence, DCO rate, proportion malignancies	12
6	Age distribution and age-specific incidence (chart)	13
6a	Age-specific incidence internationally (chart)	14
7	Standardized incidence ratio of further malignancies	15
8a	Map of cancer incidence (BRD-S) by county (chart)	17
8b	Standardized incidence ratio (SIR) by county (chart)	18
9a	Pts incident cohorts and mortality / yr	19
9b	Incidence and mortality by year of diagnosis	20
9c	Cancer-related deaths, death certification available / yr	21
10	Medians of age at death / yr	22
11	Mortality by year of death	24
12	Distribution of age at death	26
13	Age-specific mortality	27
14	Further malignancies in deaths	28
15	Age-specific mortality (first primaries)	30
16	Age-specific mortality (single primaries)	31
17	Age distribution and age-specific mortality (chart)	32
18a	Map of cancer mortality (BRD-S) by county (chart)	33
18b	Standardized mortality ratio (SMR) by county (chart)	34

Global Statements about the statistics on the Internet – Baseline Statistics (grey button ——), Survival (red button ——)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.69 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases### are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, December 2021

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.65 million to 4.10 in 2002, and to 4.69 million in 2007).
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- ### DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate.

ICD-10 codes (ICD-10 2015) used for specifying cancer site

Code	Description
C66	Malignant neoplasm of ureter
	Wanghant hoopidom of drotor

INCIDENCE

Table 1

Cases with invasive cancer by year of diagnosis, proportions of DCO, further malignancies, deaths, and active follow-up (ALL PATIENTS) (incl. DCO)

				Prop.			
				at least	Prop.		
				1 further	at least		
				malign.	1 further		Prop.
	All	DCO	Prop.	prior +	malign.	Prop.	actively
Year of	cases	cases	DCO	synchron.	after	deaths	followed
diagnosis	n	n	용	%	%	용	%
1998	20			45.0	38.7	90.0	100.0
1999	10			46.7	37.7	90.0	100.0
2000	12			45.2	37.6	83.3	100.0
2001	24			42.4	37.2	95.8	100.0
2002	28			40.4	37.4	92.9	100.0 #
2003	19			38.9	37.6	78.9	100.0
2004	24			38.7	37.7	83.3	91.7
2005	42			40.8	37.0	76.2	95.2
2006	35			40.7	37.0	91.4	100.0
2007	36			41.2	36.3	86.1	100.0 #
2008	40			41.0	35.7	80.0	97.5
2009	45			41.5	34.7	80.0	97.8
2010	49	1	2.0	43.0	34.5	81.6	98.0
2011	47			45.9	32.7	70.2	100.0
2012	45	1	2.2	46.6	32.2	80.0	100.0
2013	55			47.3	31.3	74.5	100.0
2014	50			48.7	31.3	74.0	98.0
2015	33	1	3.0	49.2	30.6	60.6	100.0
2016	57			50.2	26.9	75.4	98.2
2017	45	1	2.2	51.3	25.5	53.3	100.0
2018	56			51.8	22.3	48.2	100.0
2019	36			52.1	22.2	30.6	97.2
2020	24			52.4	26.1	37.5	100.0 ##
1998-2020	832	4	0.5	52.4	38.7	72.7	98.8

832 cases diagnosed 1998-2020 are related to a total of 821 patients. Currently, in 622 (75.8 %) of these 821 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 285 / 205 / 132 (34.7 % / 25.0 % / 16.1 %) patients exist having 2 / 3 / 4+ malignancies.

How to interpret:

In 2018, a subgroup of 56 cases has been diagnosed, of which 51.8 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 22.3 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

[#] The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

^{##} Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retreived from the respective headings.

Table 1a

Cases with invasive cancer by year of diagnosis, proportions of DCO, further malignancies, deaths, and active follow-up (MALES) (incl. DCO)

					Prop.			
					at least	Prop.		
					1 further	at least		
					malign.	1 further		Prop.
			DCO	Prop.	prior +	malign.	Prop.	actively
Year of	Males	Males	cases	DCO	synchron.	after	deaths	followed
diagnosis	n	8	n	%	%	%	8	%
aragnooro		· ·	/ 11		,		Ü	· ·
1998	13	65.0			53.8	41.6	92.3	100.0
1999	6	60.0			52.6	40.2	83.3	100.0
2000	7	58.3			50.0	40.3	85.7	100.0
2001	13	54.2			48.7	40.0	92.3	100.0
2002	15	53.6			48.1	40.1	93.3	100.0 #
2003	10	52.6			48.4	40.0	90.0	100.0
2004	15	62.5			46.8	40.3	86.7	86.7
2005	30	71.4			47.7	39.3	80.0	93.3
2006	23	65.7			47.0	39.2	91.3	100.0
2007	28	77.8			46.9	39.0	85.7	100.0 #
2008	20	50.0			44.4	38.3	80.0	95.0
2009	30	66.7			45.7	36.4	73.3	96.7
2010	34	69.4			47.1	36.3	94.1	100.0
2011	27	57.4			49.8	33.6	81.5	100.0
2012	28	62.2			50.2	32.7	82.1	100.0
2013	29	52.7			50.6	32.3	79.3	100.0
2014	35	70.0			51.8	32.8	74.3	97.1
2015	21	63.6			51.6	32.9	52.4	100.0
2016	37	64.9			53.0	28.8	83.8	100.0
2017	33	73.3			54.2	26.5	57.6	100.0
2018	37	66.1			54.8	21.4	54.1	100.0
2019	25	69.4			54.7	27.0	36.0	96.0
2020	17	70.8			55.3	31.3	41.2	100.0 ##
1998-2020	533	64.1			55.3	41.6	75.2	98.5

533 cases diagnosed 1998-2020 are related to a total of 524 patients. Currently, in 419 (80.0 %) of these 524 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 184 / 137 / 98 (35.1 % / 26.1 % / 18.7 %) patients exist having 2 / 3 / 4+ malignancies.

- # The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.
- ## Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retreived from the respective headings.

How to interpret:

In 2018, a subgroup of 37 cases has been diagnosed, of which 54.8 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 21.4 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

Table 1b

Cases with invasive cancer by year of diagnosis, proportions of DCO, further malignancies, deaths, and active follow-up (FEMALES) (incl. DCO)

					Prop. at least 1 further malign.	Prop. at least 1 further		Prop.
			DCO	Prop.	prior +	malign.	Prop.	actively
Year of	Females	Females	cases	DCO	synchron.	after		followed
diagnosis	n	%	n	용	ે	olo	90	%
1998	7	35.0			28.6	33.6	85.7	100.0
1999	4	40.0			36.4	33.3	100.0	100.0
2000	5	41.7			37.5	33.0	80.0	100.0
2001	11	45.8			33.3	32.4	100.0	100.0
2002	13	46.4			30.0	32.7	92.3	100.0 #
2003	9	47.4			26.5	33.2	66.7	100.0
2004	9	37.5			27.6	32.7	77.8	100.0
2005	12	28.6			30.0	32.7	66.7	100.0
2006	12	34.3			30.5	33.0	91.7	100.0
2007	8 /	22.2			31.1	31.6	87.5	100.0 #
2008	20	50.0			35.5	31.2	80.0	100.0
2009	15	33.3			34.4	31.5	93.3	100.0
2010	15	30.6	1	6.7	35.7	31.2	53.3	93.3
2011	20	42.6			39.4	31.2	55.0	100.0
2012	17	37.8	1	5.9	40.7	31.1	76.5	100.0
2013	26	47.3			41.9	29.4	69.2	100.0
2014	15	30.0			43.6	28.2	73.3	100.0
2015	12	36.4	1	8.3	45.2	26.0	75.0	100.0
2016	20	35.1			45.6	23.0	60.0	95.0
2017	12	26.7	1	8.3	46.2	23.3	41.7	100.0
2018	19	33.9			46.6	24.2	36.8	100.0
2019	11	30.6			47.6	11.8	18.2	100.0
2020	7	29.2			47.2	14.3	28.6	100.0 ##
1998-2020	299	35.9	4	1.3	47.2	33.6	68.2	99.3

299 cases diagnosed 1998-2020 are related to a total of 297 patients. Currently, in 203 (68.4 %) of these 297 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 101 / 68 / 34 (34.0 % / 22.9 % / 11.4 %) patients exist having 2 / 3 / 4+ malignancies.

- # The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.
- ## Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retreived from the respective headings.

How to interpret:

In 2018, a subgroup of 19 cases has been diagnosed, of which 46.6 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 24.2 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

Table 2

Incidence measures by year of diagnosis including DCO cases (with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.94 m as of 2007, respectively)

			Males	Fem.	Males	Fem.	Males	Fem.	Males	Fem.
Year of	Males	Females		Inc.	Inc.	Inc.	Inc.	Inc.		Inc.
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
-										
1998	13	7	1.2/	0.6	0.8	0.3	1.1	0.4	1.5	0.5
1999	6	4	0.5	0.3	0.3	0.1	0.5	0.2	0.7	0.2
2000	7	5	0.6	0.4	0.4	0.2	0.6	0.2	0.8	0.4
2001	13	11 /	1.1	0.9	0.6	0.4	1.0	0.6	1.4	0.8
2002	15	13	0.8	0.7	0.4	0.3	0.7	0.5	0.9	0.6
2003	10	9	0.5	0.5	0.2	0.2	0.4	0.3	0.6	0.4
2004	15	9	0.8	0.5	0.4	0.2	0.6	0.3	0.9	0.4
2005	30	12	1.6	0.6	0.8	0.3	1.3	0.4	1.7	0.5
2006	23	12	1.2	0.6	0.6	0.2	0.9	0.3	1.2	0.5
2007	28	8	1.3	0.3	0.6	0.1	0.9	0.2	1.2	0.3
2008	20	20	0.9	0.9	0.5	0.3	0.7	0.5	0.9	0.7
2009	30	15	1.3	0.6	0.7	0.2	1.0	0.3	1.3	0.5
2010	34	15	1.5	0.6	0.6	0.3	1.0	0.5	1.5	0.6
2011	27	20	1.2	0.9	0.6	0.3	0.9	0.4	1.2	0.6
2012	28	17	1.2	0.7	0.5	0.2	0.8	0.4	1.2	0.6
2013	29	26	1.3	1.1	0.5	0.3	0.8	0.5	1.2	0.8
2014	35	15	1.5	0.6	0.6	0.2	1.0	0.4	1.4	0.5
2015	21	12	0.9	0.5	0.4	0.2	0.6	0.2	0.8	0.3
2016	37	20	1.5	0.8	0.7	0.3	1.0	0.5	1.4	0.6
2017	33	12	1.4	0.5	0.6	0.1	0.9	0.2	1.2	0.3
2018	37	19	1.5	0.8	0.7	0.2	1.0	0.3	1.3	0.5
2019	25	11	1.0	0.4	0.5	0.1	0.7	0.2	0.9	0.3
2020	17	7	0.7	0.3	0.3	0.1	0.4	0.2	0.6	0.2
1998-2020	533	299	1.1	0.6	0.5	0.2	0.8	0.3	1.1	0.5

The computation of the incidence measures includes all cancers, irrespective of first or subsequent malignancy.

Table 3 $\label{eq:Age_age} \mbox{Age distribution parameters by year of diagnosis (ALL PATIENTS) } \mbox{(incl. DCO)}$

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	20	70.4	9.0	49.0	83.3	57.1	66.9	70.0	78.3	81.1
1999	10	76.9	10.4	55.0	89.5	62.4	72.3	76.8	87.7	88.6
2000	12	71.2	10.4	45.2	81.7	62.8	66.7	73.1	79.8	80.6
2001	24	73.6	7.4	57.2	87.5	61.8	70.4	74.6	77.9	80.5
2002	28	70.6	9.8	51.7	91.1	58.2	62.5	72.1	75.6	85.0
2003	19	74.5	8.6	56.6	87.3	60.9	69.8	77.4	80.7	85.0
2004	24	72.8	11.4	46.3	94.6	54.9	69.6	74.2	79.8	84.2
2005	42	69.6	13.0	36.1	93.3	53.1	65.8	71.0	77.6	82.7
2006	35	73.0	9.4	50.5	88.7	59.2	67.0	73.4	80.4	84.6
2007	36	72.4	8.4	53.2	87.8	60.3	67.2	73.9	76.4	83.7
2008	40	72.4	9.5	49.3	87.2	61.5	64.5	73.5	79.8	84.5
2009	45	73.3	10.5	29.1	89.0	60.2	69.8	73.8	81.3	83.9
2010	49	73.7	8.9	52.4	92.7	60.0	69.7	73.8	80.4	84.6
2011	47	72.3	9.9	50.3	89.7	58.5	64.6	72.6	80.3	84.8
2012	45	75.2	10.0	49.0	92.8	61.7	68.5	77.4	82.4	83.9
2013	55	74.5	9.6	40.0	90.2	61.7	70.3	75.7	81.7	83.8
2014	50	73.6	10.4	45.1	90.8	59.5	68.0	75.3	80.4	87.6
2015	33	75.5	10.7	52.7	98.9	59.5	69.6	75.3	83.1	90.3
2016	57	73.9	10.7	43.9	94.1	60.3	66.8	75.0	81.3	87.1
2017	45	74.5	9.1	48.5	89.2	65.5	69.0	75.7	81.1	84.8
2018	56	75.2	9.2	51.5	93.4	62.1	66.6	77.6	80.8	87.5
2019	36	74.0	8.7	55.7	91.4	62.1	68.8	73.3	80.7	85.1
2020	24	75.1	9.5	54.9	95.2	62.9	70.8	75.3	80.5	89.0
1998-2020	832	73.5	9.9	29.1	98.9	60.1	67.6	74.6	80.4	84.9

Table 3a

Age distribution parameters by year of diagnosis (MALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	13	69.5	9,1	49.0	81.7	55.6	67.8	69.9	74.9	80.4
1999	6	74.0	11.6	55.0	89.5	55.0	69.7	74.4	80.9	89.5
2000	7	68.9	11.5	45.2	80.3	45.2	66.2	73.0	76.6	80.3
2001	13	73.7	6.6	61.6	86.8	64.0	70.5	74.8	76.9	78.0
2002	15	70.1	9.9	52.0	91.1	58.2	62.3	71.4	75.3	80.3
2003	10	74.4	10.8	56.6	87.3	58.8	63.9	79.1	81.2	86.1
2004	15	73.4	9.8	54.9	86.9	55.9	69.8	74.9	80.1	84.2
2005	30	70.8	11.7	47.6	93.3	53.9	65.9	71.2	77.6	85.9
2006	23	71.4	8.0	50.5	84.3	64.0	65.8	71.2	79.7	80.5
2007	28	71.8	8.8	53.2	87.8	59.6	66.8	71.2	76.3	86.4
2008	20	70.0	10.9	49.3	87.2	55.0	62.5	70.3	78.8	85.0
2009	30	70.7	11.2	29.1	89.0	59.8	65.0	70.4	77.6	82.7
2010	34	76.3	7.6	60.0	92.7	67.4	70.4	76.8	82.1	85.1
2011	27	69.2	9.7	50.3	84.9	58.0	62.0	67.6	79.5	82.6
2012	28	73.0	9.9	49.0	89.1	57.4	67.4	75.1	81.8	83.4
2013	29	73.2	10.4	40.0	86.3	59.7	66.8	74.6	81.0	83.4
2014	35	74.2	11.1	45.1	90.8	57.3	68.2	75.9	81.2	88.1
2015	21	73.6	10.4	52.7	98.9	58.7	69.6	74.8	79.9	84.4
2016	37	74.7	9.5	49.9	94.1	62.7	69.8	75.0	80.8	86.4
2017	33 \	73.5	9.1	48.5	86.4	65.5	69.0	75.4	79.1	84.7
2018	37	73.3	9.9	51.5	93.4	60.8	65.0	75.4	78.6	87.9
2019	25	73.4	8.6	58.7	91.4	62.1	68.9	72.0	78.7	85.1
2020	17	75.4	8.4	60.7	90.3	62.9	70.9	75.1	80.9	89.0
1998-2020	533	72.7	9.8	29.1	98.9	60.0	66.8	73.8	79.6	84.4

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	7	72.2	9,1	58.5	83.3	58.5	64.7	72.4	80.2	83.3
1999	4	81.2	7.8	72.3	87.8	72.3	74.6	82.3	87.7	87.8
2000	5	74.3	8.7	62.8	81.7	62.8	67.3	79.2	80.6	81.7
2001	11	73.5	8.6	57.2	87.5	61.8	70.4	73.7	79.9	80.5
2002	13	71.1	10.1	51.7	87.0	58.6	65.3	72.7	76.0	85.0
2003	9	74.6	6.0	64.1	83.1	64.1	71.0	75.0	79.2	83.1
2004	9	71.8	14.3	46.3	94.6	46.3	69.4	73.5	79.4	94.6
2005	12	66.5	15.9	36.1	91.0	38.2	62.4	69.4	75.9	78.2
2006	12	76.0	11.4	52.1	88.7	58.1	73.1	75.8	85.4	87.4
2007	8	74.7	6.8	60.3	82.8	60.3	72.6	75.7	78.8	82.8
2008	20	74.7	7.3	63.4	85.7	64.1	68.6	74.5	81.0	84.5
2009	15	78.7	6.4	66.0	87.2	67.9	74.3	80.6	83.9	84.0
2010	15	68.1	9.3	52.4	79.8	54.6	59.4	72.2	75.6	78.5
2011	20	76.4	8.9	56.8	89.7	63.4	71.5	76.6	84.1	87.3
2012	17 /	78.9	9.3	56.7	92.8	67.2	77.7	80.0	83.3	91.5
2013	26	76.0	8.7	52.6	90.2	61.7	72.5	77.9	81.7	84.8
2014	15	72.2	8.5	59.5	88.0	59.5	65.4	73.5	76.8	85.4
2015	12	78.8	10.9	59.5	91.9	66.5	69.3	80.4	89.4	90.4
2016	20	72.6	12.7	43.9	88.0	53.4	65.5	75.1	81.5	87.8
2017	12	77.4	8.9	57.5	89.2	68.1	71.3	80.4	82.7	86.2
2018	19	78.9	6.4	61.3	88.6	69.8	77.3	79.2	82.2	87.5
2019	11	75.3	9.3	55.7	87.4	66.8	68.2	75.3	82.7	83.9
2020	7	74.3	12.6	54.9	95.2	54.9	64.9	75.4	80.0	95.2
1998-2020	299	74.8	9.9	36.1	95.2	60.3	69.2	75.9	81.6	86.7

Age at									
diagnosis	Cases			Males			Females		
Years	n	%	Cum.%	'n	용	Cum.%	n	용	Cum.%
0 - 4									
5-9									
10-14									
15-19									
20-24									
25-29	1	0.2	0.2	1	0.2	0.2			0.0
30-34	0	0.0	0.2			0.2			0.0
35-39	1	0.2	0.3	1	0.2	0.5			0.0
40 - 44	1	0.2	0.5			0.5	1	0.5	0.5
45-49	6	1.0	1.5	6	1.5	2.0			0.5
50-54	16	2.6	4.0	10	2.5	4.5	6	2.8	3.2
55-59	28	4.5	8.6	17	4.2	8.7	11	5.1	8.3
60-64	57	9.2	17.8	45	11.2	20.0	12	5.5	13.8
65-69	86	13.9	31.7	62	15.5	35.4	24	11.1	24.9
70-74	110	17.8	49.5	75	18.7	54.1	35	16.1	41.0
75-79	135	21.8	71.4	83	20.7	74.8	52	24.0	65.0
80-84	114	18.4	89.8	65	16.2	91.0	49	22.6	87.6
85+	63	10.2	100.0	36	9.0	100.0	27	12.4	100.0
All ages	618	100.0		401	100.0		217	100.0	

 $$\operatorname{\textsc{Table}}$5$$ Age-specific incidence, DCO rate and proportion of all cancers for period 2007-2020

							Males	Females
			Males	Females	Males	Females	Prop.all	
Age at			Age-	Age-	DCO rate	DCO rate	cancers	cancers
diagnosis	Males	Females	spec.	spec.	n=0	n=4	n=153686	n=155051
Years	n	n	incid.	incid.	%	%	%	%
0- 4								
5- 9								
10-14								
15-19								
20-24								
25-29	1		0.0				0.1	
30-34								
35-39	1		0.0				0.1	
40 - 44		1		0.0				0.0
45-49	6		0.2				0.1	
50-54	10	6	0.4	0.2			0.1	0.0
55-59	16	/ 11 /	0.8	0.5			0.1	0.1
60-64	44	12	2.5	0.6			0.3	0.1
65-69	62	24	3.8	1.3			0.3	0.1
70-74	73	35	4.9	2.0			0.3	0.2
75-79	83	51	6.9	3.4		2.0	0.3	0.3
80-84	65	49	9.0	4.6		2.0	0.4	0.3
85+	36	27	7.7	2.6		7.4	0.3	0.2
All ages	397	216			0.0	1.9	0.3	0.1
Incidence								
Raw			1.2	0.6				
WS			0.5	0.2				
ES			0.8	0.3				
BRD-S			1.1	0.5				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

ICD-10 C66: Malignant neoplasm of ureter

Age distribution and age-specific incidence 2007 - 2020 (Males: 397, Females: 216)




Figure 6. Age distribution (males: mean=73.2 yrs, median=74.2 yrs; females: mean=75.5 yrs, median=76.8 yrs) and age-specific incidence.



ICD-10 C66: Malignant neoplasm of ureter Age-specific incidence rates: international comparison Period Population 2007-2020 4.9 m 2007-2018 86.7 m




Figure 6a. Age-specific incidence in MCR registry areas compared to Germany (FRG, RKI estimates) and SEER (Surveillance, Epidemiology, and End Results, USA).



Reference:

Estimated age-specific patient population of Germany, latest update: 16 March 2021. German Centre for Cancer Registry Data, Robert Koch Institute (RKI), based on data of the population based cancer registries. http://www.krebsdaten.de. Last access: 08/17/2021 Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 21 Regs Research Data, released April 2021, based on the November 2020 submission. http://www.seer.cancer.gov.

Table 7a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of further malignancies for period 1998-2020

MALES

		Observed E	xpected		CI	CI		DCO
Diagnos	is	/ n /	n	SIR	95%	95%	EAR	용
C07-C08	Salivary gland	/ 1/	0.1	10.8	0.3	60.3	5.2	
C12-C13	Hypopharynx	/ 1/	0.1	7.2	0.2	40.2	4.9	
C14	ENT cancer	/ 1	0.0	113.2	2.9	630.6	# 5.7	
C15	Oesophagus	3	0.6	4.8	1.0	14.1	13.7	33.3
C16	Stomach	2	1.5	1.4	0.2	5.0	3.1	
C17	Small intestine	1	0.2	4.8	0.1	27.0	4.6	
C18	Colon	7	3.5	2.0	0.8	4.1	19.9	14.3
C19-C20	Rectum	2	1.7	1.2	0.1	4.2	1.6	
C22	Liver	2	1.0	2.1	0.2	7.4	5.9	
C25	Pancreas	4	1.4	2.8	0.8	7.2	14.8	50.0
C32	Larynx	1	0.3	3.5	0.1	19.6	4.1	
C33-C34		10	3.9	2.6	1.2	4.7	# 35.0	20.0
C43	Malign. melanoma	2	1.5	1.3	0.2	4.8	2.9	
C61	Prostate	44	9.3	4.7	3.4	6.3	# 199.0	2.3
C64	Kidney	12	1.1	10.9	5.6	19.1	# 62.6	50.0
C65	Renal pelvis	57	0.2		260.2	445.2	# 326.3	
C66	Ureter	9	0.1	87.9		166.9		
C67	Bladder	92	1.8	50.3	40.5	61.6	# 517.7	
C68	Urethra	11	0.0	295.7	147.6	529.0	# 62.9	
C68	Urinary org.	10	0.0	329.8	158.1	606.5	# 57.2	80.0
C70-C72	CNS cancer	1	0.4	2.6	0.1	14.2	3.5	
C76-C79	CUP	1	0.6	1.6	0.0	9.1	2.2	
C82-C85		1	1.5	0.7	0.0			
C90	Mult. myeloma	2	0.5	4.3	0.5	15.6	8.8	50.0
Not obs	erved	0	2.9	0.0	0.0	1.3	-16.4	
All fur	ther malignancies	277	34.3	8.1	7.2	9.1	# 1394	7.9
Patients			51	9				
Median ag	e at next maligna	ncv (vears)	74.					
Person-ye	_	1 1	174					
_	rvation time (yea:	rs)	3.					
	servation time (year		2.0					
-1001011 00				-				

The occurrence of further specified malignancy is statistically significant.


Table 7b

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of further malignancies for period 1998-2020 FEMALES

		Observed	Expected		CI	CI			DCO
Diagnosi	.S	/ n /	n	SIR	95%	95%		EAR	%
C16	Stomach	3 /	0.5	6.2	1.3	18.1	#	27.1	
C18	Colon	5 /	1.4	3.6	1.2	8.5	#	39.0	20.0
C23-C24	Bile	/ 1/	0.2	4.9	0.1	27.1		8.6	
C33-C34	Lung	5	0.9	5.5	1.8	12.8	#	44.0	
C50	Breast	6	3.4	1.8	0.6	3.8		27.8	
C54	Corpus uteri	2	0.7	3.0	0.4	10.9		14.4	
C56	Ovary	1	0.5	2.0	0.1	11.3		5.5	100.0
C64	Kidney	9	0.3	29.1	13.3	55.2	#	93.5	66.7
C65	Renal pelvis	24	0.0	533.9	342.1	794.4	#	257.9	
C66	Ureter	2	0.0	74.7	9.0	269.8	#	21.2	
C67	Bladder	38	0.3	131.4	93.0	180.4	#	405.9	2.6
C68	Urinary org.	1	0.0	158.3	4.0	882.1	#	10.7	
C90	Mult. myeloma	1	0.2	6.0	0.2	33.2		9.0	
C91-C96	Leukaemia	1	0.2	4.9	0.1	27.5		8.6	
Not obse	erved	0	3.9	0.0	0.0	0.9	#	-42.2	
All furt	her malignancies	99	12.5	7.9	6.4	9.6	\#	930.9	9.1
Patients			291	_					
Median age	e at next malignar	ncv (vears) 78.2	2					
Person-yea	_	1 11	929)					
_	rvation time (year	rs)	3.2	2					
	servation time (ye		1.5						
	(2)								

The occurrence of further specified malignancy is statistically significant.

Average incidence (Germany 1987 standard population) 2007 - 2020: Males



werage incidence (Germany 1987 standard population) 2007 - 2020: Females





Figure 8a. Map of cancer incidence (german standard population, incl. DCO cases) by county averaged for period 2007 to 2020. According to their individual incidence rates, the counties are displayed in different red and blue hues, being the fine white color attributed to the population mean (males 1.1/100,000 WS N=397, females 0.5/100,000 WS N=216).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,727 female residents (averaged) in the period from 2007 to 2020 a total of 11 women were identified with newly diagnosed ureteral cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 1.0/100,000 (german standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.4 and 2.0/100,000.

Standardized incidence ratio (SIR) 2007 - 2020: Males



Standardized incidence ratio (SIR) 2007 - 2020: Females




Figure 8b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2020. According to their individual SIR values, the counties are displayed in different red and blue hues, being the fine white color attributed to the population overall of 1.0 (males N=397, females N=216).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,153 female residents (averaged) in the period from 2007 to 2020 a total of 11 women were identified with newly diagnosed ureteral cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 1.89. Though, the value of this parameter may vary with an underlying probability of 99% between 0.74 and 3.92, and is therefore not statistically striking.

MORTALITY

Table 9a

Annual cohorts: Incident cancers, follow-up status, proportion of DCO, deaths among the annual cohorts and proportion of available death certificates (with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.94 m as of 2007, respectively)

		Prop.				Prop. deaths
	Incident	actively	Prop.		Prop.	with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	%	%	n	%	%
aragnooro		Č			_	v
1998	20	100.0		18	90.0	100.0
1999	10	100.0		9	90.0	100.0
2000	12	100.0		10	83.3	100.0
2001	24	100.0		23	95.8	100.0
2002	28	100.0		26	92.9	96.2
2003	19	100.0		15	78.9	100.0
2004	24	91.7		20	83.3	90.0
2005	42	95.2		32	76.2	96.9
2006	35	100.0		32	91.4	90.6
2007	36	100.0		31	86.1	100.0
2008	40	97.5		32	80.0	96.9
2009	45	97.8		36	80.0	100.0
2010	49	98.0	2.0	40	81.6	100.0
2011	47	100.0		33	70.2	97.0
2012	45	100.0	2.2	36	80.0	91.7
2013	55	100.0		41	74.5	95.1
2014	50	98.0		37	74.0	89.2
2015	33	100.0	3.0	20	60.6	85.0
2016	57	98.2		43	75.4	86.0
2017	45	100.0	2.2	24	53.3	87.5
2018	56	100.0		27	48.2	70.4
2019	36	97.2		11	30.6	81.8
2020	24	100.0		9	37.5	77.8
1998-2020	832	98.8	0.5	605	72.7	93.1

Table 9b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased within the same year of being diagnosed with cancer (incl. DCO)

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.94 m as of 2007, respectively)

			Prop.		
			deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n /	n	%	n	90
1998	20	13	84.6	3	15.0
1999	10	8	100.0	/ 1	10.0
2000	12	9	100.0	2	16.7
2001	24	12	100.0	3	12.5
2002	28	10	100.0	1	3.6
2003	19	24	95.8	3	15.8
2004	24	13	100.0	1	4.2
2005	42	16	100.0	4	9.5
2006	35	24	95.8	_ 2	5.7
2007	36	20	95.0	1	2.8
2008	40	34	100.0	2	5.0
2009	45	31	100.0	7	15.6
2010	49	27	100.0	3	6.1
2011	47	45	97.8	7	14.9
2012	45	26	100.0	4	8.9
2013	55	31	100.0	9	16.4
2014	50	50	100.0	3	6.0
2015	33	51	96.1	/3	9.1
2016	57	46	100.0	/ 7 /	12.3
2017	45	47	93.6	7 /	15.6
2018	56	32	71.9	5	8.9
2019	36	42	54.8	4	11.1
2020	24	44	93.2	6	25.0
1998-2020	832	655	93.6	88	10.6

Table 9c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.94 m as of 2007, respectively)

				Prop.
				cancer
		Prop.	Prop.	recorded
		cancer-	non-cancer-	on death
Year of	Deaths	related	related	certificate
death	n/	96	%	9
1998	13	61.5	38.5	81.8
1999	8	75.0	25.0	87.5
2000	9	77.8	22.2	88.9
2001	12	75.0	25.0	83.3
2002	10	80.0	20.0	100.0
2003	24	87.5	12.5	87.0
2004	13	69.2	30.8	69.2
2005	16	81.3	18.8	81.3
2006	24	75.0	25.0	82.6
2007	20	85.0	15.0	94.7
2008	34	70.6	29.4	73.5
2009	31	83.9	16.1	90.3
2010	27	81.5	18.5	88.9
2011	45	80.0	20.0	88.6
2012	26	69.2	30.8	80.8
2013	31	71.0	29.0	83.9
2014	50	84.0	16.0	94.0
2015	51	64.7	35.3	73.5
2016	46	73.9	26.1	84.8
2017	47	78.7	21.3	86.4
2018	32	78.1	21.9	91.3
2019	42	52.4	47.6	82.6
2020	44	54.5	45.5	85.4
1998-2020	655	73.4	26.6	85.0

					7
		/			Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(non-cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	4	65.2	66.6	61.3	66.6
1999	5	75.8	76.4	69.1	75.8
2000	6	71.8	67.0	71.8	70.6
2001	8	75.2	74.9	75.6	75.2
2002	6	74.6	73.4	78.8	74.6
2003	16	77.5	76.1	88.3	76.1
2004	7	75.6	74.6	94.7	74.6
2005	6	74.8	72.1	84.5	72.5
2006	15	78.8	77.6	82.0	78.2
2007	18	75.2	73.4	80.9	75.2
2008	20	76.8	75.9	79.7	76.1
2009	23	74.3	74.0	76.5	73.9
2010	16	75.8	75.2	78.1	75.8
2011	28	80.9	79.2	84.6	80.2
2012	19	77.3	75.8	79.3	77.3
2013	20	80.4	80.4	83.4	80.4
2014	28	79.0	79.0	85.3	79.0
2015	31	78.1	77.1	80.8	76.9
2016	32	79.7	78.2	91.9	78.6
2017	29	76.0	74.5	83.3	76.0
2018	26	76.5	75.8	82.5	76.5
2019	31	78.9	71.6	81.4	79.0
2020	33	81.2	80.5	82.7	80.5
1998-2020	427	77.6	76.5	81.4	77.0

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(non-cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	9	78.0	67.4	84.8	72.7
1999	3	77.2	79.5	77.2	79.5
2000	3	73.2	73.2		73.2
2001	4	75.0	69.6	83.7	69.6
2002	4	73.6	73.6		73.6
2003	8	82.4	80.7	89.2	82.4
2004	6	78.3	80.3	76.2	80.3
2005	10	76.2	75.2	77.2	74.7
2006	9	83.1	84.8	79.0	80.0
2007	2	84.1	84.1		84.1
2008	14	79.5	78.6	84.9	79.5
2009	8	78.5	80.8	65.0	78.5
2010	11	77.6	77.3	78.8	77.3
2011	17	80.8	80.3	84.4	80.8
2012	7	83.3	76.2	83.9	79.8
2013	11	80.9	80.9	82.1	80.4
2014	22	81.4	76.1	87.7	80.2
2015	20	82.0	79.8	85.1	80.9
2016	14	84.3	78.9	90.4	80.1
2017	18	85.2	86.3	79.5	84.7
2018	6	88.4	85.0	93.8	87.0
2019	11	80.4	80.4	82.8	79.3
2020	11	85.4	85.4	87.8	85.4
1998-2020	228	81.1	80.4	84.6	80.5

By 2018, Bavarians' life expectancy at birth is estimated at 79.3 years for boys and 83.8 years for girls.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 11a $\begin{tabular}{ll} Mortality measures (cancer-related death) and mortality-incidence-index \\ by year of death \\ MALES \end{tabular}$

Year of	Deaths	Mort.	MI-Index	Mort. 1	/I-Index	Mort. N	II-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
				7		\			
1998	3	0.3	0.23	0.2	0.24	0.2	0.22	0.3	0.19
1999	4	0.4	0.67	0.2	0.55	0.3	0.62	0.6	0.76
2000	4	0.4	0.57	0.2	0.46	0.3	0.54	0.4	0.54
2001	7	0.6	0.54	0.3	0.54	0.5	0.53	0.8	0.57
2002	4	0.2	0.27	0.1	0.27	0.2	0.27	0.2	0.25
2003	14	0.7	1.40	0.4	1.43	0.6	1.46	0.9	1.46
2004	6	0.3	0.40	0.2	0.40	0.3	0.42	0.4	0.41
2005	4	0.2	0.13	0.1	0.13	0.2	0.13	0.2	0.12
2006	11	0.6	0.50	0.2	0.44	0.5	0.53	0.7	0.61
2007	15	0.7	0.56	0.3	0.56	0.5	0.59	0.7	0.62
2008	12	0.5	0.60	0.2	0.50	0.4	0.56	0.6	0.65
2009	19	0.9	0.63	0.4	0.55	0.6	0.60	0.8	0.62
2010	14	0.6	0.41	0.3	0.46	0.4	0.45	0.6	0.43
2011	22	1.0	0.85	0.4	0.61	0.6	0.73	0.9	0.81
2012	15	0.7	0.56	0.3	0.53	0.4	0.56	0.6	0.52
2013	15	0.7	0.52	0.3	0.50	0.4	0.53	0.6	0.53
2014	26	1.1	0.74	0.4	0.64	0.7	0.69	1.0	0.77
2015	21	0.9	1.00	0.3	0.86	0.5	0.94	0.8	0.99
2016	26	1.1	0.72	0.4	0.66	0.7	0.70	0.9	0.69
2017	25	1.0	0.76	0.4	0.79	0.7	0.77	0.9	0.77
2018	21	0.9	0.57	0.4	0.57	0.6	0.57	0.8	0.56
2019	17	0.7	0.68	0.3	0.66	0.5	0.67	0.6	0.67
2020	17	0.7	1.00	0.2	0.87	0.4	0.96	0.6	0.95
1998-2020	322	0.7	0.61	0.3	0.56	0.5	0.60	0.7	0.62

Table 11b $\label{lem:mortality} \mbox{Mortality measures (cancer-related death) and mortality-incidence-index } \mbox{by year of death} \mbox{FEMALES}$

Year of	Deaths	Mort.	MI-Index	Mort. N	/I-Index	Mort. N	/I-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	5	0.4	0.71	0.2	0.82	0.3	0.76	0.4	0.72
1999	2	0.2	0.50	0.1	0.66	0.1	0.62	0.2	0.72
2000	3	0.2	0,60	0.1	0.70	0.2	0.67	0.2	0.52
2001	2	0.2	0.18	0.1	0.18	0.1	0.17	0.1	0.18
2002	4	0.2	0.31	0.1	0.23	0.1/	0.25	0.2	0.32
2003	7	0.4	0.78	0.1	0.67	0.2	0.67	0.2	0.64
2004	3	0.2	0.33	0.0	0.19	0.1	0.22	0.1	0.28
2005	9	0.5	0.75	0.2	0.54	0.3	0.61	0.3	0.67
2006	7	0.3	0.58	0.1	0.50	0.2	0.51	0.2	0.50
2007	2	0.1	0.25	0.0	0.11	0.0	0.14	0.1	0.19
2008	12	0.5	0.60	0.2	0.51	0.3	0.54	0.4	0.56
2009	7	0.3	0.47	0.1	0.46	0.1	0.46	0.2	0.39
2010	8	0.3	0.53	0.1	0.41	0.2	0.43	0.3	0.47
2011	14	0.6	0.70	0.1	0.51	0.3	0.58	0.4	0.70
2012	3	0.1	0.19	0.0	0.21	0.1	0.20	0.1	0.17
2013	7	0.3	0.27	0.1	0.21	0.1	0.22	0.2	0.25
2014	16	0.7	1.07	0.2	0.91	0.4	0.95	0.5	1.07
2015	12	0.5	1.00	0.2	1.05	0.3	1.02	0.4	1.10
2016	8	0.3	0.40	0.1	0.28	0.1	0.31	0.2	0.38
2017	12	0.5	1.00	0.1	0.73	0.2	0.81	0.3	0.83
2018	4	0.2	0.21	0.0	0.20	0.1	0.20	0.1	0.18
2019	5	0.2	0.45	0.0	0.35	0.1	0.38	0.1	0.39
2020	7	0.3	1.00	0.0	0.43	0.1	0.56	0.1	0.68
1998-2020	159	0.3	0.53	0.1	0.45	0.2	0.47	0.2	0.49

Table 12

Age distribution of age at death (cancer-related) for period 2007-2020 (incl. multiple malignancies)

Age at death Years	Cases n	% Cum.%	Males	oyo	Cum.%	Females n	00	Cum.%
0-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39								
40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85+	1 0 9 12 22 41 60 71 84 82	0.3 0.3 0.0 0.3 2.4 2.6 3.1 5.8 5.8 11.5 10.7 22.3 15.7 38.0 18.6 56.5 22.0 78.5 21.5 100.0	7 10 19 30 45 51 50	0.4 2.6 3.8 7.2 11.3 17.0 19.2 18.9 19.6	0.4 0.4 3.0 6.8 14.0 25.3 42.3 61.5 80.4 100.0	2 2 3 11 15 20 34 30	1.7 1.7 2.6 9.4 12.8 17.1 29.1 25.6	0.0 0.0 1.7 3.4 6.0 15.4 28.2 45.3 74.4 100.0
All ages	382	100.0	265	100.0		117	100.0	

Table 13

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2020

(incl. multiple malignancies)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males	Females	/= /		spec.		cancers	cancers
Years	n	n		MI-index		MI-index	%	%
0- 4								
5- 9								
10-14								
15-19								
20-24								
25-29								
30-34								
35-39								
40-44	1		0.0	1.00			0.2	
45-49	_		0.0	1.00			0.2	
50-54	7	2	0.3	0.70	0.1	0.33	0.3	0.1
55-59	10	2	0.5	0.70	0.1	0.18	0.3	0.1
60-64	19	3	1.1	0.43	0.2	0.25	0.3	0.1
65-69	30	11	1.8	0.43	0.6	0.46	0.3	0.1
70-74	45	15	3.0	0.40	0.6		0.3	0.2
75-79	51	20	4.2	0.62	1.3	0.43	0.4	0.2
80-84		34		0.81	3.2	0.69		
	50		6.9				0.5	0.4
85+	52	30	11.1	1.44	2.9	1.11	0.6	0.3
- 1 1	0.65	110					0.4	0 0
All ages	265	117					0.4	0.2
Mortality				0.65	0 0	0.54		
Raw			0.8	0.67	0.3			
WS			0.3	0.61	0.1	0.44		
ES			0.5	0.65	0.2	0.47		
BRD-S			0.8	0.66	0.2	0.49		
PYLL-70								
			1.7		0.4			
per 100,000								
ES			1.5		0.3			
AYLL-70			7.4		6.1			

					Syn-	Syn-		
					chron	chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n	%↓	n	-%	n	← %	n	← %
3		/						
C03-C06 Oral cavity	/ 1	0.3	1	100.0				
C07-C08 Salivary gland	/ 1	0.3					1	100.0
C09-C10 Oropharynx	/ 1 /	0.3	1	100.0				
C12-C13 Hypopharynx	2 -	0.5	1	50.0			1	50.0
C14 ENT cancer	1	0.3					1	100.0
C15 Oesophagus	4	1.1	1	25.0			3	75.0
C16 Stomach	3	0.8					3	100.0
C17 Small intestine	3	0.8	3	100.0				
C18 Colon	19	5.1	11	57.9	1	5.3	7	36.8
C19-C20 Rectum	7	1.9	4	57.1	_ 1	14.3	2	28.6
C21 Anus/canal	1	0.3	1	100.0				
C22 Liver	1	0.3					/ 1	100.0
C25 Pancreas	2	0.5					2	100.0
C32 Larynx	3	0.8	1	33.3/	1	33.3	1	33.3
C33-C34 Lung	10	2.7	2	20.0	1	10.0	7	70.0
C43 Malign. melanoma	4	1.1	3	75.0			1	25.0
C44 Skin others	9	2.4	7	77.8			2	22.2
C46,C49 Soft tissue	2	0.5	2	100.0				
C61 Prostate	48	13.0	24	50.0	11	22.9	13	27.1
C62 Testis	2	0.5	2	100.0				
C64 Kidney	22	6.0	10	45.5	4	18.2	8	36.4
C65 Renal pelvis	46	12.5	8	17.4	29	63.0	9	19.6
C66 Ureter	18	4.9			5	27.8	13	72.2
C67 Bladder	125	33.9	57	45.6	18	14.4	50	40.0
C68 Urethra	9	2.4	2	22.2	1	11.1	6	66.7
C68 Urinary org.	12	3.3	1_	8.3	3	25.0	8	66.7
C69 Eye melanoma	1	0.3	1	100.0				
C70-C72 CNS cancer	2	0.5					2	100.0
C76-C79 CUP	1	0.3	1	100.0				
C82-C85 NHL	8	2.2	6	75.0	2	25.0		
C90 Mult. myeloma	1	0.3					1	100.0
All further malignancies	369	100.0	150	40.7	77	20.9	142	38.5

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a further malignancy.

						Syn-	Syn-		
						chron	chron		
		Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis		n	%↓	n	← %	n	← %	n	← %
C15	Oesophagus	/ 1	0.6					1	100.0
C16	Stomach	3 /	1.8	1	33.3			2	66.7
C17	Small intestine	/ 1 /	0.6	1	100.0				
C18	Colon	/ 12 -	7.3	7	58.3	2	16.7	3	25.0
C19-C20	Rectum	4	2.4	3	75.0			1	25.0
C25	Pancreas	1	0.6	1	100.0				
C33-C34	• 5	6	3.6			2	33.3	4	66.7
C43	Malign. melanoma	4	2.4	4	100.0				
C44	Skin others	10	6.1	4	40.0			6	60.0
C48	Peritoneal	1	0.6	1	100.0				
C50	Breast	18	10.9	14	77.8			4	22.2
C53	Cervix uteri	8	4.8	8	100.0				
C54	Corpus uteri	6	3.6	6	100.0				
C56	Ovary	3	1.8	2	66.7			1	33.3
C64	Kidney	7	4.2	1	14.3	2	28.6	4	57.1
C65	Renal pelvis	23	13.9	5	21.7	11	47.8	7	30.4
C66	Ureter	4	2.4			1	25.0	3	75.0
C67	Bladder	44	26.7	15	34.1	4	9.1	25	56.8
C68	Urethra	1	0.6					1	100.0
C68	Urinary org.	3	1.8	1	33.3	1	33.3	1	33.3
C73	Thyroid	2	1.2	2	100.0				
C76-C79 CUP		1	0.6					1	100.0
C82-C85	NHL	2	1.2	2	100.0				
All fur	ther malignancies	165	100.0	78	47.3	23	13.9	64	38.8

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a further malignancy.

Table 15

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2020

(First primaries only *)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males F	'emales	spec.		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0- 4								
5- 9								
10-14								
15-19								
20-24								
25-29								
30-34								
35-39								
40-44								
45-49								
50-54	3 /	1	0.1	0.60	0.0	0.33	0.1	0.0
55-59	5	1/	0.2	0.71	0.0	0.25	0.1	0.0
60-64	7 /	2	0.4	0.35	0.1	1.00	0.1	0.0
65-69	14	/7	0.9	0.52	0.4	0.78	0.2	0.1
70-74	17	4	1.1	0.71	0.2	0.31	0.2	0.1
75-79	14	10	1.2	0.50	0.7	0.40	0.2	0.1
80-84	16	14	2.2	0.73	1.3	0.67	0.2	0.2
85+	15	15	3.2	2.14	1.4	1.25	0.2	0.2
All ages	91	54					0.2	0.1
							/	
Mortality								
Raw			0.3	0.64	0.2	0.60		
WS			0.1		0.0	0.54		
ES			0.2	0.62	0.1	0.55		
BRD-S			0.3	0.63	0.1	0.55		
DIAD 5			0.5	0.03	0.1	0.55		
PYLL-70								
per 100,000			0.7		0.2			
ES ES			0.6		0.2			
AYLL-70			7.0		5.7			
77771 / /			\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \		3.7			

^{*} See corresponding tables with multiple malignancies.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2020 (Single primaries only *)

			Males		Females		Males	Females
Age at			Age-		Age-			Prop.all
death		Females	spec.		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0- 4								
5- 9								
10-14								
15-19								
20-24								
25-29								
30-34								
35-39								
40-44								
45-49								
50-54	2	1	0.1	0.67	0.0	0.50	0.1	0.0
55-59	2		0.1				0.1	
60-64	6	1	0.3		0.1	1.00	0.1	0.0
65-69	8	5	0.5		0.3	0.71	0.1	0.1
70-74	7	1	0.5		0.1	0.11	0.1	0.0
75-79	5	7	0.4		0.5	0.47	0.1	0.1
80-84	12	9	1.7		0.8	0.64	0.2	0.1
85+	6	9	1.3		0.9	0.82	0.1	0.1
0.5+	O)	1.5	2.00	0.9	0.02	0.1	0.1
All ages	48	33					0.1	0.1
All ages	40	33					0.1	0.1
Mantalita								
Mortality			0.1	0 60	0 1	0 54		
Raw			0.1		0.1	0.54		
WS			0.1		0.0	0.50		
ES			0.1	0.61	0.0	0.50		
BRD-S			0.1	0.62	0.1	0.51		
PYLL-70								
per 100,000			0.4		0.1			
ES			0.4		0.1			
AYLL-70			6.9		5.4			

^{*} See corresponding tables with multiple malignancies.

ICD-10 C66: Malignant neoplasm of ureter

Age distribution and age-specific mortality 2007 - 2020 (Males: 265, Females: 117)





Figure 17. Distribution of age at death (bars; males: mean=73.1 yrs, median=74.6 yrs; females: mean=76.0 yrs, median=77.9 yrs) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at ureteral cancer-related death (see Table 10) should be considered.



werage mortality (Germany 1987 standard population) 2007 - 2020: Males



Average mortality (Germany 1987 standard population) 2007 - 2020: Females





Figure 18a. Map of cancer mortality (german standard population) by county averaged for period 2007 to 2020. According to their individual mortality rates, the counties are displayed in different red and blue hues, being the fine white color attributed to the population mean (males 0.8/100,000 WS N=265, females 0.2/100,000 WS N=117).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,727 female residents (averaged) in the period from 2007 to 2020 a total of 6 women died from ureteral cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.5/100,000 (german standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.1 and 1.3/100,000.

Standardized mortality ratio (SMR) 2007 - 2020: Males



Standardized mortality ratio (SMR) 2007 - 2020: Females




Figure 18b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2020. According to their individual SMR values, the counties are displayed in different red and blue hues, being the fine white color attributed to the population overall of 1.0 (males N=265, females N=117).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,153 female residents (averaged) in the period from 2007 to 2020 a total of 6 women died from ureteral cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 1.94. Though, the value of this parameter may vary with an underlying probability of 99% between 0.50 and 5.06, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases, where applicable. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head&neck and lung).

The ratio of mortality and incidence (mortality-to-incidence ratio, **MIR**, **MI-Index**) is a statistical index that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MIR. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

MCR Munich Cancer Registry (Tumorregister München)

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

SEER Surveillance, Epidemiology, and End Results (USA)

DCO Death certificate only

BRD-S German (FRG) standard population ES European standard population (old)

WS World standard population

SIR Standardized incidence ratio

CI Confidence interval EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

SMR Standardized mortality ratio

MI-index Ratio of mortality to incidence, MIR

FRG Federal Republic of Germany

Recommended Citation

Munich Cancer Registry. ICD-10 C66: Ureteral cancer - Incidence and Mortality [Internet]. 2021 [updated 2021 Dec 21; cited 2022 Feb 1]. Available from: https://www.tumorregister-muenchen.de/en/facts/base/bC66__E-ICD-10-C66-Ureteral-cancer-incidence-and-mortality.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.