Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ Deutsch

ICD-10 C73: Follicular thyroid ca.

Incidence and Mortality

Year of diagnosis	1998-2020
Patients	1,046
Diseases	1,046
Creation date	12/21/2021
Database export	12/20/2021
Population	4.95 m



Munich Cancer Registry
Cancer Registry Bavaria - Upper Bavaria Regional Center
at Klinikum Grosshadern/IBE
Marchioninistr. 15
Munich, 81377
Germany

https://www.tumorregister-muenchen.de/en

https://www.tumorregister-muenchen.de/en/facts/base/bC73F_E-ICD-10-C73-Follicular-thyroid-ca.incidence-and-mortality.pdf

Index of figures and tables

Fig./Tbl	I.	Page
1	Annual cases, DCO, mult. malignancies, follow-up / yr	4
2	Incidence by year of diagnosis	7
3	Age distribution parameters by year of diagnosis	8
4	Age distribution by 5-year age group and sex	11
5	Age-specific incidence, DCO rate, proportion malignancies	12
6	Age distribution and age-specific incidence (chart)	13
6a	Age-specific incidence internationally (chart)	14
7	Standardized incidence ratio of further malignancies	15
8a	Map of cancer incidence (BRD-S) by county (chart)	17
8b	Standardized incidence ratio (SIR) by county (chart)	18
9a	Pts incident cohorts and mortality / yr	19
9b	Incidence and mortality by year of diagnosis	20
9c	Cancer-related deaths, death certification available / yr	21
10	Medians of age at death / yr	22
11	Mortality by year of death	24
12	Distribution of age at death	26
13	Age-specific mortality	27
14	Further malignancies in deaths	28
15	Age-specific mortality (first primaries)	30
16	Age-specific mortality (single primaries)	31
17	Age distribution and age-specific mortality (chart)	32
18a	Map of cancer mortality (BRD-S) by county (chart)	33
18b	Standardized mortality ratio (SMR) by county (chart)	34

Global Statements about the statistics on the Internet – Baseline Statistics (grey button ——), Survival (red button ——)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.69 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases### are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, December 2021

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.65 million to 4.10 in 2002, and to 4.69 million in 2007).
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- ### DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate.

ICD-10 codes (ICD-10 2015) used for specifying cancer site

Code	Description
C73	Malignant neoplasm of thyroid gland
in c	ase of coexisting one of the following

Morphology codes (ICD-O-3 2013) used for specifying cancer site

Code	Description
8330/3 8331/3 8332/3 8335/3	Follicular adenocarcinoma, NOS Follicular adenocarcinoma, well differentiated Follicular adenocarcinoma, trabecular Follicular carcinoma, minimally invasive

Page 4 of 36

INCIDENCE

Table 1

Cases with invasive cancer by year of diagnosis, proportions of DCO, further malignancies, deaths, and active follow-up (ALL PATIENTS) (incl. DCO)

				Prop.			
				at least	Prop.		
				1 further	at least		
				malign.	1 further		Prop.
	All	DCO	Prop.	prior +	malign.	Prop.	actively
Year of	cases	cases	DCO	synchron.	after	deaths	followed
diagnosis	n	n	%	%	%	%	%
1998	43			9.3	14.2	58.1	95.3
1999	35			9.0	14.0	57.1	94.3
2000	46			9.7	13.6	47.8	95.7
2001	31			9.7	13.8	38.7	90.3
2002	52 /			9.2	13.6	42.3	92.3 #
2003	42			10.0	12.7	35.7	95.2
2004	54			9.6	12.1	38.9	94.4
2005	42			11.0	11.5	38.1	85.7
2006	55			10.5	10.4	36.4	94.5
2007	74			11.2	10.5	35.1	87.8 #
2008	67			10.9	9.8	34.3	95.5
2009	59			10.8	9.0	22.0	96.6
2010	48			11.1	8.8	39.6	97.9
2011	40			11.2	7.3	22.5	100.0
2012	43			11.2	7.2	32.6	95.3
2013	48			11.2	6.5	12.5	100.0
2014	45			11.2	5.7	13.3	95.6
2015	51			11.9	4.6	17.6	92.2
2016	41			12.0	4.8	2.4	100.0
2017	43			12.3	4.6	9.3	100.0
2018	47			12.6	1.1		100.0
2019	27			12.4	0.0	7.4	100.0
2020	13			12.5	0.0		100.0 ##
1998-2020	1046			12.5	14.2	29.2	95.2

^{1,046} cases diagnosed 1998-2020 are related to a total of 1,046 patients. Currently, in 274 (26.2 %) of these 1,046 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 221 / 43 / 10 (21.1 % / 4.1 % / 1.0 %) patients exist having 2 / 3 / 4+ malignancies.

How to interpret:

In 2018, a subgroup of 47 cases has been diagnosed, of which 12.6 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 1.1 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

[#] The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

^{##} Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retreived from the respective headings.

Table 1a

Cases with invasive cancer by year of diagnosis, proportions of DCO, further malignancies, deaths, and active follow-up (MALES) (incl. DCO)

					Prop.			
					at least	Prop.		
					1 further	at least		
					malign.	1 further		Prop.
			DCO	Prop.	prior +	malign.	Prop.	actively
Year of	Males	Males	cases	DCO	synchron.	after	deaths	followed
diagnosis	n	용	n	용	용	ଚ୍ଚ	%	%
1998	10	23.3			10.0	15.8	60.0	100.0
1999	13	37.1			13.0	15.1	76.9	100.0
2000	15	32.6			18.4	14.8	53.3	93.3
2001	11	35.5			14.3	14.8	36.4	90.9
2002	17	32.7			10.6	14.4	41.2	88.2 #
2003	14	33.3			11.3	13.5	42.9	100.0
2004	13	24.1			10.8	12.7	53.8	100.0
2005	11	26.2			11.5	12.5	45.5	90.9
2006	17	30.9			9.9	12.3	29.4	94.1
2007	27	36.5			11.5	11.9	44.4	85.2 #
2008	19	28.4			12.0	11.4	31.6	100.0
2009	25	42.4			12.5	9.8	24.0	96.0
2010	16	33.3			13.0	10.0	56.3	100.0
2011	12	30.0			12.7	7.6	25.0	100.0
2012	15	34.9			12.8	8.3	40.0	93.3
2013	24	50.0			12.4	7.7	16.7	100.0
2014	19	42.2			12.6	5.3	21.1	100.0
2015	16	31.4			12.9	5.2	31.3	93.8
2016	18	43.9			13.5	4.8		100.0
2017	15	34.9			13.5	6.5	6.7	100.0
2018	19	40.4			14.5	3.2		100.0
2019	8	29.6			14.1	0.0	25.0	100.0
2020	4	30.8			14.2	0.0		100.0 ##
1998-2020	358	34.2			14.2	15.8	32.4	96.4

358 cases diagnosed 1998-2020 are related to a total of 358 patients. Currently, in 107 (29.9 %) of these 358 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 85 / 20 / 2 (23.7 % / 5.6 % / 0.6 %) patients exist having 2 / 3 / 4+ malignancies.

- # The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.
- ## Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retreived from the respective headings.

How to interpret:

In 2018, a subgroup of 19 cases has been diagnosed, of which 14.5 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 3.2 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

Table 1b

Cases with invasive cancer by year of diagnosis, proportions of DCO, further malignancies, deaths, and active follow-up (FEMALES) (incl. DCO)

					Prop. at least 1 further	Prop. at least 1 further		Dwar
			DCO	Prop.	malign. prior +			Prop. actively
Year of	Fomalog	Females		DCO	synchron.	malign. after	Prop.	followed
diagnosis	n	remares %	n	% %	synchron.	arcer %	%	%
uragnosis	11	•	11/	-0	-0	•	-0	-0
1998	33	76.7			9.1	13.4	57.6	93.9
1999	22	62.9			7.3	13.4	45.5	90.9
2000	31	67.4			5.8	13.0	45.2	96.8
2001	20	64.5			7.5	13.2	40.0	90.0
2002	35	67.3			8.5	13.1	42.9	94.3 #
2003	28	66.7			9.5	12.3	32.1	92.9
2004	41	75.9			9.0	11.8	34.1	92.7
2005	31	73.8			10.8	10.9	35.5	83.9
2006	38	69.1			10.8	9.4	39.5	94.7
2007	47	63.5			11.0	9.8	29.8	89.4 #
2008	48	71.6			10.4	8.8	35.4	93.8
2009	34	57.6			10.0	8.5	20.6	97.1
2010	32	66.7			10.2	8.1	31.3	96.9
2011	28	70.0			10.5	7.0	21.4	100.0
2012	28	65.1			10.5	6.5	28.6	96.4
2013	24	50.0			10.6	5.8	8.3	100.0
2014	26	57.8			10.4	6.0	7.7	92.3
2015	35	68.6			11.4	4.3	11.4	91.4
2016	23	56.1			11.3	4.7	4.3	100.0
2017	28	65.1			11.7	3.6	10.7	100.0
2018	28	59.6			11.7	0.0		100.0
2019	19	70.4			11.5	0.0		100.0
2020	9	69.2			11.6	0.0		100.0 ##
1998-2020	688	65.8			11.6	13.4	27.5	94.6

688 cases diagnosed 1998-2020 are related to a total of 688 patients. Currently, in 167 (24.3 %) of these 688 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 136 / 23 / 8 (19.8 % / 3.3 % / 1.2 %) patients exist having 2 / 3 / 4+ malignancies.

- # The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.
- ## Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retreived from the respective headings.

How to interpret:

In 2018, a subgroup of 28 cases has been diagnosed, of which 11.7 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 0.0 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

Table 2

Incidence measures by year of diagnosis including DCO cases (with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.94 m as of 2007, respectively)

			Males	Fem.	Males	Fem.	Males	Fem.	Males	Fem.
Year of	Males	Females	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
,										
1998	10	33	0.9	2.8	0.6	1.5	0.8	2.1	1.0	2.6
1999	13	22	1.2	1.9	0.7	1,.2	1.1	1.5	1.6	1.8
2000	15	31 /	1.3	2.6	0.8	2.0	1.2	2.3	1.5	2.4
2001	11	20 /	0.9	1.6	0.6	1.0	0.9	1.4	0.9	1.5
2002	17	35	0.9	1.8	0.6	1.1	0.8	1.5	0.9	1.7
2003	14	28	0.7	1.4	0.4	1.0	0.7	1.2	0.8	1.4
2004	13	41	0.7	2.1	0.4	1.4	0.6	1.7	0.7	1.9
2005	11	31	0.6	1.6	0.3	1.0	0.5	1.3	0.6	1.4
2006	17	38	0.9	1.9	0.6	1.1	0.8	1.5	0.8	1.6
2007	27	47	1.2	2.0	0.7	1.4	1.0	1.7	1.2	1.8
2008	19	48	0.9	2.1	0.5	1.2	0.7	1.5	0.8	1.8
2009	25	34	1.1	1.5	0.7	1.0	0.9	1.2	1.1	1.4
2010	16	32	0.7	1.4	0.4	0.8	0.5	1.0	0.7	1.2
2011	12	28	0.5	1.2	0.3	0.8	0.4	1.0	0.5	1.1
2012	15	28	0.7	1.2	0.4	0.8	0.5	1.0	0.6	1.1
2013	24	24	1.0	1.0	0.7	0.6	0.9	0.8	1.0	0.9
2014	19	26	0.8	1.1	0.5	0.8	0.7	1.0	0.7	1.0
2015	16	35	0.7	1.4	0.4	0.9	0.6	1.2	0.6	1.3
2016	18	23	0.7	0.9	0.5	0.8	0.6	0.9	0.7	0.9
2017	15	28	0.6	1.1	0.4	0.7	0.5	0.9	0.6	1.0
2018	19	28	0.8	1.1	0.5	0.7	0.6	0.9	0.7	1.0
2019	8	19	0.3	0.8	0.2	0.5	0.3/	0.6	0.3	0.7
2020	4	9	0.2	0.4	0.1	0.3	0.1	0.3	0.1	0.4
1998-2020	358	688	0.8	1.4	0.5	0.9	0.6	1.2	0.7	1.3

The computation of the incidence measures includes all cancers, irrespective of first or subsequent malignancy.

Table 3 $\label{eq:Age_age} \mbox{Age distribution parameters by year of diagnosis (ALL PATIENTS) } \mbox{(incl. DCO)}$

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	43	61.4	14.7	31.0	83.2	39.7	50.8	64.3	76.0	77.7
1999	35	61.6	16.6	19.5	80.2	40.7	49.4	62.0	77.8	78.6
2000	46	56.4	20.7	11.5	88.5	29.1	36.5	58.9	72.4	79.8
2001	31	58.7	11.8	35.8	78.9	42.9	47.8	59.9	67.9	74.4
2002	52	59.8	14.4	23.4	86.4	41.9	49.0	59.9	69.6	76.8
2003	42	57.7	16.6	21.3	89.5	36,3	47.2	58.6	70.1	78.8
2004	54	58.2	17.1	14.8	86.0	36.7	45.5	62.2	70.0	78.9
2005	42	59.2	17.1	13.5	82.6	36.6	48.7	64.0	72.1	78.7
2006	55	60.3	14.8	20.2	94.9	39.5	53.1	61.9	69.4	77.2
2007	74	55.9	16.1	24.3	83.8	35.0	43.2	57.5	69.1	75.4
2008	67	60.9	14.7	29.7	86.4	42.4	48.2	64.0	70.9	80.6
2009	59	57.7	15.9	13.5	85.9	36.0	47.6	58.5	68.3	79.7
2010	48	61.7	17.2	17.2	87.7	35.7	48.6	63.8	76.0	83.4
2011	40	58.0	15.7	20.1	79.8	36.7	47.9	61.3	70.7	75.0
2012	43	55.7	16.5	20.2	88.5	30.9	45.0	53.2	68.1	74.2
2013	48	58.5	15.8	20.7	88.7	36.1	46.5	61.4	70.4	75.3
2014	45	54.8	17.4	7.7	92.3	33.1	45.5	56.4	63.9	75.4
2015	51	57.5	16.8	20.7	84.5	31.7	44.8	58.6	72.1	77.2
2016	41	52.2	17.1	16.8	84.9	33.9	39.3	50.8	69.4	72.4
2017	43	57.0	15.6	23.5	86.1	36.0	46.5	55.4	68.8	77.6
2018	47	58.2	17.4	23.3	82.3	31.4	43.1	62.5	74.0	78.7
2019	27	56.6	17.7	21.1	86.0	30.0	45.8	56.3	69.6	79.0
2020	13	57.9	19.8	23.3	82.4	35.4	37.3	67.2	74.5	81.2
1998-2020	1046	58.1	16.4	7.7	94.9	35.7	46.2	59.9	70.9	78.5

Table 3a

Age distribution parameters by year of diagnosis (MALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	10	58.5	9.5	46.2	76.3	47.9	52.9	55.0	64.3	73.8
1999	13	67.5	12.6	47.9	80.2	49.4	53.6	72.6	77.9	79.9
2000	15	66.0	15.0	40.5	88.5	46.6	56.0	63.6	78.8	86.7
2001	11	58.1	10.7	40.8	77.0	44.4	50.1	59.0	65.1	69.3
2002	17	58.0	14.1	23.4	81.0	39.3	53.1	59.8	65.5	76.8
2003	14	65.3	8.6	54.3	79.9	55,6	57.5	65.6	71.7	79.2
2004	13	62.7	17.2	36.7	85.2	40.5	45.8	64.1	75.2	83.1
2005	11	60.0	16.4	36.6	78.7	37.8	42.9	64.1	74.2	78.4
2006	17	59.7	12.0	27.2	74.8	41.4	53.5	61.5	68.5	73.1
2007	27	59.1	15.3	24.9	83.6	36.8	44.2	60.5	70.2	77.8
2008	19	59.4	13.0	30.0	79.4	35.0	49.5	64.0	68.9	73.3
2009	25	59.7	15.7	13.5	84.7	47.1	50.9	60.2	68.3	82.7
2010	16	65.1	13.9	38.8	84.7	46.6	54.0	68.3	75.2	84.1
2011	12	62.3	13.4	41.0	79.8	46.5	51.7	62.3	74.1	78.9
2012	15 /	61.0	13.9	39.6	88.5	45.6	48.0	63.7	73.6	75.2
2013	24	60.4	10.5	39.7	76.2	43.6	54.5	61.9	67.2	73.7
2014	19	60.5	16.0	33.1	92.3	41.7	45.9	58.9	73.7	87.6
2015	16	55.7	18.0	23.6	82.0	24.4	47.7	55.7	73.0	75.3
2016	18	54.8	16.2	28.4	82.4	32.7	40.6	52.4	71.0	72.5
2017	15 \	58.2	11.4	41.3	77.6	43.9	48.5	55.9	68.3	74.8
2018	19	57.6	17.5	23.7	82.3	27.4	43.1	62.5	71.6	78.7
2019	8	53.3	22.7	21.1	86.0	21.1	32.2	59.3	68.3	86.0
2020	4	66.0	20.0	37.3	82.4	37.3	52.7	72.1	79.3	82.4
1998-2020	358	60.2	14.5	13.5	92.3	40.8	50.4	61.0	71.6	78.0

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	33	62.2	16.0	31.0	83.2	37.6	50.8	65.4	76.3	77.7
1999	22	58.1	18.0	19.5	78.8	34.6	48.3	60.7	75.7	78.6
2000	31	51.8	21.8	11.5	88.4	28.9	33.7	55.9	67.9	74.5
2001	20	59.0	12.6	35.8	78.9	40.9	46.9	61.2	68.7	75.1
2002	35	60.7	14.6	27.3	86.4	41.9	48.6	59.9	71.7	79.4
2003	28	53.9	18.3	21.3	89.5	25.3	39.9	51.8	67.2	78.8
2004	41	56.8	17.1	14.8	86.0	33.9	45.5	61.6	69.2	75.5
2005	31	58.9	17.6	13.5	82.6	33.3	48.7	64.0	71.8	79.4
2006	38	60.6	16.1	20.2	94.9	36.8	52.0	62.9	70.6	80.7
2007	47	54.1	16.4	24.3	83.8	32.6	38.7	53.0	69.0	74.1
2008	48	61.5	15.4	29.7	86.4	42.4	48.1	64.3	72.8	82.1
2009	34	56.2	16.2	22.8	85.9	35.1	44.9	58.0	67.8	79.4
2010	32	60.0	18.7	17.2	87.7	35.3	45.2	62.1	76.1	82.3
2011	28	56.2	16.4	20.1	79.1	30.2	44.0	61.3	69.9	74.0
2012	28	52.9	17.4	20.2	81.7	28.0	41.3	52.3	67.7	73.8
2013	24	56.6	19.8	20.7	88.7	32.8	38.5	57.5	71.7	80.3
2014	26	50.7	17.5	7.7	84.0	25.7	42.1	52.3	63.0	72.5
2015	35	58.4	16.5	20.7	84.5	38.3	44.8	60.8	70.7	78.1
2016	23	50.2	17.9	16.8	84.9	33.9	38.4	46.9	68.2	71.0
2017	28	56.4	17.6	23.5	86.1	31.5	46.1	53.3	70.6	83.2
2018	28	58.7	17.6	23.3	82.0	31.4	44.1	61.7	75.4	79.4
2019	19	57.9	15.7	22.3	81.3	33.0	48.3	55.3	69.6	79.0
2020	9	54.3	19.8	23.3	81.2	23.3	37.2	54.3	67.8	81.2
1998-2020	688	57.1	17.2	7.7	94.9	33.7	44.4	59.4	70.4	78.8

Age at									
diagnosis	Cases			Males			Females		
Years	n	용	Cum.%	n	용	Cum.%	n	용	Cum.%
0 - 4									
5-9	1	0.2	0.2			0.0	1	0.2	0.2
10-14	1	0.2	0.3	1	0.4	0.4			0.2
15-19	3	0.5	0.8			0.4	3	0.7	1.0
20-24	20	3.1	3.9	5	2.1	2.5/	15	3.7	4.6
25-29	11	1.7	5.6	3	1.3	3.8	8	2.0	6.6
30-34	27	4.2	9.8	6	2.5	6.3	21	5.1	11.7
35-39	42	6.5	16.3	10	4.2	10.5	32	7.8	19.6
40 - 44	51	7.9	24.1	17	7.2	17.7	34	8.3	27.9
45-49	67	10.4	34.5	23	9.7	27.4	44	10.8	38.6
50-54	54	8.4	42.9	22	9.3	36.7	32	7.8	46.5
55-59	58	9.0	51.9	27	11.4	48.1	31	7.6	54.0
60-64	69	10.7	62.5	31	13.1	61.2	38	9.3	63.3
65-69	75	11.6	74.1	28	11.8	73.0	47	11.5	74.8
70-74	75	11.6	85.8	30	12.7	85.7	45	11.0	85.8
75-79	43	6.7	92.4	19	8.0	93.7	24	5.9	91.7
80-84	38	5.9	98.3	11	4.6	98.3	27	6.6	98.3
85+	11	1.7	100.0	4	1.7	100.0	7	1.7	100.0
All ages	646	100.0		237	100.0		409	100.0	

 $$\operatorname{\textsc{Table}}$5$$ Age-specific incidence, DCO rate and proportion of all cancers for period 2007-2020

							Males	Females
			Males	Females	Males	Females	Prop.all	Prop.all
Age at			Age-			DCO rate	_	cancers
diagnosis	Males	Females	spec.	spec.	n=0	n=0		n=155051
Years	n	n	incid.		%	%	%	%
0-4								
5- 9		1		0.1				1.0
10-14	1		0.1				0.7	
15-19		3		0.2				1.1
20-24	5	15	0.2	0.8			0.8	2.9
25-29	3	8	0.1	0.4			0.3	0.7
30-34	6	21	0.3	0.9			0.5	1.0
35-39	10	32	0.4	1.4			0.5	0.9
40 - 44	17	34	0.7	1.4			0.6	0.6
45-49	23	44	0.9	1.7			0.5	0.5
50-54	22	32	0.9	1.3			0.3	0.3
55-59	27	31	1.3	1.4			0.2	0.2
60-64	31	38	1.8	2.0			0.2	0.2
65-69	28	47	1.7	2.6			0.1	0.2
70-74	30	45	2.0	2.6			0.1	0.2
75-79	19	24	1.6	1.6			0.1	0.1
80-84	11	27	1.5	2.5			0.1	0.2
85+	4	7	0.9	0.7			0.0	0.0
All ages	237	409			0.0	0.0	0.2	0.3
Incidence								
Raw			0.7	1.2				
WS			0.5	0.8				
ES			0.6	1.0				
BRD-S			0.7	1.1				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

ICD-10 C73: Follicular thyroid carcinoma (FTC)

Age distribution and age-specific incidence 2007 - 2020 (Males: 237, Females: 409)




Figure 6. Age distribution (males: mean=59.3 yrs, median=60.8 yrs; females: mean=56.4 yrs, median=57.7 yrs) and age-specific incidence.



ICD-10 C73: Follicular thyroid carcinoma (FTC)




Figure 6a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).



Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 21 Regs Research Data, released April 2021, based on the November 2020 submission. http://www.seer.cancer.gov.

Table 7a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of further malignancies for period 1998-2020

MALES

	Observed Ex	xpected		CI	CI		DCO
Diagnosis	/ n /	n	SIR	95%	95%	EAR	용
C03-C06 Oral cavity	/ 1 /	0.2	4.5	0.1	25.1	4.5	
C16 Stomach	4 /	0.9	4.5	1.2	11.4	# 17.9	
C18 Colon	4/	2.2	1.8	0.5	4.7	10.4	
C19-C20 Rectum	7	1.3	5.5	2.2	11.4	# 33.1	
C21 Anus/canal	1	0.1	17.0	0.4	95.0	5.4	
C22 Liver	2	0.7	2.9	0.4	10.5	7.6	
C25 Pancreas	2	0.9	2.2	0.3	8.0	6.3	
C32 Larynx	1	0.3	4.0	0.1	22.2	4.3	
C33-C34 Lung	4	2.8	1.4	0.4	3.7	7.1	25.0
C61 Prostate	8	6.6	1.2	0.5	2.4	8.0	
C64 Kidney	3	0.8	3.7	0.8	10.7	12.6	
C65 Renal pelvis	1	0.1	10.0	0.3	55.7	5.2	
C67 Bladder	4	1.0	3.8	1.0	9.8	# 17.1	25.0
C69 Eye melanoma	1	0.0	37.5	0.9	209.0	5.6	
C73 Thyroid	10	0.2	56.4	27.0	103.6	# 56.8	
C74-C80 Cancer others	1	0.0	20.2	0.5	112.5	5.5	
C76-C79 CUP	1	0.4	2.6	0.1	14.5	3.6	
C82-C85 NHL	1	1.0	1.0	0.0	5.7	0.1	
C90 Mult. myeloma	1	0.3	3.3	0.1	18.4	4.0	
C91-C96 Leukaemia	2	0.3	5.8	0.7	21.0	9.6	
Not observed	0	3.6	0.0	0.0	1.0	-21.0	
All further malignancies	5 59	23.7	2.5	1.9	3.2	# 203.8	3.4
Patients		353					
Median age at next maligna	ancy (years)	69.4					
Person-years		1731					
Mean observation time (year	ars)	4.9					
Median observation time (years)	3.2					

The occurrence of further specified malignancy is statistically significant.

Table 7b


Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of further malignancies for period 1998-2020 FEMALES

Observed Expected CI CI DCO Diagnosis 95% 95% n SIR EAR n C07-C08 Salivary gland 0.1 18.2 0.5 101.5 2.6 C09-C10 Oropharynx 0.1 6.9 38.4 1 0.2 2.4 100.0 1 0.2 4.7 0.1 26.2 C15 Oesophagus 2.2 2 C16 Stomach 1.1 1.8 0.2 6.5 2.5 C18 Colon 4 3.1 1.3 0.3 3.3 2.4 C19-C20 Rectum 4 1.3 3.0 0.8 7.7 7.4 C25 4 1.5 2.7 0.7 6.9 7.0 Pancreas C33-C34 Lung 5 2.5 2.0 0.6 4.7 6.9 C37 Thymus 0.0 49.3 1.2 274.8 # 2.7 1 C38,C45 Mesothelioma 0.1 16.3 0.4 91.0 2.6 1 1.5 0.2 5.3 Malign. melanoma 1.4 1.8 2 C46,C49 Soft tissue 28.3 0.2 5.1 0.1 2.2 1 Breast 10.6 3.0 # C50 2.0 1.2 28.9 4.8 21 C51 Vulva 3 0.3 8.7 1.8 25.6 # 7.4 C52 0.1 89.5 Vagina 1 16.1 0.4 2.6 C54 5 2.7 1.9 0.9 6.2 8.7 Corpus uteri 0.7 0.0 C56 1 1.4 -1.0Ovary 4.1 3 11.2 C64 0.8 3.8 0.8 Kidney 6.1 1 0.1 18.2 0.5 101.5 C66 Ureter 2.6 3 1.0 14.1 C67 Bladder 0.6 4.8 6.6 1 22.8 0.6 127.0 C69 Eye melanoma 0.0 2.6 3 C70-C72 CNS cancer 0.5 6.6 1.4 19.3 7.0 C73 Thyroid 16 0.6 25.0 14.3 40.6/# 42.5 C82-C85 NHL 3 1.3 2.3 0.5 6.7 4.7 Mult. myeloma 1 0.4 2.5 0.1 13.7 1.6 C91-C96 Leukaemia 5 0.5 10.3 3.3 24.0 # 12.5 Not observed 0 3.3 0.0 0.0 1.1 -9.2 All further malignancies 94 33.9 2.8 2.2 3.4 # 166.3

Patients 681
Median age at next malignancy (years) 65.8
Person-years 3611
Mean observation time (years) 5.3
Median observation time (years) 3.9

The occurrence of further specified malignancy is statistically significant.

Average incidence (Germany 1987 standard population) 2007 - 2020: Males



werage incidence (Germany 1987 standard population) 2007 - 2020: Females





Figure 8a. Map of cancer incidence (german standard population, incl. DCO cases) by county averaged for period 2007 to 2020. According to their individual incidence rates, the counties are displayed in different red and blue hues, being the fine white color attributed to the population mean (males 0.7/100,000 WS N=237, females 1.1/100,000 WS N=409).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,727 female residents (averaged) in the period from 2007 to 2020 a total of 10 women were identified with newly diagnosed follicular thyroid ca.. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 1.0/100,000 (german standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.4 and 2.2/100,000.

Standardized incidence ratio (SIR) 2007 - 2020: Males



Standardized incidence ratio (SIR) 2007 - 2020: Females




Figure 8b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2020. According to their individual SIR values, the counties are displayed in different red and blue hues, being the fine white color attributed to the population overall of 1.0 (males N=237, females N=409).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,153 female residents (averaged) in the period from 2007 to 2020 a total of 10 women were identified with newly diagnosed follicular thyroid ca.. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.87. Though, the value of this parameter may vary with an underlying probability of 99% between 0.33 and 1.87, and is therefore not statistically striking.

MORTALITY

Table 9a

Annual cohorts: Incident cancers, follow-up status, proportion of DCO, deaths among the annual cohorts and proportion of available death certificates (with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.94 m as of 2007, respectively)

		Prop.				Prop. deaths
	Incident	actively	Prop.		Prop.	with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	%	%	n	용	90
1998	43	95.3		2.5	58.1	96.0
	_	94.3		25		
1999	35			20	57.1	95.0
2000	46	95.7		22	47.8	100.0
2001	31	90.3		12	38.7	100.0
2002	52	92.3		22	42.3	95.5
2003	42	95.2		15	35.7	93.3
2004	54	94.4		21	38.9	95.2
2005	42	85.7		16	38.1	100.0
2006	55	94.5		20	36.4	95.0
2007	74	87.8		26	35.1	96.2
2008	67	95.5		23	34.3	95.7
2009	59	96.6		13	22.0	100.0
2010	48	97.9		19	39.6	78.9
2011	40	100.0		9	22.5	100.0
2012	43	95.3		14	32.6	100.0
2013	48	100.0		6	12.5	100.0
2014	45	95.6		6	13.3	83.3
2015	51	92.2		9	17.6	77.8
2016	41	100.0		1	2.4	100.0
2017	43	100.0		4	9.3	100.0
2018	47	100.0				
2019	27	100.0		2	7.4	100.0
2020	13	100.0				
1998-2020	1046	95.2		305	29.2	95.1
1000 2020	1040	JJ • Z		303	49.4	JJ • ±

Table 9b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased within the same year of being diagnosed with cancer (incl. DCO)

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.94 m as of 2007, respectively)

			Prop.		
			deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n /	n	%	n	%
1998	43	15	100.0	6	14.0
1999	35	14	92.9	/ 1	2.9
2000	46	17	88.2	3	6.5
2001	31	9	66.7	1	3.2
2002	52	17	100.0		
2003	42	14	92.9		
2004	54	16	93.8	3	5.6
2005	42	24	100.0	1	2.4
2006	55	23	95.7		
2007	74	20	95.0	1	1.4
2008	67	16	100.0	1	1.5
2009	59	24	95.8	2	3.4
2010	48	34	97.1	4	8.3
2011	40	36	100.0	3	7.5
2012	43	19	100.0	1	2.3
2013	48	26	100.0	1 1	2.1
2014	45	22	100.0	1	2.2
2015	51	23	100.0		
2016	41	39	100.0		
2017	43	16	100.0		
2018	47	24	83.3		
2019	27	16	37.5		
2020	13	27	88.9		
1998-2020	1046	401	0.4 1	2.0	2.0
1990-2020	1046	491	94.1	29	2.8

Table 9c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to $4.94~\mathrm{m}$ as of 2007, respectively)

				Prop.
				cancer
		Prop.	Prop.	recorded
		cancer-	non-cancer-	on death
Year of	Deaths	related	related	certificate
death	n/	%	%	용
1998	15	86.7	13.3	93.3
1999	14	85.7	14.3	92.3
2000	17	76.5	23.5	73.3
2001	9	88.9	11.1	100.0
2002	17	94.1	5.9	100.0
2003	14	71.4	28.6	92.3
2004	16	62.5	37.5	60.0
2005	24	75.0	25.0	79.2
2006	23	65.2	34.8	77.3
2007	20	90.0	10.0	94.7
2008	16	62.5	37.5	93.8
2009	24	70.8	29.2	78.3
2010	\ 34	76.5	23.5	81.8
2011	36	61.1	38.9	72.2
2012	19	63.2	36.8	52.6
2013	26	50.0	50.0	53.8
2014	22	72.7	27.3	72.7
2015	23	60.9	39.1	65.2
2016	39	66.7	33.3	79.5
2017	16	62.5	37.5	56.3
2018	24	50.0	50.0	50.0
2019	16	43.8	56.3	16.7
2020	27	33.3	66.7	54.2
1998-2020	491	66.6	33.4	73.6


Table 10a Medians of age at death according to the grouping in Table 9MALES

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(non-cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	7	74.4	69.1	74.5	69.1
1999	5	62.6	67.1	62.1	67.1
2000	9	78.4	78.3	79.1	78.4
2001	4	75.0	72.3	77.8	74.4
2001	10	73.8	73.1	90.1	73.8
2002	4	71.0	71.0	50.1	71.0
2004	7	80.0	80.0	77.2	75.2
2005	7	81.2	82.5	68.6	83.8
2006	5	78.2	78.2	00.0	78.2
2007	9	73.4	73.4		73.4
2008	4	66.6	62.3	70.8	66.6
2009	8	78.6	76.0	84.9	76.9
2010	15	79.7	78.9	87.1	78.9
2011	8	77.2	70.8	77.2	70.8
2012	6	77.8	76.7	91.4	76.7
2013	8	76.8	76.6	77.0	76.4
2014	9	82.1	84.8	76.3	82.1
2015	8	78.7	81.4	78.7	82.9
2016	16	74.5	74.8	74.2	75.1
2017	2	76.8	68.4	85.1	
2018	9	73.9	55.6	77.5	67.5
2019	9	84.5	82.7	84.5	
2020	12	80.9	72.0	83.7	66.2
1998-2020	181	77.2	75.6	79.2	75.6

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

 $\begin{array}{c} \text{Table 10b} \\ \text{Medians of age at death according to the grouping in Table 9} \\ \text{FEMALES} \end{array}$

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(non-cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	8	79.0	76.7	81.3	79.0
1999	9	72.3	71.9	85.3	71.9
2000	8	79.3	76.3	79.9	78.1
2001	5	66.1	66.1		72.7
2002	7	75.6	75.6		75.6
2003	10	82.2	71.1	89.1	74.7
2004	9	78.9	80.0	73.6	78.7
2005	17	77.2	77.0	82.1	77.9
2006	18	77.0	72.5	85.8	72.5
2007	11/	78.0	78.0	73.2	81.7
2008	12	78.5	72.6	82.4	76.2
2009	16	85.3	85.1	87.9	85.3
2010	19	72.8	70.5	79.3	71.1
2011	28	85.1	80.3	88.0	80.3
2012	13	79.3	80.4	73.6	79.3
2013	18	80.4	78.1	80.6	78.1
2014	13	79.7	77.9	91.6	69.7
2015	15	84.9	83.4	91.1	83.2
2016	23	79.1	77.0	87.0	77.0
2017	14	82.3	81.9	86.6	81.9
2018	15	78.6	76.8	85.0	77.1
2019	7	84.7	84.7	77.1	87.9
2020	15	83.4	84.8	83.2	75.5
1998-2020	310	80.3	77.2	83.6	77.4



Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 11a $\begin{tabular}{ll} Mortality measures (cancer-related death) and mortality-incidence-index \\ by year of death \\ MALES \end{tabular}$

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	6	0.5	0.60	0.3	0.51	0.5	0.56	0.6	0.67
1999	4	0.4	0.31	0.2	0.28	0.3	0.27	0.4	0.23
2000	8	0.7	0.53	0.3	0.43	0.6	0.54	1.1	0.74
2001	3	0.3	0.27	0.1	0.24	0.2	0.27	0.3	0.35
2002	9	0.5	0.53	0.3	0.44	0.4	0.50	0.5	0.60
2003	4	0.2	0.29	0.1	0.27	0.2	0.28	0.2	0.28
2004	5	0.3	0.38	0.1	0.29	0.2	0.36	0.3	0.48
2005	6	0.3	0.55	0.1	0.37	0.2	0.49	0.4	0.68
2006	5	0.3	0.29	0.1	0.21	0.2	0.28	0.3	0.37
2007	9	0.4	0.33	0.2	0.27	0.3	0.31	0.4	0.37
2008	3	0.1	0.16	0.1	0.15	0.1	0.15	0.1	0.19
2009	6	0.3	0.24	0.1	0.16	0.2	0.20	0.3	0.26
2010	13	0.6	0.81	0.3	0.69	0.4	0.76	0.6	0.81
2011	4	0.2	0.33	0.1	0.31	0.1	0.32	0.2	0.40
2012	5	0.2	0.33	0.1	0.21	0.1	0.29	0.2	0.36
2013	5	0.2	0.21	0.1	0.13	0.1	0.16	0.2	0.21
2014	6	0.3	0.32	0.1	0.15	0.1	0.23	0.2	0.33
2015	6	0.3	0.38	0.1	0.25	0.2	0.32	0.2	0.35
2016	13	0.5	0.72	0.2	0.47	0.4	0.57	0.5	0.69
2017	1	0.0	0.07	0.0	0.06	0.0	0.07	0.0	0.07
2018	4	0.2	0.21	0.1	0.18	0.1	0.20	0.1	0.19
2019	2	0.1	0.25	0.0	0.13	0.1	0.19	0.1	0.22
2020	6	0.2	1.50	0.1	1.58	0.2	1.59	0.2	1.49
1998-2020	133	0.3	0.37	0.1	0.27	0.2	0.32	0.3	0.39

Table 11b $\label{lem:mortality} \mbox{Mortality measures (cancer-related death) and mortality-incidence-index } \mbox{by year of death} \mbox{FEMALES}$

Year of	Deaths	Mort.	MI-Index						
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	7	0.6	0.21	0.2	0.13	0.3	0.16	0.5	0.19
1999	8	0.7	0.36	0.3	0.27	0.5	0.31	0.6	0.35
2000	5	0.4	0,16	0.2	0.09	0.3	0.11	0.3	0.13
2001	5	0.4	0.25	0.2	0.20	0.3	0.21	0.4	0.24
2002	7	0.4	0.20	0.1	0.12	0.2/	0.14	0.3	0.18
2003	6	0.3	0.21	0.1	0.12	0.2	0.15	0.2	0.17
2004	5	0.3	0.12	0.1	0.05	0.1	0.06	0.2	0.09
2005	12	0.6	0.39	0.2	0.20	0.3	0.26	0.5	0.34
2006	10	0.5	0.26	0.2	0.19	0.3	0.21	0.4	0.24
2007	9	0.4	0.19	0.1	0.09	0.2	0.12	0.3	0.16
2008	7	0.3	0.15	0.1	0.10	0.2	0.11	0.2	0.13
2009	11	0.5	0.32	0.1	0.15	0.2	0.18	0.3	0.21
2010	13	0.6	0.41	0.2	0.30	0.4	0.35	0.5	0.38
2011	18	0.8	0.64	0.2	0.26	0.3	0.35	0.5	0.46
2012	7	0.3	0.25	0.1	0.07	0.1	0.12	0.2	0.17
2013	8	0.3	0.33	0.1	0.16	0.2	0.21	0.2	0.26
2014	10	0.4	0.38	0.2	0.19	0.2	0.23	0.3	0.32
2015	8	0.3	0.23	0.1	0.09	0.1	0.12	0.2	0.16
2016	13	0.5	0.57	0.2	0.21	0.3	0.29	0.4	0.42
2017	9	0.4	0.32	0.1	0.15	0.2	0.19	0.3	0.24
2018	8	0.3	0.29	0.1	0.17	0.2	0.21	0.2	0.23
2019	5	0.2	0.26	0.0	0.06	0.1	0.09	0.1	0.17
2020	3	0.1	0.33	0.0	0.13	0.1	0.19	0.1	0.21
1998-2020	194	0.4	0.28	0.1	0.15	0.2	0.18	0.3	0.23

Table 12

Age distribution of age at death (cancer-related) for period 2007-2020 (incl. multiple malignancies)

Age at	~								
death	Cases	0	a /o	Males	0		Females	0	a 0
Years	n	૾ૢ	Cum.%	n	용	Cum.%	n	양	Cum.%
0 4									
0-4									
5-9									
10-14									
15-19									
20-24									
25-29									
30-34	_								
35-39	1	0.5	0.5	1	1.2	1.2			0.0
40-44	0	0.0	0.5			1.2			0.0
45-49	5	2.4	2.8	3	3.6	4.8	2	1.6	1.6
50-54	4	1.9	4.7	3	3.6	8.4	1	0.8	2.3
55-59	5	2.4	7.1	1	1.2	9.6	4	3.1	5.4
60-64	26	12.3	19.3	13	15.7	25.3	13	10.1	15.5
65-69	14	6.6	25.9	5	6.0	31.3	9	7.0	22.5
70-74	32	15.1	41.0	14	16.9	48.2	18	14.0	36.4
75-79	40	18.9	59.9	17	20.5	68.7	23	17.8	54.3
80-84	36	17.0	76.9	9	10.8	79.5	27	20.9	75.2
85+	49	23.1	100.0	17	20.5	100.0	32	24.8	100.0
All ages	212	100.0		83	100.0		129	100.0	

Table 13

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2020 $\,$

(incl. multiple malignancies)

			Males		Females		Males	Females
Age at			Age-		Age-			Prop.all
death	Males F	'emales	spec.		spec.		cancers	cancers
Years	n	n		/I-index	- \	MI-index	%	% /
			7				-	
0- 4								
5- 9								
10-14								
15-19								
20-24								
25-29								
30-34								
35-39	1		0.0	0.10			0.4	
40-44								
45-49	3	2	0.1	0.13	0.1	0.05	0.2	0.1
50-54	3	1	0.1	0.14	0.0	0.03	0.1	0.0
55-59	1 /	4	0.0	0.04	0.2	0.13	0.0	0.1
60-64	13	13	0.7	0.42	0.7	0.34	0.2	0.3
65-69	5	9	0.3	0.18	0.5	0.19	0.1	0.1
70-74	14	18	0.9	0.47	1.0	0.40	0.1	0.2
75-79	17	23	1.4	0.89	1.5	0.96	0.1	0.2
80-84	9	27	1.2	0.82	2.5	1.00	0.1	0.3
85+	17	32	3.6	4.25	3.1	4.57	0.2	0.3
All ages	83	129					0.1	0.2
3								
Mortality								
Raw			0.3	0.35	0.4	0.32		
WS			0.1	0.25	0.1	0.15		
ES			0.2	0.30	0.2	0.19		
BRD-S			0.2	0.35	0.3	0.24		
PYLL-70								
per 100,000			1.0		0.8			
ES			0.8		0.7			
AYLL-70			10.6		8.0			

					Syn-	Syn-		
					chron	chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n	% ↓	n	← %	n	← %	n	← %
5		/						
C03-C06 Oral cavity	/ 1	1.6			1	100.0		
C15 Oesophagus	/ 1 /	1.6	1	100.0				
C16 Stomach	5 /	7.9					5	100.0
C18 Colon	/ 3 <	4.8					3	100.0
C19-C20 Rectum	4	6.3					4	100.0
C21 Anus/canal	1	1.6					1	100.0
C22 Liver	3	4.8					3	100.0
C25 Pancreas	1	1.6					1	100.0
C32 Larynx	3	4.8	2	66.7	1	33.3		
C33-C34 Lung	6	9.5					6	100.0
C43 Malign. melanoma	1	1.6	1	100.0				
C44 Skin others	4	6.3					4	100.0
C48 Peritoneal	1	1.6					/1	100.0
C61 Prostate	11	17.5	7	63.6			4	36.4
C62 Testis	1	1.6	1	100.0				
C64 Kidney	4	6.3	2	50.0			2	50.0
C65 Renal pelvis	1	1.6					1	100.0
C67 Bladder	2	3.2					2	100.0
C69 Eye melanoma	1	1.6	1	100.0				
C73 Thyroid	2	3.2			2	100.0		
C76-C79 CUP	1	1.6					1	100.0
C82-C85 NHL	1	1.6					1	100.0
C90 Mult. myeloma	1	1.6					1	100.0
C91-C96 Leukaemia	4	6.3					4	100.0
All further malignancies	63	100.0	15	23.8	4	6.3	44	69.8

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a further malignancy.

						Syn-	Syn-		
						chron	chron		
		Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnos	is	n	%↓	n	← %	n	← %	n	← %
C15	Oesophagus	2	2.0					2	100.0
C16	Stomach	4	4.0	1	25.0			3	75.0
C18	Colon	3 /	3.0	1	33.3			2	66.7
C19-C20	Rectum	2	2.0			1	50.0	1	50.0
C21	Anus/canal	1	1.0	1	100.0				
C22	Liver	2	2.0					2	100.0
C25	Pancreas	4	4.0	1	25.0			3	75.0
C33-C34	Lung	9	8.9					9	100.0
C37	Thymus	1	1.0					1	100.0
C43	Malign. melanoma	3	3.0	3	100.0				
C44	Skin others	5	5.0	2	40.0			3	60.0
C46,C49	Soft tissue	1	1.0					1	100.0
C50	Breast	31	30.7	14	45.2			17	54.8
C53	Cervix uteri	1	1.0	1	100.0				
C54	Corpus uteri	6	5.9	3	50.0			3	50.0
C56	Ovary	3	3.0	1	33.3			2	66.7
C64	Kidney	5	5.0	4	80.0			1	20.0
C66	Ureter	1	1.0					1	100.0
C67	Bladder	1	1.0					1	100.0
C70-C72	CNS cancer	2	2.0					2	100.0
C73	Thyroid	2	2.0			1	50.0	1	50.0
C76-C79	CUP	1	1.0	1	100.0				
C82-C85	NHL	3	3.0	1	33.3			2	66.7
C91-C96	Leukaemia	8	7.9					8	100.0
All fur	ther malignancies	101	100.0	34	33.7	2	2.0	65	64.4

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a further malignancy.

Table 15

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2020 (First primaries only *)

		Males		Females		Males	Females
Age at		Age-		Age-		Prop.all	Prop.all
death	Males Female	es spec.		spec.		cancers	cancers
Years	n n		MI-index	mortal.	MI-index	%	왕
0- 4							
5- 9							
10-14							
15-19							
20-24							
25-29							
30-34							
35-39	1	0.0	0.13			0.4	
40-44	_						
45-49	2 1	0.1	0.11	0.0	0.02	0.2	0.1
50-54	3 1	0.1	0.15	0.0	0.03	0.1	0.0
55-59	1 / 4	0.0	0.05	0.2	0.15	0.0	0.1
60-64	11/ 1/1	0.6	0.42	0.6	0.34	0.2	0.3
65-69	4 9	0.2	0.19	0.5	0.26	0.1	0.2
70-74	13 18	0.9	0.59	1.0	0.46	0.1	0.3
75-79	13 16	1.1	1.08	1.1	0.89	0.1	0.2
80-84	8 22	1.1	1.14	2.1	1.00	0.1	0.3
85+	14 22	3.0	4.67	2.1	7.33	0.2	0.2
001	11 \ 22	3.0	1.07	2.1	7.33	0.2	0.2
All ages	70 104					0.1	0.2
nii ages	70 101					/ 0.1	0.2
Mortality							
Raw /		0.2	0.37	0.3	0.30		
WS		0.1	0.26	0.1	0.14		
ES		0.1	0.20	0.2	0.14		
BRD-S		0.2	0.37	0.2	0.23		
DIAD 5		0.2	0.57	0.2	0.25		
PYLL-70							
per 100,000)	0.8		0.7			
ES ES	·	0.7		0.6			
AYLL-70		10.7		7.5			
171111 / 0		10.7		7.5			

^{*} See corresponding tables with multiple malignancies.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2020 (Single primaries only *)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males 1	Females	spec.		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0- 4								
5- 9								
10-14								
15-19								
20-24								
25-29								
30-34								
35-39	1		0.0	0.13			0.4	
40-44								
45-49	2		0.1	0.12			0.2	
50-54	3	1	0.1		0.0	0.04	0.1	0.0
55-59	1 /	2	0.0	0.05	0.1	0.08	0.0	0.1
60-64	8	6	0.5	0.36	0.3	0.22	0.2	0.2
65-69	1	9	0.1	0.08	0.5	0.32	0.0	0.2
70-74	8	8	0.5	0.42	0.5	0.24	0.1	0.1
75-79	9	10	0.7	0.75	0.7	0.59	0.1	0.1
80-84	4	14	0.6	0.57	1.3		0.1	0.2
85+	8	17	1.7	4.00	1.6	5.67	0.1	0.2
001	0	/ \	± • /	1.00	1.0	3.07	0.1	0.2
All ages	45	67					0.1	0.1
TITT ages	10	07					/ 0.1	0.1
Mortality								
Raw			0.1	0.27	0.2	0.21		
WS			0.1	0.19	0.1	0.10		
ES			0.1	0.13	0.1	0.13		
BRD-S			0.1	0.23	0.1	0.13		
סבחשם			0.1	0.27	0.1	0.10		
PYLL-70								
			0.7		0.4			
per 100,000								
ES			0.6		0.3			
AYLL-70			12.8		6.1			

^{*} See corresponding tables with multiple malignancies.

ICD-10 C73: Follicular thyroid carcinoma (FTC)

Age distribution and age-specific mortality 2007 - 2020 (Males: 83, Females: 129)





Figure 17. Distribution of age at death (bars; males: mean=65.1 yrs, median=65.4 yrs; females: mean=66.9 yrs, median=67.6 yrs) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at follicular thyroid ca.-related death (see Table 10) should be considered.



werage mortality (Germany 1987 standard population) 2007 - 2020: Males



Average mortality (Germany 1987 standard population) 2007 - 2020: Females





Figure 18a. Map of cancer mortality (german standard population) by county averaged for period 2007 to 2020. According to their individual mortality rates, the counties are displayed in different red and blue hues, being the fine white color attributed to the population mean (males 0.2/100,000 WS N=83, females 0.3/100,000 WS N=129).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,727 female residents (averaged) in the period from 2007 to 2020 a total of 3 women died from follicular thyroid ca.. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.2/100,000 (german standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.0/100,000.

Standardized mortality ratio (SMR) 2007 - 2020: Males



Standardized mortality ratio (SMR) 2007 - 2020: Females




Figure 18b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2020. According to their individual SMR values, the counties are displayed in different red and blue hues, being the fine white color attributed to the population overall of 1.0 (males N=83, females N=129).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,153 female residents (averaged) in the period from 2007 to 2020 a total of 3 women died from follicular thyroid ca.. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.88. Though, the value of this parameter may vary with an underlying probability of 99% between 0.10 and 3.20, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases, where applicable. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head&neck and lung).

The ratio of mortality and incidence (mortality-to-incidence ratio, **MIR**, **MI-Index**) is a statistical index that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MIR. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

MCR Munich Cancer Registry (Tumorregister München)

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

SEER Surveillance, Epidemiology, and End Results (USA)

DCO Death certificate only

BRD-S German (FRG) standard population ES European standard population (old)

WS World standard population

SIR Standardized incidence ratio

CI Confidence interval EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

SMR Standardized mortality ratio

MI-index Ratio of mortality to incidence, MIR

FRG Federal Republic of Germany

Recommended Citation

Munich Cancer Registry. ICD-10 C73: Follicular thyroid ca. - Incidence and Mortality [Internet]. 2021 [updated 2021 Dec 21; cited 2022 Feb 1]. Available from: https://www.tumorregister-muenchen.de/en/facts/base/bC73F_E-ICD-10-C73-Follicular-thyroid-ca.-incidence-and-mortality.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.