# **Munich Cancer Registry**



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ Deutsch

# ICD-10 C83.1: Mantle cell lymphoma

## **Incidence and Mortality**

Year of diagnosis	1998-2020
Patients	799
Diseases	800
Creation date	12/21/2021
Database export	12/20/2021
Population	4.95 m



Munich Cancer Registry
Cancer Registry Bavaria - Upper Bavaria Regional Center
at Klinikum Grosshadern/IBE
Marchioninistr. 15
Munich, 81377
Germany

https://www.tumorregister-muenchen.de/en

https://www.tumorregister-muenchen.de/en/facts/base/bC831\_E-ICD-10-C83.1-Mantle-cell-lymphoma-incidence-and-mortality.pdf

## Index of figures and tables

Fig./Tbl	I.	Page
1	Annual cases, mult. malignancies, follow-up / yr	4
2	Incidence by year of diagnosis	7
3	Age distribution parameters by year of diagnosis	8
4	Age distribution by 5-year age group and sex	11
5	Age-specific incidence, proportion malignancies	12
6	Age distribution and age-specific incidence (chart)	13
6a	Age-specific incidence internationally (chart)	14
7	Standardized incidence ratio of further malignancies	15
8a	Map of cancer incidence (BRD-S) by county (chart)	17
8b	Standardized incidence ratio (SIR) by county (chart)	18
9a	Pts incident cohorts and mortality / yr	19
9b	Incidence and mortality by year of diagnosis	20
9c	Cancer-related deaths, death certification available / yr	21
10	Medians of age at death / yr	22
11	Mortality by year of death	24
12	Distribution of age at death	26
13	Age-specific mortality	27
14	Further malignancies in deaths	28
15	Age-specific mortality (first primaries)	30
16	Age-specific mortality (single primaries)	31
17	Age distribution and age-specific mortality (chart)	32
18a	Map of cancer mortality (BRD-S) by county (chart)	33
18b	Standardized mortality ratio (SMR) by county (chart)	34

# Global Statements about the statistics on the Internet – Baseline Statistics (grey button ——), Survival (red button ——)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut<sup>#</sup>, with a total of 4.69 million inhabitants, account for the frequency of cancer diseases<sup>##</sup> and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases### are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, December 2021

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.65 million to 4.10 in 2002, and to 4.69 million in 2007).
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- ### DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate.

#### ICD-10 codes (ICD-10 2015) used for specifying cancer site

Code	Description
C83.1	Mantle cell lymphoma

#### **INCIDENCE**

Table 1

Cases with invasive cancer by year of diagnosis, proportions of further malignancies, deaths, and active follow-up (ALL PATIENTS)

		Prop.			
		at least	Prop.		
		1 further	at least		
		malign.	1 further		Prop.
	All	prior +	malign.	Prop.	actively
Year of	cases	synchron.	after	deaths	followed
diagnosis	n	०	િ	%	용
1998	8	25.0	14.1	87.5	100.0
1999	14	13.6	14.1	78.6	100.0
2000	18	15.0	14.4	88.9	100.0
2001	25	13.8	14.3	92.0	96.0
2002	19	13.1	13.8	78.9	100.0 #
2003	21	14.3	13.5	90.5	100.0
2004	33	14.5	13.6	81.8	100.0
2005	35	13.3	13.2	65.7	91.4
2006	48	15.4	12.8	75.0	100.0
2007	44	14.3	11.7	77.3	95.5 #
2008	43	14.3	11.0	69.8	100.0
2009	39	15.0	9.7	61.5	100.0
2010	51	16.1	9.4	76.5	100.0
2011	40	15.8	10.1	67.5	100.0
2012	46	15.9	11.2	56.5	100.0
2013	49	17.4	10.9	57.1	95.9
2014	49	17.2	11.4	55.1	100.0
2015	41	17.3	10.7	46.3	100.0
2016	48	17.7	8.7	56.3	97.9
2017	47	18.0	7.2	31.9	97.9
2018	22	18.1	3.8	31.8	100.0
2019	31	18.0	1.8	35.5	96.8
2020	29	18.3	0.0	17.2	100.0 ##
1998-2020	800	18.3	14.1	62.0	98.6

800 cases diagnosed 1998-2020 are related to a total of 799 patients. Currently, in 239 (29.9 %) of these 799 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 170 / 49 / 20 (21.3 % / 6.1 % / 2.5 %) patients exist having 2 / 3 / 4+ malignancies.

- # The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.
- ## Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retreived from the respective headings.

#### How to interpret:

In 2018, a subgroup of 22 cases has been diagnosed, of which 18.1 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 3.8 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

Table 1a

Cases with invasive cancer by year of diagnosis, proportions of further malignancies, deaths, and active follow-up (MALES)

			Prop.			
			at least	Prop.		
			1 further	at least		
			malign.	1 further		Prop.
			prior +	malign.	Prop.	actively
Year of	Males	Males	synchron.	after	deaths	followed
diagnosis	n	૦૦	%	%	%	용
1000	_	CO E	40.0	15.3	100 0	100 0
1998 1999	5 9	62.5	40.0	15.3	100.0 77.8	100.0
	-	64.3	21.4			100.0
2000	11	61.1	24.0	15.5	81.8	100.0
2001	16	64.0	17.1	15.4	100.0	100.0
2002	16	84.2	15.8	15.1	81.3	100.0 #
2003	16	76.2	16.4	14.8	93.8	100.0
2004	23	69.7	15.6	14.9	87.0	100.0
2005	28	80.0	13.7	14.3	64.3	92.9
2006	35	72.9	13.8	13.9	80.0	100.0
2007	33	75.0	13.5	13.1	81.8	97.0 #
2008	33	76.7	14.2	12.1	66.7	100.0
2009	23	59.0	13.7	10.9	60.9	100.0
2010	34	66.7	15.2	10.1	79.4	100.0
2011	28	70.0	15.5	10.6	60.7	100.0
2012	35	76.1	15.1	11.8	57.1	100.0
2013	33	67.3	17.2	10.9	51.5	93.9
2014	37	75.5	16.9	11.2	51.4	100.0
2015	32	78.0	17.2	9.9	46.9	100.0
2016	35	72.9	18.3	7.6	51.4	97.1
2017	31	66.0	18.3	6.0	32.3	96.8
2018	15	68.2	18.4	1.9	33.3	100.0
2019	20	64.5	18.2	2.6	30.0	95.0
2020	20	69.0	18.3	0.0	15.0	100.0 ##
1998-2020	568	71.0	18.3	15.3	61.8	98.6

568 cases diagnosed 1998-2020 are related to a total of 568 patients. Currently, in 178 (31.3 %) of these 568 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 130 / 32 / 16 (22.9 % / 5.6 % / 2.8 %) patients exist having 2 / 3 / 4+ malignancies.

## Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retreived from the respective headings.

## How to interpret:

In 2018, a subgroup of 15 cases has been diagnosed, of which 18.4 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 1.9 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

<sup>#</sup> The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

Table 1b

Cases with invasive cancer by year of diagnosis, proportions of further malignancies, deaths, and active follow-up (FEMALES)

			Prop.			
			at least	Prop.		
			1 further	at least		
			malign.	1 further		Prop.
			prior +	malign.	Prop.	actively
Year of	Females	Females	synchron.	after	deaths	followed
diagnosis	n	90	%	90	%	%
1998	3	37.5	0.0	11.1	66.7	100.0
1999	5	35.7	0.0	11.2	80.0	100.0
2000	7	38.9	0.0	11.5	100.0	100.0
2001	9	36.0	8.3	11.4	77.8	88.9
2002	3	15.8	7.4	10.4	66.7	100.0 #
2003	5	23.8	9.4	10.1	80.0	100.0
2004	10	30.3	11.9	10.3	70.0	100.0
2005	7 /	20.0	12.2	10.3	71.4	85.7
2006	13 /	27.1	19.4	10.1	61.5	100.0
2007	11/	25.0	16.4	8.4	63.6	90.9 #
2008	10	23.3	14.5	8.4	80.0	100.0
2009	16	41.0	18.2	6.9	62.5	100.0
2010	17	33.3	18.1	7.8	70.6	100.0
2011	12 \	30.0	16.4	8.8	83.3	100.0
2012	11	23.9	18.0	9.9	54.5	100.0
2013	16	32.7	18.1	11.0	68.8	100.0
2014	12	24.5	18.0	12.0	66.7	100.0
2015	9	22.0	17.6	12.7	44.4	100.0
2016	13	27.1	16.4	11.1	69.2	100.0
2017	16	34.0	17.1	9.8	31.3	100.0
2018	7	31.8	17.5	8.0	28.6	100.0
2019	11	35.5	17.5	0.0	45.5	100.0
2020	9	31.0	18.1	0.0	22.2	100.0 ##
1998-2020	232	29.0	18.1	11.1	62.5	98.7

232 cases diagnosed 1998-2020 are related to a total of 231 patients. Currently, in 61 (26.4 %) of these 231 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 40 / 17 / 4 (17.3 % / 7.4 % / 1.7 %) patients exist having 2 / 3 / 4+ malignancies.

## How to interpret:

In 2018, a subgroup of 7 cases has been diagnosed, of which 17.5 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 8.0 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

<sup>#</sup> The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

<sup>##</sup> Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retreived from the respective headings.

Table 2

Incidence measures by year of diagnosis (with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.94 m as of 2007, respectively)

			Males	Fem.	Males	Fem.	Males	Fem.	Males	Fem.
Year of	Males	Females		Inc.	Inc.	Inc.	Inc.	Inc.		Inc.
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
3										
1998	5	3	0.5	0.3	0.3	0.1	0.4	0.2	0.5	0.3
1999	9	5	0.8	0.4	0.5	0.2	0.7	0.2	0.8	0.3
2000	11	7 /	1.0	0.6	0.6	0.3	0.8	0.5	1.0	0.6
2001	16	9 /	1.4	0.7	0.8	0.4	1.2	0.6	1.5	0.7
2002	16	3 <	0.9	0.2	0.5	0.1	0.7	0.1	0.9	0.1
2003	16	5	0.9	0.3	0.5	0.1	0.7	0.2	0.8	0.2
2004	23	10	1.2	0.5	0.7	0.3	1.0	0.4	1.1	0.4
2005	28	7	1.5	0.4	0.8	0.2	1.2	0.2	1.6	0.3
2006	35	13	1.8	0.6	0.9	0.3	1.4	0.4	1.9	0.5
2007	33	11	1.5	0.5	0.8	0.2	1.2	0.3	1.5	0.4
2008	33	10	1.5	0.4	0.7	0.2	1.1	0.3	1.4	0.4
2009	23	16	1.0	0.7	0.5	0.3	0.7	0.5	0.9	0.6
2010	34	17	1.5	0.7	0.7	0.4	1.1	0.5	1.4	0.6
2011	28	12	1.3	0.5	0.7	0.2	0.9	0.3	1.1	0.4
2012	35	11	1.5	0.5	0.7	0.2	1.1	0.3	1.4	0.4
2013	33	16	1.4	0.7	0.8	0.3	1.1	0.4	1.3	0.5
2014	37	12	1.6	0.5	0.8	0.2	1.2	0.3	1.4	0.4
2015	32	9	1.3	0.4	0.7	0.1	1.0	0.2	1.2	0.3
2016	35	13	1.5	0.5	0.6	0.2	1.0	0.3	1.3	0.4
2017	31	16	1.3	0.6	0.6	0.3	0.9	0.4	1.1	0.5
2018	15	7	0.6	0.3	0.3	0.1	0.4	0.2	0.5	0.2
2019	20	11	0.8	0.4	0.4	0.2	0.6	0.3	0.7	0.3
2020	20	9	0.8	0.4	0.4	0.2	0.6	0.2	0.7	0.3
1998-2020	568	232	1.2	0.5	0.6	0.2	0.9	0.3	1.2	0.4

The computation of the incidence measures includes all cancers, irrespective of first or subsequent malignancy.

 $\mbox{Table 3}$  Age distribution parameters by year of diagnosis (ALL PATIENTS)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	8	64.9	10.7	49.9	83.8	49.9	57.2	65.0	70.6	83.8
1999	14	63.1	14.7	35.1	88.1	43.2	54.3	65.0	74.2	78.7
2000	18	66.5	10.2	46.4	81.4	49.9	63.4	66.7	73.5	80.1
2001	25	65.5	9.7	51.5	90.3	55.2	57.7	63.5	70.4	79.2
2002	19	60.5	10.7	39.2	82.4	45.2	54.8	60.1	65.2	82.0
2003	21	66.5	9.3	48.5	80.9	54.7	58.5	66.9	74.4	77.6
2004	33	65.2	14.2	36.4	92.0	38.8	58.2	67.1	74.1	79.7
2005	35	66.2	14.8	22.8	88.2	45.8	58.2	69.8	76.5	81.7
2006	48	69.7	11.1	38.0	87.3	51.8	63.5	71.7	77.7	81.6
2007	44	68.4	9.5	47.7	90.7	54.5	61.0	69.1	75.9	78.0
2008	43	69.5	10.3	47.2	88.5	50.7	67.0	70.4	76.3	82.1
2009	39	67.5	9.4	40.6	80.6	51.8	64.7	68.7	73.5	78.5
2010	51	66.4	12.1	42.1	88.1	48.4	55.2	70.1	75.3	80.5
2011	40	66.5	10.7	42.2	86.8	49.8	60.2	68.8	73.4	78.5
2012	46	68.8	11.7	26.0	90.2	54.1	65.0	71.9	75.7	77.7
2013	49	68.6	12.6	21.2	88.1	49.5	65.0	69.9	76.9	82.9
2014	49	68.1	11.2	41.1	96.9	52.1	60.9	67.1	75.7	82.5
2015	41	69.5	11.9	28.5	90.0	54.6	64.1	70.8	77.9	81.5
2016	48	71.6	11.3	37.2	87.4	54.5	67.3	73.1	80.5	84.5
2017	47	69.7	13.8	39.6	91.2	49.7	58.7	71.5	80.2	88.1
2018	22	69.6	11.7	41.7	85.0	53.7	63.8	71.5	80.0	82.2
2019	31	67.8	12.9	40.6	91.2	53.0	54.6	68.9	80.9	82.9
2020	29	69.1	13.0	49.1	92.0	52.2	58.5	70.2	80.4	87.5
1998-2020	800	67.9	11.8	21.2	96.9	51.8	60.8	69.4	76.5	81.9

Table 3a

Age distribution parameters by year of diagnosis (MALES)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
	_		_ /.	/_			\.			
1998	5	64.5	7.4	53.3	72.0	53.3	61.0	66.7	69.3	72.0
1999	9	59.2	9.7	43.2	74.2	43.2	54.3	59.9	65.3	74.2
2000	11	66.1	9.2	46.4	80.1	56.5	63.4	66.0	73.5	74.0
2001	16	67.3	10.2	55.2	90.3	55.9	59.4	64.4	73.8	81.8
2002	16	61.0	11.5	39.2	82.4	45.2	55.3	61.2	65.3	82.0
2003	16	65.3	8.5	48.5	79.5	54,7	57.5	66.9	71.7	74.7
2004	23	64.5	13.3	36.4	92.0	37.7	60.9	65.9	73.1	76.1
2005	28	65.7	15.3	22.8	88.2	45.7	58.4	69.9	76.0	81.7
2006	35	68.6	11.6	38.0	87.3	49.4	62.2	71.1	77.5	79.6
2007	33	67.8	9.5	47.7	90.7	54.5	61.0	68.0	74.5	78.0
2008	33	68.7	10.6	47.2	88.5	50.0	67.4	70.1	75.1	81.3
2009	23	67.5	8.3	44.3	80.2	62.4	65.2	68.0	71.0	77.0
2010	34	68.1	11.8	42.4	88.1	48.4	63.3	71.1	75.7	80.7
2011	28	64.7	11.7	42.2	86.8	46.9	56.5	66.0	71.9	81.0
2012	35	68.6	11.8	26.0	87.1	54.1	65.0	71.8	75.7	77.3
2013	33	67.5	13.2	21.2	85.6	49.5	65.0	69.8	74.3	79.7
2014	37	66.8	11.1	41.1	85.2	51.8	58.7	67.1	74.2	81.8
2015	32	67.7	12.5	28.5	90.0	54.5	60.5	68.6	76.5	81.5
2016	35	71.7	11.2	37.2	87.4	55.4	68.4	72.9	80.5	84.3
2017	31 \	69.1	13.2	40.3	88.8	55.2	56.8	70.2	78.0	85.6
2018	15	71.6	9.4	53.7	85.0	57.0	64.3	70.4	80.1	82.2
2019	20	66.1	11.1	52.8	82.9	52.9	55.7	64.0	77.4	81.6
2020	20	68.4	12.9	49.1	87.9	51.4	56.9	67.3	80.2	85.1
1998-2020	568	67.4	11.6	21.2	92.0	51.7	60.5	69.0	75.6	80.9

 $\mbox{Table 3b} \label{eq:table 3b}$  Age distribution parameters by year of diagnosis (FEMALES)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	3	65.6	17,1	49.9	83.8	49.9	49.9	63.2	83.8	83.8
1999	5	70.1	20.4	35.1	88.1	35.1	72.7	76.0	78.7	88.1
2000	7	67.1	12.3	49.9	81.4	49.9	51.8	68.4	78.9	81.4
2001	9	62.4	8.5	51.5	79.2	51.5	57.6	61.2	66.0	79.2
2002	3	58.0	5.2	52.1	61.8	52.1	52.1	60.1	61.8	61.8
2003	5	70.3	11.8	52.8	80.9	52.8	63.5	76.5	77.6	80.9
2004	10	67.0	16.7	38.8	89.0	44.1	54.8	68.7	79.7	88.6
2005	7	68.3	13.2	48.4	83.1	48.4	54.0	69.8	81.7	83.1
2006	13	72.7	9.7	57.6	85.3	62.7	64.2	74.6	81.6	83.9
2007	11	70.5	9.6	51.7	84.8	57.5	65.9	72.6	76.7	77.1
2008	10	72.2	9.4	52.5	84.6	59.3	67.0	72.6	78.9	83.5
2009	16	67.7	11.0	40.6	80.6	51.8	60.9	69.1	76.3	79.3
2010	17	63.2	12.3	42.1	81.9	42.2	52.6	63.6	72.0	80.5
2011	12	70.9	6.5	56.5	79.5	64.1	68.0	71.2	76.2	77.5
2012	11 /	69.4	12.3	44.1	90.2	57.0	61.1	73.1	77.5	77.7
2013	16	70.8	11.2	47.7	88.1	52.5	65.0	72.1	79.9	82.9
2014	12	72.0	11.1	60.9	96.9	63.4	63.6	67.8	78.8	85.1
2015	9	75.9	6.8	64.4	86.4	64.4	70.8	78.4	79.4	86.4
2016	13	71.1	11.9	47.9	84.9	54.1	63.0	73.2	81.9	84.5
2017	16	70.8	15.3	39.6	91.2	49.7	60.6	74.4	83.3	88.2
2018	7	65.3	15.5	41.7	83.4	41.7	51.2	72.6	77.8	83.4
2019	11	71.0	15.8	40.6	91.2	53.4	53.5	72.0	83.5	84.7
2020	9	70.7	13.7	54.0	92.0	54.0	58.9	71.7	82.2	92.0
1998-2020	232	69.3	12.1	35.1	96.9	52.3	62.2	71.0	78.8	83.5

 $\label{table 4}$  Age distribution by 5-year age group and sex for period 2007-2020

Age at									
diagnosis	Cases			Males			Females		
Years	n	용	Cum.%	n	%	Cum.%	n	%	Cum.%
0-4									
5-9									
10-14									
15-19									
20-24	1	0.2	0.2	1	0.2	0.2			0.0
25-29 30-34	2	0.3	0.5	2	0.5	0.7			0.0
35-39	2	0.0	0.5	1	0.2	1.0	1	0.6	0.0 0.6
40-44	15	2.6	3.5	9	2.2	3.2	6	3.5	4.1
45-49	21	3.6	7.1	18	4.4	7.6	3	1.8	5.9
50-54	41	7.1	14.2	28	6.8	14.4	13	7.6	13.5
55-59	43	7.4	21.6	32	7.8	22.2	11	6.5	20.0
60-64	64	11.1	32.6	45	11.0	33.3	19	11.2	31.2
65-69 70-74	93 116	16.1 20.0	48.7 68.7	70 85	17.1 20.8	50.4 71.1	23 31	13.5 18.2	44.7 62.9
75-79	89	15.4	84.1	59	14.4	85.6	30	17.6	80.6
80-84	61	10.5	94.6	40	9.8	95.4	21	12.4	92.9
85+	31	5.4	100.0	19	4.6	100.0	12	7.1	100.0
		\ \							
All ages	579	100.0		409	100.0		170	100.0	


Table 5  $\label{table 5} \mbox{Age-specific incidence and proportion of all cancers}$  for period 2007-2020

					Males	Females
			Males	Females	Prop.all	Prop.all
Age at			Age-	Age-	cancers	cancers
diagnosis	Males	Females	spec.	spec.	n=153686	n=155051
Years	n	n	incid.	incid.	%	%
iears	11	11/	incia.	incia.	0	•
0- 4						
5- 9						
10-14						
15-14						
20-24	1		0.0		0 0	
25-29	1 2				0.2	
	2		0.1		0.2	
30-34	1	1	0 0	0 0	0 1	0 0
35-39	1	1	0.0	0.0	0.1	0.0
40-44	9	6	0.4	0.2	0.3	0.1
45-49	18	3	0.7	0.1	0.4	0.0
50-54	28	13	1.1	0.5	0.3	0.1
55-59	32	11	1.5	0.5	0.3	0.1
60-64	45	19	2.5	1.0	0.3	0.1
65-69	70	23	4.3	1.3	0.3	0.1
70-74	85	31	5.7	1.8	0.3	0.2
75-79	59	30	4.9	2.0	0.2	0.2
80-84	40	21	5.5	2.0	0.3	0.1
85+	19	12	4.1	1.2	0.2	0.1
All ages	409	170			0.3	0.1
Incidence						
Raw			1.3	0.5		
WS			0.6	0.2		
ES			0.9	0.3		
BRD-S			1.1	0.4		

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

## ICD-10 C83.1: Mantle cell lymphoma

Age distribution and age-specific incidence 2007 - 2020 (Males: 409, Females: 170)



**Figure 6.** Age distribution (males: mean=68.1 yrs, median=69.9 yrs; females: mean=69.9 yrs, median=71.9 yrs) and age-specific incidence.



#### ICD-10 C83.1: Mantle cell lymphoma Age-specific incidence rates: international comparison Period Population Region 11 MCR 2007-2020 4.9 m ··∲·· SEER 2007-2018 10 1-year averaged incidence (per 100,000) 8 6 5 3 2 60-64 94 75-79 85+ Age at diagnosis (years) **FEMALES**

Figure 6a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).

**MALES** 



Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER\*Stat Database: Incidence - SEER 21 Regs Research Data, released April 2021, based on the November 2020 submission. http://www.seer.cancer.gov.

Table 7a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of further malignancies for period 1998-2020

MALES

	Observed	Expected		CI	CI		DCO
Diagnosis	/ n /	n	SIR	95%	95%	EAR	%
C03-C06 Oral cavity	2	0.3	7.1	0.9		7.9	
C09-C10 Oropharynx	/ 2 /	0.3	5.8	0.7	20.8	7.6	
C16 Stomach	/ 2/	1.3	1.5	0.2	5.4	3.0	
C17 Small intestine	1	0.2	4.4	0.1	24.7	3.6	
C18 Colon	5	3.4	1.5	0.5	3.4	7.4	
C22 Liver	1	1.1	0.9	0.0	5.2	-0.3	
C23-C24 Bile	1	0.4	2.5	0.1	14.2	2.8	
C25 Pancreas	1	1.4	0.7	0.0	3.9	-2.0	
C32 Larynx	1	0.3	2.9	0.1	16.2	3.0	
C33-C34 Lung	12	4.2	2.9	1.5	5.0	# 35.9	8.3
C43 Malign. melanoma	6	1.7	3.6	1.3	7.8	# 19.8	
C46,C49 Soft tissue	1	0.2	4.9	0.1	27.4	3.7	
C48 Peritoneal	1	0.0	33.7	0.9	187.8	4.5	
C61 Prostate	13	10.0	1.3	0.7	2.2	14.0	
C64 Kidney	2	1.2	1.6	0.2	5.9	3.6	
C66 Ureter	1	0.1	10.1	0.3	56.1	4.1	
C67 Bladder	3	1.7	1.8	0.4	5.3	6.1	
C69 Eye lymphoma	1	0.0	142.8	3.6	795.5	# 4.6	
C73 Thyroid	1	0.2	4.6	0.1	25.4	3.6	
C76-C79 CUP	2	0.6	3.4	0.4	12.4	6.5	
C82-C85 NHL	2	1.5	1.3	0.2	4.8	2.3	
C90 Mult. myeloma	1	0.5	2.1	0.1	11.8	2.4	
C91-C96 Leukaemia	2	0.5	3.7	0.4		6.7	
Not observed	0	4.6	0.0	0.0	0.8	# -21.1	
All further malignancies	64	35.8	1.8	1.4	2.3	# 129.5	1.6
Patients		55	7				
Median age at next maligna:	ncv (vears						
Person-years	ncy (years	2179					
Mean observation time (year	rel	3.9					
Median observation time (year		2.0					
median observacion cime (ye	ears)	۷.۰۷	U				

# The occurrence of further specified malignancy is statistically significant.

Table 7b


Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of further malignancies for period 1998-2020

FEMALES


	Observed 1	Expected		CI	CI		DCO
Diagnosis	/ n /	n	SIR	95%	95%	EAR	%
C16 Stomach	/ 1 /	0.3	3.2	0.1	18.0	8.6	
C17 Small intestine	/ 1/	0.1	19.4	0.5	108.2	11.8	
C18 Colon	/ 3/	0.9	3.4	0.7	9.8	26.1	
C22 Liver	1	0.1	8.5	0.2	47.3	10.9	
C25 Pancreas	1	0.4	2.2	0.1	12.4	6.8	
C33-C34 Lung	5	0.7	7.0	2.3	16.4	# 53.2	20.0
C43 Malign. melanoma	1	0.4	2.8	0.1	15.6	8.0	
C46,C49 Soft tissue	1	0.1	18.9	0.5	105.5	11.7	
C50 Breast	5	2.8	1.8	0.6	4.2	27.3	
C54 Corpus uteri	1	0.5	2.0	0.0	10.9	6.0	
C82-C85 NHL	1	0.4	2.8	0.1	15.4	7.9	
C90 Mult. myeloma	1	0.1	8.9	0.2	49.4	11.0	
C91-C96 Leukaemia	1	0.1	7.2	0.2	40.4	10.7	
Not observed	0	2.4	0.0	0.0	1.5	-29.7	
All further malignancies	23	9.3	2.5	1.6	3.7	# 170.4	4.3
Patients		225					
Median age at next malignan	cy (years	74.5					
Person-years		806					
Mean observation time (year	s)	3.6					
Median observation time (ye		2.5					

# The occurrence of further specified malignancy is statistically significant.

#### Average incidence (Germany 1987 standard population) 2007 - 2020: Males




## werage incidence (Germany 1987 standard population) 2007 - 2020: Females




**Figure 8a.** Map of cancer incidence (german standard population) by county averaged for period 2007 to 2020. According to their individual incidence rates, the counties are displayed in different red and blue hues, being the fine white color attributed to the population mean (males 1.1/100,000 WS N=409, females 0.4/100,000 WS N=170).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,727 female residents (averaged) in the period from 2007 to 2020 a total of 4 women were identified with newly diagnosed mantle cell lymphoma. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.3/100,000 (german standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.1 and 1.1/100,000.

## Standardized incidence ratio (SIR) 2007 - 2020: Males



#### Standardized incidence ratio (SIR) 2007 - 2020: Females



**Figure 8b.** Map of standardized incidence ratio (SIR) by county averaged for period 2007 to 2020. According to their individual SIR values, the counties are displayed in different red and blue hues, being the fine white color attributed to the population overall of 1.0 (males N=409, females N=170).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,153 female residents (averaged) in the period from 2007 to 2020 a total of 4 women were identified with newly diagnosed mantle cell lymphoma. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.86. Though, the value of this parameter may vary with an underlying probability of 99% between 0.14 and 2.70, and is therefore not statistically striking.

## **MORTALITY**

Table 9a

Annual cohorts: Incident cancers, follow-up status, and deaths among the annual cohorts

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.94 m as of 2007, respectively)

					Dmon
		Danas			Prop. deaths
	Incident	Prop.		Dman	with death
		actively	\\/,	Prop.	
Year of	cases	followed	Deaths	deaths	certific.
diagnosis	n	%	n	%	%
1998	8	100.0	7	87.5	100.0
1999	14	100.0	11	78.6	90.9
2000	18	100.0	16	88.9	100.0
2001	25	96.0	23	92.0	100.0
2001	19	100.0	15	78.9	100.0
2002	21	100.0	19	90.5	94.7
2003	33	100.0	27	81.8	100.0
2005	35	91.4	23	65.7	95.7
2005	48	100.0	36	75.0	97.2
2007	44	95.5	34	77.3	91.2
2007	43	100.0	30	69.8	100.0
2008	39	100.0	24		95.8
				61.5	
2010	51	100.0	39	76.5	89.7
2011	40	100.0	27	67.5	92.6
2012	46	100.0	26	56.5	92.3
2013	49	95.9	28	57.1	89.3
2014	49	100.0	27	55.1	92.6
2015	41	100.0	19	46.3	94.7
2016	48	97.9	27	56.3	88.9
2017	47	97.9	15	31.9	86.7
2018	22	100.0	7	31.8	71.4
2019	31	96.8	11	35.5	90.9
2020	29	100.0	5	17.2	80.0
1998-2020	800	98.6	496	62.0	93.8

Table 9b

Annual cohorts of incident cancers and deaths, and cases deceased within the same year of being diagnosed with cancer

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.94 m as of 2007, respectively)

				D	
V	T		Dankha da	Prop.	
Year of	Incident	D	Deaths in	deaths in	
diagnosis/	cases	Deaths	same year	same year	
death	/n /	n	n	<u></u>	
1000		4 =			
1998	8	15	/ _ /		
1999	14	4	/ 1/	7.1	
2000	18	11	1/1	5.6	
2001	25	9	_1	4.0	
2002	19	9	1	5.3	
2003	21	15	1	4.8	
2004	33	24	4	12.1	
2005	35	27	5	14.3	
2006	48	19	2	4.2	
2007	44	17	3 _	6.8	
2008	43	33	4	9.3	
2009	39	29	3	7.7	
2010	51	30	4	7.8	
2011	40	32	2	5.0	
2012	46	33	7	15.2	
2013	49	35	8	16.3	
2014	49	35	6	12.2	
2015	41	29	5	12.2	
2016	48	37	5	10.4	
2017	47	40	7	14.9	
2018	22	37	3	13.6	
2019	31	22	1	3.2	
2020	29	32	2	6.9	
		Ŭ <b>2</b>		0.3	
1998-2020	800	574	76	9.5	

Table 9c

Annual cohorts of deaths, and proportion of cancer-related and non-cancer-related deaths

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.94 m as of 2007, respectively)

				Prop.
		Prop.	Prop.	cancer recorded
		/ -	7	on death
Year of	Deaths	cancer- related	non-cancer- related	certificate
death	/ -	related %	related %	%
death	n	6	6	6
1998	15	73.3	26.7	100.0
1999	4	50.0	50.0	100.0
2000	11	81.8	18.2	88.9
2001	9	88.9	11.1	100.0
2002	9	88.9	11.1	100.0
2003	15	93.3	6.7	100.0
2004	24	79.2	20.8	91.7
2005	27	92.6	7.4	100.0
2006	/ 19	94.7	5.3	100.0
2007	17	76.5	23.5	100.0
2008	33	78.8	21.2	96.8
2009	29	89.7	10.3	96.6
2010	\ 30	90.0	10.0	100.0
2011	32	81.3	18.8	84.4
2012	33	87.9	12.1	97.0
2013	35	85.7	14.3	90.9
2014	35	85.7	14.3	94.3
2015	29	86.2	13.8	79.3
2016	37	83.8	16.2	91.9
2017	40	82.5	17.5	82.5
2018	37	67.6	32.4	75.0
2019	22	68.2	31.8	100.0
2020	32	75.0	25.0	80.6
1998-2020	574	82.6	17.4	91.9


 $\begin{array}{c} \text{Table 10a} \\ \text{Medians of age at death according to the grouping in Table 9} \\ \text{MALES} \end{array}$ 

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(non-cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
		/			
1998	10	67.4	64.6	84.2	64.7
1999	4	75.8	86.1	63.3	75.8
2000	6	72.2	72.3	65.3	72.2
2001	6	66.3	66.3		66.3
2002	6	78.6	82.3	75.0	78.6
2003	13	69.2	68.1	91.9	69.2
2004	11	69.4	69.4	74.3	69.4
2005	21	73.6	72.9	83.4	73.6
2006	12	65.7	66.6	59.8	65.7
2007	14	71.0	71.9	61.2	71.0
2008	26	71.8	69.5	78.8	73.2
2009	25	75.0	75.1	69.5	75.0
2010	13	71.2	71.3	70.4	71.2
2011	22	75.5	77.3	67.6	76.4
2012	24	76.1	76.4	68.9	75.8
2013	26	73.6	71.7	85.0	73.1
2014	25	73.6	73.3	74.1	73.6
2015	21	76.5	72.9	82.8	72.5
2016	25	78.3	77.3	82.5	78.2
2017	27	77.3	76.7	87.4	76.1
2018	24	79.0	74.8	79.2	74.8
2019	19	80.1	80.0	83.1	79.1
2020	19	80.9	80.0	84.7	78.0
1998-2020	399	74.7	74.2	78.8	73.7

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

 $\begin{array}{c} \text{Table 10b} \\ \text{Medians of age at death according to the grouping in Table 9} \\ \text{FEMALES} \end{array}$ 

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(non-cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	5	71.9	67.7	86.6	71.9
1999					
2000	5	85.5	84.4	85.5	84.4
2001	3	77.9	76.2	88.6	77.9
2002	3	62.2	62.2		62.2
2003	2	77.9	77.9		77.9
2004	13	80.5	77.9	90.8	77.9
2005	6	73.4	73.4		73.4
2006	7 /	78.8	78.8		78.8
2007	3/	66.7	66.7		66.7
2008	7	82.0	82.0		83.6
2009	4	76.8	71.6	81.9	74.4
2010	17	73.7	73.7	76.1	73.7
2011	10	77.8	77.2	89.1	77.2
2012	9	69.2	69.0	89.5	69.2
2013	\9	81.4	81.4	81.4	81.4
2014	10	73.8	71.6	79.2	73.8
2015	8	72.1	72.1	75.5	69.0
2016	12	78.9	78.9		78.9
2017	13	82.2	81.6	92.9	80.4
2018	13	79.4	78.1	82.8	79.4
2019	3	72.0	72.8	70.1	73.7
2020	13	82.1	81.9	91.5	83.1
1998-2020	175	78.5	77.1	85.5	78.2



Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 11a  $\begin{tabular}{ll} Mortality measures (cancer-related death) and mortality-incidence-index \\ by year of death \\ MALES \end{tabular}$ 

Year of	Deaths	Mort.	MI-Index	Mort. N	4I-Index	Mort. I	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	7	0.6	1.40	0.4	1.34	0.6	1.34	0.6	1.34
1999	2	0.2	0.22	0.1	0.16	0.2	0.23	0.2	0.28
2000	5	0.4	0.45	0.2	0.41	0.4	0.49	0.5	0.52
2001	6	0.5	0.38	0.3	0.40	0.4	0.38	0.5	0.37
2002	5	0.3	0.31	0.1	0.22	0.2	0.28	0.4	0.41
2003	12	0.6	0.75	0.4	0.68	0.5	0.72	0.7	0.79
2004	8	0.4	0.35	0.2	0.30	0.3	0.31	0.4	0.37
2005	19	1.0	0.68	0.5	0.60	0.8	0.67	1.1	0.67
2006	11	0.6	0.31	0.3	0.33	0.5	0.32	0.6	0.32
2007	10	0.5	0.30	0.2	0.25	0.3	0.27	0.5	0.31
2008	19	0.9	0.58	0.4	0.62	0.7	0.60	0.8	0.59
2009	24	1.1	1.04	0.5	0.92	0.8	1.07	1.1	1.19
2010	12	0.5	0.35	0.2	0.30	0.3	0.32	0.5	0.35
2011	18	0.8	0.64	0.3	0.49	0.6	0.59	0.8	0.72
2012	21	0.9	0.60	0.4	0.51	0.6	0.58	0.9	0.66
2013	22	1.0	0.67	0.5	0.59	0.7	0.63	0.9	0.68
2014	21	0.9	0.57	0.4	0.47	0.6	0.50	0.8	0.57
2015	19	0.8	0.59	0.4	0.54	0.5	0.55	0.8	0.61
2016	19	0.8	0.54	0.3	0.44	0.5	0.47	0.7	0.54
2017	22	0.9	0.71	0.3	0.53	0.5	0.59	0.8	0.71
2018	15	0.6	1.00	0.3	0.94	0.4	0.99	0.5	0.97
2019	13	0.5	0.65	0.2	0.51	0.3	0.56	0.5	0.65
2020	14	0.6	0.70	0.1	0.39	0.3	0.48	0.5	0.69
1998-2020	324	0.7	0.57	0.3	0.49	0.5	0.53	0.7	0.59

Table 11b  $\label{lem:mortality} \mbox{Mortality measures (cancer-related death) and mortality-incidence-index } \mbox{by year of death} \mbox{FEMALES}$ 

Year of death	Deaths n	Mort.	MI-Index raw	Mort.	MI-Index WS	Mort. ES	MI-Index ES	Mort. BRD-S	MI-Index BRD-S
acacii	11	Law	iaw /	***	WO		\	DIAD D	DIED 5
1998 1999	4	0.3	1.33	0.2	1.06	0.2	1.20	0.3	1.16
2000	4	0.3	0.57	0.1	0.30	0.2	0.35	0.2	0.38
2001	2	0.2	0.22	0.1	0.15	0.1	0.17	0.2	0.23
2002	3	0.2		0.1	0.68	0.1	0.70	0.1	0.88
2003	2	0.1	0.40	0.0	0.32	0.1	0.33	0.1	0.37
2004	11	0.6	1.10	0.2	0.75	0.3	0.88	0.5	1.08
2005	6	0.3	0.86	0.1	0.68	0.2	0.76	0.3	0.86
2006	7	0.3	0.58	0.1	0.47	0.2	0.53	0.3	0.56
2007	3	0.1	0.27	0.1	0.29	0.1	0.27	0.1	0.24
2008	7	0.3	0.70	0.1	0.54	0.2	0.60	0.2	0.59
2009	2	0.1	0.13	0.0	0.12	0.1	0.11	0.1	0.10
2010	15	0.6	0.88	0.2	0.60	0.4	0.66	0.5	0.83
2011	8	0.3	0.67	0.1	0.44	0.2	0.50	0.3	0.63
2012	8	0.3	0.73	0.2	0.74	0.2	0.74	0.3	0.76
2013	8	0.3	0.50	0.1	0.30	0.2	0.37	0.3	0.51
2014	9	0.4	0.75	0.1	0.61	0.2	0.66	0.3	0.76
2015	6	0.2	0.67	0.1	0.93	0.2	0.81	0.2	0.76
2016	12	0.5	0.92	0.1	0.53	0.2	0.65	0.3	0.71
2017	11	0.4	0.69	0.1	0.47	0.2	0.52	0.3	0.63
2018	10	0.4	1.43	0.1	0.72	0.2	0.90	0.3	1.14
2019	2	0.1	0.18	0.0	0.20	0.1	0.21	0.1	0.21
2020	10	0.4	1.11	0.1	0.92	0.2	0.93	0.3	1.00
1998-2020	150	0.3	0.65	0.1	0.49	0.2	0.54	0.2	0.61

Table 12

Age distribution of age at death (cancer-related) for period 2007-2020 (incl. multiple malignancies)

Age at									
death	Cases			Males			Females		
Years	n	용	Cum.%	'n	%	Cum.%	n	응	Cum.%
0 - 4									
5-9									
10-14									
15-19									
20-24									
25-29	1	0.3	0.3	1	0.4	0.4			0.0
30-34	0	0.0	0.3			0.4			0.0
35-39	0	0.0	0.3			0.4			0.0
40 - 44	0	0.0	0.3			0.4			0.0
45-49	8	2.2	2.5	6	2.4	2.8	2	1.8	1.8
50-54	11	3.1	5.6	6	2.4	5.2	5	4.5	6.3
55-59	18	5.0	10.6	14	5.6	10.8	4	3.6	9.9
60-64	27	7.5	18.1	19	7.6	18.5	8	7.2	17.1
65-69	43	11.9	30.0	32	12.9	31.3	11	9.9	27.0
70-74	56	15.6	45.6	42	16.9	48.2	14	12.6	39.6
75-79	79	21.9	67.5	54	21.7	69.9	25	22.5	62.2
80-84	73	20.3	87.8	49	19.7	89.6	24	21.6	83.8
85+	44	12.2	100.0	26	10.4	100.0	18	16.2	100.0
All ages	360	100.0		249	100.0		111	100.0	

Table 13

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2020

## (incl. multiple malignancies)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males	Females	spec.		spec.		cancers	cancers
Years	n	n		MI-index		MI-index	%	%
0- 4								
5- 9								
10-14								
15-19								
20-24								
25-29	1		0.0	0.50			1.1	
30-34								
35-39								
40-44								
45-49	6	2	0.2	0.33	0.1	0.67	0.4	0.1
50-54	6	5	0.2	0.21	0.2	0.38	0.2	0.2
55-59	14	4	0.7	0.44	0.2	0.36	0.3	0.1
60-64	19	8	1.1	0.42	0.4	0.42	0.3	0.2
65-69	32	11	2.0	0.46	0.6	0.48	0.3	0.2
70-74	42	14	2.8	0.49	0.8	0.45	0.4	0.2
75-79	54	25	4.5	0.92	1.7	0.83	0.4	0.3
80-84	49	24	6.8	1.22	2.3	1.14	0.5	0.3
85+	26	18	5.6	1.37	1.7	1.50	0.3	0.2
All ages	249	111					0.4	0.2
3								
Mortality								
Raw			0.8	0.61	0.3	0.65		
WS			0.3	0.51	0.1	0.50		
ES			0.5	0.55	0.2	0.54		
BRD-S			0.7	0.62	0.2	0.61		
PYLL-70								
per 100,000			2.4		1.0			
ES			2.0		0.8			
AYLL-70			8.7		9.0			

					Syn-	Syn-		
					chron	chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n	%↓	n	<b>←</b> %	n	<b>←</b> %	n	<b>←</b> %
C00 Lip	/ 1	0.8					1	100.0
C03-C06 Oral cavity	2	1.5					2	100.0
C09-C10 Oropharynx	2 /	1.5	1	50.0			1	50.0
C15 Oesophagus	1 -	0.8	1	100.0				
C16 Stomach	6	4.6	2	33.3			4	66.7
C18 Colon	13	9.9	5	38.5	4	30.8	4	30.8
C19-C20 Rectum	2	1.5	2	100.0				
C22 Liver	2	1.5					2	100.0
C23-C24 Bile	1	0.8					1	100.0
C25 Pancreas	1	0.8	1	100.0				
C32 Larynx	1	0.8					1,	100.0
C33-C34 Lung	11	8.4	2	18.2	2	18.2	7	63.6
C37 Thymus	1	0.8	1	100.0				
C43 Malign. melanoma	10	7.6	5	50.0			5	50.0
C44 Skin others	26	19.8	7	26.9	3	11.5	16	61.5
C46,C49 Soft tissue	1	0.8	1	100.0				
C48 Peritoneal	1	0.8					1	100.0
C50 Breast	1	0.8	1	100.0				
C60 Penis	1	0.8					1	100.0
C61 Prostate	22	16.8	17	77.3	1	4.5	4	18.2
C64 Kidney	3	2.3	2	66.7	1	33.3		
C65 Renal pelvis	1	0.8			1/	100.0		
C67 Bladder	7	5.3	3	42.9	1	14.3	3	42.9
C69 Eye lymphoma	1	0.8					1	100.0
C73 Thyroid	1	0.8	1	100.0				
C76-C79 CUP	3	2.3					3	100.0
C82-C85 NHL	5	3.8			1	20.0	4	80.0
C90 Mult. myeloma	2	1.5					2	100.0
C91-C96 Leukaemia	2	1.5					2	100.0
All further malignancies	131	100.0	52	39.7	14	10.7	65	49.6

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a further malignancy.

					Syn- chron	Syn- chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n	%↓	n	<b>←</b> %	n	<b>←</b> %	n	<b>←</b> %
C16 Stomach	3	4.7	1	33.3			2	66.7
C17 Small intestine	/ 1	1.6			1	100.0		
C18 Colon	6	9.4	4	66.7			2	33.3
C19-C20 Rectum	3	4.7	2	66.7			1	33.3
C25 Pancreas	1	1.6					1	100.0
C33-C34 Lung	7	10.9	1	14.3	1	14.3	5	71.4
C43 Malign. melanoma	4	6.3	2	50.0			2	50.0
C44 Skin others	11	17.2	1	9.1			10	90.9
C46,C49 Soft tissue	1	1.6					1	100.0
C50 Breast	10	15.6	6	60.0	_ 1	10.0	3	30.0
C53 Cervix uteri	1	1.6	1	100.0				
C54 Corpus uteri	2	3.1	1	50.0			1	50.0
C55,C57 Fem. genitals un	2	3.1	2	100.0				
C56 Ovary	1	1.6					1	100.0
C64 Kidney	2	3.1	2	100.0				
C81 Hodgkin lymphoma	1	1.6					1	100.0
C82-C85 NHL	8	12.5					8	100.0
All further malignancies	64	100.0	23	35.9	3	4.7	38	59.4

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a further malignancy.

Table 15

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2020 (First primaries only \*)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males Fe	emales	spec.		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0- 4								
5- 9								
10-14								
15-19								
20-24								
25-29	1		0.0	0.50			1.2	
30-34								
35-39								
40-44								
45-49	6	2	0.2	0.33	0.1	0.67	0.5	0.1
50-54	5	3	0.2	0.19	0.1	0.27	0.2	0.1
55-59	11 /	4	0.5	0.38	0.2	0.44	0.3	0.1
60-64	19	8	1.1	0.46	0.4	0.47	0.4	0.2
65-69	31	10	1.9	0.53	0.6	0.59	0.4	0.2
70-74	32	11	2.1	0.50	0.6	0.46	0.4	0.2
75-79	43	19	3.6	1.13	1.3	0.76	0.5	0.3
80-84	38	17	5.2	1.52	1.6	1.00	0.5	0.2
85+	19	15	4.1	1.58	1.4	1.88	0.3	0.2
001	13	13	1.1	1.30	1.1	1.00	0.5	0.2
All ages	205	89					0.4	0.2
TITE ages	200	03					/ 0.1	0.2
Mortality								
Raw			0.6	0.63	0.3	0.65		
WS			0.3		0.1	0.03		
ES			0.3	0.56	0.1	0.51		
BRD-S			0.4	0.65	0.2	0.60		
DKD-3			0.0	0.05	0.2	0.00		
PYLL-70								
per 100,000			2.2		0.8			
ES ES			1.8		0.8			
AYLL-70			8.5		8.6			
עוחח– ו ת			0.3		0.0			

<sup>\*</sup> See corresponding tables with multiple malignancies.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2020


(Single primaries only \*)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males Fe	males	spec.		spec.		cancers	cancers
Years	n	n i	mortal.	MI-index	mortal.	MI-index	૾ૢ	%
0- 4								
5- 9								
10-14								
15-19								
20-24								
25-29	1		0.0	0.50			1.2	
30-34								
35-39								
40-44								
45-49	6	2	0.2	0.33	0.1	0.67	0.5	0.1
50-54	5	3	0.2	0.20	0.1	0.27	0.2	0.1
55-59	10	4	0.5	0.36	0.2	0.44	0.3	0.1
60-64	17	8	1.0	0.45	0.4	0.50	0.3	0.2
65-69	25	7	1.5	0.53	0.4	0.47	0.3	0.1
70-74	29	9	1.9	0.54	0.5	0.43	0.3	0.1
75-79	37	18	3.1	1.23	1.2	0.78	0.4	0.2
80-84	28	13	3.9	1.47	1.2	0.87	0.4	0.2
85+	14	12	3.0	1.17	1.2	1.50	0.2	0.1
051	14	12	3.0	Ι•Ι/	1.2	1.50	0.2	0.1
All ages	172	76					0.3	0.2
TITE ages	1 / 2	// 0					0.3	0.2
Mortality								
Raw			0.5	0.61	0.2	0.60		
WS			0.2	0.49	0.1	0.48		
ES			0.4	0.43	0.1	0.40		
BRD-S			0.5	0.62	0.2	0.56		
DKD-2			0.5	0.02	0.2	0.50		
PYLL-70								
per 100,000			2.0		0.8			
ES ES			1.7		0.6			
AYLL-70			9.1		9.4			
AITT-/0			3.1		9.4			

<sup>\*</sup> See corresponding tables with multiple malignancies.

## ICD-10 C83.1: Mantle cell lymphoma

Age distribution and age-specific mortality 2007 - 2020 (Males: 249, Females: 111)




**Figure 17.** Distribution of age at death (bars; males: mean=69.5 yrs, median=71.0 yrs; females: mean=71.1 yrs, median=73.2 yrs) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).


The difference between age at diagnosis (Table 3) and age at mantle cell lymphoma-related death (see Table 10) should be considered.



#### werage mortality (Germany 1987 standard population) 2007 - 2020: Males




## Average mortality (Germany 1987 standard population) 2007 - 2020: Females




**Figure 18a.** Map of cancer mortality (german standard population) by county averaged for period 2007 to 2020. According to their individual mortality rates, the counties are displayed in different red and blue hues, being the fine white color attributed to the population mean (males 0.7/100,000 WS N=249, females 0.2/100,000 WS N=111).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,727 female residents (averaged) in the period from 2007 to 2020 a total of 0 women died from mantle cell lymphoma. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.0/100,000 (german standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 0.0/100,000.

## Standardized mortality ratio (SMR) 2007 - 2020: Males



#### Standardized mortality ratio (SMR) 2007 - 2020: Females



**Figure 18b.** Map of standardized mortality ratio (SMR) by county averaged for period 2007 to 2020. According to their individual SMR values, the counties are displayed in different red and blue hues, being the fine white color attributed to the population overall of 1.0 (males N=249, females N=111).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,153 female residents (averaged) in the period from 2007 to 2020 a total of 0 women died from mantle cell lymphoma. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.00. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 1.77, and is therefore not statistically striking.

#### Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases, where applicable. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

#### 1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

## 2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

#### 3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head&neck and lung).

The ratio of mortality and incidence (mortality-to-incidence ratio, **MIR**, **MI-Index**) is a statistical index that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MIR. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

#### **Shortcuts**

MCR Munich Cancer Registry (Tumorregister München)

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

SEER Surveillance, Epidemiology, and End Results (USA)

DCO Death certificate only

BRD-S German (FRG) standard population ES European standard population (old)

WS World standard population

SIR Standardized incidence ratio

CI Confidence interval EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

SMR Standardized mortality ratio

MI-index Ratio of mortality to incidence, MIR

FRG Federal Republic of Germany

#### **Recommended Citation**

Munich Cancer Registry. ICD-10 C83.1: Mantle cell lymphoma - Incidence and Mortality [Internet]. 2021 [updated 2021 Dec 21; cited 2022 Feb 1]. Available from: https://www.tumorregister-muenchen.de/en/facts/base/bC831\_E-ICD-10-C83.1-Mantle-cell-lymphoma-incidence-and-mortality.pdf

#### Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

#### Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.