

Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ *Deutsch*

ICD-10 C84: Mature T/NK-cell I.

Incidence and Mortality

Year of diagnosis	1998-2019
Patients	617
Diseases	617
Creation date	01/26/2021
Database export	01/07/2021
Population	4.92 m





Munich Cancer Registry
Cancer Registry Bavaria - Upper Bavaria Regional Center
at Klinikum Grosshadern/IBE
Marchioninstr. 15
Munich, 81377
Germany

<https://www.tumorregister-muenchen.de/en>

https://www.tumorregister-muenchen.de/en/facts/base/bC84__E-ICD-10-C84-Mature-T-NK-cell-I.-incidence-and-mortality.pdf

Index of figures and tables

Fig./Tbl.		Page
1	Annual cases, DCO, mult. malignancies, follow-up / yr	4
2	Incidence by year of diagnosis	7
3	Age distribution parameters by year of diagnosis	8
4	Age distribution by 5-year age group and sex	11
5	Age-specific incidence, DCO rate, proportion malignancies	12
6	Age distribution and age-specific incidence (chart)	13
6a	Age-specific incidence internationally (chart)	14
7	Standardized incidence ratio of further malignancies	15
8a	Map of cancer incidence (BRD-S) by county (chart)	17
8b	Standardized incidence ratio (SIR) by county (chart)	18
9a	Pts incident cohorts and mortality / yr	19
9b	Incidence and mortality by year of diagnosis	20
9c	Cancer-related deaths, death certification available / yr	21
10	Medians of age at death / yr	22
11	Mortality by year of death	24
12	Distribution of age at death	26
13	Age-specific mortality	27
14	Further malignancies in deaths	28
15	Age-specific mortality (first primaries)	30
16	Age-specific mortality (single primaries)	31
17	Age distribution and age-specific mortality (chart)	32
18a	Map of cancer mortality (BRD-S) by county (chart)	33
18b	Standardized mortality ratio (SMR) by county (chart)	34

**Global Statements about the statistics on the Internet –
Baseline Statistics** (grey button ) , **Survival** (red button )

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.69 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases^{###} are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, January 2021

[#] Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.65 million to 4.10 in 2002, and to 4.69 million in 2007).

^{##} Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.

^{###} DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate.

ICD-10 codes (ICD-10 2016) used for specifying cancer site

Code	Description
C84.-	Mature T/NK-cell lymphomas
C84.0	Mycosis fungoides
C84.1	Sézary disease
C84.4	Peripheral T-cell lymphoma, not elsewhere classified
C84.5	Other mature T/NK-cell lymphomas
C84.6	Anaplastic large cell lymphoma, ALK-positive
C84.7	Anaplastic large cell lymphoma, ALK-negative
C84.8	Cutaneous T-cell lymphoma, unspecified
C84.9	Mature T/NK-cell lymphoma, unspecified

INCIDENCE

Table 1

Cases by year of diagnosis, proportions of DCO, further malignancies, deaths, and active follow-up (ALL PATIENTS) (incl. DCO)

Year of diagnosis	All cases n	DCO cases n	Prop. DCO %	Prop. at least 1 further malign. prior + synchron. %	Prop. at least 1 further malign. after %	Prop. deaths %	Prop. actively followed %
1998	12			0.0	9.2	66.7	100.0
1999	14			3.8	8.9	78.6	100.0
2000	17			7.0	8.6	70.6	88.2
2001	15	1	6.7	15.5	8.7	66.7	100.0
2002	22			15.0	8.6	68.2	95.5 #
2003	24			15.4	8.0	62.5	100.0
2004	24	1	4.2	14.8	7.9	66.7	91.7
2005	33			13.7	8.3	66.7	93.9
2006	28			13.8	8.3	71.4	92.9
2007	31	4	12.9	13.6	8.1	77.4	100.0 #
2008	46	7	15.2	15.4	7.4	67.4	97.8
2009	34	1	2.9	15.3	7.5	55.9	100.0
2010	37	3	8.1	16.6	8.0	67.6	97.3
2011	44	7	15.9	18.6	7.3	72.7	97.7
2012	34	3	8.8	18.3	7.4	73.5	97.1
2013	46	5	10.9	18.0	8.7	69.6	95.7
2014	49	5	10.2	18.4	6.5	69.4	98.0
2015	32	3	9.4	19.4	4.8	53.1	100.0
2016	29	4	13.8	19.3	2.7	72.4	100.0
2017	23	3	13.0	19.5	2.3	39.1	100.0
2018	14			19.4	0.0	28.6	100.0
2019	9			19.6	0.0	44.4	88.9 ##
1998-2019	617	47	7.6	19.6	9.2	65.8	97.2

617 cases diagnosed 1998-2019 are related to a total of 617 patients. Currently, in 173 (28.0 %) of these 617 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 125 / 39 / 9 (20.3 % / 6.3 % / 1.5 %) patients exist having 2 / 3 / 4+ malignancies.

The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retrieved from the respective headings.

How to interpret:

In 2017, a subgroup of 23 cases has been diagnosed, of which 19.5 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 2.3 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

Table 1a

Cases by year of diagnosis, proportions of DCO, further malignancies, deaths, and active follow-up (MALES) (incl. DCO)

Year of diagnosis	Males n	Males %	DCO cases n	Prop. DCO %	Prop. at least 1 further malign. prior + synchron. %	Prop. at least 1 further malign. after %	Prop. deaths %	Prop. actively followed %
1998	9	75.0			0.0	9.2	55.6	100.0
1999	9	64.3			0.0	8.7	77.8	100.0
2000	8	47.1			7.7	8.9	62.5	87.5
2001	6	40.0	1	16.7	9.4	9.1	83.3	100.0
2002	15	68.2			10.6	9.2	66.7	93.3 #
2003	12	50.0			13.6	9.0	58.3	100.0
2004	12	50.0			14.1	9.0	58.3	91.7
2005	23	69.7			13.8	9.3	65.2	91.3
2006	17	60.7			13.5	9.4	76.5	100.0
2007	17	54.8	2	11.8	14.1	9.2	88.2	100.0 #
2008	34	73.9	5	14.7	17.9	8.3	67.6	97.1
2009	25	73.5			17.6	8.2	56.0	100.0
2010	26	70.3	3	11.5	19.2	9.1	73.1	100.0
2011	29	65.9	4	13.8	21.5	8.3	72.4	96.6
2012	23	67.6	2	8.7	20.4	7.9	69.6	95.7
2013	32	69.6	2	6.3	19.5	9.2	71.9	96.9
2014	35	71.4	2	5.7	19.9	6.3	68.6	97.1
2015	22	68.8	2	9.1	20.6	3.9	50.0	100.0
2016	23	79.3	4	17.4	20.4	0.0	73.9	100.0
2017	15	65.2	3	20.0	20.7	0.0	53.3	100.0
2018	10	71.4			20.4	0.0	30.0	100.0
2019	8	88.9			20.7	0.0	50.0	87.5 ##
1998-2019	410	66.5	30	7.3	20.7	9.2	66.3	97.3

410 cases diagnosed 1998-2019 are related to a total of 410 patients. Currently, in 122 (29.8 %) of these 410 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 90 / 27 / 5 (22.0 % / 6.6 % / 1.2 %) patients exist having 2 / 3 / 4+ malignancies.

The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retrieved from the respective headings.

How to interpret:

In 2017, a subgroup of 15 cases has been diagnosed, of which 20.7 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 0.0 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

Table 1b

Cases by year of diagnosis, proportions of DCO, further malignancies, deaths, and active follow-up (FEMALES) (incl. DCO)

Year of diagnosis	Females n	Females %	DCO cases n	Prop. DCO %	Prop. at least 1 further malign. prior + synchron. %	Prop. at least 1 further malign. after %	Prop. deaths %	Prop. actively followed %
1998	3	25.0			0.0	9.3	100.0	100.0
1999	5	35.7			12.5	9.4	80.0	100.0
2000	9	52.9			5.9	8.1	77.8	88.9
2001	9	60.0			23.1	8.0	55.6	100.0
2002	7	31.8			21.2	7.3	71.4	100.0 #
2003	12	50.0			17.8	5.8	66.7	100.0
2004	12	50.0	1	8.3	15.8	5.6	75.0	91.7
2005	10	30.3			13.4	6.1	70.0	100.0
2006	11	39.3			14.1	5.8	63.6	81.8
2007	14	45.2	2	14.3	13.0	5.5	64.3	100.0 #
2008	12	26.1	2	16.7	11.5	5.3	66.7	100.0
2009	9	26.5	1	11.1	11.5	5.9	55.6	100.0
2010	11	29.7			12.1	5.4	54.5	90.9
2011	15	34.1	3	20.0	13.7	4.9	73.3	100.0
2012	11	32.4	1	9.1	14.7	6.1	81.8	100.0
2013	14	30.4	3	21.4	15.2	7.3	64.3	92.9
2014	14	28.6	3	21.4	15.7	7.1	71.4	100.0
2015	10	31.3	1	10.0	17.0	7.1	60.0	100.0
2016	6	20.7			17.0	11.1	66.7	100.0
2017	8	34.8			17.3	8.3	12.5	100.0
2018	4	28.6			17.5	0.0	25.0	100.0
2019	1	11.1			17.4			100.0 ##
1998-2019	207	33.5	17	8.2	17.4	9.3	64.7	97.1

207 cases diagnosed 1998-2019 are related to a total of 207 patients. Currently, in 51 (24.6 %) of these 207 patients more than one malignancy of any cancer type has been registered. Hereby, groups of 35 / 12 / 4 (16.9 % / 5.8 % / 1.9 %) patients exist having 2 / 3 / 4+ malignancies.

The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be retrieved from the respective headings.

How to interpret:

In 2017, a subgroup of 8 cases has been diagnosed, of which 17.3 % previously and/or concurrently (synchronously) had at least one other malignancy of any cancer type. In 8.3 % of cases, at least one new malignancy has occurred during the follow-up period (all numbers refer to the date of the database export, see cover sheet).

Table 2

Incidence measures by year of diagnosis including DCO cases
 (with respect to registry area expansion from 2.65 to 4.10 m as of 2002,
 and from 4.10 to 4.92 m as of 2007, respectively)

Year of diagnosis	Males n	Females n	Males Inc. raw	Fem. Inc. raw	Males Inc. WS	Fem. Inc. WS	Males Inc. ES	Fem. Inc. ES	Males Inc. BRD-S	Fem. Inc. BRD-S
1998	9	3	0.8	0.3	0.6	0.2	0.7	0.2	0.9	0.2
1999	9	5	0.8	0.4	0.5	0.3	0.7	0.3	0.7	0.5
2000	8	9	0.7	0.7	0.6	0.5	0.7	0.6	0.7	0.8
2001	6	9	0.5	0.7	0.4	0.4	0.5	0.5	0.6	0.6
2002	15	7	0.8	0.4	0.5	0.2	0.7	0.3	0.8	0.3
2003	12	12	0.6	0.6	0.4	0.4	0.6	0.5	0.6	0.6
2004	12	12	0.6	0.6	0.5	0.4	0.6	0.5	0.6	0.5
2005	23	10	1.2	0.5	0.9	0.3	1.1	0.3	1.3	0.4
2006	17	11	0.9	0.5	0.5	0.2	0.7	0.3	0.9	0.5
2007	17	14	0.8	0.6	0.5	0.4	0.7	0.5	0.8	0.5
2008	34	12	1.5	0.5	0.9	0.3	1.2	0.3	1.5	0.4
2009	25	9	1.1	0.4	0.7	0.3	0.9	0.3	1.0	0.3
2010	26	11	1.2	0.5	0.6	0.3	0.9	0.3	1.1	0.4
2011	29	15	1.3	0.6	0.8	0.3	1.0	0.4	1.2	0.5
2012	23	11	1.0	0.5	0.6	0.3	0.8	0.3	0.9	0.4
2013	32	14	1.4	0.6	0.9	0.4	1.1	0.4	1.4	0.5
2014	35	14	1.5	0.6	0.9	0.3	1.2	0.4	1.4	0.5
2015	22	10	0.9	0.4	0.5	0.2	0.7	0.3	0.9	0.3
2016	23	6	1.0	0.2	0.5	0.1	0.7	0.2	0.9	0.2
2017	15	8	0.6	0.3	0.4	0.2	0.5	0.3	0.5	0.3
2018	10	4	0.4	0.2	0.3	0.0	0.3	0.1	0.4	0.1
2019	8	1	0.3	0.0	0.2	0.0	0.2	0.0	0.3	0.0
1998-2019	410	207	0.9	0.5	0.6	0.3	0.8	0.3	0.9	0.4

The computation of the incidence measures includes all cancers, irrespective of first or subsequent malignancy.

Table 3

Age distribution parameters by year of diagnosis (ALL PATIENTS)
(incl. DCO)

Year of diagnosis	Cases n	Std.		Median						
		Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	12	53.6	15.5	24.8	78.0	29.6	47.5	55.6	61.0	73.1
1999	14	60.2	16.6	20.8	79.7	38.4	50.0	62.1	70.0	78.1
2000	17	56.4	25.5	11.0	85.1	15.7	31.4	65.6	76.0	84.1
2001	15	63.0	18.9	26.0	89.0	34.6	49.4	60.3	80.4	86.4
2002	22	62.9	15.3	24.5	87.0	41.4	59.7	67.1	73.7	77.0
2003	24	57.9	19.3	10.9	82.9	27.6	46.9	62.8	74.0	76.8
2004	24	57.2	20.5	8.0	93.7	31.8	44.6	61.8	71.3	76.7
2005	33	59.7	21.3	3.4	81.0	23.3	54.8	68.0	72.9	78.9
2006	28	69.3	13.9	38.7	91.2	50.2	59.9	69.2	79.3	85.7
2007	31	60.0	16.9	10.6	84.0	41.2	47.6	61.5	72.5	78.0
2008	46	64.5	20.6	5.9	91.7	32.8	52.5	72.1	78.8	84.7
2009	34	60.3	18.6	13.8	89.3	36.7	50.1	63.7	69.8	82.4
2010	37	65.1	16.0	25.9	96.7	41.2	57.2	66.0	78.3	84.7
2011	44	64.0	20.1	10.5	99.2	35.2	53.5	70.5	78.8	85.6
2012	34	61.5	16.7	25.2	88.1	37.0	47.0	65.9	74.6	80.5
2013	46	61.4	20.6	16.0	91.8	23.6	49.8	64.9	77.2	82.6
2014	49	62.3	20.2	13.4	91.0	30.4	54.1	63.2	78.2	84.6
2015	32	66.5	17.4	20.7	90.6	49.6	57.8	69.8	79.9	87.1
2016	29	66.5	14.5	19.7	84.9	52.0	57.7	68.8	77.8	82.6
2017	23	61.0	21.1	2.4	88.8	35.0	52.2	63.5	75.6	85.7
2018	14	64.6	17.2	33.9	90.6	35.1	49.8	71.8	76.6	81.2
2019	9	62.1	22.8	26.1	83.0	26.1	47.0	70.2	82.8	83.0
1998-2019	617	62.3	18.8	2.4	99.2	34.6	52.5	66.4	76.0	82.8

Table 3a

Age distribution parameters by year of diagnosis (MALES)
(incl. DCO)

Year of diagnosis	Cases n	Std.		Median						
		Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	9	52.8	17.9	24.8	78.0	24.8	43.1	55.2	62.2	78.0
1999	9	60.1	10.2	38.4	70.0	38.4	59.6	60.9	68.2	70.0
2000	8	51.1	25.8	11.0	85.1	11.0	27.9	59.7	68.7	85.1
2001	6	60.2	22.3	26.0	89.0	26.0	47.3	60.9	77.3	89.0
2002	15	62.4	14.6	35.7	87.0	41.4	47.0	63.6	72.0	77.9
2003	12	58.0	15.1	25.7	74.5	33.4	53.3	61.3	68.2	73.5
2004	12	54.1	16.7	17.8	74.2	31.8	44.6	59.7	65.6	69.3
2005	23	57.1	21.8	3.4	79.2	23.3	45.6	66.1	75.0	76.7
2006	17	68.4	13.9	42.3	91.2	50.2	59.8	68.1	79.3	88.2
2007	17	60.0	19.3	10.6	80.8	23.1	55.9	68.9	72.5	78.0
2008	34	63.9	21.5	5.9	91.7	32.8	55.2	72.1	78.8	82.1
2009	25	59.8	14.8	18.1	89.3	45.5	50.1	63.6	68.7	72.9
2010	26	67.6	12.7	38.2	85.3	53.3	58.5	68.1	79.9	82.8
2011	29	60.6	18.9	10.5	86.0	31.5	51.7	61.7	72.9	81.6
2012	23	61.6	16.6	25.2	81.9	33.1	52.9	66.5	74.7	79.9
2013	32	61.9	20.0	16.0	88.4	23.6	54.6	64.9	77.0	82.6
2014	35	62.1	21.3	13.4	91.0	25.5	54.1	67.5	79.1	83.3
2015	22	65.4	18.4	20.7	90.6	49.6	57.2	70.6	79.0	82.9
2016	23	66.5	15.8	19.7	84.9	52.0	56.6	69.8	78.1	82.6
2017	15	63.0	21.4	2.4	88.8	45.5	52.2	69.2	75.6	85.9
2018	10	58.7	16.6	33.9	81.2	34.5	49.0	60.7	71.8	77.3
2019	8	59.7	23.1	26.1	83.0	26.1	40.9	60.4	82.9	83.0
1998-2019	410	61.7	18.4	2.4	91.7	33.7	52.9	65.6	74.7	81.6

Table 3b

Age distribution parameters by year of diagnosis (FEMALES)
(incl. DCO)

Year of diagnosis	Cases n	Mean	Std. dev.	Min.	Max.	Median				
						10%	25%	50%	75%	90%
1998	3	56.2	3.5	52.7	59.7	52.7	52.7	56.1	59.7	59.7
1999	5	60.5	26.2	20.8	79.7	20.8	46.3	77.7	78.1	79.7
2000	9	61.1	25.8	15.7	84.1	15.7	51.9	75.9	76.6	84.1
2001	9	64.8	17.5	34.6	86.4	34.6	54.0	60.3	80.4	86.4
2002	7	63.8	18.1	24.5	76.3	24.5	60.6	69.3	73.8	76.3
2003	12	57.8	23.4	10.9	82.9	27.6	40.5	66.5	76.8	78.4
2004	12	60.4	24.0	8.0	93.7	34.4	44.8	63.4	76.0	84.8
2005	10	65.5	19.9	10.7	81.0	38.2	68.0	68.7	72.9	80.0
2006	11	70.6	14.4	38.7	84.8	52.0	63.4	73.9	83.4	84.8
2007	14	60.0	14.1	39.9	84.0	41.2	47.6	61.4	71.2	79.7
2008	12	66.2	18.6	29.3	89.2	41.2	52.3	71.3	80.3	84.7
2009	9	61.6	27.6	13.8	88.4	13.8	60.6	69.2	82.4	88.4
2010	11	59.1	21.5	25.9	96.7	36.4	41.2	63.0	71.2	84.7
2011	15	70.6	21.5	20.2	99.2	35.2	57.3	75.4	85.6	91.6
2012	11	61.5	17.5	37.0	88.1	39.7	46.0	63.4	74.6	83.5
2013	14	60.0	22.5	16.3	91.8	26.9	44.9	68.1	77.2	78.1
2014	14	62.8	18.0	32.5	86.7	34.6	48.4	61.9	77.2	84.6
2015	10	68.9	15.5	44.1	87.2	48.2	58.3	67.3	85.6	87.2
2016	6	66.5	9.0	54.7	78.8	54.7	57.7	67.7	72.4	78.8
2017	8	57.3	21.5	19.6	85.7	19.6	44.5	60.9	71.3	85.7
2018	4	79.4	7.6	73.5	90.6	73.5	75.0	76.7	83.8	90.6
2019	1	81.4		81.4	81.4	81.4	81.4	81.4	81.4	81.4
1998-2019	207	63.5	19.6	8.0	99.2	34.6	52.1	68.1	77.6	84.8

Table 4

Age distribution by 5-year age group and sex for period 2007-2019
(incl. DCO)

Age at diagnosis Years	Cases			Males			Females		
	n	%	Cum.%	n	%	Cum.%	n	%	Cum.%
0-4	1	0.2	0.2	1	0.3	0.3			0.0
5-9	2	0.5	0.7	2	0.7	1.0			0.0
10-14	6	1.4	2.1	5	1.7	2.7	1	0.8	0.8
15-19	8	1.9	4.0	5	1.7	4.3	3	2.3	3.1
20-24	6	1.4	5.4	5	1.7	6.0	1	0.8	3.9
25-29	6	1.4	6.8	3	1.0	7.0	3	2.3	6.2
30-34	11	2.6	9.3	8	2.7	9.7	3	2.3	8.5
35-39	13	3.0	12.4	7	2.3	12.0	6	4.7	13.2
40-44	13	3.0	15.4	7	2.3	14.4	6	4.7	17.8
45-49	25	5.8	21.3	17	5.7	20.1	8	6.2	24.0
50-54	29	6.8	28.0	21	7.0	27.1	8	6.2	30.2
55-59	41	9.6	37.6	33	11.0	38.1	8	6.2	36.4
60-64	39	9.1	46.7	25	8.4	46.5	14	10.9	47.3
65-69	40	9.3	56.1	31	10.4	56.9	9	7.0	54.3
70-74	64	15.0	71.0	48	16.1	72.9	16	12.4	66.7
75-79	49	11.4	82.5	33	11.0	83.9	16	12.4	79.1
80-84	43	10.0	92.5	32	10.7	94.6	11	8.5	87.6
85+	32	7.5	100.0	16	5.4	100.0	16	12.4	100.0
All ages	428	100.0		299	100.0		129	100.0	

Table 5

Age-specific incidence, DCO rate and proportion of all cancers for period 2007–2019

Age at diagnosis Years	Males n	Females n	Males Age- spec. incid.	Females Age- spec. incid.	Males DCO rate n=29 %	Females DCO rate n=16 %	Males Prop.all cancers n=143063 %	Females Prop.all cancers n=144724 %
0- 4	1		0.1				0.5	
5- 9	2		0.1				1.8	
10-14	5	1	0.3	0.1			3.8	0.9
15-19	5	3	0.3	0.2			1.7	1.2
20-24	5	1	0.3	0.1			0.9	0.2
25-29	3	3	0.1	0.1			0.3	0.3
30-34	8	3	0.4	0.1	12.5		0.7	0.2
35-39	7	6	0.3	0.3			0.4	0.2
40-44	7	6	0.3	0.3			0.3	0.1
45-49	17	8	0.7	0.3			0.4	0.1
50-54	21	8	0.9	0.3	4.8		0.3	0.1
55-59	33	8	1.7	0.4	3.0	25.0	0.3	0.1
60-64	25	14	1.5	0.8	12.0	7.1	0.2	0.1
65-69	31	9	2.0	0.5	3.2	22.2	0.1	0.1
70-74	48	16	3.4	1.0	6.3	18.8	0.2	0.1
75-79	33	16	3.0	1.2	18.2	18.8	0.1	0.1
80-84	32	11	4.9	1.1	18.8	9.1	0.2	0.1
85+	16	16	3.8	1.7	43.8	25.0	0.2	0.1
All ages	299	129			9.7	12.4	0.2	0.1
Incidence								
Raw			1.0	0.4				
WS			0.6	0.2				
ES			0.8	0.3				
BRD-S			0.9	0.3				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).




Figure 6. Age distribution (males: mean=62.7 yrs, median=67.6 yrs; females: mean=63.9 yrs, median=66.4 yrs) and age-specific incidence.




Figure 6a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).

Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2019, based on the November 2018 submission. <http://www.seer.cancer.gov>.

Table 7a

Standardized incidence ratio (SIR, with 95% confidence limits),
excess absolute risk (EAR) and DCO rate of further malignancies
for period 1998–2019

MALES

Diagnosis	Observed n	Expected n	SIR	CI 95%	CI 95%	EAR	DCO %
C03-C06 Oral cavity	2	0.1	18.2	2.2	65.6 #	18.5	
C12-C13 Hypopharynx	1	0.1	13.7	0.3	76.5	9.1	
C14 ENT cancer	1	0.0	318.9	8.1	1777 #	9.8	
C18 Colon	3	1.2	2.5	0.5	7.3	17.7	
C19-C20 Rectum	1	0.7	1.5	0.0	8.2	3.1	
C22 Liver	1	0.4	2.7	0.1	15.3	6.2	
C32 Larynx	1	0.1	7.7	0.2	42.8	8.5	
C33-C34 Lung	9	1.5	6.1	2.8	11.5 #	73.7	
C37 Thymus	1	0.0	126.4	3.2	704.3 #	9.7	
C43 Malign. melanoma	2	0.6	3.4	0.4	12.2	13.8	
C61 Prostate	3	3.5	0.9	0.2	2.5	-5.1	33.3
C67 Bladder	2	0.6	3.5	0.4	12.7	14.0	
C81 Hodgkin lymphoma	4	0.0	106.2	28.9	271.9 #	38.9	
C82-C85 NHL	6	0.5	10.9	4.0	23.8 #	53.5	
C90 Mult. myeloma	2	0.2	11.9	1.4	43.0 #	18.0	
C91-C96 Leukaemia	3	0.2	15.5	3.2	45.2 #	27.5	
Not observed	0	3.1	0.0	0.0	1.2	-30.4	
All further malignancies	42	12.8	3.3	2.4	4.4 #	286.5	2.4
Patients		382					
Median age at next malignancy (years)		72.0					
Person-years		1020					
Mean observation time (years)		2.7					
Median observation time (years)		1.2					

The occurrence of further specified malignancy is statistically significant.

Table 7b


Standardized incidence ratio (SIR, with 95% confidence limits),
excess absolute risk (EAR) and DCO rate of further malignancies
for period 1998-2019

FEMALES

Diagnosis	Observed n	Expected n	SIR	CI 95%	CI 95%	EAR	DCO %
C21 Anus/canal	1	0.0	34.5	0.9	192.1	16.2	
C22 Liver	2	0.1	29.7	3.6	107.2 #	32.3	
C33-C34 Lung	3	0.4	7.5	1.5	22.0 #	43.5	
C43 Malign. melanoma	2	0.2	9.4	1.1	34.1 #	29.9	
C48 Peritoneal	1	0.0	47.1	1.2	262.7 #	16.4	
C50 Breast	4	1.6	2.4	0.7	6.2	39.5	
C54 Corpus uteri	1	0.3	3.3	0.1	18.6	11.7	
C67 Bladder	1	0.1	9.3	0.2	51.9	14.9	
C73 Thyroid	1	0.1	10.1	0.3	56.5	15.1	
C82-C85 NHL	4	0.2	18.1	4.9	46.3 #	63.2	
Not observed	0	2.5	0.0	0.0	1.5	-41.1	
All further malignancies	20	5.6	3.6	2.2	5.6 #	241.5	
Patients		187					
Median age at next malignancy (years)		71.8					
Person-years		598					
Mean observation time (years)		3.2					
Median observation time (years)		1.0					

The occurrence of further specified malignancy is statistically significant.

Average incidence (Germany 1987 standard population) 2007 - 2019: Males



Average incidence (Germany 1987 standard population) 2007 - 2019: Females





Figure 8a. Map of cancer incidence (german standard population, incl. DCO cases) by county averaged for period 2007 to 2019. According to their individual incidence rates, the counties are displayed in different red and blue hues, being the fine white color attributed to the population mean (males 0.9/100,000 WS N=299, females 0.3/100,000 WS N=129).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,462 female residents (averaged) in the period from 2007 to 2019 a total of 7 women were identified with newly diagnosed mature T/NK-cell I.. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.7/100,000 (german standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.2 and 1.7/100,000.

Standardized incidence ratio (SIR) 2007 - 2019: Males



Standardized incidence ratio (SIR) 2007 - 2019: Females




Figure 8b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2019. According to their individual SIR values, the counties are displayed in different red and blue hues, being the fine white color attributed to the population overall of 1.0 (males N=299, females N=129).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,153 female residents (averaged) in the period from 2007 to 2019 a total of 7 women were identified with newly diagnosed mature T/NK-cell I.. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 1.97. Though, the value of this parameter may vary with an underlying probability of 99% between 0.57 and 4.83, and is therefore not statistically striking.

MORTALITY

Table 9a

Annual cohorts: Incident cancers, follow-up status, proportion of DCO, deaths among the annual cohorts and proportion of available death certificates (with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.92 m as of 2007, respectively)

Year of diagnosis	Incident cases n	Prop. actively followed %	Prop. DCO %	Deaths n	Prop. deaths %	Prop. deaths with death certific. %
1998	12	100.0		8	66.7	100.0
1999	14	100.0		11	78.6	90.9
2000	17	88.2		12	70.6	91.7
2001	15	100.0	6.7	10	66.7	100.0
2002	22	95.5		15	68.2	100.0
2003	24	100.0		15	62.5	100.0
2004	24	91.7	4.2	16	66.7	100.0
2005	33	93.9		22	66.7	100.0
2006	28	92.9		20	71.4	95.0
2007	31	100.0	12.9	24	77.4	95.8
2008	46	97.8	15.2	31	67.4	96.8
2009	34	100.0	2.9	19	55.9	89.5
2010	37	97.3	8.1	25	67.6	96.0
2011	44	97.7	15.9	32	72.7	96.9
2012	34	97.1	8.8	25	73.5	100.0
2013	46	95.7	10.9	32	69.6	90.6
2014	49	98.0	10.2	34	69.4	91.2
2015	32	100.0	9.4	17	53.1	94.1
2016	29	100.0	13.8	21	72.4	95.2
2017	23	100.0	13.0	9	39.1	66.7
2018	14	100.0		4	28.6	25.0
2019	9	88.9		4	44.4	100.0
1998-2019	617	97.2	7.6	406	65.8	94.3

Table 9b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased within the same year of being diagnosed with cancer (incl. DCO)

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002, and from 4.10 to 4.92 m as of 2007, respectively)

Year of diagnosis/ death	Incident cases n	Deaths n	Prop. deaths with death certific. %	Prop. deaths in same year	
				n	%
1998	12	2	50.0	1	8.3
1999	14	5	100.0	1	7.1
2000	17	10	90.0	3	17.6
2001	15	13	100.0	6	40.0
2002	22	9	100.0	3	13.6
2003	24	8	100.0	2	8.3
2004	24	19	100.0	6	25.0
2005	33	12	91.7	4	12.1
2006	28	21	100.0	10	35.7
2007	31	22	100.0	9	29.0
2008	46	29	100.0	16	34.8
2009	34	26	96.2	9	26.5
2010	37	19	100.0	8	21.6
2011	44	33	100.0	16	36.4
2012	34	31	96.8	13	38.2
2013	46	31	100.0	13	28.3
2014	49	33	93.9	14	28.6
2015	32	36	100.0	8	25.0
2016	29	30	100.0	13	44.8
2017	23	22	95.5	6	26.1
2018	14	13	7.7	1	7.1
2019	9	6	33.3	1	11.1
1998–2019	617	430	94.4	163	26.4

Table 9c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates
(incl. DCO)

(with respect to registry area expansion from 2.65 to 4.10 m as of 2002,
and from 4.10 to 4.92 m as of 2007, respectively)

Year of death	Deaths n	Prop. cancer- related %	Prop. non-cancer- related %	Prop. cancer recorded on death certificate %
1998	2	50.0	50.0	100.0
1999	5	80.0	20.0	100.0
2000	10	80.0	20.0	88.9
2001	13	84.6	15.4	92.3
2002	9	100.0		100.0
2003	8	87.5	12.5	87.5
2004	19	89.5	10.5	100.0
2005	12	83.3	16.7	90.9
2006	21	90.5	9.5	100.0
2007	22	90.9	9.1	90.9
2008	29	82.8	17.2	89.7
2009	26	76.9	23.1	88.0
2010	19	73.7	26.3	84.2
2011	33	84.8	15.2	93.9
2012	31	83.9	16.1	90.0
2013	31	83.9	16.1	93.5
2014	33	72.7	27.3	90.3
2015	36	86.1	13.9	91.7
2016	30	96.7	3.3	93.3
2017	22	72.7	27.3	90.5
2018	13	38.5	61.5	100.0
2019	6	50.0	50.0	50.0
1998–2019	430	81.9	18.1	91.9

Table 10a

Medians of age at death according to the grouping in Table 9
MALES

Year of death	Deaths n	Age at death (all causes) Years	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate) Years
1998	1	78.1		78.1	78.1
1999	4	61.0	61.0		61.0
2000	5	70.3	70.3	65.6	70.3
2001	8	63.1	61.6	69.4	61.4
2002	6	76.0	76.0		76.0
2003	4	61.3	61.3		61.3
2004	6	73.7	74.9	70.7	73.7
2005	10	68.7	67.3	70.1	67.3
2006	8	68.5	68.5	74.5	68.5
2007	14	69.5	68.9	75.7	68.9
2008	20	76.7	74.3	87.6	75.4
2009	21	73.5	73.5	74.9	73.5
2010	9	61.9	61.9	54.9	61.9
2011	20	73.8	73.6	82.8	73.7
2012	23	72.8	69.9	85.5	71.8
2013	19	73.4	73.4	79.4	73.4
2014	25	73.7	73.0	75.4	73.0
2015	25	73.8	72.6	84.3	72.9
2016	24	72.1	72.1		70.2
2017	15	75.5	75.2	83.2	78.0
2018	9	71.5	58.9	74.1	
2019	5	50.5	46.1	73.9	47.8
1998-2019	281	72.8	71.7	77.3	72.0

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 10b

Medians of age at death according to the grouping in Table 9
FEMALES

Year of death	Deaths n	Age at death (all causes) Years	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate) Years
1998	1	70.6	70.6		
1999	1	100.0		100.0	100.0
2000	5	76.1	70.3	78.4	70.3
2001	5	77.8	77.8		77.8
2002	3	68.7	68.7		68.7
2003	4	64.7	78.6	50.9	78.6
2004	13	76.9	75.3	76.9	76.9
2005	2	70.4	51.4	89.4	51.4
2006	13	77.9	77.9		77.9
2007	8	69.0	68.7	82.3	68.7
2008	9	84.6	84.7	76.5	84.6
2009	5	83.7	83.7		83.7
2010	10	76.7	73.6	87.6	73.6
2011	13	83.6	81.4	83.6	81.4
2012	8	76.6	74.1	87.4	76.6
2013	12	74.0	72.9	88.4	72.2
2014	8	76.4	63.0	87.0	77.5
2015	11	71.9	72.6	65.1	66.9
2016	6	82.5	78.7	90.3	78.7
2017	7	73.1	73.1		73.1
2018	4	77.9	81.7	70.7	89.1
2019	1	81.9	81.9		
1998-2019	149	76.5	74.4	83.6	75.9

By 2018, Bavarians' life expectancy at birth is estimated at 79.3 years for boys and 83.8 years for girls.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 11a

Mortality measures (cancer-related death) and mortality-incidence-index by year of death
 MALES

Year of death	Deaths n	Mort. raw	MI-Index raw	Mort. WS	MI-Index WS	Mort. ES	MI-Index ES	Mort. BRD-S	MI-Index BRD-S
1998									
1999	4	0.4	0.44	0.2	0.41	0.3	0.41	0.3	0.43
2000	4	0.4	0.50	0.2	0.37	0.3	0.47	0.4	0.54
2001	6	0.5	1.00	0.3	0.87	0.5	0.98	0.6	0.89
2002	6	0.3	0.40	0.2	0.37	0.3	0.42	0.4	0.48
2003	4	0.2	0.33	0.1	0.26	0.2	0.28	0.2	0.34
2004	5	0.3	0.42	0.1	0.26	0.2	0.34	0.3	0.51
2005	9	0.5	0.39	0.2	0.27	0.4	0.34	0.5	0.37
2006	6	0.3	0.35	0.2	0.35	0.2	0.32	0.3	0.32
2007	13	0.6	0.76	0.4	0.75	0.5	0.75	0.6	0.73
2008	16	0.7	0.47	0.3	0.38	0.6	0.46	0.8	0.52
2009	15	0.7	0.60	0.3	0.44	0.5	0.52	0.7	0.62
2010	7	0.3	0.27	0.2	0.29	0.3	0.28	0.3	0.28
2011	18	0.8	0.62	0.4	0.46	0.6	0.54	0.7	0.61
2012	20	0.9	0.87	0.4	0.77	0.6	0.83	0.8	0.90
2013	17	0.7	0.53	0.3	0.37	0.5	0.44	0.7	0.48
2014	20	0.9	0.57	0.4	0.39	0.6	0.49	0.8	0.54
2015	22	0.9	1.00	0.5	1.03	0.7	1.01	0.9	1.00
2016	24	1.0	1.04	0.5	0.94	0.7	1.00	0.9	1.03
2017	9	0.4	0.60	0.1	0.39	0.2	0.48	0.3	0.61
2018	3	0.1	0.30	0.1	0.22	0.1	0.26	0.1	0.26
2019	2	0.1	0.25	0.1	0.40	0.1	0.37	0.1	0.28
1998-2019	230	0.5	0.56	0.3	0.46	0.4	0.52	0.5	0.56

Table 11b

Mortality measures (cancer-related death) and mortality-incidence-index
by year of death
FEMALES

Year of death	Deaths n	Mort. raw	MI-Index raw	Mort. WS	MI-Index WS	Mort. ES	MI-Index ES	Mort. BRD-S	MI-Index BRD-S
1998	1	0.1	0.33	0.0	0.26	0.1	0.27	0.1	0.35
1999									
2000	4	0.3	0.44	0.1	0.27	0.2	0.37	0.3	0.36
2001	5	0.4	0.56	0.2	0.63	0.3	0.56	0.4	0.61
2002	3	0.2	0.43	0.1	0.46	0.1	0.47	0.2	0.44
2003	3	0.2	0.25	0.1	0.15	0.1	0.19	0.1	0.22
2004	12	0.6	1.00	0.3	0.72	0.4	0.83	0.5	0.96
2005	1	0.1	0.10	0.0	0.14	0.1	0.16	0.1	0.14
2006	13	0.6	1.18	0.2	1.04	0.4	1.09	0.5	1.11
2007	7	0.3	0.50	0.1	0.39	0.2	0.42	0.2	0.45
2008	8	0.3	0.67	0.1	0.36	0.2	0.43	0.2	0.49
2009	5	0.2	0.56	0.1	0.19	0.1	0.32	0.1	0.44
2010	7	0.3	0.64	0.1	0.41	0.2	0.50	0.2	0.58
2011	10	0.4	0.67	0.1	0.46	0.2	0.54	0.3	0.70
2012	6	0.3	0.55	0.1	0.38	0.1	0.42	0.2	0.47
2013	9	0.4	0.64	0.2	0.46	0.2	0.53	0.3	0.58
2014	4	0.2	0.29	0.1	0.29	0.1	0.30	0.1	0.26
2015	9	0.4	0.90	0.2	0.91	0.2	0.90	0.3	0.95
2016	5	0.2	0.83	0.1	0.59	0.1	0.64	0.1	0.65
2017	7	0.3	0.88	0.1	0.49	0.2	0.63	0.2	0.77
2018	2	0.1	0.50	0.0	0.60	0.0	0.55	0.0	0.46
2019	1	0.0	1.00	0.0	1.00	0.0	1.00	0.0	1.00
1998-2019	122	0.3	0.59	0.1	0.43	0.2	0.49	0.2	0.53

Table 12

Age distribution of age at death (cancer-related) for period 2007-2019
(incl. multiple malignancies)

Age at death Years	Cases			Males			Females		
	n	%	Cum.%	n	%	Cum.%	n	%	Cum.%
0-4									
5-9									
10-14	1	0.4	0.4	1	0.5	0.5			0.0
15-19	1	0.4	0.8	1	0.5	1.1			0.0
20-24	1	0.4	1.1	1	0.5	1.6			0.0
25-29	2	0.8	1.9	1	0.5	2.2	1	1.3	1.3
30-34	3	1.1	3.0	3	1.6	3.8			1.3
35-39	4	1.5	4.5	2	1.1	4.8	2	2.5	3.8
40-44	3	1.1	5.6	2	1.1	5.9	1	1.3	5.0
45-49	8	3.0	8.6	6	3.2	9.1	2	2.5	7.5
50-54	11	4.1	12.8	8	4.3	13.4	3	3.8	11.3
55-59	22	8.3	21.1	19	10.2	23.7	3	3.8	15.0
60-64	22	8.3	29.3	17	9.1	32.8	5	6.3	21.3
65-69	29	10.9	40.2	19	10.2	43.0	10	12.5	33.8
70-74	50	18.8	59.0	37	19.9	62.9	13	16.3	50.0
75-79	45	16.9	75.9	31	16.7	79.6	14	17.5	67.5
80-84	35	13.2	89.1	24	12.9	92.5	11	13.8	81.3
85+	29	10.9	100.0	14	7.5	100.0	15	18.8	100.0
All ages	266	100.0		186	100.0		80	100.0	

Table 13

Age-specific mortality (cancer-related) and proportion of all cancers
for period 2007-2019
(incl. multiple malignancies)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	Males MI-index	Females Age- spec. mortal.	Females MI-index	Males Prop.all cancers %	Females Prop.all cancers %
0- 4								
5- 9								
10-14	1		0.1	0.20			3.7	
15-19	1		0.1	0.20			2.1	
20-24	1		0.1	0.20			1.5	
25-29	1	1	0.0	0.33	0.0	0.33	1.2	1.1
30-34	3		0.1	0.38			2.3	
35-39	2	2	0.1	0.29	0.1	0.33	0.8	0.5
40-44	2	1	0.1	0.29	0.0	0.17	0.3	0.1
45-49	6	2	0.2	0.35	0.1	0.25	0.4	0.1
50-54	8	3	0.3	0.38	0.1	0.38	0.3	0.1
55-59	19	3	1.0	0.58	0.2	0.38	0.5	0.1
60-64	17	5	1.0	0.68	0.3	0.36	0.3	0.1
65-69	19	10	1.2	0.61	0.6	1.11	0.2	0.2
70-74	37	13	2.6	0.77	0.8	0.81	0.3	0.2
75-79	31	14	2.8	0.94	1.0	0.88	0.3	0.2
80-84	24	11	3.7	0.75	1.1	1.00	0.3	0.1
85+	14	15	3.3	0.88	1.6	0.94	0.2	0.1
All ages	186	80					0.3	0.1
Mortality								
Raw			0.6	0.62	0.3	0.62		
WS			0.3	0.51	0.1	0.43		
ES			0.4	0.57	0.1	0.50		
BRD-S			0.6	0.62	0.2	0.55		
PYLL-70								
per 100,000			4.2		1.3			
ES			3.9		1.1			
AYLL-70			14.0		12.3			

Table 14a

Further malignancies in deaths in period 1998-2019
MALES

Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C03-C06 Oral cavity	1	1.1					1	100.0
C09-C10 Oropharynx	1	1.1	1	100.0				
C12-C13 Hypopharynx	1	1.1					1	100.0
C14 ENT cancer	1	1.1			1	100.0		
C16 Stomach	3	3.2	2	66.7			1	33.3
C17 Small intestine	1	1.1	1	100.0				
C18 Colon	8	8.6	4	50.0	1	12.5	3	37.5
C19-C20 Rectum	2	2.2	1	50.0			1	50.0
C22 Liver	1	1.1			1	100.0		
C33-C34 Lung	10	10.8	3	30.0			7	70.0
C37 Thymus	1	1.1			1	100.0		
C40-C41 Bone	1	1.1	1	100.0				
C43 Malign. melanoma	1	1.1	1	100.0				
C44 Skin others	13	14.0	4	30.8			9	69.2
C46,C49 Soft tissue	1	1.1					1	100.0
C48 Peritoneal	1	1.1	1	100.0				
C61 Prostate	16	17.2	11	68.8	3	18.8	2	12.5
C67 Bladder	3	3.2	2	66.7			1	33.3
C68 Urethra	1	1.1	1	100.0				
C73 Thyroid	2	2.2	2	100.0				
C81 Hodgkin lymphoma	4	4.3	2	50.0			2	50.0
C82-C85 NHL	10	10.8			1	10.0	9	90.0
C90 Mult. myeloma	7	7.5	4	57.1	2	28.6	1	14.3
C91-C96 Leukaemia	3	3.2			1	33.3	2	66.7
All further malignancies	93	100.0	41	44.1	11	11.8	41	44.1

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a further malignancy.

Table 14b

Further malignancies in deaths in period 1998-2019
FEMALES

Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C03-C06 Oral cavity	1	1.9	1	100.0				
C16 Stomach	1	1.9			1	100.0		
C17 Small intestine	1	1.9					1	100.0
C18 Colon	4	7.5	2	50.0			2	50.0
C21 Anus/canal	1	1.9					1	100.0
C22 Liver	2	3.8					2	100.0
C33-C34 Lung	3	5.7					3	100.0
C43 Malign. melanoma	2	3.8	1	50.0			1	50.0
C44 Skin others	7	13.2	1	14.3			6	85.7
C48 Peritoneal	1	1.9			1	100.0		
C50 Breast	15	28.3	13	86.7			2	13.3
C54 Corpus uteri	2	3.8	2	100.0				
C56 Ovary	2	3.8	2	100.0				
C67 Bladder	2	3.8					2	100.0
C82-C85 NHL	7	13.2					7	100.0
C91-C96 Leukaemia	2	3.8	1	50.0			1	50.0
All further malignancies	53	100.0	23	43.4	2	3.8	28	52.8

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a further malignancy.

Table 15

Age-specific mortality (cancer-related) and proportion of all cancers
for period 2007-2019
(First primaries only *)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	Males MI-index	Females Age- spec. mortal.	Females MI-index	Males Prop.all cancers %	Females Prop.all cancers %
0- 4								
5- 9								
10-14	1		0.1	0.20			3.7	
15-19	1		0.1	0.20			2.2	
20-24	1		0.1	0.20			1.7	
25-29	1	1	0.0	0.33	0.0	0.33	1.3	1.2
30-34	3		0.1	0.38			2.4	
35-39	2	2	0.1	0.29	0.1	0.33	0.9	0.6
40-44	2	1	0.1	0.33	0.0	0.25	0.4	0.1
45-49	5	2	0.2	0.33	0.1	0.29	0.4	0.1
50-54	6	3	0.3	0.33	0.1	0.43	0.3	0.1
55-59	15	3	0.8	0.56	0.2	0.50	0.4	0.1
60-64	14	4	0.9	0.74	0.2	0.33	0.3	0.1
65-69	15	7	1.0	0.58	0.4	0.78	0.2	0.1
70-74	28	8	2.0	0.85	0.5	0.89	0.3	0.1
75-79	24	10	2.2	1.20	0.7	0.83	0.3	0.1
80-84	16	9	2.4	0.89	0.9	1.13	0.2	0.1
85+	7	10	1.6	1.00	1.0	0.91	0.1	0.1
All ages	141	60					0.3	0.1
Mortality								
Raw			0.5	0.63	0.2	0.59		
WS			0.2	0.50	0.1	0.40		
ES			0.3	0.56	0.1	0.47		
BRD-S			0.4	0.62	0.1	0.52		
PYLL-70								
per 100,000			3.7		1.2			
ES			3.4		1.0			
AYLL-70			14.8		13.8			

* See corresponding tables with multiple malignancies.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers
for period 2007-2019
(**Single primaries only** *)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	Males MI-index	Females Age- spec. mortal.	Females MI-index	Males Prop.all cancers %	Females Prop.all cancers %
0- 4								
5- 9								
10-14	1		0.1	0.20			3.7	
15-19	1		0.1	0.20			2.2	
20-24	1		0.1	0.20			1.7	
25-29	1	1	0.0	0.33	0.0	0.33	1.3	1.2
30-34	2		0.1	0.25			1.6	
35-39	2	1	0.1	0.29	0.0	0.17	0.9	0.3
40-44	2	1	0.1	0.33	0.0	0.25	0.4	0.1
45-49	5	2	0.2	0.36	0.1	0.29	0.4	0.1
50-54	5	3	0.2	0.28	0.1	0.50	0.2	0.1
55-59	15	2	0.8	0.65	0.1	0.33	0.4	0.1
60-64	10	3	0.6	0.67	0.2	0.33	0.2	0.1
65-69	14	6	0.9	0.58	0.4	0.67	0.2	0.1
70-74	22	7	1.6	0.76	0.4	0.78	0.3	0.1
75-79	20	8	1.8	1.05	0.6	0.73	0.3	0.1
80-84	16	9	2.4	1.14	0.9	1.13	0.3	0.1
85+	3	9	0.7	0.50	0.9	0.82	0.1	0.1
All ages	120	52					0.3	0.1
Mortality								
Raw			0.4	0.59	0.2	0.54		
WS			0.2	0.46	0.1	0.36		
ES			0.3	0.52	0.1	0.42		
BRD-S			0.4	0.58	0.1	0.47		
PYLL-70								
per 100,000			3.4		1.0			
ES			3.1		0.9			
AYLL-70			15.1		13.8			

* See corresponding tables with multiple malignancies.

ICD-10 C84: Mature T/NK-cell lymphomas
 Age distribution and age-specific mortality 2007 - 2019 (Males: 186, Females: 80)





Figure 17. Distribution of age at death (bars; males: mean=66.2 yrs, median=69.4 yrs; females: mean=69.8 yrs, median=72.4 yrs) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at mature T/NK-cell I.-related death (see Table 10) should be considered.

Average mortality (Germany 1987 standard population) 2007 - 2019: Males



Average mortality (Germany 1987 standard population) 2007 - 2019: Females





Figure 18a. Map of cancer mortality (german standard population) by county averaged for period 2007 to 2019. According to their individual mortality rates, the counties are displayed in different red and blue hues, being the fine white color attributed to the population mean (males 0.6/100,000 WS N=186, females 0.2/100,000 WS N=80).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,462 female residents (averaged) in the period from 2007 to 2019 a total of 4 women died from mature T/NK-cell I.. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.4/100,000 (german standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.1 and 1.3/100,000.

Standardized mortality ratio (SMR) 2007 - 2019: Males



Standardized mortality ratio (SMR) 2007 - 2019: Females




Figure 18b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2019. According to their individual SMR values, the counties are displayed in different red and blue hues, being the fine white color attributed to the population overall of 1.0 (males N=186, females N=80).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 67,153 female residents (averaged) in the period from 2007 to 2019 a total of 4 women died from mature T/NK-cell I.. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 1.86. Though, the value of this parameter may vary with an underlying probability of 99% between 0.31 and 5.85, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases, where applicable. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head&neck and lung).

The ratio of mortality and incidence (mortality-to-incidence ratio, **MIR, MI-Index**) is a statistical index that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MIR. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

MCR	Munich Cancer Registry (Tumorregister München)
GEKID	Association of Population-based Cancer Registries in Germany (Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)
SEER	Surveillance, Epidemiology, and End Results (USA)
DCO	Death certificate only
BRD-S	German (FRG) standard population
ES	European standard population (old)
WS	World standard population
SIR	Standardized incidence ratio
CI	Confidence interval
EAR	Excess absolute risk = excess cancer cases (O - E) per 10,000 person-years
PYLL-70	Potential years of life lost prior to age 70 given a person dies before that age
AYLL-70	Average years of life lost prior to age 70 given a person dies before that age
SMR	Standardized mortality ratio
MI-index	Ratio of mortality to incidence, MIR
FRG	Federal Republic of Germany

Recommended Citation

Munich Cancer Registry. ICD-10 C84: Mature T/NK-cell I. - Incidence and Mortality [Internet]. 2021 [updated 2021 Jan 26; cited 2021 Mar 1]. Available from: https://www.tumorregister-muenchen.de/en/facts/base/bC84__E-ICD-10-C84-Mature-T-NK-cell-I.-incidence-and-mortality.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.