Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage

Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

Cancer statistics: Baseline statistics

C01: Base of tongue cancer

Year of diagnosis	1998-2011
Patients	550
Diseases	551
Creation date	04/02/2013
Export date	01/03/2013
Population	4.5 m



http://www.tumorregister-muenchen.de/en/facts/base/base_C01__E.pdf

Global Statements about the statistics on the Internet – Baseline Statistics (grey button ——), Survival (red button ——)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.5 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases**** are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR. The time-delayed acquisition of data and the occasionally high DCO-rates indicate optimizing reserves, among others, because of current financial and legal conditions that hinder the analyses.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, April 2013

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007). Death certificates from 2011 are incorporated into these analyses.
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate. A high proportion of DCO cases (≥5%) in particular cancer types indicate insufficient participation of specific cancer specializations.

Some remarks regarding this cancer type

As a general rule, these few results from the TRM form the basis of sophisticated analyses. For head and neck tumors this is not the case. Therefore the results for head and neck tumors should be interpreted with caution. In part this is due to problems of classification because of limited specific details of locality. Additionally, with advanced tumors in a close topographic location it is often not possible to determine the exact ICD localization of a tumor.

ICD-O-3 codes (topography) used for specifying cancer site

ICD-O-3 Description

C01.9 Base of tongue (posterior third of tongue, root of tongue)

INCIDENCE

Table 1

Patient cohorts by year of diagnosis including DCO cases and multiple primaries, and with proportion of deaths and active follow-up

				Prop.		Prop.
		DCO	Prop.	mult.	Prop.	actively
Year of	Cases #	cases	DCO	primaries	deaths	followed
diagnosis	n	n	%	90	%	%
1998	14			21.4	71.4	100.0
1999	22			36.4	90.9	100.0
2000	13	1	7.7	38.5	76.9	100.0
2001	19	2	10.5	31.6	84.2	100.0
2002	34			17.6	67.6	97.1
2003	42	5	11.9	54.8	83.3	100.0
2004	48	4	8.3	20.8	75.0	93.8
2005	53	3	5.7	30.2	54.7	94.3
2006	51	1	2.0	27.5	60.8	96.1
2007	52	1	1.9	26.9	61.5	82.7 ##
2008	/51	4	7.8	19.6	58.8	84.3
2009	44			25.0	43.2	90.9
2010	54	1	1.9	25.9	46.3	92.6
2011	54	2	3.7	18.5	27.8	68.5 ###
1998-2011	551	24	4.4	27.2	60.1	90.7

[#] The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

^{##} Since 2007 the percentage of actively followed patients sharply declined compared to the previous years. This is a consequence of ambiguous data protection rules that currently forbid cancer registries in Bavaria to obtain the essential life status informations from competent registration offices.

^{###} Please be aware that data of recent annual patient cohorts may not yet be fully processed. Therefore, the presented figures and tables are potentially related to different time periods as pointed out in the respective headlines or legends.

Table 1a

Patient cohorts by year of diagnosis and gender including DCO cases

	Year of	All	Males	Females	Prop. males	
C	diagnosis	n	n	n	%	
	1998	14	10	4	71.4	
	1999	22	18	4	81.8	
	2000	13	9	4	69.2	
	2001	19	12	7	63.2	
	2002	34	27	7	79.4	
	2003	42	37	5	88.1	
	2004	48	33	15	68.8	
	2005	53	40	13	75.5	
	2006	51	38	1/3	74.5	
	2007	52	42	10	80.8	
	2008	51	40	11 /	78.4	
	2009	44	29	15	65.9	
	2010	54	38	16	70.4	
	2011	54	42	12	77.8	
_						
]	L998-2011	551	415	136	75.3	

Table 2

Incidence measures by year of diagnosis and gender including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

			Males	Fem.	Males	Fem.	Males	Fem.	Males	Fem.
Year of	Males	Females	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	10	4	0.9	0.3	0.6	0.2	0.8	0.2	0.9	0.3
1999	18	4	1.6	0.3	1.1	0.2	1.4	0.3	1.5	0.3
2000	9	4	0.8	0.3	0.5	0.2	0.7	0.3	0.9	0.3
2001	12	7	1.0	0.6	0.7	0.3	1.0	0.4	1.1	0.5
2002	27	7	1.4	0.4	1.0	0.2	1.3	0.3	1.4	0.3
2003	37	5	2.0	0.3	1.2	0.1	1.7	0.2	2.0	0.2
2004	33	15 /	1.8	0.8	1.0	0.4	1.4	0.6	1.6	0.7
2005	40	13 <	2.1	0.7	1.4	0.3	1.9	0.5	2.1	0.6
2006	38	13	2.0	0.6	1.2	0.4	1.7	0.5	1.9	0.6
2007	42	10	1.9	0.4	1.2	0.3/	1.7	0.4	1.8	0.4
2008	40	11	1.8	0.5	1.1	0.3	1.5	0.4	1.8	0.4
2009	29	15	1.3	0.6	0.8	0.3	1.1	0.5	1.3	0.5
2010	38	16	1.7	0.7	1.0	0.3	1.4	0.5	1.6	0.6
2011	42	12	1.9	0.5	1.1	0.3	1.5	0.4	1.7	0.4
1998-2011	415	136	1.7	0.5	1.0	0.3	1.4	0.4	1.6	0.5



The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

Table 3

Age distribution parameters by year of diagnosis (All) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	14	57.7	13.2	31.1	78.2	44.9	50.3	58.0	67.5	77.8
1999	22	58.6	9.0	40.4	74.9	48.5	52.2	59.7	64.3	70.5
2000	13	60.3	14.0	35,9	78.3	42.5	47.6	65.5	68.7	78.2
2001	19	63.2	13.6	48.6	92.5	49.0	50.6	61.3	71.2	85.1
2002	34	60.4	8.9	43.8	81.5	50.3	54.8	58.2	65.1	72.1
2003	42	63.1	9.1	45.1	83.3	52.1	56.0	62.7	69.4	76.9
2004	48	63.8	10.0	38.6	93.3	52.0	58.5	62.6	69.7	75.4
2005	53	61.3	13.4	4.1	87.2	50.1	54.8	61.1	65.9	78.5
2006	51	60.3	12.0	19.0	84.8	46.0	52.6	60.1	68.0	76.2
2007	52	57.9	9.7	35.2	76.9	45.0	53.5	58.0	63.7	69.5
2008	51	64.2	10.2	38.3	87.7	51.7	58.7	63.6	71.1	77.4
2009	44	63.4	10.3	36.7	87.3	51.7	57.1	62.9	72.2	75.7
2010	54	64.6	11.4	38.0	92.1	50.1	57.6	65.6	72.7	77.9
2011	54	64.5	11.5	40.0	93.8	49.9	55.9	63.9	72.0	78.3
1998-2011	551	62.1	11.1	4.1	93.8	48.6	55.2	61.5	69.4	76.5

Table 3a

Age distribution parameters by year of diagnosis (MALES)
(incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	10	57.3	10.8	44.9	78.2	46.1	50.3	54.9	60.4	75.2
1999	18	56.9	8.8	40.4	72.5	43.1	50.5	58.4	63.4	70.5
2000	9	59.7	15.8	35.9	78.3	35.9	47.6	59.2	73.3	78.3
2001	12	61.1	12.1	48.6	85.1	49.0	50.4	59.2	68.2	78.7
2002	27	59.7	8.2	43.8	80.2	48.0	54.6	58.2	64.2	70.8
2003	37	62.3	8.9	45.1	83.3	50.6	56.0	62.4	66.7	76.9
2004	33	62.9	8.8	38.6	80.3	54.8	58.9	62.5	68.3	72.5
2005	40	59.5	14.1	4.1	87.1	46.3	52.9	59.8	65.6	78.2
2006	38	61.0	10.5	38.7	84.8	46.7	54.4	60.1	66.8	77.4
2007	42	58.8	9.5	37.1	76.9	45.6	55.5	58.9	64.5	69.5
2008	40	64.4	9.3	38.3	85.9	53.8	59.9	63.2	71.1	77.7
2009	29	62.1	10.2	36.7	77.3	48.0	56.6	61.9	70.7	76.2
2010	38	63.8	12.4	38.0	92.1	49.5	54.0	63.9	72.7	81.3
2011	42	63.9	11.1	40.0	86.5	49.9	54.8	64.1	71.8	77.5
1998-2011	415	61.5	10.7	4.1	92.1	48.6	54.8	61.1	68.4	76.3

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	4	58.8	20.0	31.1	77.8	31.1	45.0	63.2	72.6	77.8
1999	4	66.1	6.6	59.0	74.9	59.0	61.6	65.3	70.6	74.9
2000	4	61.8	10.6	46.0	68.2	46.0	55.7	66.5	67.9	68.2
2001	7	66.8	16,2	50.2	92.5	50.2	50.6	63.0	83.0	92.5
2002	7	63.3	11.4	50.3	81.5	50.3	55.2	58.3	73.9	81.5
2003	5	68.8	9.6	53.1	77.3	53.1	68.5	69.4	75.7	77.3
2004	15	66.0	12.3	48.0	93.3	50.3	57.0	63.1	75.2	81.4
2005	13	66.9	9.3	57.1	87.2	57.2	60.9	64.4	72.9	79.3
2006	13	58.1	15.9	19.0	76.4	45.9	48.1	65.4	70.1	71.3
2007	10	53.9	10.1	35.2	70.5	40.1	47.5	55.1	61.3	66.3
2008	11	63.7	13.4	45.6	87.7	49.7	51.5	64.5	76.3	77.4
2009	15	66.0	10.4	45.8	87.3	54.2	57.8	66.8	72.6	75.1
2010	16	66.5	8.7	47.1	77.9	53.4	60.6	67.8	73.2	77.8
2011	12	66.5	13.2	48.6	93.8	55.9	57.1	62.3	75.1	84.0
1998-2011	136	64.0	12.2	19.0	93.8	48.6	57.1	64.8	72.2	77.8

Age at									
diagnosis	Cases			Males			Females		
Years	n	%	Cum.%	n	%	Cum.%	n	%	Cum.%
0-4	1	0.2	0.2	1	0.2	0.2			0.0
5-9	0	0.0	0.2			0.2			0.0
10-14	0	0.0	0.2			0.2			0.0
15-19	1	0.2	0.4			0.2	1	0.7	0.7
20-24	0	0.0	0.4			0.2			0.7
25-29	0	0.0	0.4			0.2			0.7
30-34	1	0.2	0.5			0.2	1	0.7	1.5
35-39	11	2.0	2.5	10	2.4	2.7/	1	0.7	2.2
40-44	14	2.5	5.1	14	3.4	6.0			2.2
45-49	38	6.9	12.0	24	5.8	11.8	14	10.3	12.5
50-54	68	12.3	24.3	58	14.0	25.8	10	7.4	19.9
55-59	101	18.3	42.6	78	18.8	44.6	23	16.9	36.8
60-64	105	19.1	61.7	87	21.0	65.5	18	13.2	50.0
65-69	81	14.7	76.4	56	13.5	79.0	25	18.4	68.4
70-74	56	10.2	86.6	38	9.2	88.2	18	13.2	81.6
75-79	48	8.7	95.3	33	8.0	96.1	15	11.0	92.6
80-84	14	2.5	97.8	10	2.4	98.6	4	2.9	95.6
85+	12	2.2	100.0	6	1.4	100.0	6	4.4	100.0
All ages	551	100.0		415	100.0		136	100.0	
65-69 70-74 75-79 80-84 85+	81 56 48 14 12	14.7 10.2 8.7 2.5 2.2	76.4 86.6 95.3 97.8	56 38 33 10 6	13.5 9.2 8.0 2.4 1.4	79.0 88.2 96.1 98.6	25 18 15 4 6	18.4 13.2 11.0 2.9 4.4	68. 81. 92. 95.

Included in the statistics are 36.7% multiple primaries in males and 32.4% in females.

Table 5

Age-specific incidence, DCO rate and proportion of all cancers for period 1998-2011

Age at diagnosis Years 0- 4	Males n	Females n	Age- spec. incid.	incid.		Females DCO rate n=6 %		Females Prop.all cancers n=129521 %
5- 9			0.0	0.0				
10-14			0.0	0.0				
15-19		1	0.0	0.1				0.4
20-24			0.0	0.0				
25-29			0.0	0.0				
30-34		1	0.0	0.1				0.1
35-39	10	1	0.5	0.0			0.5	0.0
40-44	14		0.6	0.0			0.5	
45-49	24	14	1.2	0.7	4.2		0.5	0.2
50-54	58	10	3.5	0.6	1.7		0.8	0.1
55-59	78	23	5.0	1.4	3.8	4.3	0.6	0.2
60-64	87	18	5.7	1.1	2.3	1.3	0.5	0.1
65-69	55	25	4.0	1.7	7.3		0.2	0.2
70-74	38	18	3.7	1.5	5.3	_5.6	0.2	0.1
75-79	33	15	4.9	1.5	12.1	13.3	0.2	0.1
80-84	10	4	2.5	0.5	12.1	13.5	0.1	0.0
85+	6	6	2.2	0.8	16.7	33.3	0.1	0.0
037	O	O	۷.۷	0.0	10.7	33.3	0.1	0.0
All ages	414	136			4.3	4.4	0.3	0.1
Incidence								
Raw			1.6	0.5				
WS			1.0					
ES			1.4					
BRD-S								
DICD D				ט ט				
			1.6	0.5				
			1.0	0.5				
			1.0	0.5				
			1.0	0.5				
			1.0	0.5				
			1.0	0.5				
			1.0	0.5				
			1.0	0.5				
			1.0	0.5				
			1.0	0.5				
			1.0	0.5				
			1.0	0.5				
			1.0	0.5				
			1.0	0.5				
			1.0	0.5				
			1.0	0.5				
			1.0	0.5				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

Table 6a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2011

MALES

	Observed	Expected		LCL	UCL		DCO
Diagnosis	n /	n	SIR	95%	95%	EAR	%
C03-C06 Oral cavity	3	0.1	24.7	5.1	72.1 #	36.1	
C09-C10 Oropharynx	2	0.2	12.0	1.5	43.5 #	23.0	
C12-C13 Hypopharynx	2 /	0.1	21.9	2.7	79.2 #	24.0	
C15 Oesophagus	4	0.2	19.1	5.2	48.9 #	47.6	25.0
C18 Colon	2	0.9	2.2	0.3	7.9	13.6	
C19-C20 Rectum	2	0.6	3.3	0.4	11.8	17.4	
C22 Liver	2	0.3	7.1	0.9	25.8	21.6	50.0
C32 Larynx	3	0.1	22.6	4.7	66.1 #	36.0	66.7
C33-C34 Lung	10	1.2	8.0	3.8	14.7 #	109.8	10.0
C61 Prostate	2	2.8	0.7	0.1	2.6	-10.2	
C64 Kidney	3	0.4	8.0	1.7	23.4 #	32.9	
C91-C96 Leukaemia	2	0.1	14.8	1.8	53.3 #	23.4	
Other primaries	5	1.2	4.2	1.4	9.9 #	47.9	
Not observed	0	1.8	0.0	0.0	2.0	-23.1	
All mult. primaries	42	10.1	4.1	3.0	5.6 #	399.9	11.9

Patients	306
Mean age at second malignancy (years)	63.4
Person-years	797
Mean observation time (years)	2.6
Median observation time (years)	1.5


The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries".

Table 6b

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2011 FEMALES

	Observed	Expected		LCL	UCL		DCO
Diagnosis	n	n	SIR	95%	95%	EAR	%
C15 Oesophagus	2	0.0	107.4	13.0	387.8 #	61.2	
C33-C34 Lung	5	0.3	20.0	6.5	46.6 #	146.8	40.0
C50 Breast	2	/ 1.1	1.8	0.2	6.6	28.0	50.0
C53 Cervix uteri	2	0.0	42.4	5.1	153.1 #	60.3	
Other primaries	/5	0.3	15.6	5.1	36.4 #	144.6	
Not observed	0	1.8	0.0	0.0	2.1	-55.0	
All mult. primaries	16	3.5	4.6	2.6	7.4 #	385.8	18.8



The occurrence of second malignancy is statistically significant.

Observed second malignancy with count 1 are pooled in category "Other primaries".






Figure 7. Age distribution and age-specific incidence



C01: Malignant neoplasm of base of tongue Age-specific incidence in international comparison




Figure 7a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).



Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2012, based on the November 2011 submission. http://www.seer.cancer.gov.






Figure 8. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.



Average incidence (world standard population) 2003 - 2008: Males



Average incidence (world standard population) 2003 - 2008: Females





Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 1.2/100,000 WS N=218, females 0.3/100,000 WS N=65). Since cancer data are not available in some counties until 2007, the local incidence rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 1 women were identified with newly diagnosed base of tongue cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.2/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.3/100,000.

Standardized incidence ratio (SIR) 2003 - 2008: Males



Standardized incidence ratio (SIR) 2003 - 2008: Females




Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=218, females N=65). Since cancer data are not available in some counties until 2007, the local SIR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 1 women were identified with newly diagnosed base of tongue cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.50. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 3.73, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts, and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

		Prop.				Prop. deaths
	Incident	actively	Prop.		Prop.	with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	%	%	/ n /	8	8
1998	14	100.0		10	71.4	100.0
1999	22	100.0		20	90.9	90.0
2000	13	100.0	7.7	10	76.9	100.0
2001	19	100.0	10.5	16	84.2	93.8
2002	34	97.1		23	67.6	95.7
2003	42	100.0	11.9	35	83.3	94.3
2004	48	93.8	8.3	36	75.0	100.0
2005	53	94.3	5.7	29	54.7	100.0
2006	51	96.1	2.0	31	60.8	96.8
2007	52	82.7	1.9	32	61.5	100.0
2008	51	84.3	7.8	30	58.8	100.0
2009	44	90.9		19	43.2	100.0
2010	54	92.6	1.9	25	46.3	100.0
2011	54	68.5	3.7	15	27.8	93.3
1998-2011	551	90.7	4.4	331	60.1	97.6

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

			Prop. deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	n	%	n	%
1998	14	12	91.7		
1999	22	11	90.9	1	4.5
2000	13	18	100.0	3	23.1
2001	19	16	93.8	4	21.1
2002	34	20	100.0		
2003	42	24	91.7	10	23.8
2004	48	29	100.0	10	20.8
2005	53	26	100.0	7	13.2
2006	51	47	97.9	12	23.5
2007	52	33	97.0	7	13.5
2008	51	29	96.6	12	23.5
2009	44	35	100.0	4	9.1
2010	54	38	100.0	10	18.5
2011	54	45	100.0	4	7.4
1998-2011	551	383	97.9	84	15.2

Table 10c

Annual cohorts of deaths, proportion of cancer-related and not cancer-related deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

		Prop.	Prop.	Prop. cancer recorded on death	
Year of	Deaths	related	related	certificate	
death	n	%	90	8	
1998	12	75.0	25.0	90.9	
1999	11	72.7	27.3	90.0	
2000	18	66.7	33.3	72.2	
2001	16	75.0	25.0	93.3	
2002	20	95.0	5.0	100.0	
2003	24	62.5	37.5	90.9	
2004	29	93.1	6.9	96.6	
2005	26	96.2	3.8	96.2	
2006	47	89.4	10.6	95.7	
2007	33	81.8	18.2	90.6	
2008	29	82.8	17.2	96.4	
2009	/ 35	85.7	14.3	91.4	
2010	38	89.5	10.5	94.7	
2011	45	84.4	15.6	91.1	
1998-2011	383	84.1	15.9	92.8	

Table 11a $\begin{tabular}{ll} \begin{tabular}{ll} \begin{tabula$

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(not cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	8	60.6	58.9	65.7	59.8
1999	8	61.1	60.1	64.2	60.0
2000	16	66.2	64.3	69.3	63.9
2001	12	61.1	58.0	76.6	59.5
2002	15	62.1	61.7	67.3	62.1
2003	21	61.7	62.4	60.2	62.2
2004	23	65.0	64.1	86.1	64.1
2005	19	64.2	63.9	70.1	63.9
2006	33	66.0	65.4	69.4	65.6
2007	26	61.9	61.6	64.9	62.2
2008	24	63.3	63.6	61.4	62.5
2009	29	63.7	63.4	65.7	63.8
2010	27	66.9	67.1	61.7	66.9
2011	37	66.9	65.3	76.9	65.9
1998-2011	298	64.2	63.7	67.6	63.8




Table 11b Means of age at death according to the grouping in Table 10 FEMALES

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(not cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	4	64.8	58.0	85.2	64.8
1999	3	60.6	55.9	70.0	60.6
2000	2	71.4	71.4		71.4
2001	4	72.0	59.3	84.7	70.4
2002	5	70.3	70.3		70.3
2003	3	63.0	85.4	51.8	68.4
2004	6	64.7	61.4	81.6	64.7
2005	7	61.9	61.9		61.9
2006	14	67.2	67.2		67.2
2007	7	72.0	70.5	74.1	68.7
2008	5	65.2	60.7	71.9	62.2
2009	6	61.7	62.0	60.0	62.0
2010	11 /	71.2	66.8	83.2	67.6
2011	8/	73.2	74.8	68.5	71.5
1998-2011	85	67.6	66.0	73.5	66.6



Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 12a $\begin{tabular}{ll} Mortality measures (cancer-related death) and mortality-incidence-index \\ by year of death \\ \hline MALES \\ \end{tabular}$

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	6	0.5	0.60	0.4	0.60	0.4	0.54	0.5	0.53
1999	6	0.5	0.33	0.3	0.32	0.5	0.33	0.5	0.33
2000	10	0.9	1.11	0.5	1.02	0.8	1.08	1.1	1.28
2001	10	0.9	0.83	0.6	0.83	0.8	0.79	0.8	0.72
2002	14	0.8	0.52	0.5	0.53	0.7	0.54	0.9	0.62
2003	14	0.7	0.39	0.5	0.38	0.6	0.38	0.7	0.36
2004	22	1.2	0.67	0.7	0.68	1.0	0.69	1.2	0.76
2005	18	1.0	0.45	0.6	0.39	0.8	0.42	0.9	0.45
2006	28	1.5	0.74	0.8	0.67	1.2	0.67	1.4	0.71
2007	23	1.0	0.55	0.6	0.52	0.9	0.54	1.0	0.56
2008	21	0.9	0.53	0.6	0.50	0.8	0.50	0.9	0.50
2009	25	1.1	0.86	0.7	0.84	0.9	0.83	1.0	0.83
2010	26	1.2	0.68	0.6	0.67	0.9	0.68	1.1	0.70
2011	32	1.4	0.76	0.8	0.78	1.2	0.80	1.4	0.80
1998-2011	255	1.0	0.62	0.6	0.59	0.9	0.60	1.0	0.63

Table 12b

Mortality measures (cancer-related death) and mortality-incidence-index by year of death

FEMALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	3	0.3	0.75	0.2	1.00	0.2	0.94	0.3	0.90
1999	2	0.2	0.50	0.1	0.56	0.2	0.58	0.2	0.51
2000	2	0.2	0.50	0.1	0.27	0.1	0.31	0.1	0.42
2001	2	0.2	0.29	0.1	0.40	0.2	0.35	0.2	0.37
2002	5	0.3	0.71	0.1	0.60	0.2	0.59	0.2	0.68
2003	1	0.1	0.20	0.0	0.09	0.0	0.11	0.0	0.11
2004	5	0.3	0.33	0.2	0.39	0.2	0.36	0.2	0.36
2005	7	0.4	0.54	0.2	0.64	0.3	0.64	0.3	0.61
2006	14	0.7	1.08	0.4	0.83	0.5	0.95	0.6	0.94
2007	4	0.2	0.40	0.1	0.26	0.1	0.28	0.1	0.30
2008	3	0.1	0.27	0.1	0.29	0.1	0.29	0.1	0.28
2009	5	0.2	0.33	0.1	0.39	0.2	0.36	0.2	0.38
2010	8	0.3	0.50	0.2	0.51	0.3	0.51	0.3	0.50
2011	6	0.3	0.50	0.1	0.36	0.1	0.37	0.2	0.37
1998-2011	67	0.3	0.49	0.1	0.47	0.2	0.48	0.2	0.48

base_C01__E.pdf

Table 13

Age distribution of age at death (cancer-related) for period 1998-2011 (incl. multiple primaries)

Age at									
death	Cases			Males			Females		
Years	n	%	Cum.%	n	96	Cum.%	n	%	Cum.%
35-39	2	0.6	0.6	2	0.8	0.8			0.0
40-44	7	2.2	2.8	/ 7	2.7	3.5			0.0
45-49	23	7.1	9.9	17	6.6	10.2	6	9.0	9.0
50-54	35	10.8	20.7	28	10.9	21.1	7	10.4	19.4
55-59	45	13.9	34.7	39	15.2	36.3	6	9.0	28.4
60-64	70	21.7	56.3	56	21.9	58.2	14	20.9	49.3
65-69	48	14.9	71.2	41	16.0	74.2	7	10.4	59.7
70-74	36	11.1	82.4	22	8.6	82.8	14	20.9	80.6
75-79	33	10.2	92.6	27	10.5	93.4	6	9.0	89.6
80-84	12	3.7	96.3	10	3.9	97.3	2	3.0	92.5
85+	12	3.7	100.0	7	2.7	100.0	5	7.5	100.0
All ages	s 323	100.0		256	100.0		67	100.0	

Included in the statistics are 36.7% multiple primaries in males and 32.4% in females.

Table 14

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2011 (incl. multiple primaries)

Age at			Males Age-		Females Age-		Males Prop.all	Females Prop.all
death	Males	Females	spec.		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0- 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29			0.0		0.0			
30-34			0.0		0.0			
35-39	2		0.1		0.0		0.5	
40-44	7		0.3	0.50	0.0		0.9	
45-49	17	6	0.9	0.71	0.3	0.43	1.1	0.3
50-54	28	7	1.7		0.4	0.70	1.0	0.3
55-59	39	6	2.5		0.4		0.8	0.1
60-64	56	14	3.7		0.9	0.78	0.7	0.3
65-69	41	7	3.0		0.5	0.28	0.4	0.1
70-74	22	14	2.1		1.1	0.78	0.2	0.2
75-79	27	6	4.0		0.6		0.2	0.1
80-84	10	2	2.5		0.3		0.1	0.0
85+	7	5	2.5	1.17	0.7	0.83	0.1	0.0
All ages	256	67					0.4	0.1
Mortality								
Raw			1.0	0.62	0.3			
WS			0.6	0.59	0.1	0.47		
ES			0.9		0.2			
BRD-S			1.0	0.63	0.2	0.48		
PYLL-70								
per 100,000			9.4		2.0			
ES			8.4		1.8			
AYLL-70			11.3		11.4			

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

	Total	Total	Pre	Pre	Syn- chron ±30d	Syn- chron ±30d	Post	Post
Diagnosis	n /	%↓	n	←%	n	~%	n	←%
C00 Lip	1	0.8					1	100.0
C03-C06 Oral cavity	/5	4.0			\ 1	20.0	4	80.0
C09-C10 Oropharynx	10	7.9			1	10.0	9	90.0
C12-C13 Hypopharynx	6	4.8	3	50.0	2	33.3	1	16.7
C15 Oesophagus	12	9.5	5	41.7	1	8.3	6	50.0
C16 Stomach	4 /	3.2	2	50.0			2	50.0
C17 Small intestine	/ 1 -	0.8					1	100.0
C18 Colon	5	4.0	1	20.0	/ 1	20.0	3	60.0
C21 Anus/canal	1	0.8	1	100.0				
C22 Liver	2	1.6					2	100.0
C25 Pancreas	5	4.0	2	40.0			3	60.0
C30-C31 Sinuses	1	0.8	1	100.0				
C32 Larynx	11	8.7	7	63.6	1	9.1	3	27.3
C33-C34 Lung	21	16.7	3	14.3	1	4.8	17	81.0
C38,C45 Mesothelioma	1	0.8					1	100.0
C43 Malign. melanoma	1	0.8	1	100.0				
C44 Skin others	7	5.6	2	28.6			5	71.4
C46,C49 Soft tissue	2	1.6	2	100.0				
C50 Breast	1	0.8	1	100.0				
C61 Prostate	6	4.8	3	50.0			3	50.0
C62 Testis	1	0.8	1	100.0				
C64 Kidney	5	4.0	1	20.0	1 \	20.0	3	60.0
C67 Bladder	5	4.0	3	60.0			2	40.0
C70-C72 CNS cancer	1	0.8	1	100.0				
C73 Thyroid	1	0.8					1	100.0
C76-C79 CUP	8	6.3	6	75.0	2	25.0		
C82-C85 NHL	1	0.8					1	100.0
C91-C96 Leukaemia	1	0.8					1	100.0
All mult. primaries	126	100.0	46	36.5	11	8.7	69	54.8

Multiple primaries with number of cases n<1 are pooled in category "Other primaries".

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 15b

Multiple primaries in deaths in period 1998-2011

FEMALES

						Syn- chron	Syn- chron		
		Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosi	S	n	%↓	n	← %	n	~ %	n	← %
C03-C06	Oral cavity	1 /	3,6					1	100.0
C09-C10	Oropharynx	5	17.9			1	20.0	4	80.0
C12-C13	Hypopharynx	2	7.1			1	50.0	1	50.0
C15	0esophagus	3	10.7	1	33.3			2	66.7
C25	Pancreas	/1	3.6					1	100.0
C32	Larynx	4	14.3	1	25.0	2	50.0	1	25.0
C33-C34	Lung	6	21.4			1	16.7	5	83.3
C44	Skin others	1	3.6	_1	100.0				
C50	Breast	1	3.6					1	100.0
C54	Corpus uteri	1	3.6	1	100.0				
C76-C79	CUP	3	10.7	3	100.0				
All mult	. primaries	28	100.0	7	25.0	5	17.9	16	57.1



ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2011

(Singular primaries only *)

Age at death Years	Males n	Females n	_ /	MI-index	Females Age- spec. mortal.	MI-index	cancers	Females Prop.all cancers
0 4								
0 – 4 5 – 9			0.0		0.0			
5- 9 10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29			0.0		0.0			
30-34			0.0		0.0			
35-39	1		0.0	0.13	0.0		0.3	
40-44	6		0.3		0.0		0.9	
45-49	12	6	0.6	0.60	0.3	0.46	0.9	0.4
50-54	27	5	1.6	0.50	0.3	0.71	1.1	0.2
55-59	31	6	2.0	0.46	0.4	0.29	0.7	0.2
60-64	44	10	2.9	0.65	0.6	0.63	0.7	0.2
65-69	35	4	2.6	0.73	0.3	0.21	0.4	0.1
70-74	15	11/	1.5	0.52	0.9	0.69	0.2	0.2
75-79	17	6	2.5	1.00	0.6	0.43	0.2	0.1
80-84	8	/ 1	2.0		0.1	0.33	0.1	0.0
85+	5	4	1.8	1.25	0.5	2.00	0.1	0.0
All ages	201	53					0.4	0.1
Mortality			0.0	0.60	0 0	0.45		
Raw			0.8		0.2	0.46		
WS			0.5	0.58	0.1	0.44		
ES			0.7 0.8	0.59 0.62	0.2	0.45		
BRD-S			0.0	0.62	0.2	0.45		
PYLL-70								
per 100,000			7.7		1.7			
ES			6.9		1.5			
AYLL-70			11.2		12.3			

^{*} See corresponding tables with multiple primaries.

Table 17

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2011

(Single primaries only *)

Age at death Years	Males n	Females	_ /	MI-index	Females Age- spec. mortal.	MI-index	cancers	Females Prop.all cancers %
0- 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29			0.0		0.0			
30-34			0.0		0.0			
35-39	1		0.0	0.13	0.0		0.3	
40-44	5		0.2	0.50	0.0		0.8	
45-49	8	4	0.4	0.57	0.2	0.33	0.6	0.3
50-54	23	5	1.4	0.48	0.3	0.83	1.0	0.3
55-59	26	5	1.7	0.43	0.3	0.26	0.6	0.2
60-64	37	8	2.4	0.63	0.5	0.57	0.6	0.2
65-69	27	4	2.0		0.3	0.24	0.4	0.1
70-74	11	8	1.1		0.6	0.53	0.1	0.1
75-79	13	5	1.9		0.5	0.36	0.2	0.1
80-84	5	/ 1	1.2		0.1	1.00	0.1	0.0
85+	5	3	1.8	1.25	0.4	1.50	0.1	0.0
All ages	161	43					0.3	0.1
1111 0300		\ -					\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	0.1
Mortality								
Raw			0.6	0.54	0.2	0.42		
WS			0.4	0.52	0.1	0.39		
ES			0.5	0.53	0.1	0.40		
BRD-S			0.6	0.55	0.1	0.40		
PYLL-70								
per 100,000			6.3		1.4			
ES ES			5.6		1.2			
AYLL-70			11.2		11.9			
					==.,			

^{*} See corresponding tables with multiple primaries.





Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at base of tongue cancer-related death (see Table 10) should be considered.



Average mortality (world standard population) 2003 - 2008: Males



Average mortality (world standard population) 2003 - 2008: Females




Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2003 to 2008. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.6/100,000 WS N=121, females 0.1/100,000 WS N=33). Since cancer data are not available in some counties until 2007, the local mortality rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 0 women died from base of tongue cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.0/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.4/100,000.

Standardized mortality ratio (SMR) 2003 - 2008: Males




Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=121, females N=33). Since cancer data are not available in some counties until 2007, the local SMR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 0 women died from base of tongue cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.00. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 5.32, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the tumor-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

base_C01__E.pdf

Shortcuts

AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

BRD-S German standard population

DCO Death certificate only EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

ES European standard population (old) FRG Federal Republic of Germany

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

LCL Lower confidence limit

MI-index Ratio between mortality and incidence

MCR Munich Cancer Registry (Tumorregister München)

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age

SEER Surveillance, Epidemiology, and End Results (USA)

SIR Standardized incidence ratio
SMR Standardized mortality ratio
UCL Upper confidence limit
WS World standard population

Recommended Citation

Munich Cancer Registry. Baseline statistics C01: Base of tongue cancer [Internet]. 2013 [updated 2013 Apr 2; cited 2013 Jun 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/base/base_C01__E.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

Fig./Tbl	l.	Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	3
1a	Gender distribution by year of diagnosis	4
2	Incidence by year of diagnosis	5
3	Age distribution parameters by year of diagnosis	6
4	Age distribution by 5-year age group and gender	8
5	Age-specific incidence and DCO rate	9
6	Standardized incidence ratio of second primaries	10
7	Age distribution and age-specific incidence (chart)	12
7a	Age-specific incidence internationally (chart)	13
8	Cumulative follow-up years (chart)	14
9a	Map of cancer incidence (WS) by county (chart)	15
9b	Standardized incidence ratio (SIR) by county (chart)	16
10a	Pts incident cohorts and mortality / yr	17
10b	Incidence and mortality by year of diagnosis	18
10c	Cancer-related deaths, death certification available / yr	19
11	Means of age at death / yr	20
12	Mortality by year of death	22
13	Distribution of age at death	23
14	Age-specific mortality	24
15	Multiple primaries in deaths	25
16	Age-specific mortality (first primaries)	27
17	Age-specific mortality (single primaries)	28
18	Age distribution and age-specific mortality (chart)	29
19a	Map of cancer mortality (WS) by county (chart)	30
19b	Standardized mortality ratio (SMR) by county (chart)	31