Munich Cancer Registry



- Survival
- Selection Matrix
- ▶ Homepage

Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

Cancer statistics: Baseline statistics

Year of diagnosis	1998-2011
Patients	1094
Diseases	1097
Creation date	04/02/2013
Export date	01/03/2013
Population (females)	2.3 m



C51: Vulva cancer

Global Statements about the statistics on the Internet -

Baseline Statistics (grey button ___), Survival (red button ___)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.5 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases^{###} are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR. The time-delayed acquisition of data and the occasionally high DCO-rates indicate optimizing reserves, among others, because of current financial and legal conditions that hinder the analyses.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, April 2013

- [#] Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007). Death certificates from 2011 are incorporated into these analyses.
- ^{##} Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- ### DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate. A high proportion of DCO cases (≥5%) in particular cancer types indicate insufficient participation of specific cancer specializations.

INCIDENCE

Table 1

Patient cohorts by year of diagnosis including DCO cases and multiple primaries, and with proportion of deaths and active follow-up

				Prop.		Prop.
		DCO	Prop.	mult.	Prop.	actively
Year of	Cases #	cases	DCO	primaries	deaths	followed
diagnosis	n	n	olo	90	90	<u>0</u>
1998	42	2	4.8	26.2	78.6	97.6
1999	37	3	8.1	8.1	78.4	100.0
2000	40	5	12.5	22.5	67.5	100.0
2001	36	4	11.1	13.9	63.9	100.0
2002	69	5	7.2	15.9	60.9	100.0
2003	75	5	6.7	30.7	68.0	97.3
2004	73	5	6.8	31.5	65.8	94.5
2005	90	6	6.7	27.8	55.6	97.8
2006	86	1	1.2	23.3	44.2	95.3
2007	107	4	3.7	19.6	43.9	83.2 ##
2008	110	4	3.6	23.6	46.4	65.5
2009	100	3	3.0	20.0	35.0	68.0
2010	132	4	3.0	23.5	34.8	96.2
2011	100	6	6.0	19.0	27.0	80.0 ###
1998-2011	1097	57	5.2	22.5	49.9	88.5

The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

- ## Since 2007 the percentage of actively followed patients sharply declined compared to the previous years. This is a consequence of ambiguous data protection rules that currently forbid cancer registries in Bavaria to obtain the essential life status informations from competent registration offices.
- ### Please be aware that data of recent annual patient cohorts may not yet be fully processed. Therefore, the presented figures and tables are potentially related to different time periods as pointed out in the respective headlines or legends.

Incidence measures by year of diagnosis and gender including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

Year of	Cases	Incidence	Incidence WS	Incidence ES	Incidence BRD-S	
diagnosis	n	raw	WS	E2	BRD-5	
1	10					
1998	42	3.6	1.5	2.3	3.0	
1999	37	3.1	1.2	1.9	2.5	
2000	40	3.3	1.6	2.2	2.8	
2001	36	3.0	1.3	2.0	2.5	
2002	69	3.5	1.5	2.2	2.9	
2003	75	3.8	1.4	2.2	2.9	
2004	73	3.7	1.5	2.3	3.0	
2005	90	4.5	1.6	2.5	3.5	
2006	86	4.3	1.7	2.5	3.3	
2007	107	4.6	1.9	2.9	3.7	
2008	110	4.7	1.9	2.8	3.6	
2009	100	4.3	1.7	2.5	3.3	
2010	132	5.6	2.2	3.3	4.3	
2011	100	4.3	1.6	2.4	3.2	
1998-2011	1097	4.2	1.7	2.5	3.3	

The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

Year of	Cases		Std.					Median			
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%	
1998	42	72.9	13.4	36.3	93.3	52.1	68.6	75.1	80.9	87.2	
1999	37	71.8	15.5	34.3	94.0	48.7	62.4	75.6	84.8	88.0	
2000	40	68.1	17.6	26,1	92.3	42.4	58.1	73.2	81.8	88.2	
2001	36	69.9	13.5	33.3	89.1	49.6	60.6	70.8	80.1	86.2	
2002	69	70.7	16.1	26.9	96.1	39.3	64.9	74.6	81.5	89.8	
2003	75	74.1	14.9	34.3	94.8	50.2	63.8	78.9	85.7	91.0	
2004	73	72.3	14.6	32.7	95.5	51.7	63.4	76.3	83.8	88.6	
2005	90	75.1	13.0	40.8	96.9	54.2	68.7	78.8	83.7	89.6	
2006	86	73.0	14.4	34.6	97.2	47.7	65.3	74.0	84.6	88.7	
2007	107	71.0	14.8	29.4	96.7	47.7	60.7	74.6	82.6	86.7	
2008	110	73.2	14.0	28.7	96.9	52.8	65.9	74.9	83.5	89.1	
2009	100	73.0	15.1	24.9	100	51.3	64.5	76.7	84.7	88.3	
2010	132	72.3	13.3	36.5	95.1	52.1	65.8	73.5	82.2	88.1	
2011	100	73.3	14.2	30.3	98.5	51.2	69.0	75.8	83.3	88.8	
1998-2011	1097	72.5	14.4	24.9	100	50.1	64.5	75.4	83.0	88.7	

Age distribution parameters by year of diagnosis (incl. DCO)

Age distribution by 5-year age group for period 1998-2011 (incl. DCO)

Age at diagnosis	Cases	00	Charme &		
Years	n	10	Cum.%		
20-24	1	0.1	0.1		
25-29	4	0.4	0.5		
30-34	11	1.0	1.5		
35-39	18	1.6	3.1		
40-44	25	2.3	5.4		
45-49	50	4.6	9.9		
50-54	40	3.6	13.6		
55-59	66	6.0	19.6		
60-64	73	6.7	26.3		
65-69	109	9.9	36.2		
70-74	134	12.2	48.4		
75-79	168	15.3	63.7		
80-84	180	16.4	80.1		
85+	218	19.9	100.0		
All ages	1097	100.0			

Included in the statistics are 29.1% multiple primaries.

Prop. all DCO rate Age at cancers diagnosis Cases Age-spec. n=57 n=129521 Years incidence % ° n 0- 4 0.0 5-9 0.0 10 - 140.0 15-19 0.0 20-24 1 0.1 0.2 25-29 4 0.2 0.4 0.6 30-34 11 0.6 35-39 18 0.9 0.6 40 - 4425 1.2 0.5 0.7 45-49 50 2.6 50-54 40 2.3 0.4 55-59 3.0 0.6 66 4.0 60-64 73 0.5 4.6 65-69 7.3 0.9 0.7 109 70-74 10.9 3.0 134 0.9 16.9 75-79 3.6 168 1.1 22.6 80-84 5.0 180 1.3 85+ 218 29.4 16.1 1.5 5.2 All ages 1097 0.8 Incidence 4.2 Raw 1.7 WS 2.5 ES BRD-S 3.3

Age-specific incidence, DCO rate and proportion of all cancers for period 1998-2011

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2011

	Observed E	xpected		LCL	UCL		DCO
Diagnosis	n	n	SIR	95%	95%	EAR	00
C18 Colon	9	3.6	2.5	1.1	4.8 #	21.8	33.3
C21 Anus/canal	6	0.2	37.3	13.7	81.1 #	23.5	
C33-C34 Lung	5	2.0	2.5	0.8	5.8	12.1	
C43 Malign. melanoma	. 4	1.0	4.1	1.1	10.5 #	12.2	25.0
C50 Breast	14	8.5	1.6	0.9	2.8	22.1	14.3
C51 Vulva	3	0.3	8.8	1.8	25.6 #	10.7	
C52 Vagina	2	0.1	29.5	3.6	106.7 #	7.8	
C53 Cervix uteri	3	0.4	7.5	1.6	22.1 #	10.5	
C54 Corpus uteri	3	1.6	1.9	0.4	5.6	5.7	
C56 Ovary	5	1.3	3.8	1.2	8.9 #	14.9	40.0
C70-C72 CNS cancer	2	0.4	4.6	0.6	16.7	6.3	50.0
C76-C79 CUP	2	0.6	3.2	0.4	11.4	5.5	
C82-C85 NHL	2	1.2	1.6	0.2	5.9	3.1	
Other primaries	10	6.0	1.7	0.8	3.1	16.2	10.0
Not observed	0	3.9	0.0	0.0	0.9 #	-15.6	
All mult. primaries	70	31.1	2.3	1.8	2.8 #	156.6	14.3

Patients	762
Mean age at second malignancy (years)	73.6
Person-years	2485
Mean observation time (years)	3.3
Median observation time (years)	2.2

The occurrence of second malignancy is statistically significant.

Observed second malignancy with count 1 are pooled in category "Other primaries".




Figure 7. Age distribution and age-specific incidence







Figure 7a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).



Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2012, based on the November 2011 submission. http://www.seer.cancer.gov.




C51: Malignant neoplasm of vulva

Figure 8. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.





Average incidence (world standard population) 2003 - 2008

Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (1.7/100,000 WS N=518). Since cancer data are not available in some counties until 2007, the local incidence rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 12 women were identified with newly diagnosed vulva cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.9/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.4 and 2.1/100,000.

Standardized incidence ratio (SIR) 2003 - 2008




Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (N=518). Since cancer data are not available in some counties until 2007, the local SIR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 12 women were identified with newly diagnosed vulva cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.81. Though, the value of this parameter may vary with an underlying probability of 99% between 0.33 and 1.63, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts, and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

		Prop.				Prop. deaths
	Incident	actively	Prop.		Prop.	with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	90	00	n	00	00
1998	42	97.6	4.8	33	78.6	87.9
1999	37	100.0	8.1	29	78.4	89.7
2000	40	100.0	12.5	27	67.5	85.2
2001	36	100.0	11.1	23	63.9	100.0
2002	69	100.0	7.2	42	60.9	95.2
2003	75	97.3	6.7	51	68.0	98.0
2004	73	94.5	6.8	48	65.8	97.9
2005	90	97.8	6.7	50	55.6	98.0
2006	86	95.3	1.2	38	44.2	97.4
2007	107	83.2	3.7	47	43.9	100.0
2008	110	65.5	3.6	51	46.4	96.1
2009	100	68.0	3.0	35	35.0	100.0
2010	132	96.2	3.0	46	34.8	93.5
2011	100	80.0	6.0	27	27.0	100.0
1998-2011	1097	88.5	5.2	547	49.9	96.0




Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

			Prop.		
			deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	n	00	n	00
1998	42	23	78.3	5	11.9
1999	37	26	80.8	6	16.2
2000	40	31	96.8	4	10.0
2001	36	28	89.3	4	11.1
2002	69	40	95.0	12	17.4
2003	75	41	97.6	11	14.7
2004	73	52	100.0	11	15.1
2005	90	50	100.0	12	13.3
2006	86	48	95.8	5	5.8
2007	107	60	95.0	14	13.1
2008	110	66	100.0	17	15.5
2009	100	59	98.3	7	7.0
2010	132	66	100.0	17	12.9
2011	100	82	98.8	15	15.0
1998-2011	1097	672	96.4	140	12.8

Munich Cancer Registry

Table 10c

Annual cohorts of deaths, proportion of cancer-related and not cancerrelated deaths, and cancer recorded on death certificates (incl. DCO) (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

		Prop. cancer-	Prop. not cancer-	Prop. cancer recorded on death	
Year of	Deaths	related	related	certificate	
death	n	00	8	ક	
1998	23	69.6	30.4	83.3	
1999	26	61.5	38.5	95.2	
2000	31	61.3	38.7	76.7	
2001	28	60.7	39.3	80.0	
2002	40	72.5	27.5	73.7	
2003	41	75.6	24.4	80.0	
2004	52	76.9	23.1	86.5	
2005	50	66.0	34.0	76.0	
2006	48	60.4	39.6	73.9	
2007	60	70.0	30.0	80.7	
2008	66	68.2	31.8	72.7	
2009	59	62.7	37.3	77.6	
2010	66	63.6	36.4	74.2	
2011	82	68.3	31.7	77.8	
1998-2011	672	67.3	32.7	78.1	

Munich Cancer Registry

Table	11

Means of age at death according to the grouping in Table 10

		Ngo at	Ngo at	Ago at	Age at death
		Age at	Age at	Age at	
		death	death	death	(according
		(all	(cancer-	(not cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	23	79.3	76.7	85.3	80.0
1999	26	77.1	73.9	82.3	75.5
2000	31	82.5	82.9	81.8	83.3
2001	28	78.3	77.6	79.4	78.4
2002	40	81.3	80.0	84.6	80.5
2003	41	77.7	74.6	87.3	75.0
2004	52	81.6	80.6	84.9	81.0
2005	50	80.9	78.1	86.1	79.7
2006	48	79.6	77.0	83.5	78.0
2007	60	81.3	79.1	86.3	79.9
2008	66	81.5	79.3	86.4	79.4
2009	59	81.4	80.0	83.8	80.8
2010	66	80.0	77.9	83.8	78.5
2011	82	80.6	78.4	85.2	78.6
1998-2011	672	80.5	78.5	84.5	79.2

By 2010, life expectancy for a newborn male in Germany is 77.5 years compared with 82.6 years for his female counterpart.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

			b	y year	of death				
Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	16	1.4	0.38	0.5	0.33	0.8	0.34	1.0	0.34
1999	16	1.3	0.43	0.6	0.45	0.8	0.45	1.1	0.44
2000	19	1.6	0.48	0.4	0.26	0.7	0.33	1.1	0.38
2001	17	1.4	0.47	0.5	0.36	0.8	0.40	1.1	0.45
2002	29	1.5	0.42	0.4	0.28	0.7	0.32	1.1	0.37
2003	31	1.6	0.41	0.6	0.38	0.9	0.40	1.2	0.41
2004	40	2.0	0.55	0.6	0.38	0.9	0.41	1.3	0.45
2005	33	1.7	0.37	0.5	0.32	0.8	0.33	1.2	0.33
2006	29	1.4	0.34	0.5	0.26	0.8	0.30	1.2	0.36
2007	42	1.8	0.39	0.5	0.27	0.9	0.30	1.3	0.34
2008	45	1.9	0.41	0.6	0.31	0.9	0.33	1.3	0.35
2009	37	1.6	0.37	0.5	0.28	0.8	0.30	1.1	0.33
2010	42	1.8	0.32	0.5	0.24	0.9	0.27	1.3	0.30
2011	56	2.4	0.56	0.8	0.51	1.2	0.51	1.6	0.50
1998-2011	452	1.7	0.41	0.5	0.32	0.9	0.35	1.2	0.37

Mortality measures (cancer-related death) and mortality-incidence-index

Munich Cancer Registry

Age distribution of age at death (cancer-related) for period 1998-2011 (incl. multiple primaries)

Age at death Years	Cases	00	Cum.%	
30-34	3	0.7	0.7	
35-39	2	0.4	1.1	
40-44	4	0.9	2.0	
45-49	3	0.7	2.7	
50-54	6	1.3	4.0	
55-59	13	2.9	6.9	
60-64	23	5.1	11.9	
65-69	46	10.2	22.1	
70-74	47	10.4	32.5	
75-79	62	13.7	46.2	
80-84	98	21.7	67.9	
85+	145	32.1	100.0	
All ages	452	100.0		

Included in the statistics are 29.1% multiple primaries.

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2011 (incl. multiple primaries)

Age at death	Cases /	Age-spec.		Prop. all cancers	
Years	n n	nortality	MI-index	90	
0- 4 5- 9 10-14 15-19 20-24		0.0 0.0 0.0 0.0 0.0 0.0			
25-29		0.0			
30-34	3	0.2	0.27	1.5	
35-39	2	0.1	0.11	0.4	
40 - 44	4	0.2	0.16	0.4	
45-49	3	0.2	0.06	0.2	
50-54	6	0.3	0.15	0.2	
55-59	13	0.8	0.20	0.3	
60-64	23	1.4	0.32	0.4	
65-69	46	3.1	0.42	0.7	
70-74	47	3.8	0.35	0.6	
75-79 80-84	62 98	6.2 12.3	0.37 0.54	0.7 1.0	
85+	145	12.5	0.54	1.3	
01+	THD	19.5	0.07	1.5	
All ages	452			0.7	
Mortality					
Raw		1.7	0.41		
WS		0.5	0.32		
ES		0.9	0.35		
BRD-S		1.2	0.37		
PYLL-70		1 0			
per 100,000		4.0			
ES		3.4			
AYLL-70		9.1			

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Multiple primaries in deaths in period 1998-2011

					Syn- chron	Syn- chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n	% ↓	n	\$→	n	olo	n	00
C00 Lip	1	0.5	1	100.0				
C03-C06 Oral cavity	1	0.5					1	100.0
C12-C13 Hypopharynx	/1	0.5	1	100.0				
C15 Oesophagus	/ 1	0.5					1	100.0
C16 Stomach	6	3.2	1	16.7			5	83.3
C18 Colon	14	7.4	4	28.6	1	7.1	9	64.3
C19-C20 Rectum	8	4.3	5	62.5	1	12.5	2	25.0
C21 Anus/canal	5	2.7	3	60.0			2	40.0
C22 Liver	3	1.6					3	100.0
C23-C24 Bile	1	0.5					1	100.0
C25 Pancreas	2	1.1	1	50.0			1	50.0
C32 Larynx	2	1.1	2	100.0				
C33-C34 Lung	11	5.9	2	18.2	1	9.1	8	72.7
C43 Malign. melanoma	8	4.3	2	25.0	1	12.5	5	62.5
C44 Skin others	9	4.8	2	22.2	3	33.3	4	44.4
C46,C49 Soft tissue	1	0.5	1	100.0				
C48 Peritoneal	1	0.5	1	100.0				
C50 Breast	26	13.8	17	65.4	1	3.8	8	30.8
C51 Vulva	5	2.7			1	20.0	4	80.0
C52 Vagina	4	2.1	1	25.0	1	25.0	2	50.0
C53 Cervix uteri	17	9.0	12	70.6	3	17.6	2	11.8
C54 Corpus uteri	15	8.0	14	93.3	1	6.7		
C55,C57 Fem. genitals un	2	1.1	1	50.0			1	50.0
C56 Ovary	11	5.9	6	54.5	2	18.2	3	27.3
C64 Kidney	3	1.6	2	66.7			1	33.3
C67 Bladder	7	3.7	2	28.6			5	71.4
C68 Urethra	2	1.1	1	50.0			1	50.0
C70-C72 CNS cancer	3	1.6	2	66.7			1	33.3
C76-C79 CUP	3	1.6			2	66.7	1	33.3
C81 Hodgkin lymphoma	1	0.5	1	100.0				
C82-C85 NHL	7	3.7	1	14.3			6	85.7
C90 Mult. myeloma	2	1.1					2	100.0
C91-C96 Leukaemia	5	2.7	1	20.0			4	80.0
All mult. primaries	188	100.0	87	46.3	18	9.6	83	44.1

Multiple primaries with number of cases n<1 are pooled in category "Other primaries".

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2011 (Singular primaries only *)

Age at death Years		ge-spec. ortality	MI-index	Prop. all cancers %	
0- 4 5- 9 10-14 15-19 20-24		0.0 0.0 0.0 0.0 0.0 0.0			
25-29		0.0			
30-34	2 2	0.1	0.20	1.1	
35-39 40-44	2	0.1	0.12 0.10	0.5 0.2	
45-49	2	0.1	0.04	0.1	
50-54	6	0.3	0.16	0.3	
55-59	11	0.7	0.19	0.3	
60-64	17	1.1	0.29	0.4	
65-69	33	2.2	0.38	0.6	
70-74 75-79	39 49	3.2 4.9	0.36 0.35	0.6	
80-84	49 76	4.9 9.6	0.53	1.0	
85+	118	15.9	0.65	1.3	
All ages	357			0.7	
Mortality					
Raw		1.4	0.39		
WS		0.4	0.30		
ES		0.7	0.33		
BRD-S		1.0	0.35		
PYLL-70					
per 100,000		3.1			
ES		2.6			
AYLL-70		9.2			

* See corresponding tables with multiple primaries.

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2011 (Single primaries only *)

Age at				Prop. all	
death		ge-spec.		cancers	
Years	n mo	ortality	MI-index	00	
0- 4 5- 9 10-14		0.0 0.0 0.0			
15-19		0.0			
20-24		0.0			
25-29		0.0			
30-34	1	0.1	0.10	0.6	
35-39	2	0.1	0.12	0.5	
40-44	2	0.1	0.11	0.3	
45-49	2	0.1	0.05	0.1	
50-54	6	0.3	0.17	0.3	
55-59	8	0.5	0.15	0.3	
60-64	16	1.0	0.32	0.4	
65-69	23	1.5	0.29	0.5	
70-74	35	2.8	0.34	0.6	
75-79	35	3.5	0.27	0.6	
80-84	62	7.8	0.47	1.0	
85+	106	14.3	0.60	1.3	
All ages	298			0.7	
Mortality					
Raw		1.1	0.35		
WS		0.3	0.27		
ES		0.6	0.29		
BRD-S		0.8	0.31		
PYLL-70		2 6			
per 100,000		2.6			
ES DVIII 70		2.2			
AYLL-70		9.8			

* See corresponding tables with multiple primaries.





Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at vulva cancer-related death (see Table 10) should be considered.



Average mortality (world standard population) 2003 - 2008

Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2003 to 2008. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (0.5/100,000 WS N=207). Since cancer data are not available in some counties until 2007, the local mortality rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 7 women died from vulva cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.5/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.1 and 1.5/100,000.

Standardized mortality ratio (SMR) 2003 - 2008




Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (N=207). Since cancer data are not available in some counties until 2007, the local SMR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 7 women died from vulva cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 1.23. Though, the value of this parameter may vary with an underlying probability of 99% between 0.36 and 3.01, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the tumor-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

AYLL-70 BRD-S DCO	Average years of life lost prior to age 70 given a person dies before that age German standard population Death certificate only
EAR	Excess absolute risk
	= excess cancer cases (O - E) per 10,000 person-years
ES	European standard population (old)
FRG	Federal Republic of Germany
GEKID	Association of Population-based Cancer Registries in Germany
	(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)
LCL	Lower confidence limit
MI-index	Ratio between mortality and incidence
MCR	Munich Cancer Registry (Tumorregister München)
PYLL-70	Potential years of life lost prior to age 70 given a person dies before that age
SEER	Surveillance, Epidemiology, and End Results (USA)
SIR	Standardized incidence ratio
SMR	Standardized mortality ratio
UCL	Upper confidence limit
WS	World standard population

Recommended Citation

Munich Cancer Registry. Baseline statistics C51: Vulva cancer [Internet]. 2013 [updated 2013 Apr 2; cited 2013 Jun 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/base/base_C51_E.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Munich Cancer Registry

Index of figures and tables

Fig./Tb	l.	Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	3
2	Incidence by year of diagnosis	4
3	Age distribution parameters by year of diagnosis	5
4	Age distribution by 5-year age group	6
5	Age-specific incidence and DCO rate	7
6	Standardized incidence ratio of second primaries	8
7	Age distribution and age-specific incidence (chart)	9
7a	Age-specific incidence internationally (chart)	10
8	Cumulative follow-up years (chart)	11
9a	Map of cancer incidence (WS) by county (chart)	12
9b	Standardized incidence ratio (SIR) by county (chart)	13
10a	Pts incident cohorts and mortality / yr	14
10b	Incidence and mortality by year of diagnosis	15
10c	Cancer-related deaths, death certification available / yr	16
11	Means of age at death / yr	17
12	Mortality by year of death	18
13	Distribution of age at death	19
14	Age-specific mortality	20
15	Multiple primaries in deaths	21
16	Age-specific mortality (first primaries)	22
17	Age-specific mortality (single primaries)	23
18	Age distribution and age-specific mortality (chart)	24
19a	Map of cancer mortality (WS) by county (chart)	25
19b	Standardized mortality ratio (SMR) by county (chart)	26

