Munich Cancer Registry



- Survival
- Selection Matrix
- ► Homepage

Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

Cancer statistics: Baseline statistics

D05: Breast cancer i.s. (women)

Year of diagnosis	1998-2011
Patients	3756
Diseases	3812
Creation date	04/02/2013
Export date	01/03/2013
Population (females)	2.3 m



http://www.tumorregister-muenchen.de/en/facts/base/base_D05f_E.pdf

Global Statements about the statistics on the Internet -

Baseline Statistics (grey button ___), Survival (red button ___)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.5 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases^{###} are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR. The time-delayed acquisition of data and the occasionally high DCO-rates indicate optimizing reserves, among others, because of current financial and legal conditions that hinder the analyses.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, April 2013

- [#] Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007). Death certificates from 2011 are incorporated into these analyses.
- ^{##} Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- ### DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate. A high proportion of DCO cases (≥5%) in particular cancer types indicate insufficient participation of specific cancer specializations.

INCIDENCE

Table 1

Patient cohorts by year of diagnosis including DCO cases and multiple primaries, and with proportion of deaths and active follow-up

		DCO	Prop.	Prop. mult.	Prop.	Prop. actively
Year of	Cases #	cases	DCO	primaries	deaths	followed
diagnosis	n	n	%	8	00	8
1998	107			46.7	17.8	91.6
1999	127			39.4	20.5	99.2
2000	137			36.5	20.4	97.1
2001	163			42.3	12.9	98.8
2002	216			35.2	13.9	94.4
2003	217			31.8	8.8	95.9
2004	268			32.1	9.0	94.8
2005	312			30.1	6.1	95.5
2006	301			34.9	5.3	91.0
2007	340			34.1	2.6	76.8 ##
2008	412			24.0	2.4	48.3
2009	402			25.4	3.2	58.5
2010	420			21.9	1.2	84.3
2011	390			20.0	0.8	63.1 ###
1998-2011	3812			29.8	6.3	80.0

- # The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.
- ## Since 2007 the percentage of actively followed patients sharply declined compared to the previous years. This is a consequence of ambiguous data protection rules that currently forbid cancer registries in Bavaria to obtain the essential life status informations from competent registration offices.
- ### Please be aware that data of recent annual patient cohorts may not yet be fully processed. Therefore, the presented figures and tables are potentially related to different time periods as pointed out in the respective headlines or legends.

Incidence measures by year of diagnosis and gender including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

Year of diagnosis	Cases n	Incidence raw	Incidence WS	Incidence ES	Incidence BRD-S	
1998	107	9.1	6.0	8.0	8.5	
1999	127	10.7	7.0	9.3	10.0	
2000	137	11.4	7.0	9.5	10.5	
2001	163	13.4	8.7	11.6	12.5	
2002	216	11.0	7.3	9.7	10.5	
2003	217	11.0	7.1	9.4	10.2	
2004	268	13.6	8.3	11.1	12.1	
2005	312	15.7	9.6	12.7	13.8	
2006	301	15.0	9.5	12.7	13.7	
2007	340	14.7	9.1	12.2	13.1	
2008	412	17.8	11.0	14.7	15.6	
2009	402	17.3	10.7	14.3	15.2	
2010	420	17.9	11.4	15.3	16.2	
2011	390	16.7	10.2	13.7	14.7	
1998-2011	3812	14.5	9.1	12.1	13.0	

The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

Voor of	Gogog		0+4					Modian		
Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	107	56.8	11.0	28.5	89.2	43.9	49.3	56.4	62.3	69.0
1999	127	58.4	10.3	40.9	88.2	45.9	50.5	57.7	63.3	72.4
2000	137	59.7	12.0	30.9	85.6	44.1	51.4	59.1	67.8	77.5
2001	163	57.9	11.8	26.0	93.0	42.6	49.8	58.1	65.5	72.6
2002	216	58.4	11.2	34.3	88.8	45.5	50.1	57.9	65.1	74.2
2003	217	59.2	11.3	33.3	91.7	44.1	50.2	58.8	67.6	73.6
2004	268	60.9	10.8	32.5	90.7	46.7	53.5	62.3	67.7	74.1
2005	312	60.0	11.5	30.0	91.4	43.2	52.5	61.5	66.8	73.3
2006	301	59.4	10.6	33.8	83.7	45.7	51.1	60.4	66.7	72.2
2007	340	60.0	11.6	26.3	90.8	43.9	51.9	60.9	67.8	73.7
2008	412	59.6	11.4	34.0	92.5	44.2	51.1	59.9	67.7	71.7
2009	402	59.9	11.2	26.2	89.6	44.7	51.9	60.9	67.4	73.1
2010	420	59.0	10.6	27.2	89.8	45.0	51.5	59.3	66.7	72.2
2011	390	60.1	11.1	30.8	92.5	45.3	51.0	60.5	68.5	73.7
1998-2011	3812	59.5	11.2	26.0	93.0	44.6	51.3	59.8	67.3	73.3

Age distribution parameters by year of diagnosis (incl. DCO)

Age distribution by 5-year age group for period 1998-2011 (incl. DCO)

Age at diagnosis Years	Cases n	୧	Cum.%	
25-29	7	0.2	0.2	
30-34	35	0.9	1.1	
35-39	98	2.6	3.7	
40-44	267	7.0	10.7	
45-49	390	10.2	20.9	
50-54	546	14.3	35.2	
55-59	586	15.4	50.6	
60-64	650	17.1	67.7	
65-69	628	16.5	84.1	
70-74	309	8.1	92.2	
75-79	161	4.2	96.5	
80-84	83	2.2	98.6	
85+	52	1.4	100.0	
All ages	3812	100.0		

Included in the statistics are 33.8% multiple primaries.

Prop. all Age at DCO rate cancers diagnosis Cases Age-spec. n=0 n=129521 Years incidence % ° n 0- 4 0.0 5-9 0.0 10 - 140.0 15-19 0.0 20 - 240.0 0.4 25-29 7 0.8 30-34 35 1.9 2.0 35-39 98 4.7 3.0 40 - 44265 12.5 5.0 45-49 388 20.3 5.4 50-54 31.6 5.9 542 55-59 35.4 4.9 580 60-64 40.3 4.3 646 65-69 41.8 3.8 622 70-74 24.8 2.0 306 75-79 16.0 159 1.1 80-84 82 10.3 0.6 85+ 52 7.0 0.4 2.9 All ages 3782 0.0 Incidence 14.4 Raw 9.0 WS 12.0 ES BRD-S 12.9

Age-specific incidence, DCO rate and proportion of all cancers for period 1998-2011

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).


Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2011

	Observe	d Expected	l	LCL	UCL		DCO
Diagnosis	n	n	SIR	95%	95%	EAR	00
C16 Stomach	3	3.4	0.9	0.2	2.6	-0.3	
C17 Small intestine	2	0.6	3.5	0.4	12.5	1.1	
C18 Colon	13	9.9	1.3	0.7	2.2	2.3	
C19-C20 Rectum	9	4.8	1.9	0.9	3.5	3.2	
C22 Liver	3	1.1	2.6	0.5	7.7	1.4	33.3
C25 Pancreas	13	4.2	3.1	1.6	5.2 #	6.7	15.4
C33-C34 Lung	20	8.9	2.3	1.4	3.5 #	8.5	5.0
C43 Malign. melanoma	a 8	4.6	1.7	0.8	3.4	2.6	
C50 Breast	306	41.9	7.3	6.5	8.2 #	201.1	
C51 Vulva	3	1.0	3.1	0.6	9.2	1.6	
C54 Corpus uteri	18	7.1	2.6	1.5	4.0 #	8.3	5.6
C56 Ovary	7	5.1	1.4	0.6	2.8	1.5	
C64 Kidney	4	2.8	1.4	0.4	3.7	0.9	
C70-C72 CNS cancer	3	1.8	1.7	0.4	5.0	0.9	33.3
C73 Thyroid	2	3.0	0.7	0.1	2.4	-0.8	
C76-C79 CUP	2	1.6	1.2	0.1	4.4	0.3	50.0
C82-C85 NHL	9	4.1	2.2	1.0	4.1	3.7	
C90 Mult. myeloma	2	1.3	1.6	0.2	5.6	0.5	
C91-C96 Leukaemia	5	1.6	3.2	1.0	7.4 #	2.6	
Other primaries	11	6.0	1.8	0.9	3.3	3.8	
Not observed	0	5.9	0.0	0.0	0.6 #	-4.5	
All mult. primaries	443	120.6	3.7	3.3	4.0 #	245.5	1.6

Patients	2674
Mean age at second malignancy (years)	64.1
Person-years	13132
Mean observation time (years)	4.9
Median observation time (years)	4.4

The occurrence of second malignancy is statistically significant.

Observed second malignancy with count 1 are pooled in category "Other primaries".



D05: Malignant neoplasm of breast in situ (women) Age distribution and age-specific incidence 1998 - 2011 (n=3782)

Figure 7. Age distribution and age-specific incidence





Figure 7a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).



Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2012, based on the November 2011 submission. http://www.seer.cancer.gov.




D05: Malignant neoplasm of breast in situ (women)

Figure 8. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.






Average incidence (world standard population) 2003 - 2008

Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (9.0/100,000 WS N=1,754). Since cancer data are not available in some counties until 2007, the local incidence rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 56 women were identified with newly diagnosed breast cancer i.s. (women). Therefore, the mean incidence rate for this cancer type in this area can be calculated at 9.1/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 6.2 and 13.0/100,000.



Standardized incidence ratio (SIR) 2003 - 2008

Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (N=1,754). Since cancer data are not available in some counties until 2007, the local SIR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 56 women were identified with newly diagnosed breast cancer i.s. (women). Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 1.02. Though, the value of this parameter may vary with an underlying probability of 99% between 0.71 and 1.43, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts, and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

		Prop.				Prop. deaths
	Incident	actively	Prop.		Prop.	with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	00	0/0	n	00	00
1998	107	91.6		19	17.8	100.0
1999	127	99.2		26	20.5	96.2
2000	137	97.1		28	20.4	96.4
2001	163	98.8		21	12.9	95.2
2002	216	94.4		30	13.9	100.0
2003	217	95.9		19	8.8	100.0
2004	268	94.8		24	9.0	100.0
2005	312	95.5		19	6.1	94.7
2006	301	91.0		16	5.3	100.0
2007	340	76.8		9	2.6	100.0
2008	412	48.3		10	2.4	90.0
2009	402	58.5		13	3.2	100.0
2010	420	84.3		5	1.2	100.0
2011	390	63.1		3	0.8	66.7
1998-2011	3812	80.0		242	6.3	97.5

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

			Prop.		
			deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	n	<u>%</u>	n	8
1998	107	17	94.1		
1999	127	21	85.7		
2000	137	19	78.9	1	0.7
2001	163	15	73.3		
2002	216	25	96.0		
2003	217	38	92.1		
2004	268	42	97.6		
2005	312	40	95.0	2	0.6
2006	301	45	88.9	1	0.3
2007	340	35	97.1		
2008	412	44	95.5	1	0.2
2009	402	56	100.0	2	0.5
2010	420	62	95.2	1	0.2
2011	390	65	100.0	1	0.3
1998-2011	3812	524	94.3	9	0.2

Table 10c

Annual cohorts of deaths, proportion of cancer-related and not cancerrelated deaths, and cancer recorded on death certificates (incl. DCO) (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

		Prop. cancer-	Prop. not cancer-	Prop. cancer recorded on death	
Year of	Deaths	related	related	certificate	
death	n	20	8	8	
1998	17	58.8	41.2	50.0	
1999	21	61.9	38.1	77.8	
2000	19	47.4	52.6	60.0	
2001	15	40.0	60.0	54.5	
2002	25	44.0	56.0	66.7	
2003	38	73.7	26.3	74.3	
2004	42	57.1	42.9	65.9	
2005	40	57.5	42.5	71.1	
2006	45	66.7	33.3	85.0	
2007	35	65.7	34.3	73.5	
2008	44	50.0	50.0	64.3	
2009	56	39.3	60.7	50.0	
2010	62	53.2	46.8	72.9	
2011	65	44.6	55.4	60.0	
1998-2011	524	54.0	46.0	66.6	

Munich Cancer Registry

Table	11

Means of age at death according to the grouping in Table 10

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(not cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	17	75.2	69.1	84.0	70.3
1999	21	71.0	68.4	75.3	70.4
2000	19	75.9	66.7	84.3	69.9
2001	15	74.5	63.8	81.7	64.1
2002	25	74.0	68.9	78.0	70.5
2003	38	71.1	66.9	82.9	67.1
2004	42	72.8	66.9	80.7	68.4
2005	40	77.2	74.2	81.4	75.2
2006	45	73.2	71.2	77.2	71.2
2007	35	71.9	68.6	78.0	68.3
2008	44	77.4	72.4	82.4	75.2
2009	56	76.6	65.7	83.6	68.8
2010	62	75.4	69.6	82.0	73.7
2011	65	77.0	71.1	81.7	72.9
1998-2011	524	74.8	69.3	81.3	71.1

By 2010, life expectancy for a newborn male in Germany is 77.5 years compared with 82.6 years for his female counterpart.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

			by	y year	of death				
Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	10	0.9	0.10	0.4	0.07	0.6	0.08	0.7	0.08
1999	13	1.1	0.10	0.5	0.07	0.7	0.08	0.9	0.09
2000	9	0.7	0.07	0.4	0.05	0.5	0.06	0.7	0.06
2001	б	0.5	0.04	0.3	0.03	0.4	0.03	0.4	0.03
2002	11	0.6	0.05	0.3	0.04	0.4	0.04	0.5	0.05
2003	28	1.4	0.13	0.7	0.10	1.0	0.11	1.2	0.12
2004	24	1.2	0.09	0.6	0.08	0.9	0.08	1.0	0.08
2005	23	1.2	0.07	0.5	0.05	0.7	0.06	0.9	0.07
2006	30	1.5	0.10	0.7	0.07	1.0	0.08	1.3	0.09
2007	23	1.0	0.07	0.5	0.05	0.7	0.06	0.9	0.07
2008	22	0.9	0.05	0.4	0.04	0.6	0.04	0.7	0.05
2009	22	0.9	0.05	0.5	0.05	0.7	0.05	0.8	0.05
2010	33	1.4	0.08	0.6	0.06	0.9	0.06	1.1	0.07
2011	29	1.2	0.07	0.5	0.05	0.8	0.06	1.0	0.07
1998-2011	283	1.1	0.07	0.5	0.06	0.7	0.06	0.9	0.07

Mortality measures (cancer-related death) and mortality-incidence-index

Age distribution of age at death (cancer-related) for period 1998-2011 (incl. multiple primaries)

Age at death Years	Cases n	96	Cum.%	
35-39	2	0.7	0.7	
40-44	6	2.1	2.8	
45-49	13	4.5	7.2	
50-54	17	5.9	13.1	
55-59	27	9.3	22.4	
60-64	49	16.9	39.3	
65-69	36	12.4	51.7	
70-74	39	13.4	65.2	
75-79	45	15.5	80.7	
80-84	17	5.9	86.6	
85+	39	13.4	100.0	
All ages	290	100.0		

Included in the statistics are 33.8% multiple primaries.

Munich Cancer Registry

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2011 (incl. multiple primaries)

Age at death	Cases Ag	0-5005		Prop. all cancers	
Years	/ =	e-spec. rtality	MI-index	calleers %	
ICALS	11 110	Itality	MI-INGEX	^т о	
0- 4 5- 9 10-14		0.0 0.0 0.0			
15-19		0.0			
20-24		0.0			
25-29		0.0			
30-34		0.0			
35-39	2	0.1	0.02	0.4	
40-44	6	0.3	0.02	0.6	
45-49	13	0.7	0.03	0.7	
50-54	17	1.0	0.03	0.6	
55-59	27	1.6	0.05	0.7	
60-64	49	3.1	0.08	0.9	
65-69	36	2.4	0.06	0.5	
70-74	39	3.2	0.13	0.5	
75-79	45	4.5	0.28	0.5	
80-84	17	2.1	0.20	0.2	
85+	39	5.3	0.75	0.3	
All ages	290			0.5	
Mortality					
Raw		1.1	0.08		
WS		0.5	0.06		
ES		0.8	0.06		
BRD-S		0.9	0.07		
PYLL-70					
per 100,000		7.2			
ES		6.1			
AYLL-70		10.8			

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.



Multiple primaries in deaths in period 1998-2011

					Syn-	Syn-		
	metel	metel	Deeg	Dree	chron	chron	Deet	Deet
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n	%↓	n	%→	n	⇔%	n	60→
	/							
C16 Stomach	4	1.3	1	25.0			3	75.0
C17 Small intestine	3	1.0					3	100.0
C18 Colon	10	3.2	1	10.0			9	90.0
C19-C20 Rectum	10	3.2	3	30.0			7	70.0
C22 Liver	3	1.0					3	100.0
C23-C24 Bile	3	1.0					3	100.0
C25 Pancreas	12	3.8					12	100.0
C33-C34 Lung	23	7.3			1	4.3	22	95.7
C43 Malign. melanoma	7	2.2	3	42.9			4	57.1
C44 Skin others	6	1.9	1	16.7	1	16.7	4	66.7
C50 Breast	158	50.2			59	37.3	99	62.7
C53 Cervix uteri	3	1.0	3	100.0				
C54 Corpus uteri	9	2.9	7	77.8			2	22.2
C56 Ovary	13	4.1	4	30.8	1	7.7	8	61.5
C70-C72 CNS cancer	5	1.6	1	20.0			4	80.0
C73 Thyroid	4	1.3	1	25.0			3	75.0
C82-C85 NHL	9	2.9	2	22.2			7	77.8
C90 Mult. myeloma	9	2.9	1	11.1			8	88.9
C91-C96 Leukaemia	8	2.5					8	100.0
Other primaries	16	5.1	4	25.0	1	6.3	11	68.8
-								
All mult. primaries	315	100.0	32	10.2	63	20.0	220	69.8

Multiple primaries with number of cases n<3 are pooled in category "Other primaries".

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2011 (Singular primaries only *)

Age at				Prop. all	
death		e-spec.		cancers	
Years	n mor	tality	MI-index	010	
0- 4 5- 9		0.0			
10-14		0.0			
15-19		0.0			
20-24		0.0			
25-29		0.0			
30-34		0.0			
35-39		0.0			
40-44	4	0.2	0.02	0.5	
45-49	5	0.3	0.02	0.3	
50-54	б	0.3	0.01	0.3	
55-59	14	0.9	0.03	0.4	
60-64	25	1.6	0.05	0.5	
65-69	19	1.3	0.04	0.3	
70-74	21	1.7	0.10	0.3	
75-79	29	2.9	0.30	0.4	
80-84	11	1.4	0.20	0.1	
85+	27	3.6	0.73	0.3	
		0.0			
All ages	161			0.3	
Mortality					
Raw		0.6	0.05		
WS		0.3	0.04		
ES		0.4	0.04		
BRD-S		0.5	0.05		
PYLL-70					
per 100,000		3.3			
ES		2.7			
AYLL-70		10.1			

* See corresponding tables with multiple primaries.



Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2011 (**Single primaries only ***)

Age at				Prop. all	
death	Cases Ag	ge-spec.		cancers	
Years	n mo	ortality	MI-index	00	
0- 4		0.0			
5-9		0.0			
10-14		0.0			
15-19		0.0			
20-24		0.0			
25-29		0.0			
30-34		0.0			
35-39		0.0			
40 - 44	3	0.1	0.01	0.4	
45-49	2	0.1	0.01	0.1	
50-54		0.0			
55-59	3	0.2	0.01	0.1	
60-64	8	0.5	0.02	0.2	
65-69	8	0.5	0.02	0.2	
70-74	5	0.4	0.03	0.1	
75-79	5	0.5	0.06	0.1	
80-84	1	0.1	0.02	0.0	
85+	12	1.6	0.33	0.2	
All ages	47			0.1	
5					
Mortality					
Raw		0.2	0.02		
WS		0.1	0.01		
ES		0.1	0.01		
BRD-S		0.1	0.01		
PYLL-70					
per 100,000	0	1.1			
ES		0.9			
AYLL-70		10.2			

* See corresponding tables with multiple primaries.







Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). "

The difference between age at diagnosis (Table 3) and age at breast cancer i.s. (women)-related death (see Table 10) should be considered.



Average mortality (world standard population) 2003 - 2008

Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2003 to 2008. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (0.6/100,000 WS N=142). Since cancer data are not available in some counties until 2007, the local mortality rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 5 women died from breast cancer i.s. (women). Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.6/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.1 and 1.9/100,000.

Standardized mortality ratio (SMR) 2003 - 2008




Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (N=142). Since cancer data are not available in some counties until 2007, the local SMR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 5 women died from breast cancer i.s. (women). Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 1.20. Though, the value of this parameter may vary with an underlying probability of 99% between 0.26 and 3.38, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the tumor-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

AYLL-70 BRD-S	Average years of life lost prior to age 70 given a person dies before that age German standard population
DCO	Death certificate only
EAR	Excess absolute risk
	= excess cancer cases (O - E) per 10,000 person-years
ES	European standard population (old)
FRG	Federal Republic of Germany
GEKID	Association of Population-based Cancer Registries in Germany
	(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)
LCL	Lower confidence limit
MI-index	Ratio between mortality and incidence
MCR	Munich Cancer Registry (Tumorregister München)
PYLL-70	Potential years of life lost prior to age 70 given a person dies before that age
SEER	Surveillance, Epidemiology, and End Results (USA)
SIR	Standardized incidence ratio
SMR	Standardized mortality ratio
UCL	Upper confidence limit
WS	World standard population

Recommended Citation

Munich Cancer Registry. Baseline statistics D05: Breast cancer i.s. (women) [Internet]. 2013 [updated 2013 Apr 2; cited 2013 Jun 1]. Available from: http://www.tumorregistermuenchen.de/en/facts/base/base_D05f_E.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Munich Cancer Registry

Index of figures and tables

Fig./Tb	l.	Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	3
2	Incidence by year of diagnosis	4
3	Age distribution parameters by year of diagnosis	5
4	Age distribution by 5-year age group	6
5	Age-specific incidence and DCO rate	7
6	Standardized incidence ratio of second primaries	8
7	Age distribution and age-specific incidence (chart)	9
7a	Age-specific incidence internationally (chart)	10
8	Cumulative follow-up years (chart)	11
9a	Map of cancer incidence (WS) by county (chart)	12
9b	Standardized incidence ratio (SIR) by county (chart)	13
10a	Pts incident cohorts and mortality / yr	14
10b	Incidence and mortality by year of diagnosis	15
10c	Cancer-related deaths, death certification available / yr	16
11	Means of age at death / yr	17
12	Mortality by year of death	18
13	Distribution of age at death	19
14	Age-specific mortality	20
15	Multiple primaries in deaths	21
16	Age-specific mortality (first primaries)	22
17	Age-specific mortality (single primaries)	23
18	Age distribution and age-specific mortality (chart)	24
19a	Map of cancer mortality (WS) by county (chart)	25
19b	Standardized mortality ratio (SMR) by county (chart)	26