Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage

Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

Cancer statistics: Baseline statistics

CML: Chronic myel. leukaemia (morph.)

Year of diagnosis	1998-2011
Patients	683
Diseases	683
Creation date	04/02/2013
Export date	01/03/2013
Population	4.5 m



http://www.tumorregister-muenchen.de/en/facts/base/base_hCML_E.pdf

Global Statements about the statistics on the Internet – Baseline Statistics (grey button ——), Survival (red button ——)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.5 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases**** are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR. The time-delayed acquisition of data and the occasionally high DCO-rates indicate optimizing reserves, among others, because of current financial and legal conditions that hinder the analyses.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, April 2013

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007). Death certificates from 2011 are incorporated into these analyses.
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate. A high proportion of DCO cases (≥5%) in particular cancer types indicate insufficient participation of specific cancer specializations.

Some remarks regarding this cancer type

The results for leukemias should be interpreted with caution. As with other primarily non-surgically or non-radiologically treated cancer diseases, the MCR hardly manages to obtain even the simplest information on this cancer. The proportion of DCO cases indicates a situation that is far away from a satisfying cooperation. In the group of institutions that potentially participate in reporting are a few hospitals that refuse any contribution to MCR.

ICD-10 codes used for specifying cancer site

ICD-10	Description
C92.1	Chronic myeloid leukaemia [CML], BCR/ABL-positive

ICD-O-3 codes (morphology) used for specifying cancer site

Code	Description	
9863/3 9875/3 9876/3	Chronic myeloid leukemia, NOS Chronic myelogenous leukemia, BCR/ABL positive Atypical chronic myeloid leukemia, BCR/ABL negative	

INCIDENCE

Table 1

Patient cohorts by year of diagnosis including DCO cases and multiple primaries, and with proportion of deaths and active follow-up

				Prop.		Prop.
		DCO	Prop.	mult.	Prop.	actively
Year of	Cases #	cases	DCO	primaries	deaths	followed
diagnosis	n	n	%	%	%	%
1998	17	/ 1 /	5.9	5.9	76.5	100.0
1999	33	5	15.2	9.1	69.7	100.0
2000	37	10	27.0	8.1	59.5	100.0
2001	38	17	44.7	18.4	73.7	97.4
2002	54	20	37.0	14.8	68.5	96.3
2003	69	26	37.7	14.5	69.6	98.6
2004	55	18	32.7	20.0	58.2	96.4
2005	51	18	35.3	23.5	58.8	98.0
2006	51	22	43.1	33.3	70.6	94.1
2007	58	20	34.5	29.3	51.7	91.4 ##
2008	65	26	40.0	30.8	55.4	83.1
2009	53	14	26.4	28.3	39.6	79.2
2010	59	15	25.4	30.5	39.0	94.9
2011	43	15	34.9	25.6	48.8	83.7 ###
1998-2011	683	227	33.2	22.4	58.6	93.1

[#] The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

^{##} Since 2007 the percentage of actively followed patients sharply declined compared to the previous years. This is a consequence of ambiguous data protection rules that currently forbid cancer registries in Bavaria to obtain the essential life status informations from competent registration offices.

^{###} Please be aware that data of recent annual patient cohorts may not yet be fully processed. Therefore, the presented figures and tables are potentially related to different time periods as pointed out in the respective headlines or legends.

Table 1a

Patient cohorts by year of diagnosis and gender including DCO cases

All	Males	Females	Prop. males
n	n	n	%
17 /	7	10	41.2
33/	18	15	54.5
37	/ 19	18	51.4
38	/ 18	20	47.4
54	23	31	42.6
69	33	36	47.8
55	34	21/	61.8
51	22	29	43.1
51	31	20	60.8
58	34	24	58.6
65	38	27	58.5
53	26	27	49.1
59	30	29	50.8
43	26	17	60.5
683	359	324	52.6
	n 17 33 37 38 54 69 55 51 58 65 53 59 43	n n 17 7 33 18 37 19 38 18 54 23 69 33 55 34 51 22 51 31 58 34 65 38 53 26 59 30 43 26	n n n n 17 7 10 33 18 15 37 19 18 38 18 20 54 23 31 69 33 36 55 34 21 51 22 29 51 31 20 58 34 24 65 38 27 53 26 27 59 30 29 43 26 17

Table 2

Incidence measures by year of diagnosis and gender including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

			Males	Fem.	Males	Fem.	Males	Fem.	Males	Fem.
Year of	Males	Females	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	7	10	0.6	0.9	0.4	0.5	0.6	0.6	0.7	0.7
1999	18	15	1.6	1.3	1.1	0.6	1.4	0.9	1.7	1.1
2000	19	18	1.7	1.5	1.2	0.7	1.5	1.0	1.8	1.2
2001	18	20	1.6	1.6	1.2	0.7	1.4	1.0	1.6	1.4
2002	23	31	1.2	1.6	0.7	0.8	1.1	1.1	1.4	1.3
2003	33	36	1.8	1.8	1.0	0.8	1.5	1.2	1.9	1.5
2004	34	21 /	1.8	1.1	1.1	0.5	1.5	0.7	1.8	0.9
2005	22	29 🤇	1.2	1.5	0.7	0.7	0.9	1.0	1.2	1.2
2006	31	20	1.6	1.0	0.9	0.4	1.3	0.6	1.7	0.8
2007	34	24	1.5	1.0	0.9	0.4	1.2	0.6	1.5	0.8
2008	38	27	1.7	1.2	0.9	0.5	1.3	0.7	1.7	0.9
2009	26	27	1.2	1.2	0.7	0.6	1.0	0.8	1.1	0.9
2010	30	29	1.3	1.2	0.7	0.6	1.0	0.8	1.3	1.0
2011	26	17	1.2	0.7	0.7	0.3	0.9	0.4	1.1	0.6
1998-2011	359	324	1.4	1.2	0.9	0.6	1.2	0.8	1.5	1.0



The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

base_hCML_E.pdf

Table 3 Age distribution parameters by year of diagnosis (All) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	17	59.9	22.9	27.0	87.4	27.0	36.6	69.2	81.5	86.5
1999	33	61.3	18.5	1.5	88.5	35.3	54.9	68.2	72.5	79.7
2000	37	63.5	19.9	16,5	97.6	34.7	49.8	66.7	78.1	88.0
2001	38	62.4	20.9	9.8	96.4	32.9	51.2	64.7	79.1	85.9
2002	54	66.1	16.2	27.4	96.1	41.9	55.4	69.7	77.8	83.8
2003	69	65.0	18.8	21.0	98.9	37.8	51.6	70.3	80.8	85.5
2004	55	65.5	15.9	19.4	93.3	44.6	55.2	67.3	76.5	84.7
2005	51	66.1	18.1	9.3	92.9	42.5	56.7	70.3	79.1	84.3
2006	51	69.3	15.9	28.0	92.5	42.0	61.0	73.0	79.4	86.5
2007	58	65.7	19.8	10.6	94.2	37.4	47.2	69.8	81.9	86.0
2008	65	70.7	16.0	26.6	98.1	47.2	62.5	74.4	81.5	87.2
2009	53	65.7	16.5	28.3	100	43.0	54.2	65.9	77.2	86.3
2010	59	67.0	16.8	15.7	90.9	47.5	54.9	71.7	78.2	84.4
2011	43	65.8	18.2	16.6	91.4	42.8	53.3	69.6	81.4	84.1
1998-2011	683	65.9	17.9	1.5	100	39.6	54.4	69.7	79.4	85.6

Table 3a Age distribution parameters by year of diagnosis (MALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	7	60.3	21.0	32.4	86.5	32.4	36.6	69.2	76.4	86.5
1999	18	56.3	20.5	1.5	80.3	30.7	43.4	60.2	69.8	78.3
2000	19	59.6	19.0	16.5	97.6	34.7	48.2	60.5	72.4	79.6
2001	18	51.6	22.8	9.8	96.4	23.3	37.7	51.5	64.0	81.6
2002	23	62.0	15.9	32.7	83.8	38.0	49.7	62.4	75.9	78.7
2003	33	60.7	19.4	21.0	91.1	35.2	41.9	64.5	77.3	83.7
2004	34	64.1	16.2	19.4	90.1	41.8	51.5	65.8	75.5	84.6
2005	22	62.3	20.8	9.3	91.0	38.1	43.7	71.5	78.2	81.5
2006	31	68.4	15.1	37.3	92.5	48.2	57.3	70.7	77.2	86.5
2007	34	62.3	20.5	10.6	91.8	33.9	46.8	68.0	77.4	84.0
2008	38	69.6	14.8	28.0	98.1	47.6	62.5	72.6	80.9	84.2
2009	26	63.4	17.9	28.3	92.2	38.2	51.2	64.9	77.2	86.3
2010	30	67.2	17.8	15.7	89.2	43.9	64.8	73.2	78.2	81.1
2011	26	62.8	19.6	16.6	91.4	41.3	52.7	67.0	75.7	84.1
1998-2011	359	63.1	18.5	1.5	98.1	37.5	50.3	67.3	77.2	84.0

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	10	59.5	25.3	27.0	87.4	27.0	33.6	65.8	81.7	86.5
1999	15	67.3	14.2	35.3	88.5	46.3	55.6	70.4	79.5	79.8
2000	18	67.6	20.5	27.4	93.9	32.2	54.5	73.4	86.2	89.0
2001	20	72.1	13.4	33.6	88.7	57.1	63.9	73.9	83.0	86.3
2002	31	69.1	16.0	27.4	96.1	53.0	60.5	70.9	82.5	88.4
2003	36	69.0	17.6	25.2	98.9	42.2	54.9	74.6	82.5	87.8
2004	21	67.8	15.4	28.0	93.3	49.3	57.1	69.0	80.2	84.7
2005	29	69.0	15.4	25.2	92.9	46.6	62.9	70.3	79.5	85.6
2006	20	70.7	17.4	28.0	92.2	37,1	68.8	74.3	80.4	87.8
2007	24	70.7	17.9	35.1	94.2	43.6	58.8	78.0	85.2	86.4
2008	27	72.1	17.8	26.6	94.9	41.4	60.7	77.1	85.4	88.6
2009	27	67.9	15.1	38.7	100	46.3	59.2	66.9	79.8	89.0
2010	29	66.8	15.9	29.0	90.9	47.5	53.9	70.9	75.9	87.8
2011	17	70.4	15.1	41.3	87.1	45.7	58.1	73.7	82.3	85.7
1998-2011	324	69.0	16.6	25.2	100	43.7	58.0	72.6	82.2	87.3

Table 4 Age distribution by 5-year age group and gender for period 1998-2011 (incl. DCO)

Age at									
diagnosis	Cases			Males			Females		
Years	n	%	Cum.%	n	010	Cum.%	n	%	Cum.%
0-4	1	0.1	0.1	1	0.3	0.3			0.0
5-9	2	0.3	0.4	2	0.6	0.8			0.0
10-14	1	0.1	0.6	/ 1	0.3	1.1			0.0
15-19	7	1.0	1.6	/ 7	1.9	3.1			0.0
20-24	2	0.3	1.9	2	0.6	3.6			0.0
25-29	15	2.2	4.1	5	1.4	5.0	10	3.1	3.1
30-34	15	2.2	6.3	11	3.1	8.1	4	1.2	4.3
35-39	27	4.0	10.2	18	5.0	13.1	9	2.8	7.1
40 - 44	33	4.8	15.1	23	6.4	19.5	10	3.1	10.2
45-49	36	5.3	20.4	19	5.3	24.8	17	5.2	15.4
50-54	41	6.0	26.4	22	6.1	30.9	19	5.9	21.3
55-59	39	5.7	32.1	21	5.8	36.8	18	5.6	26.9
60-64	49	7.2	39.2	28	7.8	44.6	21	6.5	33.3
65-69	81	11.9	51.1	48	13.4	57.9	33	10.2	43.5
70-74	88	12.9	64.0	37	10.3	68.2	51	15.7	59.3
75-79	88	12.9	76.9	57	15.9	84.1	31	9.6	68.8
80-84	81	11.9	88.7	30	8.4	92.5	51	15.7	84.6
85+	77	11.3	100.0	27	7.5	100.0	50	15.4	100.0
All ages	683	100.0		359	100.0		324	100.0	

Included in the statistics are 27.9% multiple primaries in males and 27.8% in females.

Table 5 Age-specific incidence, DCO rate and proportion of all cancers for period 1998-2011

			101 1	eriou i	770 2011			
Age at	Ma la m	nl.	Age-	Females Age-	DCO rate	Females DCO rate	cancers	Females Prop.all cancers
diagnosis			- /	spec.	n=112	n=115		n=129521
Years	n	n	incid.	incid.	%	%	%	%
0- 4	1		0.1	0.0			0.4	
5- 9	2		0.2	0.0			1.3	
10-14	1		0.1	0.0			0.8	
15-19	7		0.5	0.0	14.3		2.4	
20-24	2		0.1	0.0			0.4	
25-29	5	10	0.3	0.6			0.6	1.1
30-34	11	4	0.6	0.0			0.8	0.2
35-39	18	9	0.8	0.4			0.9	0.2
40-44	23	10	1.0	0.4	8.7		0.8	0.3
					0.7			
45-49	19	17	1.0	0.9	4 =	5 0	0.4	0.2
50-54	22	19	1.3	1.1	4.5	5.3	0.3	0.2
55-59	21	18	1.3	1.1	9.5	16.7	0.2	0.2
60-64	28	21	1.8	1.3	21.4	19.0	0.1	0.1
65-69	48	/33	3.5	2.2	27.1	24.2	0.2	0.2
70-74	37	51	3.6	4.1	43.2	39.2	0.2	0.3
75-79	57	31 /	8.4	3.1	43.9	48.4	0.3	0.2
80-84	30	51	7.4	6.4	83.3	51.0	0.3	0.4
85+	27	50	9.7	6.7	77.8	76.0	0.3	0.3
All ages	359	324			31.2	35.5	0.3	0.3
Incidence								
Raw			1.4	1.2				
WS			0.9	0.6				
ES			1.2	0.8				
BRD-S			1.5	1.0				


The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

Table 6a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2011

MALES

	Observed Ex	xpected		LCL	UCL		DCO
Diagnosis	'n	n	SIR	95%	95%	EAR	%
C18 Colon	2	0.9	2.3	0.3	8.2	14.2	
C43 Malign. melanoma	2	0.4	5.5	0.7	19.8	20.6	
C61 Prostate	7 /	2.6	2.7	1.1	5.6 #	56.0	
C67 Bladder	2 /	0.4	5.5	0.7	20.0	20.7	
C82-C85 NHL	2	0.4	5.7	0.7	20.6	20.8	
C91-C96 Leukaemia	3	0.1	21.7	4.5	63.5 #	36.1	66.7
Other primaries	8	2.6	3.1	1.3	6.1 #	68.3	
Not observed	0	1.8	0.0	0.0	2.1	-22.3	
All mult. primaries	26	9.0	2.9	1.9	4.2 #	214.4	7.7
-							




The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries".

Table 6b


Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2011 FEMALES

		Observed	Expected		LCL	UCL		DCO
Diagnosi	.s	n	n	SIR	95%	95%	EAR	%
C18	Colon	2	0.6	3.5	0.4	12.8	22.3	
C50	Breast	4	1.8	2.2	0.6	5.7	34.4	
C54	Corpus uteri	2	0.3	6.3	0.8	22.6	26.1	
Other pr	rimaries	5	1.1	4.4	1.4	10.2 #	60.0	60.0
Not obse	erved	0	2.0	0.0	0.0	1.9	-30.7	
All mult	. primaries	13	5.8	2.2	1.2	3.8 #	112.1	23.1



The occurrence of second malignancy is statistically significant.

Observed second malignancy with count 1 are pooled in category "Other primaries".







Figure 7. Age distribution and age-specific incidence








Figure 7a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).



Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2012, based on the November 2011 submission. http://www.seer.cancer.gov.





Figure 8. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.



Average incidence (world standard population) 2003 - 2008: Males



Average incidence (world standard population) 2003 - 2008: Females




Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.9/100,000 WS N=183, females 0.6/100,000 WS N=151). Since cancer data are not available in some counties until 2007, the local incidence rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 4 women were identified with newly diagnosed chronic myel. leukaemia (morph.). Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.5/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.1 and 2.0/100,000.

Standardized incidence ratio (SIR) 2003 - 2008: Males




Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=183, females N=151). Since cancer data are not available in some counties until 2007, the local SIR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 4 women were identified with newly diagnosed chronic myel. leukaemia (morph.). Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.92. Though, the value of this parameter may vary with an underlying probability of 99% between 0.15 and 2.89, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts, and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

		Prop.				Prop. deaths
	Incident	actively	Prop.		Prop.	with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	%	%	/ n /	%	%
1998	17	100.0	5.9	13	76.5	92.3
1999	33	100.0	15.2	23	69.7	95.7
2000	37	100.0	27.0	22	59.5	100.0
2001	38	97.4	44.7	28	73.7	96.4
2002	54	96.3	37.0	37	68.5	100.0
2003	69	98.6	37.7	48	69.6	100.0
2004	55	96.4	32.7	32	58.2	96.9
2005	51	98.0	35.3	30	58.8	100.0
2006	51	94.1	43.1	36	70.6	100.0
2007	58	91.4	34.5	30	51.7	100.0
2008	65	83.1	40.0	36	55.4	100.0
2009	53	79.2	26.4	21	39.6	100.0
2010	59	94.9	25.4	23	39.0	100.0
2011	43	83.7	34.9	21	48.8	100.0
1998-2011	683	93.1	33.2	400	58.6	99.0

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

(with respect to registry area expansion from $2.51\ \mathrm{to}\ 3.96\ \mathrm{m}$ as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

			Prop.		
			deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	/ n /	%	n	%
1998	17	14	100.0	2	11.8
1999	33	14	92.9	1	3.0
2000	37	27	92.6	10	27.0
2001	38	30	96.7	13	34.2
2002	54	37	100.0	24	44.4
2003	69	31	100.0	26	37.7
2004	55	37	97.3	20	36.4
2005	51	36	100.0	20	39.2
2006	51	35	100.0	28	54.9
2007	58	27	100.0	23	39.7
2008	65	27	96.3	27	41.5
2009	53	25	100.0	18	34.0
2010	59	21	100.0	18	30.5
2011	43	35	100.0	19	44.2
1998-2011	683	396	98.5	249	36.5

Table 10c

Annual cohorts of deaths, proportion of cancer-related and not cancerrelated deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

				Prop.	
				cancer	
		Prop.	Prop.	recorded	
		cancer-	not cancer-	on death	
Year of	Deaths	related	related	certificate	
death	n /	8	%	%	
1998	14	57.1	42.9	85.7	
1999	14	57.1	42.9	92.3	
2000	27	77.8	22.2	100.0	
2001	30	70.0	30.0	93.1	
2002	37	83.8	16.2	97.3	
2003	31	77.4	22.6	100.0	
2004	37	81.1	18.9	94.4	
2005	36	88.9	11.1	100.0	
2006	35	94.3	5.7	100.0	
2007	27	85.2	14.8	100.0	
2008	27	59.3	40.7	88.5	
2009	25	72.0	28.0	96.0	
2010	21	76.2	23.8	90.5	
2011	35	80.0	20.0	94.3	
1998-2011	396	78.0	22.0	95.9	

Table 11a Means of age at death according to the grouping in Table 10 MALES

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(not cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	8	58.5	59.8	57.2	60.2
1999	11	62.5	52.1	74.9	60.1
2000	15	70.1	69.1	76.7	72.8
2001	6	61.9	61.5	64.0	61.9
2002	13	63.6	65.4	57.6	64.5
2003	15	66.9	66.3	76.5	66.9
2004	20	69.8	69.3	71.8	69.7
2005	22	67.1	66.3	74.7	67.1
2006	22	73.2	73.0	75.9	73.2
2007	11	71.9	70.5	85.5	71.9
2008	15	67.5	71.3	59.8	69.9
2009	14	77.0	76.2	79.1	77.5
2010	15	77.6	76.4	79.9	77.6
2011	22	73.6	71.4	83.8	73.6
1998-2011	209	69.6	69.1	71.9	70.0

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 11b Means of age at death according to the grouping in Table 10 FEMALES

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(not cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	6	49.6	48.2	52.3	56.2
1999	3	79.8	79.1	81.2	79.1
2000	12	80.7	77.8	86.6	80.7
2001	24	76.7	72.3	85.5	76.6
2002	24	73.8	72.2	85.4	73.8
2003	16	79.8	77.0	84.3	79.8
2004	17	70.7	72.4	62.8	70.0
2005	14	74.1	71.9	87.5	74.1
2006	13	72.2	72.6	67.0	72.2
2007	16	77.1	75.2	85.1	77.1
2008	12	76.8	69.4	84.2	75.2
2009	11	80.6	85.9	66.5	80.6
2010	6/	78.0	78.0		78.0
2011	1,3	76.2	73.8	84.2	75.7
1998-2011	187	75.2	73.6	80.5	75.3



Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 12a Mortality measures (cancer-related death) and mortality-incidence-index by year of death MALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	4	0.4	0.57	0.2	0.51	0.3	0.52	0.4	0.53
1999	6	0.5	0.33	0.4	0.34	0.5	0.33	0.5	0.33
2000	13	1.1	0.68	0.7	0.60	1.0	0.68	1.4	0.78
2001	5	0.4	0.28	0.2	0.21	0.3	0.25	0.4	0.26
2002	10	0.5	0.43	0.3	0.44	0.4	0.42	0.6	0.39
2003	14	0.7	0.42	0.4	0.40	0.6	0.42	0.8	0.44
2004	16	0.9	0.47	0.4	0.41	0.7	0.48	1.0	0.52
2005	20	1.1	0.91	0.5	0.78	0.8	0.86	1.1	0.93
2006	21	1.1	0.68	0.5	0.63	0.9	0.66	1.2	0.72
2007	10	0.5	0.29	0.2	0.23	0.3	0.28	0.5	0.30
2008	10	0.4	0.26	0.2	0.25	0.3	0.28	0.5	0.27
2009	10	0.4	0.38	0.2	0.25	0.3	0.32	0.5	0.43
2010	10	0.4	0.33	0.2	0.26	0.3	0.31	0.4	0.33
2011	18	0.8	0.69	0.4	0.56	0.6	0.63	0.8	0.72
1998-2011	167	0.7	0.47	0.3	0.40	0.5	0.45	0.7	0.48

Table 12b Mortality measures (cancer-related death) and mortality-incidence-index by year of death FEMALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	4	0.3	0.40	0.2	0.50	0.3	0.50	0.3	0.43
1999	2	0.2	0.13	0.0	0.06	0.1	0.09	0.2	0.14
2000	8	0.7	0.44	0.2	0.29	0.4	0.36	0.5	0.44
2001	16	1.3	0.80	0.5	0.73	0.8	0.76	1.1	0.83
2002	21	1.1	0.68	0.5	0.60	0.7	0.63	0.9	0.69
2003	10	0.5	0.28	0.2	0.19	0.3	0.23	0.4	0.28
2004	14	0.7	0.67	0.3	0.55	0.5	0.62	0.6	0.68
2005	12	0.6	0.41	0.2	0.36	0.4	0.38	0.5	0.38
2006	12	0.6	0.60	0.2	0.58	0.4	0.64	0.5	0.63
2007	13	0.6	0.54	0.2	0.50	0.3	0.54	0.4	0.54
2008	6	0.3	0.22	0.1	0.25	0.2	0.26	0.2	0.26
2009	8	0.3	0.30	0.1	0.12	0.1	0.17	0.2	0.21
2010	6	0.3	0.21	0.1	0.12	0.1	0.15	0.2	0.17
2011	10	0.4	0.59	0.2	0.53	0.2	0.55	0.3	0.61
1998-2011	142	0.5	0.44	0.2	0.36	0.3	0.40	0.4	0.43

Table 13 Age distribution of age at death (cancer-related) for period 1998-2011 (incl. multiple primaries)

Age at								
death	Cases		Mal	es		Females		
Years	n	% Cu	m.% n	્ર	Cum.%	n	%	Cum.%
15-19	1	0.3	0.3	1 0.6	0.6			0.0
20-24	0	0.0	0.3		0.6			0.0
25-29	2	0.6	1.0	2 1.2	1.8			0.0
30-34	4	1.3	2.3	2 1.2	3.0	2	1.4	1.4
35-39	7	2.3	4.5	5 3.0	6.0	2	1.4	2.8
40 - 44	9	2.9	7.4	9 5.4	11.4			2.8
45-49	13	4.2 / 1	1.7	7 4.2	15.6	6	4.2	7.0
50-54	6	1.9 / 1	3.6	4 2.4	18.0	2	1.4	8.5
55-59	23	7.4 - 2	1.0 1	7.2	25.1	11	7.7	16.2
60-64	18	5.8 2	6.9 1	6.0	31.1	8	5.6	21.8
65-69	33	10.7 3	7.5 2	1 12.6	43.7	12	8.5	30.3
70-74	43	13.9 5	1.5 1	9 11.4	55.1	24	16.9	47.2
75-79	64	20.7 7	2.2 3	3 22.8	77.8	26	18.3	65.5
80-84	43	13.9 8	6.1 1	7 10.2	88.0	26	18.3	83.8
85+	43	13.9 10	0.0 2	0 12.0	100.0	23	16.2	100.0
All ages	309	100.0	16	7 100.0		142	100.0	

Included in the statistics are 27.9% multiple primaries in males and 27.8% in females.

Table 14

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2011 (incl. multiple primaries)

		Males		Females		Males	Females
Age at		Age-		Age-			Prop.all
death	Males Femal	_ /		spec.		cancers	cancers
Years	n n	mortal.	MI-index	mortal.	MI-index	%	%
0- 4		0.0		0.0			
5- 9		0.0		0.0			
10-14		0.0		0.0			
15-19	1	0.1		0.0		2.6	
20-24		0.0		0.0			
25-29	2	0.1		0.0		2.3	
30-34	2 2			0.1	0.50	1.2	1.0
35-39	5 2			0.1	0.22	1.4	0.4
40-44	9	0.4		0.0		1.2	
45-49	7 6	0.4		0.3		0.5	0.3
50-54	4 2			0.1	0.11	0.1	0.1
55-59	12 11	0.8		0.7		0.2	0.3
60-64	10 8			0.5	0.38	0.1	0.1
65-69	21 12		0.44	0.8	0.36	0.2	0.2
70-74	19 24			1.9	0.47	0.2	0.3
75-79	38 26			2.6	0.84	0.3	0.3
80-84	17 26			3.3	0.51	0.2	0.3
85+	20 23	7.2	0.74	3.1	0.46	0.3	0.2
All ages	167 142					0.2	0.2
Mortality							
Raw		0.7	0.47	0.5	0.44		
WS		0.3		0.2	0.36		
ES		0.5		0.3	0.40		
BRD-S		0.7	0.48	0.4	0.43		
PYLL-70							
per 100,000		5.0		2.4			
ES		4.3		2.0			
AYLL-70		15.4		12.5			

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15a $\begin{tabular}{ll} Multiple primaries in deaths in period 1998-2011 \\ \hline MALES \\ \end{tabular}$

					Syn- chron	Syn- chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n	%↓	n	~%	n	← %	n	~%
C16 Stomach	3	6,7	2	66.7			1	33.3
C18 Colon	4	8.9	1	25.0			3	75.0
C19-C20 Rectum	2	4.4	1	50.0	1	50.0		
C22 Liver	3	6.7	1	33.3			2	66.7
C25 Pancreas	/1	2.2	1	100.0				
C33-C34 Lung	4	8.9	1	25.0	1	25.0	2	50.0
C44 Skin others	3	6.7					3	100.0
C61 Prostate	7	15.6	6	85.7			1	14.3
C62 Testis	1	2.2	1	100.0				
C64 Kidney	3	6.7	3	100.0				
C65 Renal pelvis	1	2.2	1	100.0				
C67 Bladder	5	11.1	5	100.0				
C76-C79 CUP	2	4.4	1	50.0	1	50.0		
C82-C85 NHL	2	4.4	1	50.0			1	50.0
C91-C96 Leukaemia	4	8.9			1	25.0	3 /	75.0
All mult. primaries	45	100.0	25	55.6	4	8.9	16	35.6
_ /								




Table 15b Multiple primaries in deaths in period 1998-2011 FEMALES

Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C12-C13 Hypopharynx	1	2.5	1	100.0				
C18 Colon	/1	2.5	1	100.0				
C19-C20 Rectum	2	5.0	2	100.0				
C23-C24 Bile	1	2.5	1	100.0				
C25 Pancreas	2	5.0	_	100.0	2	100.0		
C33-C34 Lung	1 /	2.5			1	100.0		
C43 Malign. melanoma	/ 2	5.0	2	100.0	_	100.0		
C44 Skin others	3	7.5	1	33.3			2	66.7
C50 Breast	8	20.0	4	50.0	1	12.5	3	37.5
C51 Vulva	2	5.0	1	50.0	_	12.5	1	50.0
C52 Vagina	2	5.0	1	50.0			1	50.0
C53 Cervix uteri	3	7.5	2	66.7	1	33.3	_	
C54 Corpus uteri	5	12.5	4	80.0	_		1	20.0
C67 Bladder	1	2.5	1	100.0				
C70-C72 CNS cancer	1	2.5	1	100.0				
C74-C80 Cancer others	1	2.5					$\sqrt{1}$	100.0
C82-C85 NHL	1	2.5					1	100.0
C91-C96 Leukaemia	3	7.5			1	33.3	2	66.7
All mult. primaries	40	100.0	22	55.0	6	15.0	12	30.0




Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2011

(Singular primaries only *)

Age at death Years	Males n	Females	_ /	MI-index	Females Age- spec. mortal.	MI-index	cancers	Females Prop.all cancers %
0- 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19	1		0.1	0.14	0.0		2.9	
20-24			0.0		0.0			
25-29	2		0.1	0.40	0.0		2.5	
30-34	2	2 <	0.1	0.18	0.1	0.67	1.2	1.1
35-39	4	2	0.2		0.1	0.22	1.2	0.5
40-44	9		0.4		0.0		1.3	
45-49	7	6	0.4		0.3	0.38	0.5	0.4
50-54	3	1	0.2		0.1	0.07	0.1	0.0
55-59	10	8	0.6		0.5	0.57	0.2	0.2
60-64	10	8	0.7		0.5	0.47	0.2	0.2
65-69	19	10	1.4		0.7	0.37	0.2	0.2
70-74	18	19	1.7		1.5	0.42	0.2	0.3
75-79	31	22	4.6		2.2	0.92	0.4	0.3
80-84	13	19	3.2		2.4	0.54	0.2	0.2
85+	17	19	6.1	0.77	2.6	0.49	0.3	0.2
All ages	146	116					0.3	0.2
							\	
Mortality								
Raw			0.6		0.4	0.44		
WS			0.3	0.41	0.2	0.36		
ES			0.5	0.46	0.3	0.39		
BRD-S			0.6	0.50	0.4	0.43		
PYLL-70								
per 100,000			4.6		2.1			
ES ES			4.0		1.8			
AYLL-70			15.6		12.9			
VITTI-/0			13.0		14.9			

^{*} See corresponding tables with multiple primaries.

Table 17 Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2011 (Single primaries only *)

Age at			Males Age-		Females Age-		Males	Females Prop.all
death	Males	Females			spec.		cancers	cancers
Years	n	n		MI-index		MI-index		%
0- 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19	1		0.1	0.14	0.0		2.9	
20-24			0.0		0.0			
25-29	2		0.1	0.40	0.0		2.7	
30-34	2	2 <	0.1	0.20	0.1	0.67	1.3	1.2
35-39	4	2	0.2		0.1	0.22	1.2	0.5
40-44	8		0.4	0.38	0.0		1.2	
45-49	7	6	0.4	0.37	0.3	0.46	0.5	0.4
50-54	3		0.2	0.15	0.0		0.1	
55-59	10	7	0.6	0.50	0.4	0.54	0.2	0.2
60-64	10	7	0.7	0.56	0.4	0.41	0.2	0.2
65-69	17	10	1.2	0.55	0.7	0.38	0.2	0.2
70-74	15	19	1.5	0.56	1.5	0.43	0.2	0.3
75-79	29	19	4.3		1.9		0.4	0.3
80-84	10	17	2.5		2.1	0.55	0.2	0.3
85+	16	18	5.8	0.73	2.4	0.47	0.4	0.2
All ages	134	107					0.3	0.2
Mortality								
Raw			0.5		0.4			
WS			0.3		0.2			
ES			0.4		0.2	0.38		
BRD-S			0.6	0.49	0.3	0.42		
PYLL-70								
per 100,000			4.5		2.0			
ES			3.9		1.7			
AYLL-70			15.9		12.9			

^{*} See corresponding tables with multiple primaries.





Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at chronic myel. leukaemia (morph.)-related death (see Table 10) should be considered.



Average mortality (world standard population) 2003 - 2008: Males



Average mortality (world standard population) 2003 - 2008: Females





Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2003 to 2008. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.4/100,000 WS N=89, females 0.2/100,000 WS N=67). Since cancer data are not available in some counties until 2007, the local mortality rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 2 women died from chronic myel. leukaemia (morph.). Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.3/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.2/100,000.

Standardized mortality ratio (SMR) 2003 - 2008: Males



Standardized mortality ratio (SMR) 2003 - 2008: Females




Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=89, females N=67). Since cancer data are not available in some counties until 2007, the local SMR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 2 women died from chronic myel. leukaemia (morph.). Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 1.04. Though, the value of this parameter may vary with an underlying probability of 99% between 0.05 and 4.84, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the tumor-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

BRD-S German standard population

DCO Death certificate only EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

ES European standard population (old)
FRG Federal Republic of Germany

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

LCL Lower confidence limit

MI-index Ratio between mortality and incidence

MCR Munich Cancer Registry (Tumorregister München)

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age

SEER Surveillance, Epidemiology, and End Results (USA)

SIR Standardized incidence ratio
SMR Standardized mortality ratio
UCL Upper confidence limit
WS World standard population

Recommended Citation

Munich Cancer Registry. Baseline statistics CML: Chronic myel. leukaemia (morph.) [Internet]. 2013 [updated 2013 Apr 2; cited 2013 Jun 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/base/base_hCML_E.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

Fig./Tbl	l.	Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	4
1a	Gender distribution by year of diagnosis	5
2	Incidence by year of diagnosis	6
3	Age distribution parameters by year of diagnosis	7
4	Age distribution by 5-year age group and gender	9
5	Age-specific incidence and DCO rate	10
6	Standardized incidence ratio of second primaries	11
7	Age distribution and age-specific incidence (chart)	13
7a	Age-specific incidence internationally (chart)	14
8	Cumulative follow-up years (chart)	15
9a	Map of cancer incidence (WS) by county (chart)	16
9b	Standardized incidence ratio (SIR) by county (chart)	17
10a	Pts incident cohorts and mortality / yr	18
10b	Incidence and mortality by year of diagnosis	19
10c	Cancer-related deaths, death certification available / yr	20
11	Means of age at death / yr	21
12	Mortality by year of death	23
13	Distribution of age at death	24
14	Age-specific mortality	25
15	Multiple primaries in deaths	26
16	Age-specific mortality (first primaries)	28
17	Age-specific mortality (single primaries)	29
18	Age distribution and age-specific mortality (chart)	30
19a	Map of cancer mortality (WS) by county (chart)	31
19b	Standardized mortality ratio (SMR) by county (chart)	32