# **Munich Cancer Registry**



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage

Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

# **Cancer statistics: Baseline statistics**

# **GI-NET: Gastroint. neuroend. tumor**

Year of diagnosis	1998-2011
Patients	1341
Diseases	1351
Creation date	04/02/2013
Export date	01/03/2013
Population	4.5 m



http://www.tumorregister-muenchen.de/en/facts/base/base\_hGNETE.pdf

# Global Statements about the statistics on the Internet – Baseline Statistics (grey button —), Survival (red button —)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut<sup>#</sup>, with a total of 4.5 million inhabitants, account for the frequency of cancer diseases<sup>##</sup> and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases\*\*\*\* are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.


The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR. The time-delayed acquisition of data and the occasionally high DCO-rates indicate optimizing reserves, among others, because of current financial and legal conditions that hinder the analyses.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, April 2013

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007). Death certificates from 2011 are incorporated into these analyses.
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate. A high proportion of DCO cases (≥5%) in particular cancer types indicate insufficient participation of specific cancer specializations.



# ICD-10 codes used for specifying cancer site

Code	Description
C16 C17 C18 C19 C20	Stomach Small intestine Colon Rectosigmoid junction Rectum

# ICD-O-3 codes (morphology) used for specifying cancer site

Code	Description
8013/3	Large cell neuroendocrine carcinoma
8041/3	Small cell carcinoma, NOS
8152/1	Glucagonoma, NOS
8153/3	Gastrinoma, malignant
8156/3	Somatostatinoma, malignant
8240/3	Carcinoid tumor, NOS
8241/3	Enterochromaffin cell carcinoid
8243/3	Goblet cell carcinoid
8244/3	Composite carcinoid
8245/1	Tubular carcinoid
8246/3	Neuroendocrine carcinoma, NOS
8249/3	Atypical carcinoid tumor
8683/0	Gangliocytic paraganglioma

#### Reference:

Bosman FT, Carneiro F, Hruban RH, Theise ND, editors. WHO Classification of Tumours of the Digestive System 4th edition, IARC, Lyon (2010).

## **INCIDENCE**

Table 1

Patient cohorts by year of diagnosis including DCO cases and multiple primaries, and with proportion of deaths and active follow-up

				Prop.		Prop.
		DCO	Prop.	mult.	Prop.	actively
Year of	Cases #	cases	DCO	primaries	deaths	followed
diagnosis	n	n	%	90	୪	90
1998	45			24.4	55.6	100.0
1999	53			20.8	47.2	94.3
2000	41			26.8	41.5	97.6
2001	47			36.2	44.7	100.0
2002	75			22.7	38.7	96.0
2003	76			27.6	43.4	94.7
2004	108			28.7	38.9	92.6
2005	98			30.6	42.9	91.8
2006	127			27.6	37.0	89.8
2007	137			30.7	35.8	77.4 ##
2008	140			23.6	32.9	65.7
2009	139			23.7	29.5	71.9
2010	145			26.9	20.0	93.8
2011	120			20.8	11.7	70.0 ###
1998-2011	1351			26.4	34.0	85.0

<sup>#</sup> The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

<sup>##</sup> Since 2007 the percentage of actively followed patients sharply declined compared to the previous years. This is a consequence of ambiguous data protection rules that currently forbid cancer registries in Bavaria to obtain the essential life status informations from competent registration offices.

<sup>###</sup> Please be aware that data of recent annual patient cohorts may not yet be fully processed. Therefore, the presented figures and tables are potentially related to different time periods as pointed out in the respective headlines or legends.

Table 1a

Patient cohorts by year of diagnosis and gender including DCO cases

Year of	All	Males	Females	Prop. males	
diagnosis	n	n	n	%	
1998	45	24	21	53.3	
1999	53/	25	28	47.2	
2000	41	19	22	46.3	
2001	47	26	21	55.3	
2002	75	37	38	49.3	
2003	76	42	34	55.3	
2004	108	58	50	53.7	
2005	98	54	44	55.1	
2006	127	75	52	59.1	
2007	137	88	49	64.2	
2008	140	72	68	51.4	
2009	139	74	65	53.2	
2010	145	73	72	50.3	
2011	120	70	50	58.3	
1998-2011	1351	737	614	54.6	

Table 2

Incidence measures by year of diagnosis and gender including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

			Males	Fem.	Males	Fem.	Males	Fem.	Males	Fem.
Year of	Males	Females	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	24	21	2.2	1.8	1.3	1.0	1.9	1.3	2.2	1.6
1999	25	28	2.2	2.4	1.6	1.5	2.0	1.9	2.2	2.1
2000	19	22	1.7	1.8	1.0	1.0	1.5	1.4	1.8	1.6
2001	26	21	2.2	1.7	1.4	1,1	2.0	1.4	2.3	1.6
2002	37	38	2.0	1.9	1.2	1.2	1.7	1.5	2.0	1.8
2003	42	34	2.2	1.7	1.3	1.0	1.9	1.4	2.2	1.6
2004	58	50	3.1	2.5	1.9	/1.5	2.6	1.9	3.2	2.3
2005	54	44	2.9	2.2	1.7	/ 1.1	2.4	1.6	2.9	1.9
2006	75	52	3.9	2.6	2.1	1.5	3.1	1.9	3.7	2.3
2007	88	49	4.0	2.1	2.4	1.3	3.3	1.6	3.8	1.9
2008	72	68	3.2	2.9	1.9	1.6	2.6	2.1	3.2	2.5
2009	74	65	3.3	2.8	1.9	1.7	2.7	2.2	3.2	2.6
2010	73	72	3.2	3.1	1.9	1.9	2.6	2.5	3.0	2.8
2011	70	50	3.1	2.1	1.6	1.4	2.3	1.7	2.9	2.0
1998-2011	737	614	2.9	2.3	1.7	1.4	2.4	1.8	2.9	2.1



The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

Table 3

Age distribution parameters by year of diagnosis (All) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	45	63.5	15.2	13.2	87.9	46.6	56.0	64.1	72.5	83.3
1999	53	57.8	16.3	24.9	87.9	29.8	49.4	59.9	67.7	75.0
2000	41	61.8	16.8	24,7	92.1	39.8	49.5	61.0	75.5	80.6
2001	47	61.8	12.8	29.6	88.5	47.7	53.9	61.2	71.8	79.8
2002	75	61.7	16.4	17.7	90.9	35.0	51.2	64.1	73.3	80.3
2003	76	62.9	13.1	23.5	85.0	50.4	56.5	63.6	71.6	79.1
2004	108	61.5	15.0	13.8	93.3	39.1	52.3	63.9	72.3	78.3
2005	98	65.1	14.8	16.1	90.8	46.9	57.6	66.8	75.5	82.5
2006	127	64.6	12.5	29.9	91.9	46.1	55.3	66.3	73.7	80.0
2007	137	62.0	15.4	15.8	91.2	40.4	55.2	64.3	71.2	80.0
2008	140	64.0	14.7	18.9	93.9	44.7	56.3	66.4	73.4	80.1
2009	139	63.0	15.8	15.9	92.6	38.6	54.7	63.9	74.6	81.9
2010	145	61.2	14.1	14.9	89.6	42.3	54.1	61.9	71.6	78.5
2011	120	62.7	15.9	17.1	92.0	41.3	51.7	66.1	74.6	81.4
1998-2011	1351	62.6	14.9	13.2	93.9	42.3	54.3	64.3	72.9	80.0

Table 3a

Age distribution parameters by year of diagnosis (MALES)
(incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	24	62.1	11.4	32.1	81.7	46.6	56.9	63.8	66.5	76.6
1999	25	55.4	17.1	24.9	85.3	27.4	43.0	59.2	66.5	73.7
2000	19	62.1	15.1	37.4	92.1	39.0	49.5	60.1	74.4	80.6
2001	26	62.0	12.1	34.3	88.5	50.5	54.0	60.0	69.1	79.8
2002	37	61.8	13.2	27.1	88.3	42.3	57.9	64.1	70.6	76.2
2003	42	64.5	10.0	32.4	85.0	55.0	58.8	63.9	70.9	76.7
2004	58	62.2	12.0	27.8	78.9	47.7	53.9	63.9	71.4	76.5
2005	54	64.4	13.7	19.0	87.6	48.7	57.8	66.5	74.8	77.6
2006	75	65.5	10.4	38.5	85.7	49.5	59.0	67.4	72.8	76.9
2007	88	61.9	14.6	15.8	91.2	39.7	56.4	63.6	70.2	80.0
2008	72	64.5	11.9	19.3	85.6	49.9	58.6	65.7	73.4	79.1
2009	74	65.0	12.4	31.4	89.0	49.6	56.9	65.4	73.5	81.4
2010	73	62.4	12.4	26.3	86.8	47.1	55.2	63.2	71.6	78.0
2011	70	65.9	12.7	39.7	86.8	45.7	56.6	67.9	76.0	81.5
1998-2011	737	63.4	12.8	15.8	92.1	47.0	56.5	64.6	72.3	78.7

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	21	65.0	18.8	13.2	87.9	47.0	55.5	65.4	80.7	84.2
1999	28	59.9	15.4	26.9	87.9	38.4	51.0	62.0	72.2	76.9
2000	22	61.5	18.4	24.7	92.0	39.8	46.0	62.0	78.5	81.0
2001	21	61.6	14.0	29.6	84.4	47.7	53.9	62.6	71.8	75.7
2002	38	61.6	19.2	17.7	90.9	30.6	49.1	64.5	74.8	84.7
2003	34	60.9	16.1	23.5	84.0	33.0	53.0	63.0	73.6	79.4
2004	50	60.7	18.0	13.8	93.3	35.0	51.7	63.9	75.5	79.1
2005	44	65.9	16.1	16.1	90.8	40.7	56.6	68.7	78.4	83.0
2006	52	63.3	15.1	29.9	91.9	43.5	50.3	63.9	75.1	82.8
2007	49	62.0	16.7	17.8	86.8	41.0	53.5	65.2	73.9	81.7
2008	68	63.5	17.2	18.9	93.9	39.2	52.2	68.2	73.3	83.0
2009	65	60.7	18.8	15.9	92.6	35.0	49.4	62.3	76.3	84.2
2010	72	60.0	15.7	14.9	89.6	39.7	51.3	61.2	71.2	78.8
2011	50	58.3	18.8	17.1	92.0	27.9	46.9	60.2	71.3	80.9
1998-2011	614	61.7	17.1	13.2	93.9	37.0	51.7	63.9	73.9	81.7

Table 4

Age distribution by 5-year age group and gender for period 1998-2011 (incl. DCO)

Age at									
diagnosis	Cases			Males			Females		
Years	n	%	Cum.%	n	%	Cum.%	n	용	Cum.%
10-14	3	0.2	0.2			0.0	3	0.5	0.5
15-19	13	1.0	1,2	/ 3	0.4	0.4	10	1.6	2.1
20-24	11	0.8	2.0	/ 1	0.1	0.5	10	1.6	3.7
25-29	21	1.6	3.6	/ 10	1.4	1.9	11	1.8	5.5
30-34	32	2.4	5.9	14	1.9	3.8	18	2.9	8.5
35-39	36	2.7	8.6	12	1.6	5.4	24	3.9	12.4
40-44	49	3.6	12.2	24	3.3	8.7	25	4.1	16.4
45-49	75	5.6	17.8	37	5.0	13.7	38	6.2	22.6
50-54	111	8.2	26.0	58	7.9	21.6	53	8.6	31.3
55-59	139	10.3	36.3	88	11.9	33.5	51	8.3	39.6
60-64	206	15.2	51.5	127	17.2	50.7	79	12.9	52.4
65-69	211	15.6	67.1	135	18.3	69.1	76	12.4	64.8
70-74	163	12.1	79.2	90	12.2	81.3	73	11.9	76.7
75-79	147	10.9	90.1	83	11.3	92.5	64	10.4	87.1
80-84	87	6.4	96.5	39	5.3	97.8	48	7.8	95.0
85+	47	3.5	100.0	16	2.2	100.0	31	5.0	100.0
All ages	1351	100.0		737	100.0		614	100.0	

Included in the statistics are 37.5% multiple primaries in males and 26.5% in females.

Table 5

Age-specific incidence, DCO rate and proportion of all cancers for period 1998-2011

Age at diagnosis Years	Males n	Females n	Age- spec. incid.			Females DCO rate n=0 %	cancers	Females Prop.all cancers n=129521
0- 4 5- 9			0.0	0.0				
10-14		3	0.0	0.2				2.0
15-19	3	10	0.2	0.8			1.0	4.2
20-24	1	10	0.1	0.7			0.2	2.3
25-29	10	11	0.6	0.6			1.2	1.2
30-34	14	18	0.7	1.0			1.1	1.0
35-39	12	24	0.5	1.2			0.6	0.7
40-44	24	25	1.1	1.2			0.9	0.5
45-49	37	38	1.9	2.0			0.8	0.5
50-54	58	53	3.5	3.1			0.8	0.6
55-59	88	51	5.6	3.1			0.7	0.4
60-64	125	78	8.2	4.9			0.7	0.5
65-69	134	76	9.8	5.1			0.6	0.5
70-74	90	73	8.7	5.9			0.4	0.5
75-79 80-84	82 38	63 48	12.1 9.4	6.3			0.5 0.3	0.4
85+	36 16	31	5.8	6.0 4.2			0.3	0.4 0.2
037	10	31	5.0	4.2			0.2	0.2
All ages	732	612			0.0	0.0	0.6	0.5
Incidence								
Incidence Raw			2.9	2.3				
Raw WS			1.7	1.4				
Raw WS ES			1.7 2.4	1.4 1.8				
Raw WS			1.7	1.4				
Raw WS ES			1.7 2.4	1.4 1.8				
Raw WS ES			1.7 2.4	1.4 1.8				
Raw WS ES			1.7 2.4	1.4 1.8				
Raw WS ES			1.7 2.4	1.4 1.8				
Raw WS ES			1.7 2.4	1.4 1.8				
Raw WS ES			1.7 2.4	1.4 1.8				
Raw WS ES			1.7 2.4	1.4 1.8				
Raw WS ES			1.7 2.4	1.4 1.8				
Raw WS ES			1.7 2.4	1.4 1.8				
Raw WS ES			1.7 2.4	1.4 1.8				
Raw WS ES			1.7 2.4	1.4 1.8				
Raw WS ES			1.7 2.4	1.4 1.8				
Raw WS ES			1.7 2.4	1.4 1.8				
Raw WS ES			1.7 2.4	1.4 1.8				
Raw WS ES			1.7 2.4	1.4 1.8				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

Table 6a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries

for period 1998-2011 MALES

	Observed Ex	xpected		LCL	UCL		DCO
Diagnosis	n	n	SIR	95%	95%	EAR	%
C16 Stomagh	4	1.0	3.9	1 1	10.1 #	16 0	
C16 Stomach					**		
C17 Small intestine	2	0.1	16.6		60.0 #	10.2	
C18 Colon	/ 16 /	2.5	6.5	3.7	10.6 #	73.8	
C19-C20 Rectum	/ 10/	1.5	6.6	3.2	12.2 #	46.2	
C22 Liver	4	0.7	5.6	1.5	14.3 #	17.9	25.0
C23-C24 Bile	2	0.2	8.6	1.0	30.9 #	9.6	
C25 Pancreas	6	0.9	6.9	2.5	14.9 #	27.9	
C33-C34 Lung	5	3.1	1.6	0.5	3.7	10.1	20.0
C43 Malign. melanoma	3	1.0	2.9	0.6	8.4	10.6	
C61 Prostate	14	7.5	1.9	1.0	3.1 #	35.2	
C64 Kidney	6	0.9	6.4	2.3	13.9 #	27.5	
C67 Bladder	2	1.0	2.0	0.2	7.4	5.5	
C73 Thyroid	2	0.2	10.1	1.2	36.5 #	9.8	
C76-C79 CUP	2	0.4	4.8	0.6	17.4	8.6	
C82-C85 NHL	4	1.0	4.1	1.1	10.5 #	16.5	
C91-C96 Leukaemia	2	0.4	5.4	0.7	19.6	8.9	100.0
Other primaries	6	0.9	7.0	2.6	15.1 #	28.0	
Not observed	0	2.5	0.0	0.0	1,5	-13.6	
All mult. primaries	90	25.9	3.5	2.8	4.3 #	349.1	4.4

Patients	534
Mean age at second malignancy (years)	68.6
Person-years	1836
Mean observation time (years)	3.4
Median observation time (years)	2.6

# The occurrence of second malignancy is statistically significant.


Observed second primaries with count 1 are pooled in category "Other primaries".

base\_hGNETE.pdf

Table 6b

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2011 FEMALES

	Observed E	Expected		LCL U	DCO		
Diagnosis	n	n	SIR	95% 9	15%	EAR	%
C16 Stomach	4	0.6	6.7	1.8 17	.2 #	20.3	25.0
C18 Colon	9 /	1.6	5.5	2.5 10	.4 #	43.8	
C19-C20 Rectum	5 /	0.7	6.9	2.2 16	.1 #	25.4	20.0
C25 Pancreas	/ 4/	0.7	5.9	1.6 15	.0 #	19.7	
C33-C34 Lung	3	1.2	2.6	0.5 7	.6	11.0	
C43 Malign. melanoma	a 2	0.6	3.4	0.4 12	. 4	8.4	
C50 Breast	8	5.2	1.5	0.7 3	.0	16.8	
C54 Corpus uteri	2	0.9	/ 2.1 /	0.3 7	.8	6.4	
C56 Ovary	2	0.7	2.8	0.3 10	.2	7.7	
C64 Kidney	3	0.4	7.4	1.5 21	.6 #	15.4	33.3
C82-C85 NHL	3	0.6	4.9	1.0 14	.3 #	14.2	
Other primaries	4	0.6	6.2	1.7 15	.8 #	19.9	
Not observed	0	2.8	0.0	0.0 1	. 3	-16.6	
All mult. primaries	49	16.7	2.9	2.2 3	.9 #	192.4	6.1



# The occurrence of second malignancy is statistically significant.

Observed second malignancy with count 1 are pooled in category "Other primaries".

## GI-NET: Gastrointestinal neuroendocrine tumor





Figure 7. Age distribution and age-specific incidence



# GI-NET: Gastrointestinal neuroendocrine tumor



**Figure 7a.** Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).



#### Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER\*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2012, based on the November 2011 submission. http://www.seer.cancer.gov.





Figure 8. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.



#### Average incidence (world standard population) 2003 - 2008: Males




## Average incidence (world standard population) 2003 - 2008: Females



**Figure 9a.** Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 1.9/100,000 WS N=373, females 1.3/100,000 WS N=289). Since cancer data are not available in some counties until 2007, the local incidence rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 10 women were identified with newly diagnosed gastroint. neuroend. tumor. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 1.2/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.4 and 3.0/100,000.

#### Standardized incidence ratio (SIR) 2003 - 2008: Males



**Figure 9b.** Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=373, females N=289). Since cancer data are not available in some counties until 2007, the local SIR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 10 women were identified with newly diagnosed gastroint. neuroend. tumor. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 1.14. Though, the value of this parameter may vary with an underlying probability of 99% between 0.43 and 2.45, and is therefore not statistically striking.

## **MORTALITY**

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts, and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

		Prop.				Prop. deaths
	Incident	actively	Prop.		Prop.	with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	%	%	/ n /	%	%
1998	45	100.0		25	55.6	92.0
1999	53	94.3		25	47.2	100.0
2000	41	97.6		17	41.5	88.2
2001	47	100.0		21	44.7	95.2
2002	75	96.0		29	38.7	100.0
2003	76	94.7		33	43.4	100.0
2004	108	92.6		42	38.9	97.6
2005	98	91.8		42	42.9	97.6
2006	127	89.8		47	37.0	97.9
2007	137	77.4		49	35.8	95.9
2008	140	65.7		46	32.9	97.8
2009	139	71.9		41	29.5	100.0
2010	145	93.8		29	20.0	93.1
2011	120	70.0		14	11.7	92.9
1998-2011	1351	85.0		460	34.0	97.0

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

			Prop.		
			deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	/ n /	%	n	%
					/
1998	45	/ /9	88.9	4	8.9
1999	53	/ 9	100.0	4	7.5
2000	41	20	95.0	5	12.2
2001	47	10	100.0	3	6.4
2002	75	19	100.0	8	10.7
2003	76	33	97.0	/ 11	14.5
2004	108	36	100.0	10	9.3
2005	98	46	93.5	16	16.3
2006	127	41	95.1	11	8.7
2007	137	53	98.1	12	8.8
2008	140	64	95.3	17	12.1
2009	139	63	100.0	18	12.9
2010	145	59	98.3	18	12.4
2011	120	48	95.8	9	7.5
1998-2011	1351	510	97.1	146	10.8

#### Table 10c

Annual cohorts of deaths, proportion of cancer-related and not cancer-related deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

				Prop.	
				cancer	
		Prop.	Prop.	recorded	
		cancer-	not cancer-	on death	
Year of	Deaths	related	related	certificate	
death	n /	%	%	%	
1998	9 9	77.8	22.2	62.5	
1999	9	88.9	11.1	88.9	
2000	20	65.0	35.0	73.7	
2001	10	80.0	20.0	80.0	
2002	19	78.9	21.1	89.5	
2003	33	63.6	36.4	78.1	
2004	36	72.2	27.8	77.8	
2005	46	65.2	34.8	76.7	
2006	41	82.9	17.1	87.2	
2007	53	79.2	20.8	80.8	
2008	64	84.4	15.6	93.4	
2009	63	66.7	33.3	76.2	
2010	59	69.5	30.5	75.9	
2011	48	83.3	16.7	87.0	
1998-2011	510	74.7	25.3	81.4	

Table 11a  $\begin{tabular}{ll} Means of age at death according to the grouping in Table 10 \\ \hline MALES \end{tabular}$ 

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(not cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	3	67.0	70.9	59.3	71.3
1999	5	74.3	71.9	83.7	71.9
2000	14	69.7	66.5	81.2	66.5
2001	9	70.4	68.3	87.5	68.3
2002	13	68.0	66.0	92.4	68.0
2003	16	73.2	71.7	75.1	72.5
2004	16	70.9	68.3	73.5	69.0
2005	27	73.2	72.9	74.0	73.6
2006	19	72.7	71.4	77.8	71.4
2007	30	71.4	71.9	69.0	71.9
2008	43	69.0	67.8	76.1	68.1
2009	40	71.2	69.7	74.2	69.4
2010	35	70.6	68.6	76.3	68.6
2011	29	68.2	67.6	72.2	66.4
1000 2011	200	70.7	60.3	75.0	60 E
1990-2011	299	70.7	09.3	75.0	09.5
2005 2006 2007 2008 2009 2010	27 19 30 43 40 35	73.2 72.7 71.4 69.0 71.2 70.6	72.9 71.4 71.9 67.8 69.7 68.6	74.0 77.8 69.0 76.1 74.2 76.3	73.6 71.4 71.9 68.1 69.4 68.6




Table 11b  $\label{eq:means} \mbox{Means of age at death according to the grouping in Table 10 }$ 

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(not cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	6	82.4	81.2	88.1	82.2
1999	4	78.9	78.9		78.9
2000	6	73.9	71.6	75.0	77.2
2001	1	92.2		92.2	
2002	6	77.1	78.7	75.4	74.7
2003	17	74.9	73.3	78.6	73.8
2004	20	72.8	72.0	79.4	73.4
2005	19	76.0	70.5	83.5	72.5
2006	22	75.5	75.4	75.6	75.7
2007	23	71.1	68.9	77.3	68.9
2008	21	74.4	71.3	87.5	74.5
2009	23	74.7	71.5	82.0	73.3
2010	24	73.0	67.0	82.9	70.5
2011	19	71.6	68.2	84.3	70.3
1998-2011	211	74.2	71.7	81.1	73.1



Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 12a Mortality measures (cancer-related death) and mortality-incidence-index by year of death MALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	2	0.2	0.08	0.1	0.09	0.2	0.09	0.2	0.10
1999	4	0.4	0.16	0.2	0.15	0.3	0.17	0.5	0.22
2000	11	1.0	0.58	0.6	0.58	0.9	0.60	1.1	0.58
2001	8	0.7	0.32	0.4	0.29	0.6	0.32	0.9	0.39
2002	12	0.6	0.32	0.4	0.30	0.5	0.32	0.7	0.34
2003	9	0.5	0.21	0.3	0.20	0.4	0.22	0.6	0.25
2004	8	0.4	0.14	0.2	0.13	0.4	0.14	0.5	0.15
2005	19	1.0	0.36	0.5	0.29	0.8	0.33	1.1	0.38
2006	15	0.8	0.20	0.4	0.19	0.6	0.20	0.9	0.23
2007	25	1.1	0.29	0.5	0.22	0.8	0.25	1.1	0.29
2008	37	1.7	0.51	0.9	0.48	1.3	0.50	1.6	0.50
2009	26	1.2	0.36	0.6	0.30	0.9	0.33	1.1	0.36
2010	26	1.2	0.36	0.6	0.32	0.9	0.33	1.2	0.38
2011	25	1.1	0.36	0.6	0.35	0.8	0.37	1.0	0.36
1998-2011	227	0.9	0.31	0.5	0.28	0.7	0.30	0.9	0.32

Table 12b Mortality measures (cancer-related death) and mortality-incidence-index by year of death FEMALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	5	0.4	0.24	0.1	0.10	0.2	0.15	0.4	0.25
1999	4	0.3	0.14	0.1	0.08	0.2	0.10	0.2	0.10
2000	2	0.2	0.09	0.1	0.06	0.1	0.08	0.1	0.07
2001									
2002	3	0.2	0.08	0.1	0.04	0.1	0.05	0.1	0.06
2003	12	0.6	0.35	0.3	0.24	0.4	0.28	0.5	0.31
2004	18	0.9	0.36	0.4	0.23	0.6	0.29	0.8	0.33
2005	11	0.6	0.25	0.2	0.21	0.3	0.22	0.5	0.24
2006	19	0.9	0.37	0.3	0.24	0.5	0.28	0.7	0.33
2007	17	0.7	0.35	0.3	0.27	0.5	0.29	0.6	0.31
2008	17	0.7	0.25	0.3	0.17	0.4	0.19	0.6	0.22
2009	16	0.7	0.25	0.3	0.16	0.4	0.19	0.5	0.20
2010	15	0.6	0.21	0.3	0.17	0.5	0.18	0.6	0.20
2011	15	0.6	0.30	0.3	0.22	0.4	0.25	0.5	0.27
1998-2011	154	0.6	0.25	0.2	0.17	0.4	0.20	0.5	0.22

Table 13

Age distribution of age at death (cancer-related) for period 1998-2011 (incl. multiple primaries)

Age at									
death	Cases			Males			Females		
Years	n	%	Cum.%	n	96	Cum.%	n	%	Cum.%
25-29	1	0.3	0.3	1	0.4	0.4			0.0
30-34	2	0.5	0.8			0.4	2	1.3	1.3
35-39	4	1.0	1.8			0.4	4	2.6	3.9
40-44	7	1.8	3.7	/ 3	1.3	1.8	4	2.6	6.5
45-49	6	1.6	5.2	3	1.3	3.1	3	1.9	8.4
50-54	19	5.0	10.2	14	6.1	9.2	5	3.2	11.7
55-59	32	8.4	18.6	21	9.2	18.4	11	7.1	18.8
60-64	39	10.2	28.8	27	11.8	30.3	12	7.8	26.6
65-69	69	18.1	46.9	48	21.1	51.3	21	13.6	40.3
70-74	60	15.7	62.6	42	18.4	69.7	18	11.7	51.9
75-79	61	16.0	78.5	36	15.8	85.5	25	16.2	68.2
80-84	48	12.6	91.1	21	9.2	94.7	27	17.5	85.7
85+	34	8.9	100.0	12	5.3	100.0	22	14.3	100.0
All ages	382	100.0		228	100.0		154	100.0	

Included in the statistics are 37.5% multiple primaries in males and 26.5% in females.

Table 14

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2011 (incl. multiple primaries)

			Males		Females		Males	Females
Age at			Age-		Age-		_	Prop.all
death		Females	_ /		spec.	! ]	cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0- 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29	1		0.1	0.10	0.0		1.1	
30-34		2 <	0.0		0.1	0.11		1.0
35-39		4	0.0		0.2	0.17		0.9
40-44	3	4	0.1	0.13	0.2	0.16	0.4	0.4
45-49	3	3	0.2		0.2	0.08	0.2	0.2
50-54	14	5	0.8	0.24	0.3	0.09	0.5	0.2
55-59	21	11	1.3	0.24	0.7	0.22	0.4	0.3
60-64	27	12	1.8	0.21	0.7	0.15	0.4	0.2
65-69	48	21	3.5	0.36	1.4	0.28	0.5	0.3
70-74	42	18	4.1	0.47	1.5	0.25	0.4	0.2
75-79	36	25	5.3	0.43	2.5	0.39	0.3	0.3
80-84	21	27	5.2	0.54	3.4	0.56	0.2	0.3
85+	12	22	4.3	0.75	3.0	0.71	0.2	0.2
All ages	228	154					0.3	0.3
Mortality								
Raw			0.9		0.6	0.25		
WS			0.5	0.28	0.2	0.17		
ES			0.7		0.4	0.20		
BRD-S			0.9	0.32	0.5	0.22		
PYLL-70								
per 100,000			4.5		3.3			
ES			4.0		2.8			
AYLL-70			8.7		12.1			

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15a

Multiple primaries in deaths in period 1998-2011

MALES

Diagnos	is	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C15	Oesophagus	3	3.0	2	66.7			1	33.3
C16	Stomach	/2	2.0					2	100.0
C17	Small intestine	/ 9	8.9			4	44.4	5	55.6
C18	Colon	/ 11	10.9			10	90.9	1	9.1
C19-C20	Rectum	6	5.9			4	66.7	2	33.3
C21	Anus/canal	/ 1 /	1.0					1	100.0
C22	Liver	4	4.0			1	25.0	3	75.0
C23-C24	Bile	2	2.0			/ 1	50.0	1	50.0
C25	Pancreas	5	5.0	1	20.0	3	60.0	1	20.0
C33-C34	Lung	8	7.9	1	12.5	2	25.0	5	62.5
C38,C45	Mesothelioma	1	1.0					1	100.0
C43	Malign. melanoma	1	1.0					1	100.0
C44	Skin others	4	4.0	4	100.0				
C46,C49	Soft tissue	2	2.0	1	50.0			1	50.0
C50	Breast	1	1.0	1	100.0				
C61	Prostate	17	16.8	12	70.6	_ 2	11.8	3	17.6
C62	Testis	2	2.0	2	100.0				
C64	Kidney	5	5.0	2	40.0	2	40.0	1	20.0
C65	Renal pelvis	2	2.0					2	100.0
C66	Ureter	1	1.0					1	100.0
C67	Bladder	5	5.0	3	60.0			2	40.0
C70-C72	CNS cancer	1	1.0			1	100.0		
C76-C79	CUP	1	1.0			1	100.0		
C82-C85	NHL	4	4.0	1	25.0	1	25.0	2	50.0
C91-C96	Leukaemia	3	3.0					3	100.0
All mul	t. primaries	101	100.0	30	29.7	32	31.7	39	38.6

Multiple primaries with number of cases n<1 are pooled in category "Other primaries".

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 15b

Multiple primaries in deaths in period 1998-2011
FEMALES

						Syn- chron	Syn- chron		
		Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnos	is	n /	%↓	n	<b>←%</b>	n	<b>~</b> %	n	<b>←</b> %
C15	Oesophagus	1	/1.5					1	100.0
C16	Stomach	/3	4.6					3	100.0
C17	Small intestine	3	4.6			3	100.0		
C18	Colon	8	12.3			3	37.5	5	62.5
C19-C20	Rectum	4	6.2			2	50.0	2	50.0
C21	Anus/canal	/ 1 /	1.5	1	100.0				
C25	Pancreas	/ 1 -	1.5					1	100.0
C33-C34	Lung	6	9.2	3	50.0			3	50.0
C43	Malign. melanoma	2	3.1	2	100.0				
C50	Breast	15	23.1	13	86.7			2	13.3
C51	Vulva	2	3.1	2	100.0				
C54	Corpus uteri	4	6.2	2	50.0	2	50.0		
C56	Ovary	6	9.2	2	33.3	4	66.7		
C64	Kidney	2	3.1	1	50.0			1	50.0
C67	Bladder	2	3.1	2	100.0				
C70-C72	CNS cancer	1	1.5					$\sqrt{1}$	100.0
C76-C79	CUP	1	1.5					1	100.0
C82-C85	NHL	2	3.1	1	50.0			1	50.0
C91-C96	Leukaemia	1	1.5					1	100.0
All mul	t. primaries	65	100.0	29	44.6	14	21.5	22	33.8




Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2011

(Singular primaries only \*)

Age at			Males Age-		Females Age-		Males Prop.all	Females Prop.all
death	Males	Females	spec.		spec.		cancers	cancers
Years	n	n		MI-index		MI-index	%	%
0- 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29	1		0.1	0.10	0.0		1.2	
30-34		2	0.0		0.1	0.12		1.1
35-39		2	0.0		0.1	0.09		0.5
40-44	2	4	0.1		0.2	0.17	0.3	0.5
45-49	3	3	0.2		0.2		0.2	0.2
50-54	14	4	0.8	0.30	0.2	0.09	0.6	0.2
55-59	16	9	1.0		0.5	0.22	0.4	0.3
60-64	21	/11	1.4		0.7	0.17	0.3	0.2
65-69	38	13	2.8		0.9	0.26	0.5	0.2
70-74	30	12	2.9		1.0	0.22	0.3	0.2
75-79	24	1/7	3.6		1.7		0.3	0.2
80-84	12	20	3.0		2.5	0.51	0.2	0.3
85+	8	15	2.9	0.80	2.0	0.58	0.1	0.2
	\	_ \ _					\	
All ages	169	112					0.3	0.2
76 1 7 1								
Mortality			0.7	0.20	0 4	0.00		
Raw			0.7		0.4			
WS			0.4		0.2	0.15		
ES			0.5	0.29	0.3	0.18		
BRD-S			0.7	0.31	0.3	0.20		
PYLL-70								
per 100,000			3.8		2.7			
ES			3.4		2.3			
AYLL-70			9.1		12.8			

<sup>\*</sup> See corresponding tables with multiple primaries.


Table 17

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2011

(Single primaries only \*)

Age at			Males Age-		Females Age-		Males	Females Prop.all
death	Males	Females	spec.		spec.		cancers	cancers
Years	n	n		MI-index		MI-index	%	%
0- 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0,0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29	1		0.1	0.10	0.0		1.3	
30-34		2 <	0.0		0.1	0.12		1.2
35-39		2	0.0		0.1	0.10		0.5
40-44	2	4	0.1		0.2	0.17	0.3	0.5
45-49	3	3	0.2	0.09	0.2		0.2	0.2
50-54	14	4	0.8	0.32	0.2	0.09	0.6	0.2
55-59	16	7	1.0		0.4		0.4	0.2
60-64	21	10	1.4	0.24	0.6	0.17	0.4	0.2
65-69	32	/ 11	2.3		0.7	0.23	0.4	0.2
70-74	21	12	2.0	0.35	1.0	0.23	0.3	0.2
75-79	20	16	3.0	0.45	1.6	0.34	0.3	0.3
80-84	12	18	3.0	0.71	2.3	0.47	0.2	0.3
85+	6	14	2.2	0.67	1.9	0.54	0.1	0.2
	\	\						
All ages	148	103					0.3	0.2
Mortality			0.5	0.00	0.4	0.01		
Raw			0.6	0.29	0.4			
WS			0.3	0.26	0.2	0.15		
ES			0.5	0.28	0.2	0.17		
BRD-S			0.6	0.30	0.3	0.19		
PYLL-70								
per 100,000			3.7		2.6			
ES			3.4		2.2			
AYLL-70			9.5		13.4			

<sup>\*</sup> See corresponding tables with multiple primaries.




**Figure 18.** Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at gastroint. neuroend. tumor-related death (see Table 10) should be considered.



#### Average mortality (world standard population) 2003 - 2008: Males




## Average mortality (world standard population) 2003 - 2008: Females



**Figure 19a.** Map of cancer mortality (world standard population) by county averaged for period 2003 to 2008. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.5/100,000 WS N=106, females 0.3/100,000 WS N=91). Since cancer data are not available in some counties until 2007, the local mortality rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 2 women died from gastroint. neuroend. tumor. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.1/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 0.7/100,000.

#### Standardized mortality ratio (SMR) 2003 - 2008: Males








Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=106, females N=91). Since cancer data are not available in some counties until 2007, the local SMR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 2 women died from gastroint. neuroend. tumor. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.76. Though, the value of this parameter may vary with an underlying probability of 99% between 0.04 and 3.53, and is therefore not statistically striking.

#### **Statistical Notes**

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

## 1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the tumor-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

**3. Single primary** (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

#### **Shortcuts**

AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

BRD-S German standard population

DCO Death certificate only EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

ES European standard population (old) FRG Federal Republic of Germany

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

LCL Lower confidence limit

MI-index Ratio between mortality and incidence

MCR Munich Cancer Registry (Tumorregister München)

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age

SEER Surveillance, Epidemiology, and End Results (USA)

SIR Standardized incidence ratio
SMR Standardized mortality ratio
UCL Upper confidence limit
WS World standard population

#### **Recommended Citation**

Munich Cancer Registry. Baseline statistics GI-NET: Gastroint. neuroend. tumor [Internet]. 2013 [updated 2013 Apr 2; cited 2013 Jun 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/base/base\_hGNETE.pdf

#### Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

#### Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

base\_hGNETE.pdf

# Index of figures and tables

ol.	Page
Pts cohorts, DCO, mult. prim., follow-up / yr	4
Gender distribution by year of diagnosis	5
Incidence by year of diagnosis	6
Age distribution parameters by year of diagnosis	7
Age distribution by 5-year age group and gender	9
Age-specific incidence and DCO rate	10
Standardized incidence ratio of second primaries	11
Age distribution and age-specific incidence (chart)	13
Age-specific incidence internationally (chart)	14
Cumulative follow-up years (chart)	15
Map of cancer incidence (WS) by county (chart)	16
Standardized incidence ratio (SIR) by county (chart)	17
Pts incident cohorts and mortality / yr	18
Incidence and mortality by year of diagnosis	19
Cancer-related deaths, death certification available / yr	20
Means of age at death / yr	21
Mortality by year of death	23
Distribution of age at death	24
Age-specific mortality	25
Multiple primaries in deaths	26
Age-specific mortality (first primaries)	28
Age-specific mortality (single primaries)	29
Age distribution and age-specific mortality (chart)	30
Map of cancer mortality (WS) by county (chart)	31
Standardized mortality ratio (SMR) by county (chart)	32
	Pts cohorts, DCO, mult. prim., follow-up / yr Gender distribution by year of diagnosis Incidence by year of diagnosis Age distribution parameters by year of diagnosis Age distribution by 5-year age group and gender Age-specific incidence and DCO rate Standardized incidence ratio of second primaries Age distribution and age-specific incidence (chart) Age-specific incidence internationally (chart) Cumulative follow-up years (chart) Map of cancer incidence (WS) by county (chart) Standardized incidence ratio (SIR) by county (chart) Pts incident cohorts and mortality / yr Incidence and mortality by year of diagnosis Cancer-related deaths, death certification available / yr Means of age at death / yr Mortality by year of death Distribution of age at death Age-specific mortality Multiple primaries in deaths Age-specific mortality (first primaries) Age-specific mortality (single primaries) Age-specific mortality (ws) by county (chart) Map of cancer mortality (WS) by county (chart)