Munich Cancer Registry



- Survival
- Selection Matrix
- ► Homepage

Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

Cancer statistics: Baseline statistics

C07, C08: Salivary gland cancer

Year of diagnosis	1998-2012
Patients	598
Diseases	598
Creation date	03/20/2014
Export date	02/12/2014
Population	4.5 m



http://www.tumorregister-muenchen.de/en/facts/base/base_C0708E.pdf

Global Statements about the statistics on the Internet -

Baseline Statistics (grey button ___), Survival (red button ___)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.5 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases^{###} are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR. The time-delayed acquisition of data and the occasionally high DCO-rates indicate optimizing reserves, among others, because of current financial and legal conditions that hinder the analyses.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, March 2014

- [#] Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007). Death certificates from 2013 are incorporated into these analyses.
- ^{##} Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- ^{###} DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate. A high proportion of DCO cases (≥5%) in particular cancer types indicate insufficient participation of specific cancer specializations.

Some remarks regarding this cancer type

As a general rule, these few results from the TRM form the basis of sophisticated analyses. For head and neck tumors this is not the case. Therefore the results for head and neck tumors should be interpreted with caution. In part this is due to problems of classification because of limited specific details of locality. Additionally, with advanced tumors in a close topographic location it is often not possible to determine the exact ICD localization of a tumor.

ICD-10 codes used for specifying cancer site

ICD-10	Description
C07	Malignant neoplasm of parotid gland
C08	Malignant neoplasm of other and unspecified major salivary glands
C08.0	Submandibular gland
C08.1	Sublingual gland
C08.8	Overlapping lesion of major salivary glands
C08 9	Major salivary gland unspecified

C08.9 Major salivary gland, unspecified

INCIDENCE

Table 1

Patient cohorts by year of diagnosis including DCO cases and multiple primaries, and with proportion of deaths and active follow-up

Year of	Cases	DCO cases	Prop. DCO	Prop. mult. primaries %	Prop. deaths %	Prop. actively followed
diagnosis	n	n	6	6	6	6
1998	25	6	24.0	28.0	60.0	100.0
1999	22	2	9.1	27.3	63.6	90.9
2000	29	4	13.8	27.6	72.4	96.6
2001	25	5	20.0	16.0	64.0	92.0
2002	46	3	6.5	39.1	60.9	95.7 #
2003	31	3	9.7	29.0	48.4	93.5 #
2004	39	5	12.8	33.3	56.4	100.0 #
2005	41	3	7.3	39.0	58.5	95.1 #
2006	37	1	2.7	27.0	48.6	89.2 #
2007	49	2	4.1	38.8	46.9	85.7 # ##
2008	53	2	3.8	24.5	47.2	64.2
2009	48			43.8	58.3	83.3
2010	62	б	9.7	46.8	41.9	77.4
2011	53	3	5.7	20.8	20.8	69.8
2012	38	1	2.6	28.9	18.4	97.4 ###
1998-2012	598	46	7.7	32.6	49.0	86.6

The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

- ## Since 2007 the percentage of actively followed patients sharply declined compared to the previous years. This is a consequence of ambiguous data protection rules that currently forbid cancer registries in Bavaria to obtain the essential life status informations from competent registration offices.
- ### Please be aware that data of recent annual patient cohorts may not yet be fully processed. Therefore, the presented figures and tables are potentially related to different time periods as pointed out in the respective headlines or legends.

Table 1a

Patient cohorts by year of diagnosis and gender including DCO cases

Year of diagnosis	All n	Males n	Females n	Prop. males %	
1998	25	14	11	56.0	
1999	22	9	13	40.9	
2000	29	16	13	55.2	
2001	25	11	14	44.0	
2002	46	23	23	50.0	
2003	31	15	16	48.4	
2004	39	23	16	59.0	
2005	41	25	16	61.0	
2006	37	20	17	54.1	
2007	49	28	21	57.1	
2008	53	35	18	66.0	
2009	48	28	20	58.3	
2010	62	44	18	71.0	
2011	53	31	22	58.5	
2012	38	16	22	42.1	
1998-2012	598	338	260	56.5	

Table 2

Incidence measures by year of diagnosis and gender including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

Year of diagnosis	Males n	Females n	Males Inc. raw	Fem. Inc. raw	Males Inc. WS	Fem. Inc. WS	Males Inc. ES	Fem. Inc. ES	Males Inc. BRD-S	Fem. Inc. BRD-S
1998	14	11	1.3	0.9	0.8	0.6	1.1	0.7	1.6	0.8
1999	9	13	0.8	1.1	0.5	0.6	0.7	0.7	0.9	0.9
2000	16	13	1.4	1.1	0.8	0.4	1.3	0.7	1.6	0.9
2001	11	14	0.9	1.2	0.6	0.7	0.8	0.9	1.1	1.0
2002	23	23	1.2	1.2	0.7	0.5	1.0	0.7	1.3	1.0
2003	15	16	0.8	0.8	0.5	0.4	0.7	0.6	0.8	0.7
2004	23	16	1.2	0.8	0.7	0.4	1.0	0.6	1.4	0.7
2005	25	16	1.3	0.8	0.7	0.4	1.0	0.6	1.3	0.7
2006	20	17	1.0	0.8	0.6	0.6	0.8	0.7	1.1	0.8
2007	28	21	1.3	0.9	0.7	0.5	1.0	0.6	1.3	0.7
2008	35	18	1.6	0.8	0.9	0.4	1.2	0.5	1.5	0.6
2009	28	20	1.3	0.9	0.7	0.5	0.9	0.6	1.2	0.7
2010	44	18	2.0	0.8	1.0	0.4	1.5	0.5	1.9	0.6
2011	31	22	1.4	0.9	0.7	0.5	1.0	0.7	1.2	0.8
2012	16	22	0.7	0.9	0.3	0.4	0.5	0.6	0.6	0.7
1998-2012	338	260	1.2	0.9	0.7	0.5	1.0	0.6	1.3	0.7

The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

Table 3

Year of	Cases		Std.					Median		
		Meere		Min	Mass	1 0 9.	<u>ог</u> е.	50%	7	0.0.%
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	508	75%	90%
1998	25	63.1	22.1	9.5	97.4	31.9	51.4	62.3	81.9	85.6
1999	22	68.2	18.2	13.9	90.9	54.0	63.6	70.3	80.7	85.3
2000	29	70.6	13.2	48.8	91.9	51.7	60.0	72.7	80.7	88.3
2001	25	65.2	18.5	16.4	95.8	42.4	56.2	65.5	78.6	84.0
2002	46	67.6	14.7	31.4	96.4	46.6	60.1	68.8	78.6	84.6
2003	31	61.7	17.7	22.9	90.5	36.2	51.4	61.8	75.5	81.1
2004	39	66.9	19.2	24.7	94.9	37.7	50.1	70.5	81.9	90.6
2005	41	65.3	15.7	31.9	93.1	43.5	54.3	65.9	79.0	82.0
2006	37	60.1	17.1	21.6	89.7	38.8	45.1	63.5	73.9	83.9
2007	49	64.3	18.4	7.7	92.9	33.2	55.9	67.9	76.1	86.6
2008	53	67.5	17.5	19.8	98.4	47.4	60.1	68.5	81.1	86.4
2009	48	66.6	18.7	16.6	96.1	38.7	61.0	68.8	80.6	85.8
2010	62	67.5	18.7	18.2	95.3	42.7	53.0	71.1	81.0	89.1
2011	53	68.1	17.2	14.4	95.5	46.1	60.8	70.6	78.9	87.1
2012	38	68.7	13.9	37.1	89.7	45.0	60.6	72.3	77.3	85.7
1998-2012	598	66.2	17.4	7.7	98.4	41.8	55.1	69.0	79.0	86.2

Age distribution parameters by year of diagnosis (All) (incl. DCO)

Table 3a

Age distribution parameters by year of diagnosis (MALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	14	62.1	18.0	31.9	85.6	34.1	51.4	60.4	82.4	84.2
1999	9	65.8	16.4	32.0	90.4	32.0	60.9	69.0	72.3	90.4
2000	16	67.1	12.6	48.8	89.7	50.0	58.7	64.8	75.1	88.3
2001	11	66.8	12.4	48.6	84.0	52.0	55.1	65.0	78.8	83.9
2002	23	64.1	12.0	40.1	84.6	46.6	55.0	65.2	73.9	78.6
2003	15	58.3	13.7	29.1	81.1	36.2	51.4	61.3	69.0	72.6
2004	23	66.7	19.0	26.7	92.4	44.8	46.3	70.5	81.9	90.2
2005	25	67.0	14.7	31.9	87.8	43.8	62.2	70.4	77.8	82.0
2006	20	63.5	14.3	39.1	84.3	42.1	54.4	64.0	77.7	81.7
2007	28	63.0	17.1	15.7	84.6	33.1	54.0	69.3	75.2	77.9
2008	35	67.4	14.4	19.8	89.1	49.4	60.1	68.0	78.1	84.3
2009	28	68.6	15.6	16.6	86.8	48.2	63.5	69.9	80.6	85.5
2010	44	68.4	18.2	18.2	95.3	42.7	56.2	71.8	80.8	89.1
2011	31	71.6	15.5	14.4	95.5	55.7	67.2	72.8	82.0	87.1
2012	16	68.4	12.3	45.0	87.6	45.1	61.5	70.9	76.8	82.4
1998-2012	338	66.5	15.5	14.4	95.5	45.5	56.7	68.8	77.9	84.3

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	11	64.4	27.3	9.5	97.4	29.6	42.5	73.6	81.9	93.7
1999	13	69.9	19.8	13.9	90.9	55.1	64.3	75.7	81.6	85.3
2000	13	74.8	13.2	51.7	91.9	52.9	66.4	79.1	84.7	87.6
2001	14	63.9	22.5	16.4	95.8	27.1	56.2	66.7	78.6	90.7
2002	23	71.1	16.5	31.4	96.4	48.2	63.5	71.5	82.4	89.4
2003	16	65.0	20.6	22.9	90.5	30.3	50.1	74.1	79.9	85.2
2004	16	67.1	20.2	24.7	94.9	37.1	57.1	70.1	79.4	93.2
2005	16	62.5	17.4	33.8	93.1	38.9	50.4	60.3	79.6	83.7
2006	17	56.1	19.6	21.6	89.7	26.6	41.3	54.5	69.9	84.0
2007	21	65.9	20.4	7.7	92.9	40.5	58.5	67.5	85.0	88.1
2008	18	67.6	22.8	25.6	98.4	28.7	51.8	74.3	82.8	90.7
2009	20	63.8	22.4	16.8	96.1	25.1	51.3	67.8	81.0	87.4
2010	18	65.3	20.1	27.9	90.4	29.5	46.5	70.7	82.4	89.8
2011	22	63.3	18.7	17.2	94.6	40.3	48.5	66.0	75.3	84.8
2012	22	68.9	15.2	37.1	89.7	44.2	57.2	74.0	77.7	85.7
1998-2012	260	65.9	19.6	7.7	98.4	37.8	54.3	69.4	80.7	88.0

Age at	~						_ 1		
diagnosis	Cases			Males			Females		
Years	n	00	Cum.%	n	olo	Cum.%	n	olo	Cum.%
5-9	2	0.3	0.3			0.0	2	0.8	0.8
10-14	2	0.3	0.7	1	0.3	0.3	1	0.4	1.2
15-19	8	1.3	2.0	4	1.2	1.5	4	1.5	2.7
20-24	3	0.5	2.5			1.5	3	1.2	3.8
25-29	10	1.7	4.2	3	0.9	2.4	7	2.7	6.5
30-34	13	2.2	6.4	8	2.4	4.7	5	1.9	8.5
35-39	13	2.2	8.5	4	1.2	5.9	9	3.5	11.9
40-44	25	4.2	12.7	10	3.0	8.9	15	5.8	17.7
45-49	29	4.8	17.6	21	6.2	15.1	8	3.1	20.8
50-54	41	6.9	24.4	26	7.7	22.8	15	5.8	26.5
55-59	26	4.3	28.8	15	4.4	27.2	11	4.2	30.8
60-64	62	10.4	39.1	41	12.1	39.3	21	8.1	38.8
65-69	78	13.0	52.2	47	13.9	53.3	31	11.9	50.8
70-74	83	13.9	66.1	54	16.0	69.2	29	11.2	61.9
75-79	68	11.4	77.4	40	11.8	81.1	28	10.8	72.7
80-84	62	10.4	87.8	34	10.1	91.1	28	10.8	83.5
85+	73	12.2	100.0	30	8.9	100.0	43	16.5	100.0
All ages	598	100.0		338	100.0		260	100.0	
-									

Age distribution by 5-year age group and gender for period 1998-2012 (incl. DCO)

Table 4

Included in the statistics are 55.9% multiple primaries in males and 34.6% in females.

Table 5

Males Females Males Females Males Females Prop.all Prop.all Age at DCO rate DCO rate cancers cancers Age- Agediagnosis Males Females n=17 n=29 n=146755 n=142297 spec. spec. Years incid. incid. % n n % % % 0- 4 0.0 0.0 5-9 2 0.0 0.2 1.8 10 - 140.1 0.1 0.7 0.6 1 1 15-19 4 0.3 0.3 1.5 4 1.3 20-24 3 0.0 0.2 0.6 25-29 3 7 0.2 0.4 0.3 0.7 30-34 8 5 0.4 0.2 0.6 0.3 35-39 4 9 0.2 0.4 0.2 0.3 40 - 4410 15 0.4 0.7 0.3 0.3 45-49 21 0.4 0.4 0.1 8 1.0 4.8 12.5 50-54 0.1 26 15 1.4 0.8 6.7 0.3 55-59 0.9 0.1 15 11 0.6 0.1 60-64 0.1 41 21 2.5 1.2 4.8 0.2 65-69 47 0.2 31 3.2 1.9 2.1 0.2 70-74 3.7 54 29 4.7 2.1 0.2 0.2 3.4 75-79 40 5.3 2.6 10.0 0.2 28 10.7 0.2 80-84 34 7.5 3.2 14.7 7.1 0.2 28 0.3 9.7 5.2 0.3 85+ 30 43 13.3 46.5 0.3 260 5.0 11.2 0.2 0.2 All ages 338 Incidence 0.9 Raw 1.2 0.7 WS 0.5 ES 1.0 0.6 BRD-S 1.3 0.7

Age-specific incidence, DCO rate and proportion of all cancers for period 1998-2012

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

Table 6a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2012 MALES

Diamagia	Observed	Expected	GTD	LCL	UCL		DCO
Diagnosis	n	n	SIR	95%	95%	EAR	olo
Cl6 Stomach	2	0.5	4.0	0.5	14.4	18.1	
C18 Colon C19-C20 Rectum	2 2	1.2 0.7	1.7 2.9	$0.2 \\ 0.4$	6.1 10.5	10.0 15.9	
C32 Larynx	2	0.1	15.1	1.8		22.6	
C33-C34 Lung	11	1.4	7.7	3.9		115.8	9.1
C61 Prostate	8	3.6	2.2		4.4	53.2	25.0
C67 Bladder	3 2	0.5	5.8	1.2	16.9 #	30.0	
C82-C85 NHL	Z	0.5	4.2	0.5	15.2	18.4	
Other primaries	8	1.3	5.9	2.6	11.7 #	80.4	25.0
Not observed	0	2.4	0.0	0.0	1.5	-29.3	
All mult. primaries	40	12.3	3.3	2.3	4.4 #	335.2	12.5
Patients		2	22				
Mean age at second mal	ignancy (ye		.1				
Person-years Mean observation time	(vearg)		27				
Median observation time							
-	· - · ·						

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries".

Table 6b

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2012 FEMALES

	Observed	Expected		LCL	UCL		DCO
Diagnosis	n	n	SIR	95%	95%	EAR	00
G10 G 1 - -	2	0.0	0 1	0.2		14 0	
C18 Colon	2	0.9	2.1	0.3	7.7	14.2	
C25 Pancreas	2	0.4	5.2	0.6	18.7	21.5	100.0
C33-C34 Lung	6	0.5	11.5	4.2	24.9 #	73.0	50.0
C50 Breast	4	2.3	1.7	0.5	4.5	22.8	25.0
C73 Thyroid	2	0.1	14.8	1.8	53.6 #	24.9	50.0
Other primaries	8	1.1	7.3	3.2	14.4 #	92.1	
Not observed	0	2.9	0.0	0.0	1.3	-38.6	
All mult. primaries	24	8.3	2.9	1.9	4.3 #	209.9	29.2

Patients	178	
Mean age at second malignancy (years)	72.2	
Person-years	750	
Mean observation time (years)	4.2	
Median observation time (years)	3.3	

The occurrence of second malignancy is statistically significant.

Observed second malignancies with count 1 are pooled in category "Other primaries".




Figure 7. Age distribution and age-specific incidence







Figure 7a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).



Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2012, based on the November 2011 submission. http://www.seer.cancer.gov.




C07, C08: Malignant neoplasm of salivary glands

Figure 8. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.






Average incidence (world standard population) 2003 - 2008: Males

Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.7/100,000 WS N=136, females 0.4/100,000 WS N=99). Since cancer data are not available in some counties until 2007, the local incidence rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 1 women were identified with newly diagnosed salivary gland cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.2/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.8/100,000.



Standardized incidence ratio (SIR) 2003 - 2008: Males

Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=136, females N=99). Since cancer data are not available in some counties until 2007, the local SIR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 1 women were identified with newly diagnosed salivary gland cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.34. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 2.54, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts, and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

		Prop.				Prop. deaths
	Incident	actively	Prop.		Prop.	with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	%	olo	n	00	90
1998	25	100.0	24.0	15	60.0	100.0
1999	22	90.9	9.1	14	63.6	100.0
2000	29	96.6	13.8	21	72.4	95.2
2001	25	92.0	20.0	16	64.0	93.8
2002	46	95.7	6.5	28	60.9	96.4
2003	31	93.5	9.7	15	48.4	93.3
2004	39	100.0	12.8	22	56.4	86.4
2005	41	95.1	7.3	24	58.5	100.0
2006	37	89.2	2.7	18	48.6	100.0
2007	49	85.7	4.1	23	46.9	95.7
2008	53	64.2	3.8	25	47.2	96.0
2009	48	83.3		28	58.3	100.0
2010	62	77.4	9.7	26	41.9	100.0
2011	53	69.8	5.7	11	20.8	100.0
2012	38	97.4	2.6	7	18.4	100.0
1998-2012	598	86.6	7.7	293	49.0	96.9

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

			Prop.		
			deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	n	8	n	8
1998	25	16	87.5	7	28.0
1999	22	16	100.0	3	13.6
2000	29	17	100.0	6	20.7
2001	25	13	92.3	5	20.0
2002	46	27	92.6	4	8.7
2003	31	20	95.0	4	12.9
2004	39	31	90.3	10	25.6
2005	41	18	100.0	5	12.2
2006	37	20	95.0	6	16.2
2007	49	23	95.7	3	6.1
2008	53	25	100.0	4	7.5
2009	48	35	100.0	6	12.5
2010	62	33	97.0	8	12.9
2011	53	39	100.0	3	5.7
2012	38	42	95.2	4	10.5
1998-2012	598	375	96.3	78	13.0

Table 10c

Annual cohorts of deaths, proportion of cancer-related and not cancerrelated deaths, and cancer recorded on death certificates (incl. DCO) (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

		Prop. cancer-	Prop. not cancer-	Prop. cancer recorded on death	
Year of	Deaths	related	related	certificate	
death	n	90	90	રુ	
1998	16	43.8	56.3	85.7	
1999	16	50.0	50.0	68.8	
2000	17	94.1	5.9	100.0	
2001	13	69.2	30.8	75.0	
2002	27	63.0	37.0	80.0	
2003	20	80.0	20.0	84.2	
2004	31	77.4	22.6	100.0	
2005	18	50.0	50.0	72.2	
2006	20	65.0	35.0	84.2	
2007	23	65.2	34.8	72.7	
2008	25	84.0	16.0	84.0	
2009	35	74.3	25.7	82.9	
2010	33	87.9	12.1	90.6	
2011	39	61.5	38.5	74.4	
2012	42	69.0	31.0	87.5	
1998-2012	375	70.1	29.9	83.4	

Munich Cancer Registry

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(not cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	4	79.1	75.4	80.4	80.5
1999	8	71.0	75.8	63.0	72.7
2000	7	75.1	75.1		75.1
2001	б	70.8	71.7	66.2	71.7
2002	17	71.9	70.8	73.7	72.7
2003	11	74.5	74.5		75.8
2004	19	71.4	71.9	70.1	69.6
2005	5	76.6	79.5	74.6	81.5
2006	12	74.8	73.9	76.6	74.2
2007	15	80.1	79.1	82.5	78.6
2008	15	67.6	66.9	78.1	64.8
2009	27	71.8	69.4	80.0	70.5
2010	21	70.3	70.3		70.8
2011	27	77.0	73.7	85.0	76.2
2012	22	78.5	75.8	85.4	76.6
1998-2012	216	73.8	72.6	78.0	73.1

Table 11a

Means of age at death according to the grouping in Table 10 $$\rm MALES$$

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(not cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	12	82.4	82.1	82.7	82.4
1999	8	83.5	74.0	89.2	81.8
2000	10	78.1	79.1	70.0	78.1
2001	7	82.2	81.4	83.4	81.4
2002	10	80.0	77.7	83.4	76.7
2003	9	76.9	71.6	83.5	73.9
2004	12	81.1	80.9	82.5	81.8
2005	13	74.3	68.1	81.4	71.3
2006	8	86.3	88.3	83.0	88.3
2007	8	88.3	85.5	91.0	89.2
2008	10	74.1	69.2	85.5	70.6
2009	8	81.6	78.3	87.2	77.3
2010	12	74.7	68.7	86.8	69.5
2011	12	84.4	76.0	90.5	79.8
2012	20	75.3	75.6	74.8	76.1
1998-2012	159	79.6	76.6	84.2	77.8

Table 11b

Means of age at death according to the grouping in Table 10 FEMALES

By 2010, life expectancy for a newborn male in Germany is 77.5 years compared with 82.6 years for his female counterpart.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 12a

Mortality measures (cancer-related death) and mortality-incidence-index by year of death MALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	1	0.1	0.07	0.0	0.05	0.1	0.07	0.1	0.10
1999	5	0.4	0.56	0.3	0.54	0.4	0.58	0.6	0.68
2000	7	0.6	0.44	0.3	0.40	0.6	0.44	0.8	0.52
2001	5	0.4	0.45	0.2	0.43	0.4	0.46	0.6	0.52
2002	11	0.6	0.48	0.3	0.43	0.5	0.49	0.7	0.55
2003	11	0.6	0.73	0.3	0.54	0.5	0.70	0.7	0.92
2004	14	0.7	0.61	0.4	0.56	0.6	0.62	0.9	0.65
2005	2	0.1	0.08	0.0	0.05	0.1	0.07	0.1	0.11
2006	8	0.4	0.40	0.2	0.33	0.3	0.37	0.4	0.40
2007	11	0.5	0.39	0.2	0.27	0.4	0.35	0.6	0.43
2008	14	0.6	0.40	0.3	0.37	0.5	0.40	0.6	0.41
2009	21	0.9	0.75	0.5	0.68	0.7	0.73	0.9	0.72
2010	21	0.9	0.48	0.4	0.43	0.7	0.46	0.9	0.48
2011	19	0.8	0.61	0.4	0.54	0.6	0.59	0.8	0.63
2012	16	0.7	1.00	0.3	0.87	0.5	1.00	0.7	1.01
1998-2012	166	0.6	0.49	0.3	0.42	0.5	0.47	0.7	0.52

Table 12b

Mortality measures (cancer-related death) and mortality-incidence-index by year of death FEMALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	б	0.5	0.55	0.1	0.19	0.2	0.33	0.4	0.47
1999	3	0.3	0.23	0.1	0.20	0.2	0.22	0.2	0.26
2000	9	0.7	0.69	0.2	0.57	0.4	0.60	0.6	0.64
2001	4	0.3	0.29	0.1	0.13	0.2	0.18	0.2	0.21
2002	б	0.3	0.26	0.1	0.22	0.2	0.22	0.2	0.23
2003	5	0.3	0.31	0.1	0.27	0.2	0.28	0.2	0.29
2004	10	0.5	0.63	0.2	0.38	0.3	0.47	0.4	0.54
2005	7	0.4	0.44	0.1	0.34	0.2	0.38	0.3	0.41
2006	5	0.2	0.29	0.0	0.08	0.1	0.13	0.1	0.17
2007	4	0.2	0.19	0.0	0.07	0.1	0.11	0.1	0.17
2008	7	0.3	0.39	0.1	0.36	0.2	0.40	0.2	0.36
2009	5	0.2	0.25	0.1	0.13	0.1	0.17	0.1	0.19
2010	8	0.3	0.44	0.2	0.40	0.2	0.43	0.3	0.46
2011	5	0.2	0.23	0.1	0.12	0.1	0.16	0.2	0.19
2012	13	0.6	0.59	0.2	0.50	0.3	0.51	0.4	0.52
1998-2012	97	0.3	0.37	0.1	0.25	0.2	0.29	0.2	0.33

Table 13

Age at										
death	Cases			Males			Females			
Years	n	00	Cum.%	n	olo	Cum.%	n	00	Cum.%	
30-34	1	0.4	0.4			0.0	1	1.0	1.0	
35-39	0	0.0	0.4			0.0			1.0	
40 - 44	7	2.7	3.0	5	3.0	3.0	2	2.1	3.1	
45-49	11	4.2	7.2	7	4.2	7.2	4	4.1	7.2	
50-54	8	3.0	10.3	7	4.2	11.4	1	1.0	8.2	
55-59	13	4.9	15.2	10	6.0	17.5	3	3.1	11.3	
60-64	18	6.8	22.1	9	5.4	22.9	9	9.3	20.6	
65-69	26	9.9	31.9	20	12.0	34.9	6	6.2	26.8	
70-74	39	14.8	46.8	28	16.9	51.8	11	11.3	38.1	
75-79	46	17.5	64.3	30	18.1	69.9	16	16.5	54.6	
80-84	38	14.4	78.7	25	15.1	84.9	13	13.4	68.0	
85+	56	21.3	100.0	25	15.1	100.0	31	32.0	100.0	
All ages	263	100.0		166	100.0		97	100.0		

Age distribution of age at death (cancer-related) for period 1998-2012 (incl. multiple primaries)

Included in the statistics are 55.9% multiple primaries in males and 34.6% in females.



Males Females Males Females Age at Age-Age-Prop.all Prop.all death Males Females spec. cancers cancers spec. Years mortal. MI-index mortal. MI-index % n n 8 0- 4 0.0 0.0 5-9 0.0 0.0 10 - 140.0 0.0 15-19 0.0 0.0 20 - 240.0 0.0 25-29 0.0 0.0 30-34 1 0.0 0.0 0.20 0.5 35-39 0.0 0.0 2 40 - 445 0.2 0.50 0.1 0.13 0.6 0.2 45-49 7 0.3 0.33 0.2 0.50 0.4 0.2 4 0.07 50-54 7 0.27 0.1 0.2 0.0 1 0.4 55-59 0.27 3 0.2 0.2 0.1 10 0.6 0.67 60-64 9 0.5 0.1 0.1 9 0.5 0.22 0.43 65-69 20 0.4 0.2 0.1 6 1.4 0.43 0.19 70-74 28 0.8 0.2 0.1 2.4 0.52 0.38 11 75-79 30 4.0 1.5 0.2 16 0.75 0.57 0.2 80-84 25 13 5.5 0.74 1.5 0.46 0.1 0.3 31 3.8 0.2 85+ 25 8.1 0.83 0.72 0.3 All ages 97 0.2 0.1 166 Mortality 0.3 Raw 0.6 0.49 0.37 WS 0.3 0.42 0.1 0.25 ES 0.5 0.47 0.2 0.29 BRD-S 0.7 0.52 0.2 0.33 PYLL-70 per 100,000 2.7 1.3 ES 2.3 1.1 AYLL-70 11.4 12.3

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2012 (incl. multiple primaries)

Table 14

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15a

Multiple primaries in deaths in period 1998-2012 $$\rm MALES$$

					Syn- chron	Syn- chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n	i0tai %↓	n	++E ←%	±50u n	+30a ←%	n	JaOi %→
Diagnosis		0↓		ۥ		← 0	11	← 0
C00 Lip	2	1.6	2	100.0				
C03-C06 Oral cavity	/1	0.8	1	100.0				
C09-C10 Oropharynx	3	2.5	2	66.7			1	33.3
C11 Nasopharynx	1	0.8	1	100.0				
C12-C13 Hypopharynx	1	0.8	1	100.0				
C16 Stomach	2	1.6	1	50.0			1	50.0
C18 Colon	2	1.6					2	100.0
C19-C20 Rectum	3	2.5	1	33.3			2	66.7
C22 Liver	3	2.5			2	66.7	1	33.3
C25 Pancreas	2	1.6					2	100.0
C30 Middle/inner ear	2	1.6	2	100.0				
C30-C31 Sinuses	1	0.8					1	100.0
C32 Larynx	2	1.6	1	50.0			1	50.0
C33-C34 Lung	10	8.2			1	10.0	9	90.0
C38,C45 Mesothelioma	1	0.8					1	100.0
C43 Malign. melanoma	2	1.6	1	50.0	_ 1	50.0		
C44 Skin others	47	38.5	24	51.1	6	12.8	17	36.2
C46,C49 Soft tissue	1	0.8					1	100.0
C61 Prostate	14	11.5	11	78.6	2	14.3	1	7.1
C62 Testis	1	0.8	1	100.0				
C64 Kidney	1	0.8					1	100.0
C67 Bladder	7	5.7	4	57.1			3	42.9
C70-C72 CNS cancer	2	1.6	1	50.0			1	50.0
C76-C79 CUP	2	1.6	1	50.0	1	50.0		
C82-C85 NHL	7	5.7	5	71.4	1/	14.3	1	14.3
C91-C96 Leukaemia	2	1.6					2	100.0
All mult. primaries	122	100.0	60	49.2	14	11.5	48	39.3

Multiple primaries with number of cases n<1 are pooled in category "Other primaries".

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 15b

Multiple primaries in deaths in period 1998-2012 FEMALES

					Syn- chron	Syn- chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n	&↓	n	6→	n	~%	n	÷
COO Lip	1 /	1.9			1	100.0		
C12-C13 Hypopharynx	1/	1.9					1	100.0
C16 Stomach	3 1	5.7	1	33.3			2	66.7
C19-C20 Rectum	í	1.9					1	100.0
C23-C24 Bile	1	1.9					1	100.0
C25 Pancreas	3	5.7					3	100.0
C26 GI cancer	/ 1	1.9			1	100.0		
C33-C34 Lung	8	15.1					8	100.0
C44 Skin others	10	18.9	4	40.0			6	60.0
C50 Breast	10	18.9	5	50.0			5	50.0
C54 Corpus uteri	2	3.8	1	50.0	1	50.0		
C56 Ovary	2	3.8	1	50.0			1	50.0
C64 Kidney	2	3.8					2	100.0
C69 Eye carcinoma	1	1.9	1	100.0				
C73 Thyroid	2	3.8	1	50.0			1 /	50.0
C82-C85 NHL	4	7.5	1	25.0			3	75.0
C91-C96 Leukaemia	1	1.9					1	100.0
All mult. primaries	53	100.0	15	28.3	3	5.7	35	66.0

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2012 (Singular primaries only *)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	MI-index	Females Age- spec. mortal.	MI-index	cancers	Females Prop.all cancers %
0- 4			0.0		0.0			
5-9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29			0.0		0.0			
30-34		1	0.0		0.0	0.33		0.5
35-39			0.0		0.0			
40 - 44	5	2	0.2	0.56	0.1	0.14	0.7	0.2
45-49	7	4	0.3	0.33	0.2	0.50	0.5	0.2
50-54	4	1	0.2	0.19	0.1	0.07	0.1	0.0
55-59	10	1	0.6		0.1	0.09	0.2	0.0
60-64	6	9	0.4		0.5	0.45	0.1	0.2
65-69	11	4	0.7		0.2	0.18	0.1	0.1
70-74	18	8	1.6	0.50	0.6	0.33	0.2	0.1
75-79	20	13	2.7		1.2	0.65	0.2	0.2
80-84	17	10	3.7		1.2	0.45	0.2	0.1
85+	11	26	3.5	0.92	3.2	0.70	0.2	0.3
777	100	70					0.0	0 1
All ages	109	79					0.2	0.1
Mortality								
Raw			0.4	0.45	0.3	0.36		
WS			0.2		0.1	0.23		
ES			0.3		0.2	0.28		
BRD-S			0.4	0.48	0.2	0.32		
PYLL-70								
per 100,000			2.3		1.2			
ES			2.0		1.0			
AYLL-70			13.1		13.2			

* See corresponding tables with multiple primaries.



Table 17

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2012 (Single primaries only *)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	MI-index	Females Age- spec. mortal.	MI-index	cancers	Females Prop.all cancers %
0- 4			0.0		0.0			
5-9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29			0.0		0.0			
30-34			0.0		0.0			
35-39		_	0.0		0.0			
40-44	5	2	0.2		0.1	0.18	0.7	0.2
45-49	6	2	0.3	0.30	0.1	0.29	0.4	0.1
50-54	3 7	1	0.2	0.16	0.1	0.08	0.1	0.0
55-59		1 6	0.4		0.1	0.11	0.2	0.0
60-64 65-69	6 8	3	0.4 0.5		0.3 0.2	0.35 0.16	0.1 0.1	0.1 0.1
70-74	。 12	3 7	1.0	0.33	0.2	0.10	0.1	0.1
75-79	12	8	1.0		0.5	0.53	0.1	0.1
80-84	13	6	2.9		0.7	0.32	0.2	0.1
85+	9	23	2.9		2.8	0.52	0.2	0.3
0.01	2	25	2.9	0.75	2.0	0.00	0.2	0.5
All ages	82	59					0.2	0.1
Mortality								
Raw			0.3	0.39	0.2	0.31		
WS			0.2	0.32	0.1	0.19		
ES			0.2	0.37	0.1	0.23		
BRD-S			0.3	0.41	0.1	0.26		
DVIII 70								
PYLL-70			1.9		0 7			
per 100,000			1.9		0.7			
ES AYLL-70			13.6		0.6 12.2			
AIDD-10			13.0		12.2			

* See corresponding tables with multiple primaries.





Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).


The difference between age at diagnosis (Table 3) and age at salivary gland cancer-related death (see Table 10) should be considered.



Average mortality (world standard population) 2003 - 2008: Males

Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2003 to 2008. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.2/100,000 WS N=55, females 0.1/100,000 WS N=38). Since cancer data are not available in some counties until 2007, the local mortality rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 0 women died from salivary gland cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.0/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.4/100,000.






Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=55, females N=38). Since cancer data are not available in some counties until 2007, the local SMR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 0 women died from salivary gland cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.00. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 4.99, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the tumor-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

AYLL-70 BRD-S	Average years of life lost prior to age 70 given a person dies before that age German standard population
DCO	Death certificate only
EAR	Excess absolute risk
	= excess cancer cases (O - E) per 10,000 person-years
ES	European standard population (old)
FRG	Federal Republic of Germany
GEKID	Association of Population-based Cancer Registries in Germany
	(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)
LCL	Lower confidence limit
MI-index	Ratio between mortality and incidence
MCR	Munich Cancer Registry (Tumorregister München)
PYLL-70	Potential years of life lost prior to age 70 given a person dies before that age
SEER	Surveillance, Epidemiology, and End Results (USA)
SIR	Standardized incidence ratio
SMR	Standardized mortality ratio
UCL	Upper confidence limit
WS	World standard population

Recommended Citation

Munich Cancer Registry. Baseline statistics C07, C08: Salivary gland cancer [Internet]. 2014 [updated 2014 Mar 20; cited 2014 May 1]. Available from: http://www.tumorregistermuenchen.de/en/facts/base/base_C0708E.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Munich Cancer Registry

Index of figures and tables

Fig./Tb	l.	Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	4
1a	Gender distribution by year of diagnosis	5
2	Incidence by year of diagnosis	6
3	Age distribution parameters by year of diagnosis	7
4	Age distribution by 5-year age group and gender	9
5	Age-specific incidence and DCO rate	10
6	Standardized incidence ratio of second primaries	11
7	Age distribution and age-specific incidence (chart)	13
7a	Age-specific incidence internationally (chart)	14
8	Cumulative follow-up years (chart)	15
9a	Map of cancer incidence (WS) by county (chart)	16
9b	Standardized incidence ratio (SIR) by county (chart)	17
10a	Pts incident cohorts and mortality / yr	18
10b	Incidence and mortality by year of diagnosis	19
10c	Cancer-related deaths, death certification available / yr	20
11	Means of age at death / yr	21
12	Mortality by year of death	23
13	Distribution of age at death	24
14	Age-specific mortality	25
15	Multiple primaries in deaths	26
16	Age-specific mortality (first primaries)	28
17	Age-specific mortality (single primaries)	29
18	Age distribution and age-specific mortality (chart)	30
19a	Map of cancer mortality (WS) by county (chart)	31
19b	Standardized mortality ratio (SMR) by county (chart)	32