Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage

Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

Cancer statistics: Baseline statistics

C15: Oesophagus cancer

Year of diagnosis	1998-2012
Patients	3,552
Diseases	3,553
Creation date	03/20/2014
Export date	02/12/2014
Population	4.5 m



http://www.tumorregister-muenchen.de/en/facts/base/base_C15__E.pdf

Global Statements about the statistics on the Internet – Baseline Statistics (grey button —), Survival (red button —)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.5 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases**** are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.


The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR. The time-delayed acquisition of data and the occasionally high DCO-rates indicate optimizing reserves, among others, because of current financial and legal conditions that hinder the analyses.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, March 2014

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007). Death certificates from 2013 are incorporated into these analyses.
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- ^{###} DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate. A high proportion of DCO cases (≥5%) in particular cancer types indicate insufficient participation of specific cancer specializations.



ICD-10 codes used for specifying cancer site

ICD-10 Description


Note

Two subclassifications are available:

- .0 .2 according to the anatomical segment
- .3 .5 according to the division into three parts

The deviation from the principle of mutual exclusiveness of particular categories is intentional, because both classifications are in use and although the associated anatomical structures do not match.

C15 C15.0 C15.1 C15.2 C15.3 C15.4 C15.5 C15.8	Malignant neoplasm of oesophagus Cervical part of oesophagus Thoracic part of oesophagus Abdominal part of oesophagus Upper third of oesophagus Middle third of oesophagus Lower third of oesophagus Overlapping lesion of oesophagus
C15.8 C15.9	Overlapping lesion of oesophagus Oesophagus, unspecified



INCIDENCE

Table 1

Patient cohorts by year of diagnosis including DCO cases and multiple primaries, and with proportion of deaths and active follow-up

				Prop.		Prop.
		DCO	Prop.	mult.	Prop.	actively
Year of	Cases	cases	DCO	primaries	deaths	followed
diagnosis	n	'n	%	90	%	%
1998	138	15	10.9	18.1	97.8	99.3
1999	132	9	6.8	18.2	89.4	99.2
2000	130	11	8.5	23.1	90.0	99.2
2001	144	6	4.2	25.7	91.0	100.0
2002	268	31	11.6	25.4	90.3	100.0 #
2003	218	24	11.0	23.9	89.9	99.1 #
2004	218	19	8.7	26.1	88.5	99.1 #
2005	265	21	7.9	29.8	87.5	98.9 #
2006	232	7	3.0	28.4	82.3	98.3 #
2007	292	9	3.1	26.7	83.2	93.2 # ##
2008	284	11	3.9	26.4	77.8	85.2
2009	304	15	4.9	26.0	74.3	88.8
2010	307	16	5.2	20.8	70.4	85.7
2011	337	20	5.9	31.8	66.8	91.4
2012	284	19	6.7	27.8	45.8	96.5 ###
1998-2012	3553	233	6.6	25.9	79.3	94.6

[#] The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

^{##} Since 2007 the percentage of actively followed patients sharply declined compared to the previous years. This is a consequence of ambiguous data protection rules that currently forbid cancer registries in Bavaria to obtain the essential life status informations from competent registration offices.

^{###} Please be aware that data of recent annual patient cohorts may not yet be fully processed. Therefore, the presented figures and tables are potentially related to different time periods as pointed out in the respective headlines or legends.

Table 1a

Patient cohorts by year of diagnosis and gender including DCO cases

Year of	All	Males	Females	Prop. males	
diagnosis	n	n	n	%	
1998	138	107	31	77.5	
1999	132	/111	21	84.1	
2000	130	101	29	77.7	
2001	144	112	32	77.8	
2002	268	212	56	79.1	
2003	218	176	42	80.7	
2004	218	174	44	79.8	
2005	265	218	47	82.3	
2006	232	179	53	77.2	
2007	292	237	55	81.2	
2008	284	230	54	81.0	
2009	304	239	65	78.6	
2010	307	224	83	73.0	
2011	337	278	59	82.5	
2012	284	211	73	74.3	
1000 0010	0.5.5.0				
1998-2012	3553	2809	744	79.1	

base_C15__E.pdf

Table 2

Incidence measures by year of diagnosis and gender including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

			Males	Fem.	Males	Fem.	Males	Fem.	Males	Fem.
Year of	Males	Females	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	107	31	9.7	2.6	6.0	1.2	8.7	1.7	10.4	2.3
1999	111	21	9.9	1.8	6.0	1.0	8.7	1.4	10.2	1.6
2000	101	29	8.9	2.4	5.4	1.3	7.8	1.9	9.6	2.1
2001	112	32	9.7/	2.6	5.9	1.2	8.6	1.8	10.4	2.2
2002	212	56	11.4	2.9	7.0	1.3	9.8	1.9	11.5	2.4
2003	176	42	9.4	2.1	5.6	1.1	7.9	1.6	9.4	1.8
2004	174	44	9.2	2.2	5.4	/1.1	7.7	1.6	9.1	1.8
2005	218	47	11.5	2.4	6.4	1.0	9.3	1.5	11.4	1.9
2006	179	53	9.3	2.6	5.2	1.3	7.5	1.9	9.0	2.2
2007	237	55	10.7	2.4	6.0	1.2	8.7	1.7	10.4	2.0
2008	230	54	10.3	2.3	5.8	1.1	8.3	1.6	10.0	2.0
2009	239	65	10.7	2.8	5.7	1.3	8.3	1.9	10.1	2.3
2010	224	83	9.9	3.5	5.5	1.5	7.9	2.2	9.5	2.7
2011	278	59	12.2	2.5	6.4	1.1	9.2	1.6	11.3	2.1
2012	211	73	9.2	3.1	5.0	1.6	7.2	2.2	8.6	2.5
1998-2012	2809	744	10.2	2.6	5.8	1.2	8.4	1.8	10.0	2.2



The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

Table 3 $\label{eq:Age_distribution_parameters} \mbox{ Age distribution parameters by year of diagnosis (All) } \\ \mbox{ (incl. DCO)}$

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	138	64.3	12.5	35.7	93.8	49.5	56.1	63.0	74.9	80.4
1999	132	63.6	10.4	37.6	89.6	52.6	56.7	61.7	71.7	77.2
2000	130	64.0	11.7	39.6	92.2	49.7	56.0	61.8	72.6	79.7
2001	144	65.4	11,2	38.9	97.2	52.6	57.1	63.7	73.4	82.1
2002	268	65.2	11.7	33.5	95.5	50.1	57.9	64.1	73.5	80.8
2003	218	65.5	11.1	39.0	92.5	50.5	57.8	64.8	73.1	81.3
2004	218	65.8	11.0	36.5	97.2	52.4	59.1	64.9	71.9	80.3
2005	265	66.6	10.7	34.8	96.0	54.1	58.7	66.0	74.8	80.6
2006	232	66.2	9.8	38.4	94.3	54.1	59.6	65.8	72.2	80.3
2007	292	66.1	10.6	33.4	89.9	52.8	59.6	65.9	73.3	80.2
2008	284	67.3	10.5	32.2	96.2	54.1	60.3	66.3	74.4	82.0
2009	304	67.1	10.8	35.6	94.4	52.0	59.3	68.0	74.3	80.7
2010	307	67.1	11.8	32.0	96.3	53.1	59.7	67.2	75.5	83.3
2011	337	68.3	10.4	44.0	94.6	54.5	60.9	68.8	75.5	82.3
2012	284	66.9	10.5	39.2	91.3	51.9	60.2	67.2	73.7	80.6
1998-2012	3553	66.3	11.0	32.0	97.2	52.4	58.7	66.0	73.8	81.2

Table 3a

Age distribution parameters by year of diagnosis (MALES)
(incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	107	62.4	12.1	35.7	93.8	47.9	53.2	60.4	70.9	78.1
1999	111	63.5	10.8	37.6	89.6	51.9	56.2	62.2	71.7	77.2
2000	101	64.1	10.9	39.6	92.2	49.8	56.8	62.5	72.3	78.1
2001	112	64.2	10.7	38.9	97.2	51.5	56.7	62.6	70.3	79.8
2002	212	63.8	11.0	33.5	92.4	50.1	57.3	63.4	71.4	77.5
2003	176	65.1	10.4	39.0	92.5	50.4	57.9	64.9	71.8	78.7
2004	174	64.9	10.6	36.5	94.7	52.3	58.4	64.1	71.5	79.7
2005	218	65.7	10.4	34.8	96.0	53.6	58.1	65.4	74.0	79.6
2006	179	66.0	9.1	38.4	94.3	54.4	59.7	66.0	71.8	77.8
2007	237	65.8	10.4	38.7	89.9	52.6	59.2	66.0	73.2	79.8
2008	230	66.8	10.4	32.2	91.6	53.6	60.1	65.7	73.7	81.6
2009	239	66.7	10.3	35.6	89.0	52.7	59.3	68.0	73.7	80.2
2010	224	65.8	11.5	32.0	91.0	51.6	57.5	65.5	74.1	81.7
2011	278	67.9	10.1	44.0	94.6	54.7	60.9	68.5	74.4	81.8
2012	211	66.7	9.9	39.2	90.1	52.8	60.2	66.5	73.6	78.7
1998-2012	2809	65.6	10.6	32.0	97.2	52.1	58.2	65.5	73.0	79.7

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	31	71.0	11.7	48.1	91.6	56.4	61.4	72.4	81.0	87.2
1999	21	63.9	8.4	52.6	80.1	54.0	58.7	61.5	70.5	74.8
2000	29	63.7	14.1	40.6	89.8	46.8	54.8	58.7	78.1	85.2
2001	32	69.9	12,1	52.6	91.4	54.3	60.4	68.5	81.0	86.3
2002	56	70.6	13.0	44.7	95.5	49.8	61.2	72.2	80.5	87.7
2003	42	67.2	13.5	42.8	92.4	52.6	56.9	63.7	78.8	84.4
2004	44	69.1	11.8	46.4	97.2	56.5	61.9	66.8	75.7	87.2
2005	47	70.8	11.4	40.6	91.4	55.9	62.2	71.3	79.2	85.9
2006	53	66.8	11.8	44.7	92.5	51,6	59.4	65.4	75.9	82.7
2007	55	67.4	11.1	33.4	85.5	52.9	61.6	65.8	78.2	83.6
2008	54	69.5	10.9	46.3	96.2	57.9	61.5	68.1	79.5	83.2
2009	65	68.4	12.5	44.1	94.4	51.4	59.2	67.9	77.9	86.1
2010	83	70.8	11.9	33.3	96.3	57.0	63.7	71.1	78.9	85.8
2011	59	70.4	11.6	47.1	91.5	53.4	60.8	72.0	80.0	83.9
2012	73	67.5	12.1	40.9	91.3	50.5	60.5	68.2	74.1	85.7
1998-2012	744	68.8	12.0	33.3	97.2	52.9	60.4	68.2	78.1	84.9

03/20/2014

Munich Cancer Registry

Table 4

Age distribution by 5-year age group and gender for period 1998-2012 (incl. DCO)

Age at									
diagnosis	Cases			Males			Females		
Years	n	%	Cum.%	n	96	Cum.%	n	%	Cum.%
30-34	6	0.2	0.2	4	0.1	0.1	2	0.3	0.3
35-39	23	0.6	0.8	23	0.8	1.0			0.3
40-44	53	1.5	2.3	40	1.4	2.4	13	1.7	2.0
45-49	155	4.4	6.7	128	4.6	6.9	27	3.6	5.6
50-54	307	8.6	15.3	244	8.7	15.6	63	8.5	14.1
55-59	470	13.2	28.5	400	14.2	29.9	70	9.4	23.5
60-64	634	17.8	46.4	504	17.9	47.8	130	17.5	41.0
65-69	624	17.6	63.9	514	18.3	66.1	110	14.8	55.8
70-74	491	13.8	77.8	399	14.2	80.3	92	12.4	68.1
75-79	364	10.2	88.0	288	10.3	90.6	76	10.2	78.4
80-84	255	7.2	95.2	168	6.0	96.5	87	11.7	90.1
85+	171	4.8	100.0	97	3.5	100.0	74	9.9	100.0
All ages	3553	100.0		2809	100.0		744	100.0	

Included in the statistics are 31.8% multiple primaries in males and 33.6% in females.

Table 5

Age-specific incidence, DCO rate and proportion of all cancers for period 1998-2012

Age at diagnosis Years 0- 4 5- 9 10-14 15-19	Males n	Females n	Age- spec.	Females Age- spec. incid. 0.0 0.0 0.0 0.0 0.0		Females DCO rate n=64 %	cancers	Females Prop.all cancers n=142297 %
20-24			0.0	0.0				
25-29	4	_ /	0.0	0.0			0 0	0 1
30-34	4	2	0.2	0.1			0.3	0.1
35-39	23		1.0	0.0			1.1	
40-44	40	13	1.7	0.6			1.3	0.2
45-49	128	27	5.9	1.3	4.7	3.7	2.6	0.3
50-54	244	63	13.2	3.3	5.3	3.2	3.0	0.6
55-59	400	70	23.5	3.9	2.3	2.9	3.0	0.5
60-64	504	130	30.6	7.5	5.2	0.8	2.5	0.8
65-69	513	110	35.0	6.9	5.1	3.6	2.0	0.6
70-74	399	92	34.4	6.7	5.5	5.4	1.6	0.5
75-79	288	76	38.2	6.9	6.9	9.2	1.5	0.5
80-84	168	87	37.0	10.1	14.3	20.7	1.3	0.6
85+	97	74	31.3	9.0	23.7	32.4	1.1	0.5
03+	91	/ 1	31.3	9.0	23.7	32.4	1.1	0.5
All ages	2808	744			6.0	8.6	1.9	0.5
Incidence								
Raw			10.2	2.6				
WS			5.8	1.2				
ES			8.4	1.8				
BRD-S			10.0	2.2				
BRD B			10.0	2.2				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

Table 6a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2012

MALES

	Observed	Expected		LCL	UCL		DCO
Diagnosis	n	n	SIR	95%	95%	EAR	%
C03-C06 Oral cavity	13	0.5	25.8	13.7	44.1 #	37.4	7.7
C09-C10 Oropharynx	16	0.6	24.9	14.2	40.4 #	45.9	
C12-C13 Hypopharynx	6 /	0.4	16.3	6.0	35.5 #	16.8	
C16 Stomach	/10 /	2.1	4.8	2.3	8.9 #	23.7	10.0
C17 Small intestine	/ 3 /	0.2	12.0	2.5	35.1 #	8.2	
C18 Colon	/ 12 /	4.9	2.4	1.3	4.3 #	21.2	8.3
C19-C20 Rectum	4	3.0	1.3	0.4	3.4	3.1	
C21 Anus/canal	2	0.1	18.1	2.2	65.4 #	5.6	
C22 Liver	9	1.4	6.4	2.9	12.1 #	22.7	33.3
C25 Pancreas	3	1.8	1.7	0.4	5.0	3.7	
C32 Larynx	6	0.6	10.1	3.7	22.0 #	16.2	
C33-C34 Lung	34	6.2	5.5	3.8	7.7 #	83.1	8.8
C50 Breast	3	0.1	23.1	4.8	67.4 #	8.6	66.7
C61 Prostate	25	15.5	1.6	1.0	2.4 #	28.5	16.0
C64 Kidney	6	1.9	3.2	1.2	7.1 #	12.4	
C67 Bladder	6	2.1	2.9	_1.1	6.2 #	11.7	16.7
C76-C79 CUP	2	0.9	2.3	0.3	8.4	3.4	
C82-C85 NHL	4	2.0	2.0	0.6	5.2	6.0	25.0
C90 Mult. myeloma	3	0.6	4.8	1.0	14.0	7.1	33.3
Other primaries	4	3.3	1.2	0.3	3.1	2.2	25.0
Not observed	0	4.1	0.0	0.0	0.9 #	-12.2	
All mult. primaries	171	52.1	3.3	2.8	3.8 #	355.3	11.1

Patients	1918
Mean age at second malignancy (years)	66.3
Person-years	3345
Mean observation time (years)	1.7
Median observation time (years)	0.8

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries".

Table 6b

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2012

FEMALES

	Observed	Expected		LCL	UCL		DCO		
Diagnosis	n	n	SIR	95%	95%	EAR	%		
	/_		<u></u>	\					
C03-C06 Oral cavity	2	0.1			123.9 #				
C09-C10 Oropharynx	5 /	0.0	115.5		269.5 #				
C16 Stomach	2	0.3	6.7	- 1 -		20.5			
C18 Colon	5	0.9			13.5 #		20.0		
C32 Larynx	2 /	0.0	108.0		390.1 #				
C33-C34 Lung	/ 7/	0.7	10.6	4.3					
C43 Malign. melanoma		0.3	9.2		•				
C50 Breast	8	3.0	2.7		5.3 #		12.5		
C76-C79 CUP	2	0.1	13.6	1.7	49.3 #	22.3			
0.1			5 4		11 0 1		16 5		
Other primaries	6	1.1	5.4			58.9	16.7		
Not observed	0	2.7	0.0	0.0	1.4	-32.9			
711 11 1	4.0	0 0	4 6	2.2	c 0 "	1 204 0	п 1		
All mult. primaries	42	9.2	4.6	3.3	6.2 #	394.9	7.1		

Patients	530
Mean age at second malignancy (years)	67.1
Person-years	830
Mean observation time (years)	1.6
Median observation time (years)	0.8

The occurrence of second malignancy is statistically significant.

Observed second malignancies with count 1 are pooled in category "Other primaries".




Figure 7. Age distribution and age-specific incidence



C15: Malignant neoplasm of oesophagus




Figure 7a. Age-specific incidence in MCR registry areas compared to Germany (FRG, GEKID extrapolation) and SEER (Surveillance, Epidemiology, and End Results, USA).



Reference:

Extrapolated age-specific patient population of Germany, data status middle of 2010. Association of Population-based Cancer Registries in Germany (GEKID e.V.). Berlin, 2011. http://www.gekid.de. Last access: 05/12/2011

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2012, based on the November 2011 submission. http://www.seer.cancer.gov.





Figure 8. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.



Average incidence (world standard population) 2003 - 2008: Males



Average incidence (world standard population) 2003 - 2008: Females




Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 5.7/100,000 WS N=1,152, females 1.2/100,000 WS N=285). Since cancer data are not available in some counties until 2007, the local incidence rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 7 women were identified with newly diagnosed oesophagus cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.8/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.2 and 2.4/100,000.

Standardized incidence ratio (SIR) 2003 - 2008: Males




Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=1,152, females N=285). Since cancer data are not available in some counties until 2007, the local SIR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 7 women were identified with newly diagnosed oesophagus cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.83. Though, the value of this parameter may vary with an underlying probability of 99% between 0.24 and 2.03, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts, and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

	Incident	Prop. actively	Prop.		Prop.	Prop. deaths with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	%	%	n	%	%
aragnobib	11		Ů	/ 11 /	0	o
1998	138	99.3	10.9	135	97.8	91.9
1999	132	99.2	6.8	118	89.4	94.9
2000	130	99.2	8.5	117	90.0	95.7
2001	144	100.0	4.2	131	91.0	96.2
2002	268	100.0	11.6	242	90.3	97.5
2003	218	99.1	11.0	196	89.9	98.0
2004	218	99.1	8.7	193	88.5	97.9
2005	265	98.9	7.9	232	87.5	98.3
2006	232	98.3	3.0	191	82.3	98.4
2007	292	93.2	3.1	243	83.2	98.4
2008	284	85.2	3.9	221	77.8	99.1
2009	304	88.8	4.9	226	74.3	99.6
2010	307	85.7	5.2	216	70.4	99.1
2011	337	91.4	5.9	225	66.8	97.8
2012	284	96.5	6.7	130	45.8	96.2
1998-2012	3553	94.6	6.6	2816	79.3	97.6

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

			Prop. deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	n	96	n	8
1998	138	110	91.8	60	43.5
1999	132	106	91.5	38	28.8
2000	130	99	97.0	30	23.1
2001	144	136	94.9	56	38.9
2002	268	219	98.6	/ 111	41.4
2003	218	186	98.4	83	38.1
2004	218	189	97.4	71	32.6
2005	265	214	98.1	94	35.5
2006	232	199	97.0	77	33.2
2007	292	228	97.8	85	29.1
2008	284	217	98.6	78	27.5
2009	304	238	99.2	83	27.3
2010	307	241	98.8	77	25.1
2011	337	286	98.6	119	35.3
2012	284	254	98.8	91	32.0
1998-2012	3553	2922	97.6	1153	32.5

Table 10c

Annual cohorts of deaths, proportion of cancer-related and not cancer-related deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

				Prop. cancer	
		Drop	Prop.	recorded	
		Prop.			
	/	cancer-	not cancer-	on death	
Year of	Deaths	related	related	certificate	
death	n	%	8	%	
1998	110	82.7	17.3	94.1	
1999	106	90.6	9.4	97.9	
2000	99	87.9	12.1	95.8	
2001	136	80.1	19.9	96.9	
2002	219	92.2	7.8	97.2	
2003	186	90.3	9.7	95.6	
2004	189	92.1	7.9	97.3	
2005	214	95.8	4.2	98.6	
2006	199	94.0	6.0	98.4	
2007	228	88.2	11.8	94.2	
2008	217	91.2	8.8	96.3	
2009	238	88.2	11.8	93.2	
2010	241	89.6	10.4	95.4	
2011	286	86.0	14.0	94.3	
2012	254	90.9	9.1	94.8	
1998-2012	2922	89.7	10.3	95.9	

Table 11a $\begin{tabular}{ll} \begin{tabular}{ll} \begin{tabula$

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(not cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	79	64.1	62.7	70.3	64.4
1999	86	65.4	65.0	69.6	65.6
2000	83	64.3	62.6	75.4	63.5
2001	102	65.4	65.5	65.2	65.1
2002	169	66.4	66.0	71.6	66.2
2003	158	65.5	65.5	66.0	65.6
2004	148	65.0	65.1	63.3	65.0
2005	170	66.6	66.2	75.3	66.7
2006	159	67.0	67.0	66.9	67.1
2007	186	67.3	66.6	72.9	66.8
2008	166	68.7	68.3	72.4	68.6
2009	191	68.3	67.9	71.4	68.1
2010	184	67.9	67.7	70.3	67.6
2011	226	69.2	68.2	76.0	68.7
2012	193	68.9	68.5	73.0	68.6
1998-2012	2300	67.1	66.6	71.0	66.9




Table 11b Means of age at death according to the grouping in Table 10 FEMALES

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(not cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	31	71.2	70.2	76.4	71.4
1999	20	71.5	70.3	82.6	70.3
2000	16	67.7	68.4	58.0	67.7
2001	34	71.2	70.2	73.5	72.0
2002	50	71.9	70.8	84.1	71.9
2003	28	69.0	68.2	78.5	68.2
2004	41	68.2	68.1	69.2	68.0
2005	44	69.6	69.6	70.2	69.9
2006	40	72.7	73.3	66.2	73.2
2007	42	69.2	68.1	80.1	68.6
2008	51	68.5	68.5	68.2	68.7
2009	47	69.4	68.8	71.9	69.2
2010	57	72.8	72.3	76.0	72.8
2011	60	72.3	72.6	70.5	72.3
2012	61	70.4	69.9	74.4	70.6
1998-2012	622	70.5	70.1	73.8	70.6



Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 12a $\begin{tabular}{ll} Mortality measures (cancer-related death) and mortality-incidence-index \\ by year of death \\ \hline MALES \\ \end{tabular}$

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	65	5.9	0.61	3.6	0.60	5.2	0.60	6.3	0.60
1999	78	7.0	0.70	4.2	0.69	6.1	0.70	7.4	0.73
2000	72	6.3	0.71	3.9	0.73	5.6	0.73	6.6	0.69
2001	85	7.3	0.76	4.4	0.75	6.5	0.75	7.9	0.75
2002	156	8.4	0.74	5.0	0.71	7.1	0.72	8.5	0.74
2003	142	7.6	0.81	4.4	0.79	6.4	0.81	7.8	0.82
2004	136	7.2	0.78	4.2	0.77	6.0	0.78	7.3	0.80
2005	162	8.6	0.74	4.8	0.75	7.0	0.75	8.5	0.75
2006	150	7.8	0.84	4.3	0.82	6.2	0.83	7.6	0.84
2007	163	7.4	0.69	4.1	0.68	5.9	0.68	7.2	0.69
2008	151	6.8	0.66	3.6	0.62	5.3	0.64	6.5	0.65
2009	171	7.7	0.72	4.0	0.70	5.8	0.70	7.2	0.71
2010	167	7.4	0.75	3.9	0.72	5.7	0.73	7.0	0.74
2011	195	8.5	0.70	4.4	0.69	6.5	0.70	8.0	0.71
2012	177	7.7	0.84	4.0	0.80	5.9	0.82	7.2	0.84
1998-2012	2070	7.5	0.74	4.2	0.72	6.1	0.73	7.4	0.74

Table 12b

Mortality measures (cancer-related death) and mortality-incidence-index by year of death

FEMALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	26	2.2	0.84	1.0	0.85	1.5	0.86	2.0	0.87
1999	18	1.5	0.86	0.7	0.72	1.0	0.72	1.3	0.82
2000	15	1.2	0.52	0.6	0.45	0.9	0.47	1.1	0.53
2001	24	2.0	0.75	0.9	0.72	1.3	0.72	1.6	0.73
2002	46	2.3	0.82	1.1	0.79	1.6	0.83	2.0	0.83
2003	26	1.3	0.62	0.6	0.58	0.9	0.58	1.1	0.61
2004	38	1.9	0.86	0.9	0.85	1.3	0.86	1.6	0.89
2005	43	2.2	0.91	1.0	0.98	1.5	0.98	1.8	0.90
2006	37	1.8	0.70	0.7	0.53	1.1	0.58	1.5	0.66
2007	38	1.6	0.69	0.8	0.65	1.1	0.68	1.4	0.70
2008	47	2.0	0.87	1.0	0.88	1.4	0.86	1.7	0.85
2009	39	1.7	0.60	0.8	0.63	1.2	0.60	1.4	0.61
2010	49	2.1	0.59	0.9	0.57	1.3	0.58	1.6	0.57
2011	51	2.2	0.86	0.9	0.79	1.3	0.80	1.6	0.78
2012	54	2.3	0.74	1.1	0.68	1.5	0.71	1.8	0.73
1998-2012	551	1.9	0.74	0.9	0.71	1.3	0.72	1.6	0.73

Table 13

Age distribution of age at death (cancer-related) for period 1998-2012 (incl. multiple primaries)

Age at									
death	Cases			Males			Females		
Years	n	%	Cum.%	n	96	Cum.%	n	%	Cum.%
30-34	3	0.1	0.1	2	0.1	0.1	1	0.2	0.2
35-39	5	0.2	0.3	5	0.2	0.3			0.2
40 - 44	29	1.1	1.4	27	1.3	1.6	2	0.4	0.5
45-49	108	4.1	5.5	92	4.4	6.1	16	2.9	3.4
50-54	173	6.6	12.1	136	6.6	12.7	37	6.7	10.2
55-59	345	13.2	25.3	296	14.3	26.9	49	8.9	19.1
60-64	457	17.4	42.7	359	17.3	44.3	98	17.8	36.8
65-69	467	17.8	60.5	389	18.8	63.1	78	14.2	51.0
70-74	399	15.2	75.7	324	15.6	78.7	75	13.6	64.6
75-79	280	10.7	86.4	215	10.4	89,1	65	11.8	76.4
80-84	212	8.1	94.5	142	6.9	95.9	70	12.7	89.1
85+	144	5.5	100.0	84	4.1	100.0	60	10.9	100.0
All ages	2622	100.0		2071	100.0		551	100.0	

Included in the statistics are 31.8% multiple primaries in males and 33.6% in females.

Table 14

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2012 (incl. multiple primaries)

Age at			Males Age-		Females Age-		Males Prop.all	Females Prop.all
death	Males	Females	spec.		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0 - 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29			0.0		0.0			
30-34	2	1 <	0.1		0.0	0.50	1.1	0.5
35-39	5		0.2		0.0		1.3	
40-44	27	2	1.1		0.1	0.15	3.3	0.2
45-49	92	16	4.3	0.72	0.8	0.59	5.4	0.9
50-54	136	37	7.4	0.56	2.0	0.59	4.4	1.3
55-59	296	49	17.4		2.8	0.70	5.3	1.1
60-64	359	98	21.8		5.6	0.75	4.3	1.6
65-69	389	78	26.5	0.76	4.9	0.71	3.5	1.0
70-74	324	75	28.0	0.81	5.4	0.82	2.6	0.8
75-79	215	65	28.5	0.75	5.9	0.86	1.8	0.7
80-84	142	70	31.3	0.85	8.1	0.80	1.4	0.7
85+	84	60	27.1	0.87	7.3	0.81	1.0	0.5
All ages	2071	551					2.8	0.8
Mortality								
Raw			7.5		1.9	0.74		
WS			4.2	0.72	0.9	0.71		
ES			6.1	0.73	1.3	0.72		
BRD-S			7.4	0.74	1.6	0.73		
PYLL-70								
per 100,000			51.6		10.8			
ES			45.7		9.2			
AYLL-70			9.8		9.4			

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15a $\begin{tabular}{ll} Multiple primaries in deaths in period 1998-2012 \\ \hline MALES \end{tabular}$

					Syn- chron	Syn- chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n /	%↓	n	← %	n	← %	n	← %
C03-C06 Oral cavity	56	8.1	39	69.6	9	16.1	8	14.3
C09-C10 Oropharynx	78	11.3	52	66.7	9	11.5	17	21.8
C12-C13 Hypopharynx	48	6.9	33	68.8	7	14.6	8	16.7
C16 Stomach	25	3.6	7	28.0	14	56.0	4	16.0
C18 Colon	34	4.9	22	64.7	5	14.7	7	20.6
C19-C20 Rectum	24	3.5	16	66.7	4	16.7	4	16.7
C22 Liver	16	2.3	2	12.5	9	56.3	5	31.3
C25 Pancreas	11	1.6	2	18.2	4	36.4	5	45.5
C32 Larynx	37	5.3	28	75.7	5	13.5	4	10.8
C33-C34 Lung	86	12.4	27	31.4	27	31.4	32	37.2
C43 Malign. melanoma	10	1.4	9	90.0	1	10.0		
C44 Skin others	29	4.2	20	69.0	2	6.9	7	24.1
C61 Prostate	98	14.2	74	75.5	7	7.1	17	17.3
C64 Kidney	19	2.7	13	68.4	1	5.3	5	26.3
C67 Bladder	36	5.2	26	72.2	1	2.8	9	25.0
C73 Thyroid	7	1.0	4	57.1	_ 1	14.3	2	28.6
C76-C79 CUP	12	1.7	7	58.3	4	33.3	1	8.3
C81 Hodgkin lymphoma	6	0.9	6	100.0				
C82-C85 NHL	11	1.6	8	72.7	1	9.1	2	18.2
C91-C96 Leukaemia	10	1.4	4	40.0	3	30.0	3	30.0
Other primaries	39	5.6	20	51.3	5	12.8	14	35.9
All mult. primaries	692	100.0	419	60.5	119	17.2	154	22.3

Multiple primaries with number of cases n<6 are pooled in category "Other primaries".

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 15b $\label{eq:multiple primaries in deaths in period 1998-2012 }$ FEMALES

	_	_			Syn- chron	Syn- chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n /	%↓	n	←%	n	← %	n	-%
G02 G06 O1		- 7	11	100 0				
C03-C06 Oral cavity	11	5.7	11	100.0 66.7	1	16 7	4	16 7
C09-C10 Oropharynx	24	12.5	16		4	16.7	4	16.7
C12-C13 Hypopharynx	8	4.2	3	37.5	5	62.5	1	100 0
C15 Oesophagus	1	0.5				05 0	1	100.0
C16 Stomach	4	2.1	-		1	25.0	3	75.0
C17 Small intestine	2 /	1.0	1	50.0	1	50.0	_	
C18 Colon	/ 11 -	5.7	7	63.6	2	18.2	2	18.2
C19-C20 Rectum	5	2.6	4	80.0	/ 1	20.0		
C21 Anus/canal	2	1.0	1	50.0	1	50.0		
C23-C24 Bile	2	1.0			2	100.0		
C25 Pancreas	4	2.1	2	50.0			2	50.0
C32 Larynx	4	2.1	2	50.0			2	50.0
C33-C34 Lung	11	5.7	3	27.3	2	18.2	6	54.5
C43 Malign. melanoma	5	2.6	4	80.0			1	20.0
C44 Skin others	2	1.0	2	100.0				
C50 Breast	60	31.3	52	86.7	_ 3	5.0	5	8.3
C51 Vulva	1	0.5	1	100.0				
C53 Cervix uteri	5	2.6	4	80.0			1	20.0
C54 Corpus uteri	4	2.1	4	100.0				
C56 Ovary	1	0.5	1	100.0				
C64 Kidney	1	0.5	1	100.0				
C65 Renal pelvis	1	0.5	1	100.0				
C66 Ureter	1	0.5	1	100.0				
C67 Bladder	7	3.6	6	85.7	1	14.3		
C70-C72 CNS cancer	3	1.6	3	100.0				
C73 Thyroid	4	2.1	4	100.0				
C76-C79 CUP	2	1.0	1	50.0	1	50.0		
C82-C85 NHL	3	1.6	2	66.7	1	33.3		
C90 Mult. myeloma	1	0.5					1	100.0
C91-C96 Leukaemia	2	1.0	2	100.0				
All mult. primaries	All mult. primaries 192 100.0 139 72.4 25 13.0 28 14.6							

Multiple primaries with number of cases n<1 are pooled in category "Other primaries".

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2012

(Singular primaries only *)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males	Females			spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
			/					
0 - 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29			0.0		0.0			
30-34	2	1 <	0.1	0.50	0.0/	0.50	1.2	0.5
35-39	4		0.2	0.21	0.0		1.1	
40-44	24	2	1.0	0.62	0.1	0.18	3.2	0.2
45-49	79	10	3.7	0.68	0.5	0.48	5.1	0.6
50-54	111	29	6.0	0.54	1.5	0.58	4.1	1.2
55-59	240	39	14.1	0.72	2.2	0.72	5.0	1.0
60-64	284	74	17.2		4.3	0.75	4.0	1.5
65-69	303	57	20.7		3.6	0.77	3.3	0.9
70-74	257		22.2		3.8	0.78	2.6	0.7
75-79	165	49	21.9		4.5	0.84	1.8	0.6
80-84	103	52	22.7		6.0	0.73	1.4	0.6
85+	61	50	19.7		6.1	0.81	1.0	0.5
	<u> </u>			0.00	0.1	3.32		0.0
All ages	1633	415					2.8	0.8
5							\	
Mortality								
Raw			5.9	0.73	1.4	0.73		
WS			3.3		0.7	0.70		
ES			4.8		1.0	0.71		
BRD-S			5.8		1.2	0.72		
DKD 5			3.0	0.71	1.2	0.72		
PYLL-70								
per 100,000			42.3		8.2			
ES			37.4		7.0			
AYLL-70			10.0		9.5			

^{*} See corresponding tables with multiple primaries.

Table 17

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2012

(Single primaries only *)

			Males		Females		Males	Females
Age at	_	_	Age-		Age-		_	Prop.all
death		Females	_ /		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0- 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29			0.0		0.0			
30-34	2	1 <	0.1	0.50	0.0	1.00	1.2	0.6
35-39	4		0.2	0.21	0.0		1.2	
40-44	23	2	0.9	0.64	0.1	0.18	3.2	0.2
45-49	74	10	3.4	0.67	0.5	0.48	5.2	0.7
50-54	107	28	5.8	0.57	1.5	0.58	4.4	1.3
55-59	229	37	13.5	0.72	2.1	0.74	5.2	1.1
60-64	267	70	16.2	0.71	4.0	0.79	4.3	1.6
65-69	283	53	19.3	0.76	3.3	0.73	3.6	1.0
70-74	237	50	20.5	0.81	3.6	0.78	2.9	0.8
75-79	151	46	20.0	0.74	4.2	0.81	2.0	0.7
80-84	90	49	19.8	0.85	5.7	0.74	1.5	0.7
85+	56	50	18.1	0.80	6.1	0.83	1.1	0.6
711	1500	206					3.0	0.0
All ages	1523	396					3.0	0.8
Mortality								
Raw			5.5	0.73	1.4	0.73		
WS			3.1	0.71	0.6	0.70		
ES			4.5		0.9	0.71		
BRD-S			5.4		1.1	0.72		
PYLL-70								
per 100,000			40.2		7.9			
ES			35.6		6.7			
AYLL-70			10.1		9.6			

^{*} See corresponding tables with multiple primaries.





Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at oesophagus cancer-related death (see Table 10) should be considered.



Average mortality (world standard population) 2003 - 2008: Males




Average mortality (world standard population) 2003 - 2008: Females



Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2003 to 2008. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 4.2/100,000 WS N=864, females 0.8/100,000 WS N=215). Since cancer data are not available in some counties until 2007, the local mortality rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 6 women died from oesophagus cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.7/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.1 and 2.0/100,000.

Standardized mortality ratio (SMR) 2003 - 2008: Males



Standardized mortality ratio (SMR) 2003 - 2008: Females



Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=864, females N=215). Since cancer data are not available in some counties until 2007, the local SMR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 6 women died from oesophagus cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.95. Though, the value of this parameter may vary with an underlying probability of 99% between 0.24 and 2.47, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the tumor-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

BRD-S German standard population

DCO Death certificate only EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

ES European standard population (old) FRG Federal Republic of Germany

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

LCL Lower confidence limit

MI-index Ratio between mortality and incidence

MCR Munich Cancer Registry (Tumorregister München)

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age

SEER Surveillance, Epidemiology, and End Results (USA)

SIR Standardized incidence ratio
SMR Standardized mortality ratio
UCL Upper confidence limit
WS World standard population

Recommended Citation

Munich Cancer Registry. Baseline statistics C15: Oesophagus cancer [Internet]. 2014 [updated 2014 Mar 20; cited 2014 May 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/base/base_C15__E.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

Fig./Tbl	l.	Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	4
1a	Gender distribution by year of diagnosis	5
2	Incidence by year of diagnosis	6
3	Age distribution parameters by year of diagnosis	7
4	Age distribution by 5-year age group and gender	9
5	Age-specific incidence and DCO rate	10
6	Standardized incidence ratio of second primaries	11
7	Age distribution and age-specific incidence (chart)	13
7a	Age-specific incidence internationally (chart)	14
8	Cumulative follow-up years (chart)	15
9a	Map of cancer incidence (WS) by county (chart)	16
9b	Standardized incidence ratio (SIR) by county (chart)	17
10a	Pts incident cohorts and mortality / yr	18
10b	Incidence and mortality by year of diagnosis	19
10c	Cancer-related deaths, death certification available / yr	20
11	Means of age at death / yr	21
12	Mortality by year of death	23
13	Distribution of age at death	24
14	Age-specific mortality	25
15	Multiple primaries in deaths	26
16	Age-specific mortality (first primaries)	28
17	Age-specific mortality (single primaries)	29
18	Age distribution and age-specific mortality (chart)	30
19a	Map of cancer mortality (WS) by county (chart)	31
19b	Standardized mortality ratio (SMR) by county (chart)	32