Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage

Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

Cancer statistics: Baseline statistics

C23-C24: Gallbladder cancer

Year of diagnosis	1998-2012
Patients	3,058
Diseases	3,061
Creation date	03/20/2014
Export date	02/12/2014
Population	4.5 m



http://www.tumorregister-muenchen.de/en/facts/base/base_C2324E.pdf

Global Statements about the statistics on the Internet – Baseline Statistics (grey button —), Survival (red button —)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.5 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases**** are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR. The time-delayed acquisition of data and the occasionally high DCO-rates indicate optimizing reserves, among others, because of current financial and legal conditions that hinder the analyses.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, March 2014

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007). Death certificates from 2013 are incorporated into these analyses.
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- **** DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate. A high proportion of DCO cases (≥5%) in particular cancer types indicate insufficient participation of specific cancer specializations.

ICD-10 codes used for specifying cancer site

ICD-10	Description
C23	Malignant neoplasm of gallbladder
C24,-	Malignant neoplasm of other and unspecified parts of biliary tract
C24.0	Extrahepatic bile duct
C24.1	Ampulla of Vater
C24.8	Overlapping lesion of biliary tract
C24.9	Biliary tract, unspecified

INCIDENCE

and multiple primaries, and with proportion of deaths and active follow-up

Table 1

Patient cohorts by year of diagnosis including DCO cases

				Prop.		Prop.
		DCO	Prop.	mult.	Prop.	actively
Year of	Cases	cases	DCO	primaries	deaths	followed
diagnosis	n	'n	%	%	%	%
1998	106	24	22.6	8.5	91.5	100.0
1999	113	17	15.0	19.5	95.6	100.0
2000	114	27	23.7	11.4	95.6	100.0
2001	125	40	32.0	11.2	89.6	97.6
2002	237	80	33.8	16.9	94.9	100.0 #
2003	229	72	31.4	12.2	93.0	99.6 #
2004	232	60	25.9	13.4	87.5	96.6 #
2005	212	58	27.4	17.5	87.7	98.1 #
2006	249	54	21.7	18.9	88.8	98.0 #
2007	259	47	18.1	20.8	90.7	96.1 # ##
2008	260	60	23.1	21.2	86.9	92.3
2009	254	40	15.7	16.5	81.1	87.4
2010	230	43	18.7	20.4	79.6	89.6
2011	240	35	14.6	23.3	78.8	87.1
2012	201	28	13.9	19.9	59.7	97.0 ###
1998-2012	3061	685	22.4	17.5	86.0	95.3

[#] The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

^{##} Since 2007 the percentage of actively followed patients sharply declined compared to the previous years. This is a consequence of ambiguous data protection rules that currently forbid cancer registries in Bavaria to obtain the essential life status informations from competent registration offices.

^{###} Please be aware that data of recent annual patient cohorts may not yet be fully processed. Therefore, the presented figures and tables are potentially related to different time periods as pointed out in the respective headlines or legends.

Table 1a

Patient cohorts by year of diagnosis and gender including DCO cases

6					
Year of	All	Males	Females	Prop. males	
diagnosis	n	n	n	%	
1998	106	37	69	34.9	
1999	113/	45	68	39.8	
2000	114	53	61	46.5	
2001	125	48	77	38.4	
2002	237	103	134	43.5	
2003	229	91	138	39.7	
2004	232	82	150	35.3	
2005	212	91	121	42.9	
2006	249	96	153	38.6	
2007	259	104	155	40.2	
2008	260	123	137	47.3	
2009	254	126	128	49.6	
2010	230	102	128	44.3	
2011	240	115	125	47.9	
2012	201	90	111	44.8	
1998-2012	3061	1306	1755	42.7	

Table 2

Incidence measures by year of diagnosis and gender including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

			Males	Fem.	Males	Fem.	Males	Fem.	Males	Fem.
Year of	Males	Females	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	37	69	3.3	5.9	1.9	2.5	3.1	3.7	4.5	4.7
1999	45	68	4.0	5.7	2.3	2.3	3.6	3.5	4.6	4.8
2000	53	61	4.7	5.1	2.7	1.6	4.1	2.7	5.6	4.1
2001	48	77	4.1	6.3	2.3	2.6	3.7	4.0	5.2	5.3
2002	103	134	5.5	6.8	3.1	2.3	4.7	3.8	6.0	5.4
2003	91	138	4.9	7.0	2.6	2.5	4.0	3.9	5.4	5.5
2004	82	150	4.4	7.6	2.4	2.8	3.6	4.3	4.7	5.8
2005	91	121	4.8	6.1	2.5	2.3	3.8	3.5	4.9	4.7
2006	96	153	5.0	7.6	2.6	2.5	3.9	4.1	5.1	5.6
2007	104	155	4.7	6.7	2.6	2.5	3.7	3.9	4.7	5.2
2008	123	137	5.5	5.9	2.6	2.1	4.1	3.2	5.6	4.3
2009	126	128	5.6	5.5	2.7	2.0	4.1	3.0	5.6	4.0
2010	102	128	4.5	5.5	2.1	2.0	3.3	3.0	4.4	4.1
2011	115	125	5.0	5.3	2.3	1.8	3.6	2.9	4.9	3.8
2012	90	111	3.9	4.7	1.9	1.7	2.8	2.6	3.7	3.6
1998-2012	1306	1755	4.8	6.1	2.4	2.2	3.7	3.4	4.9	4.7



The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

Table 3 $\label{eq:Age_distribution_parameters} \mbox{ Age distribution parameters by year of diagnosis (All)} \mbox{ (incl. DCO)}$

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	106	73.2	12.7	28.5	99.5	55.5	64.8	74.7	82.6	88.8
1999	113	72.2	12.3	35.3	96.5	56.5	66.8	73.2	79.5	87.0
2000	114	74.9	11.0	44,5	94.0	60.0	66.8	76.2	82.2	88.7
2001	125	72.8	11,9	40.2	99.1	56.5	65.3	74.6	80.5	87.8
2002	237	73.7	11.6	34.1	94.5	57.8	65.9	74.6	82.1	88.1
2003	229	74.3	10.3	37.2	96.8	60.0	66.8	75.0	81.9	87.3
2004	232	73.7	11.8	34.0	100	57.0	67.1	74.2	82.7	88.0
2005	212	72.7	11.6	44.5	98.0	57.5	63.9	72.1	81.8	86.8
2006	249	74.4	12.0	36.7	99.2	57.9	67.4	74.8	83.6	89.2
2007	259	71.9	12.0	35.2	97.1	55.8	64.5	72.4	80.3	87.0
2008	260	74.0	11.6	32.9	99.3	59.1	67.6	74.7	83.1	87.3
2009	254	73.1	11.9	26.5	97.7	56.0	67.0	73.7	81.6	87.7
2010	230	73.2	11.1	43.5	93.8	58.4	65.9	74.3	81.8	87.2
2011	240	73.8	11.5	36.1	100	58.6	66.0	75.2	83.0	88.1
2012	201	73.2	10.9	29.3	99.8	60.4	66.3	74.3	80.4	86.8
1998-2012	3061	73.4	11.6	26.5	100	58.0	66.2	74.3	82.1	87.6

Table 3a

Age distribution parameters by year of diagnosis (MALES)

(incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	37	71.9	10.5	51.0	88.9	55.5	62.2	74.7	79.2	83.6
1999	45	68.1	13.0	35.3	89.8	54.1	59.2	68.7	77.5	86.0
2000	53	71.6	12.1	44.5	94.0	56.4	62.3	72.5	80.5	86.0
2001	48	71.9	11.6	42.1	92.8	56.5	63.0	73.4	79.6	88.0
2002	103	70.3	10.3	44.7	93.5	58.1	62.9	70.0	77.7	85.2
2003	91	72.1	9.7	52.5	95.2	58.9	63.9	72.0	79.7	84.8
2004	82	69.9	11.4	34.0	91.5	54.2	63.9	71.1	76.7	83.7
2005	91	70.0	11.1	47.0	98.0	57.0	63.0	69.4	78.9	86.0
2006	96	69.8	12.0	36.7	94.5	53.8	62.3	69.3	79.0	84.2
2007	104	67.9	11.3	35.2	93.1	53.7	61.4	66.5	76.2	82.6
2008	123	72.3	10.4	37.0	93.1	59.2	65.7	72.6	80.6	84.6
2009	126	71.5	11.2	43.3	97.7	53.4	64.9	72.0	80.0	84.6
2010	102	71.2	10.5	43.5	93.7	58.0	64.2	72.8	78.6	84.2
2011	115	71.4	11.1	38.9	92.1	56.4	64.3	73.6	79.4	84.0
2012	90	71.1	9.5	49.8	93.5	57.6	64.5	72.2	76.2	83.1
1998-2012	1306	70.8	11.0	34.0	98.0	56.5	63.5	71.3	78.9	84.6

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	69	73.9	13.7	28.5	99.5	54.5	65.8	75.2	84.5	90.1
1999	68	75.0	11.1	43.7	96.5	61.6	69.2	75.7	81.9	90.1
2000	61	77.9	9.1	51,1	92.4	66.4	74.0	78.3	82.5	89.7
2001	77	73.3	12.2	40.2	99.1	54.9	65.7	75.7	81.6	87.8
2002	134	76.4	11.8	34.1	94.5	57.8	71.9	79.2	83.6	89.7
2003	138	75.7	10.5	37.2	96.8	61.8	68.1	77.2	83.0	88.1
2004	150	75.7	11.6	44.2	100	60.0	68.5	76.7	84.0	90.1
2005	121	74.7	11.7	44.5	98.0	59.4	65.3	76.5	82.8	89.3
2006	153	77.2	/11.1	43.2	99.2	64.8	71.2	78.5	85.7	91.2
2007	155	74.6	11.7	39.2	97.1	57.9	67.9	75.6	83.0	89.2
2008	137	75.4	12.5	32.9	99.3	59.1	68.9	78.0	84.7	88.1
2009	128	74.7	12.4	26.5	96.6	58.9	67.8	76.2	84.3	88.5
2010	128	74.9	11.4	45.3	93.8	59.1	68.1	75.6	83.7	88.9
2011	125	76.1	11.4	36.1	100	61.1	69.4	76.3	85.2	89.4
2012	111	75.0	11.7	29.3	99.8	62.4	68.2	76.5	83.2	89.6
1998-2012	1755	75.4	11.6	26.5	100	59.9	68.5	76.8	83.9	89.2

Table 4 $\label{eq:Age} \mbox{Age distribution by 5-year age group and gender for period 1998-2012 } \\ (incl. DCO)$

Age at									
diagnosis	Cases			Males			Females		
Years	n	%	Cum.%	n	00	Cum.%	n	%	Cum.%
25-29	3	0.1	0.1			0.0	3	0.2	0.2
30-34	4	0.1	0.2	/ 1	0.1	0.1	3	0.2	0.3
35-39	12	0.4	0.6	/ 7	0.5	0.6	5	0.3	0.6
40-44	24	0.8	1.4	/ 11	0.8	1.5	13	0.7	1.4
45-49	62	2.0	3.4	28	2.1	3.6	34	1.9	3.3
50-54	111	3.6	7.1	59	4.5	8.1	52	3.0	6.3
55-59	173	5.7	12.7	106	8.1	16.2	67	3.8	10.1
60-64	293	9.6	22.3	171	13.1	29.3	122	7.0	17.0
65-69	432	14.1	36.4	221	16.9	46.2	211	12.0	29.1
70-74	487	15.9	52.3	222	17.0	63.2	265	15.1	44.2
75-79	496	16.2	68.5	203	15.5	78.8	293	16.7	60.9
80-84	469	15.3	83.8	157	12.0	90.8	312	17.8	78.6
85+	495	16.2	100.0	120	9.2	100.0	375	21.4	100.0
All ages	3061	100.0		1306	100.0		1755	100.0	

Included in the statistics are 21.9% multiple primaries in males and 19.1% in females.

Table 5

Age-specific incidence, DCO rate and proportion of all cancers for period 1998-2012

Age at diagnosis Years 0- 4	Males n	Females n	Age- spec.	Females Age- spec. incid.		Females DCO rate n=459 %		Females Prop.all cancers n=142297
5- 9			0.0	0.0				
10-14			0.0	0.0				
15-19			0.0	0.0				
20-24			0.0	0.0				
25-29		3	0.0	0.2				0.3
30-34	1	3	0.0	0.1			0.1	0.2
35-39	7	5	0.3	0.2		40.0	0.3	0.1
40 - 44	11	13	0.5	0.6		7.7	0.4	0.2
45-49	28	34	1.3	1.6		8.8	0.6	0.4
50-54	59	52	3.2	2.8	10.2	7.7	0.7	0.5
55-59	106	67	6.2	3.8	5.7	11.9	0.8	0.5
60-64	171	122	10.4	7.0	7.0	9.0	0.8	0.8
65-69	221	211	15.1	13.2	11.8	7.6	0.9	1.2
70-74	222	264	19.2	19.1	16.2	13.3	0.9	1.6
75-79	203	293	26.9	26.8	17.2	24.6	1.1	1.8
80-84	157	312	34.6	36.1	29.9	35.6	1.3	2.1
85+	119	375	38.4	45.8	47.9	52.3	1.3	2.3
All ages	1305	1754			17.2	26.2	0.9	1.2
Incidence								
Raw			4.8	6.1				
WS			2.4					
ES			3.7	3.4				
BRD-S			4.9	4.7				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

Table 6a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2012

MALES

	Observed Ex	spected		LCL UCL		DCO
Diagnosis	'n	n	SIR	95% 95%	EAR	%
C15 Oesophagus	2	0.4	5.1	0.6 18.5	12.7	
C16 Stomach	3	1.0	3.1	0.6 9.0	15.9	33.3
C18 Colon	6	2.3	2.6	1.0 5.7	29.1	16.7
C22 Liver	3 /	0.6	4.8	1.0 14.0	18.7	
C25 Pancreas	3	0.8	3.8	0.8 11.0	17.4	33.3
C33-C34 Lung	3	2.7	1.1	0.2 3.3	2.6	
C43 Malign. melanoma	3	0.8	3.5	0.7 10.3	16.9	
C61 Prostate	10	6.9	1.4	0.7 2.7	24.3	20.0
C67 Bladder	2	1.0	2.0	0.2 7.2	7.9	
C82-C85 NHL	2	0.9	2.3	0.3 8.1	8.8	50.0
Other primaries	4	1.1	3.5	0.9 8.9	22.5	50.0
Not observed	0	4.5	0.0	0.0 0.8 #	-35.4	
All mult. primaries	41	23.0	1.8	1.3 2.4 #	141.4	19.5

Patients	794
Mean age at second malignancy (years)	68.8
Person-years	1270
Mean observation time (years)	1.6
Median observation time (years)	0.8

The occurrence of second malignancy is statistically significant.


Observed second primaries with count 1 are pooled in category "Other primaries".

Table 6b

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2012

FEMALES

	Observed	Expected		LCL	UCL		DCO
Diagnosis	n	n	SIR	95%	95%	EAR	%
C16 Stomach	2	0.9	2.3	0.3	8.2	7.3	100.0
C18 Colon	8	2.3	3.4	1.5	6.7 #	36.8	50.0
C19-C20 Rectum	3 /	1.0	3.1	0.6	8.9	13.2	33.3
C22 Liver	2	0.3	7.9	1.0	28.5	11.4	
C23-C24 Bile	2	0.4	5.7	0.7	20.4	10.7	
C25 Pancreas	/7	1.0	7.1	2.9	14.6 #	39.2	14.3
C33-C34 Lung	5	1.4	3.6	1.2	8.4 #	23.6	
C50 Breast	2	5.6	0.4	0.0	1.3	-23.6	
C56 Ovary	8	0.9	9.2	4.0	18.0 #	46.5	25.0
C64 Kidney	3	0.5	5.6	1.2	16.3 #	16.1	
C82-C85 NHL	5	0.8	6.0	2.0	14.1 #	27.2	40.0
Other primaries	6	1.8	3.3	1.2	7.1 #	27.1	16.7
Not observed	0	3.7	0.0	0.0	1.0 #	-24.4	
All mult. primaries	53	20.6	2.6	1.9	3.4 #	211.0	24.5
_ /							



The occurrence of second malignancy is statistically significant.

Observed second malignancies with count 1 are pooled in category "Other primaries".

C23-C24: Malignant neoplasm of gallbladder and other parts of biliary tract




Figure 7. Age distribution and age-specific incidence



C23-C24: Malignant neoplasm of gallbladder and other parts of biliary tract




Figure 7a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).



Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2012, based on the November 2011 submission. http://www.seer.cancer.gov.







Figure 8. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.



Average incidence (world standard population) 2003 - 2008: Males



Average incidence (world standard population) 2003 - 2008: Females





Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 2.5/100,000 WS N=555, females 2.4/100,000 WS N=801). Since cancer data are not available in some counties until 2007, the local incidence rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 18 women were identified with newly diagnosed gallbladder cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 2.2/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 1.0 and 4.3/100,000.

Standardized incidence ratio (SIR) 2003 - 2008: Males



Standardized incidence ratio (SIR) 2003 - 2008: Females




Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=555, females N=801). Since cancer data are not available in some counties until 2007, the local SIR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 18 women were identified with newly diagnosed gallbladder cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.80. Though, the value of this parameter may vary with an underlying probability of 99% between 0.40 and 1.42, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts, and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

		Prop.				Prop. deaths
	Incident	actively	Prop.		Prop.	with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	%	%	/ n /	%	%
1998	106	100.0	22.6	97	91.5	93.8
1999	113	100.0	15.0	108	95.6	96.3
2000	114	100.0	23.7	109	95.6	97.2
2001	125	97.6	32.0	112	89.6	96.4
2002	237	100.0	33.8	225	94.9	97.8
2003	229	99.6	31.4	213	93.0	98.6
2004	232	96.6	25.9	203	87.5	97.5
2005	212	98.1	27.4	186	87.7	98.9
2006	249	98.0	21.7	221	88.8	98.6
2007	259	96.1	18.1	235	90.7	98.7
2008	260	92.3	23.1	226	86.9	99.6
2009	254	87.4	15.7	206	81.1	99.0
2010	230	89.6	18.7	183	79.6	99.5
2011	240	87.1	14.6	189	78.8	98.4
2012	201	97.0	13.9	120	59.7	95.0
1998-2012	3061	95.3	22.4	2633	86.0	98.1

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

			Prop. deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	n	%	n	ૄ
1998	106	94	93.6	57	53.8
1999	113	104	94.2	54	47.8
2000	114	121	95.9	62	54.4
2001	125	114	97.4	55	44.0
2002	237	191	99.0	146	61.6
2003	229	148	96.6	115	50.2
2004	232	155	98.7	108	46.6
2005	212	165	97.6	96	45.3
2006	249	201	98.0	125	50.2
2007	259	191	98.4	102	39.4
2008	260	202	98.0	123	47.3
2009	254	208	98.6	100	39.4
2010	230	208	100.0	94	40.9
2011	240	206	99.5	100	41.7
2012	201	212	98.1	91	45.3
1998-2012	3061	2520	97.9	1428	46.7

Table 10c

Annual cohorts of deaths, proportion of cancer-related and not cancer-related deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

				Prop. cancer	
		Prop.	Prop.	recorded	
		cancer-	not cancer-	on death	
Year of	Deaths	related	related	certificate	
death	n	%	%	%	
1998	94	87.2	12.8	95.5	
1999	104	80.8	19.2	94.9	
2000	121	91.7	8.3	98.3	
2001	114	91.2	8.8	96.4	
2002	191	90.6	9.4	93.7	
2003	148	89.2	10.8	95.8	
2004	155	90.3	9.7	96.1	
2005	165	91.5	8.5	95.7	
2006	201	87.6	12.4	91.9	
2007	191	93.2	6.8	95.7	
2008	202	94.6	5.4	96.5	
2009	208	88.9	11.1	93.2	
2010	208	90.9	9.1	94.7	
2011	206	91.3	8.7	95.6	
2012	212	87.3	12.7	92.8	
1998-2012	2520	90.0	10.0	94.9	

Table 11a $\begin{tabular}{ll} Means of age at death according to the grouping in Table 10 \\ \hline MALES \end{tabular}$

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(not cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	36	73.5	73.2	74.7	73.7
1999	36	72.3	71.1	75.5	70.8
2000	42	73.9	74.1	72.2	74.0
2001	52	71.7	71.2	74.9	71.8
2002	81	71.6	71.0	76.2	71.3
2003	61	73.7	73.4	75.9	74.2
2004	59	71.8	71.5	74.3	71.7
2005	84	70.0	69.8	71.9	69.9
2006	78	73.3	72.8	76.4	73.3
2007	68	69.4	69.3	71.2	70.0
2008	77	72.0	71.9	73.5	71.8
2009	103	71.1	71.1	71.2	71.5
2010	89	73.3	72.7	79.7	73.1
2011	106	72.6	72.5	73.2	72.4
2012	96	72.1	71.6	76.1	72.0
1998-2012	1068	72.0	71.7	74.7	72.0




Table 11b Means of age at death according to the grouping in Table 10 FEMALES

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(not cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	58	74.8	73.9	82.5	74.6
1999	68	76.7	76.3	79.3	76.8
2000	79	76.7	76.2	81.6	76.6
2001	62	77.7	77.1	89.1	77.8
2002	110	76.0	75.7	79.5	76.0
2003	87	75.7	75.2	80.2	75.9
2004	96	77.2	76.9	79.5	77.1
2005	81	75.8	75.0	84.3	75.5
2006	123	78.0	76.8	86.1	76.9
2007	123	76.5	76.0	84.4	76.1
2008	125	75.9	75.4	85.2	76.0
2009	105	78.5	77.5	86.2	77.7
2010	119	76.6	75.9	82.6	76.2
2011	100	75.6	75.1	79.9	75.4
2012	116	76.9	76.1	82.1	76.1
1998-2012	1452	76.6	76.0	82.6	76.3



Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 12a $\begin{tabular}{ll} Mortality measures (cancer-related death) and mortality-incidence-index \\ by year of death \\ \hline MALES \\ \end{tabular}$

Year of					MI-Index		MI-Index ES		MI-Index
death	n	raw	raw	WS	WS	ES	ŁS	BRD-S	BRD-S
1998	30	2.7	0.81	1.5	0.82	2.5	0.82	3.9	0.86
1999	26	2.3	0.51	1.3	0.52	2.1	0.52	2.7	0.59
2000	39		0.75		0.72		0.77		
		3.4	- 7	1.9		3.1		4.3	0.79
2001	45	3.9	0.94	2.2	0.95	3.5	0.93	4.6	0.88
2002	71	3.8	0.69	2.1	0.67	3.2	0.69	4.3	0.71
2003	54	2.9	0.59	1.5	0.58	2.4	0.59	3.3	0.61
2004	54	2.9	0.66	1.5	0.64	2.4	0.65	3.2	0.67
2005	77	4.1	0.85	2.2	0.85	3.3	0.85	4.2	0.85
2006	68	3.6	0.71	1.8	0.68	2.7	0.70	3.7	0.72
2007	63	2.8	0.61	1.5	0.57	2,1	0.58	2.7	0.58
2008	72	3.2	0.59	1.6	0.60	2.5	0.60	3.4	0.60
2009	92	4.1	0.73	2.0	0.76	3.1	0.75	4.1	0.74
2010	82	3.6	0.80	1.7	0.77	2.6	0.79	3.5	0.79
2011	98	4.3	0.85	1.9	0.80	3.0	0.83	4.1	0.83
2012	84	3.7	0.93	1.7	0.93	2.6	0.94	3.6	0.96
1998-2012	955	3.5	0.73	1.7	0.72	2.7	0.73	3.6	0.74

Table 12b

Mortality measures (cancer-related death) and mortality-incidence-index by year of death

FEMALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	52	4.4	0.75	1.8	0.71	2.7	0.74	3.6	0.75
1999	58	4.9	0.85	1.8	0.77	2.8	0.79	3.9	0.82
2000	72	6.0	1.18	2.1	1.33	3.4	1.25	4.8	1.16
2001	59	4.9	0.77	1.7	0.63	2.8	0.69	4.0	0.75
2002	102	5.2	0.76	1.9	0.80	2.9	0.77	4.1	0.76
2003	78	4.0	0.57	1.5	0.59	2.3	0.59	3.2	0.58
2004	86	4.4	0.57	1.5	0.54	2.4	0.56	3.4	0.58
2005	74	3.7	0.61	1.3	0.56	2.0	0.57	2.9	0.62
2006	108	5.4	0.71	1.8	0.72	2.9	0.71	4.0	0.71
2007	115	5.0	0.74	1.7	0.69	2.8	0.71	3.8	0.74
2008	119	5.1	0.87	1.8	0.85	2.8	0.87	3.8	0.90
2009	93	4.0	0.73	1.2	0.60	2.0	0.66	2.9	0.72
2010	107	4.6	0.84	1.6	0.80	2.4	0.81	3.3	0.80
2011	90	3.8	0.73	1.4	0.76	2.1	0.73	2.8	0.73
2012	101	4.3	0.91	1.4	0.84	2.3	0.88	3.1	0.88
1998-2012	1314	4.6	0.75	1.6	0.72	2.5	0.73	3.5	0.75

Table 13

Age distribution of age at death (cancer-related) for period 1998-2012 (incl. multiple primaries)

Age at									
death	Cases			Males			Females		
Years	n	%	Cum.%	n	96	Cum.%	n	%	Cum.%
25-29	2	0.1	0.1			0.0	2	0.2	0.2
30-34	2	0.1	0.2	/ 1	0.1	0.1	1	0.1	0.2
35-39	6	0.3	0.4	2	0.2	0.3	4	0.3	0.5
40-44	10	0.4	0.9	/ 7	0.7	1.0	3	0.2	0.8
45-49	38	1.7	2.6	12	1.3	2.3	26	2.0	2.7
50-54	68	3.0	5.5	38	4.0	6.3	30	2.3	5.0
55-59	138	6.1	11.6	78	8.2	14.4	60	4.6	9.6
60-64	195	8.6	20.2	118	12.3	26.8	77	5.9	15.4
65-69	291	12.8	33.0	147	15.4	42.1	144	11.0	26.4
70-74	367	16.2	49.2	175	18.3	60.4	192	14.6	41.0
75-79	396	17.4	66.6	153	16.0	76.4	243	18.5	59.5
80-84	360	15.8	82.4	118	12.3	88.7	242	18.4	77.9
85+	399	17.6	100.0	108	11.3	100.0	291	22.1	100.0
All ages	2272	100.0		957	100.0		1315	100.0	

Included in the statistics are 21.9% multiple primaries in males and 19.1% in females.

Table 14

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2012 (incl. multiple primaries)

			Males		Females		Males	Females
Age at	_	_	Age-		Age-		_	Prop.all
death		Females	_ /		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0- 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29		2	0.0		0.1	0.67		1.8
30-34	1	1	0.0	1.00	0.0	0.33	0.6	0.5
35-39	2	4	0.1		0.2	0.80	0.5	0.8
40-44	7	3	0.3		0.1	0.23	0.9	0.3
45-49	12	26	0.6	0.43	1.2		0.7	1.4
50-54	38	30	2.1		1.6	0.58	1.2	1.0
55-59	78	60	4.6		3.4		1.4	1.3
60-64	118	77	7.2		4.4		1.4	1.3
65-69	147	144	10.0		9.0	0.68	1.3	1.9
70-74	175	192	15.1	0.79	13.9		1.4	2.1
75-79	153	243	20.3	0.75	22.2	0.83	1.3	2.5
80-84	118	242	26.0	0.75	28.0	0.78	1.2	2.3
85+	108	291	34.8	0.90	35.5	0.78	1.3	2.3
All ages	957	1315					1.3	2.0
Mortality								
Raw			3.5		4.6	0.75		
WS			1.7	0.72	1.6	0.72		
ES			2.7	0.73	2.5			
BRD-S			3.6	0.74	3.5	0.75		
PYLL-70								
per 100,000			14.0		12.8			
ES			12.3		10.8			
AYLL-70			8.6		9.0			
-								

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15a $\begin{tabular}{ll} Multiple primaries in deaths in period 1998-2012 \\ \hline MALES \end{tabular}$

	Total	Total	Pre	Pre	Syn- chron ±30d	Syn- chron ±30d	Post	Post
Diagnosis	n /	%↓	n	-%	n	±30α ←%	n	-%
C03-C06 Oral cavity	2	0.9	1	50.0			1	50.0
C12-C13 Hypopharynx	2	0.9	1	50.0			1	50.0
C15 Oesophagus	/ 2	0.9			1	50.0	1	50.0
C16 Stomach	/ 11	5.0	7	63.6	2	18.2	2	18.2
C17 Small intestine	/ 3	1.4	1	33.3	1	33.3	1	33.3
C18 Colon	20 /	9.1	14	70.0	4	20.0	2	10.0
C19-C20 Rectum	/ 12	5.5	12	100.0				
C22 Liver	3	1.4			2	66.7	1	33.3
C23-C24 Bile	3	1.4			1	33.3	2	66.7
C25 Pancreas	7	3.2			3	42.9	4	57.1
C26 GI cancer	2	0.9			1	50.0	1	50.0
C32 Larynx	5	2.3	4	80.0	1	20.0		
C33-C34 Lung	10	4.5	3	30.0	2	20.0	5	50.0
C43 Malign. melanoma	9	4.1	6	66.7			3	33.3
C44 Skin others	10	4.5	5	50.0	1	10.0	4	40.0
C61 Prostate	58	26.4	45	77.6	_ 4	6.9	9	15.5
C62 Testis	5	2.3	5	100.0				
C64 Kidney	11	5.0	8	72.7	1	9.1	2	18.2
C67 Bladder	17	7.7	13	76.5	2	11.8	2	11.8
C69 Eye melanoma	2	0.9	1	50.0			1	50.0
C70-C72 CNS cancer	3	1.4	2	66.7	1	33.3		
C76-C79 CUP	4	1.8	2	50.0	2	50.0		
C82-C85 NHL	5	2.3	2	40.0	1	20.0	2	40.0
C90 Mult. myeloma	3	1.4					3	100.0
Other primaries	11	5.0	6	54.5	3	27.3	2	18.2
All mult. primaries	220	100.0	138	62.7	33	15.0	49	22.3

Multiple primaries with number of cases n<2 are pooled in category "Other primaries".

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

base_C2324E.pdf

Table 15b

Multiple primaries in deaths in period 1998-2012
FEMALES

Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C03-C06 Oral cavity	3	1.3	3	100.0				
C07-C08 Salivary gland	/2	0.8	2	100.0				
C16 Stomach	6	2.5	1	16.7	1	16.7	4	66.7
C17 Small intestine	2	0.8	1	50.0			1	50.0
C18 Colon	26	10.9	15	57.7	2	7.7	9	34.6
C19-C20 Rectum	12 /	5.0	7	58.3	3	25.0	2	16.7
C22 Liver	/ 3 4	1.3	1	33.3	1	33.3	1	33.3
C25 Pancreas	7	2.9	2	28.6	4	57.1	1	14.3
C33-C34 Lung	8	3.4	3	37.5	1	12.5	4	50.0
C43 Malign. melanoma	11	4.6	10	90.9			1	9.1
C44 Skin others	8	3.4	4	50.0	3	37.5	1	12.5
C50 Breast	58	24.4	55	94.8	2	3.4	1	1.7
C51 Vulva	3	1.3	3	100.0				
C52 Vagina	2	0.8	1	50.0	1	50.0		
C53 Cervix uteri	3	1.3	2	66.7			1	33.3
C54 Corpus uteri	17	7.1	17	100.0				
C56 Ovary	16	6.7	5	31.3	3	18.8	8	50.0
C64 Kidney	11	4.6	8	72.7	3	27.3		
C67 Bladder	9	3.8	9	100.0				
C70-C72 CNS cancer	3	1.3	2	66.7	1	33.3		
C73 Thyroid	2	0.8	2	100.0				
C76-C79 CUP	5	2.1	2	40.0	2	40.0	1	20.0
C82-C85 NHL	10	4.2	7	70.0	2	20.0	1	10.0
C91-C96 Leukaemia	2	0.8	1	50.0			1	50.0
Other primaries	9	3.8	3	33.3	4	44.4	2	22.2
All mult. primaries	238	100.0	166	69.7	33	13.9	39	16.4

Multiple primaries with number of cases n<2 are pooled in category "Other primaries".

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2012

(Singular primaries only *)

			Males		Females		Males	Females
Age at			Age-		Age-		_	Prop.all
death		Females	_ /		spec.	! 1	cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0- 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29		2	0.0		0.1	0.67		1.9
30-34	1	1	0.0	1.00	0.0	0.33	0.6	0.5
35-39	2	4	0.1		0.2	0.80	0.6	0.9
40-44	7	3	0.3		0.1	0.23	0.9	0.3
45-49	12	25	0.6	0.44	1.2		0.8	1.5
50-54	35	28	1.9	0.66	1.5	0.57	1.3	1.2
55-59	70	53	4.1	0.74	3.0	0.93	1.5	1.4
60-64	95	69	5.8	0.65	4.0	0.63	1.4	1.4
65-69	132	124	9.0	0.67	7.7	0.73	1.4	2.0
70-74	147	164	12.7	0.79	11.9	0.75	1.5	2.3
75-79	125	205	16.6	0.77	18.7	0.82	1.4	2.6
80-84	95	198	20.9	0.77	22.9	0.76	1.3	2.4
85+	80	265	25.8	0.84	32.4	0.82	1.3	2.6
- 7 7	0.01	1 1 4 1					1 1	0 1
All ages	801	1141					1.4	2.1
Mortality								
Raw			2.9	0.73	4.0	0.76		
WS			1.5	0.73	1.4			
ES			2.3		2.2			
BRD-S			3.0	0.74	3.0	0.76		
BRD B			3.0	0.71	3.0	0.70		
PYLL-70								
per 100,000			12.5		11.7			
ES			11.0		10.0			
AYLL-70			8.7		9.3			

^{*} See corresponding tables with multiple primaries.

Table 17 Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2012 (Single primaries only *)

Age at death	Males	Females	Males Age- spec.		Females Age- spec.		Males Prop.all cancers	Females Prop.all cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0 - 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29		2	0.0		0.1	0.67		2.1
30-34	1	1	0.0	1.00	0.0	0.33	0.6	0.6
35-39	2	4	0.1	0.40	0.2	0.80	0.6	1.0
40-44	7	2	0.3		0.1	0.15	1.0	0.2
45-49	12	25	0.6	0.48	1.2	0.76	0.8	1.7
50-54	30	27	1.6	0.57	1.4	0.57	1.2	1.2
55-59	68	49	4.0	0.76	2.8	0.88	1.6	1.5
60-64	92	69	5.6	0.66	4.0	0.64	1.5	1.6
65-69	129	122	8.8	0.69	7.6	0.73	1.6	2.3
70-74	139	159	12.0	0.77	11.5		1.7	2.6
75-79	118	195	15.7		17.8		1.6	2.9
80-84	92	192	20.3		22.2	0.75	1.6	2.8
85+	78	260	25.2	0.83	31.7	0.81	1.6	3.0
777	7.60	1107					1 -	2 4
All ages	768	1107					1.5	2.4
Mortality								
Raw			2.8	0.72	3.9	0.76		
WS			1.4		1.3	0.73		
ES			2.2		2.1	0.74		
BRD-S			2.2	0.72	2.9			
DKD 5			۵.۶	0.75	۵. ۶	0.75		
PYLL-70								
per 100,000			11.9		11.3			
ES			10.4		9.6			
AYLL-70			8.7		9.2			

^{*} See corresponding tables with multiple primaries.







Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at gallbladder cancer-related death (see Table 10) should be considered.



Average mortality (world standard population) 2003 - 2008: Males



Average mortality (world standard population) 2003 - 2008: Females




Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2003 to 2008. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 1.7/100,000 WS N=374, females 1.6/100,000 WS N=553). Since cancer data are not available in some counties until 2007, the local mortality rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 15 women died from gallbladder cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 1.8/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.8 and 3.6/100,000.

Standardized mortality ratio (SMR) 2003 - 2008: Males




Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=374, females N=553). Since cancer data are not available in some counties until 2007, the local SMR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 15 women died from gallbladder cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.96. Though, the value of this parameter may vary with an underlying probability of 99% between 0.44 and 1.81, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the tumor-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

BRD-S German standard population

DCO Death certificate only EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

ES European standard population (old) FRG Federal Republic of Germany

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

LCL Lower confidence limit

MI-index Ratio between mortality and incidence

MCR Munich Cancer Registry (Tumorregister München)

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age

SEER Surveillance, Epidemiology, and End Results (USA)

SIR Standardized incidence ratio
SMR Standardized mortality ratio
UCL Upper confidence limit
WS World standard population

Recommended Citation

Munich Cancer Registry. Baseline statistics C23-C24: Gallbladder cancer [Internet]. 2014 [updated 2014 Mar 20; cited 2014 May 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/base/base_C2324E.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

Fig./Tbl	l.	Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	3
1a	Gender distribution by year of diagnosis	4
2	Incidence by year of diagnosis	5
3	Age distribution parameters by year of diagnosis	6
4	Age distribution by 5-year age group and gender	8
5	Age-specific incidence and DCO rate	9
6	Standardized incidence ratio of second primaries	10
7	Age distribution and age-specific incidence (chart)	12
7a	Age-specific incidence internationally (chart)	13
8	Cumulative follow-up years (chart)	14
9a	Map of cancer incidence (WS) by county (chart)	15
9b	Standardized incidence ratio (SIR) by county (chart)	16
10a	Pts incident cohorts and mortality / yr	17
10b	Incidence and mortality by year of diagnosis	18
10c	Cancer-related deaths, death certification available / yr	19
11	Means of age at death / yr	20
12	Mortality by year of death	22
13	Distribution of age at death	23
14	Age-specific mortality	24
15	Multiple primaries in deaths	25
16	Age-specific mortality (first primaries)	27
17	Age-specific mortality (single primaries)	28
18	Age distribution and age-specific mortality (chart)	29
19a	Map of cancer mortality (WS) by county (chart)	30
19b	Standardized mortality ratio (SMR) by county (chart)	31