

Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage

Munich Cancer Registry at Munich Cancer Center
Marchioninstr. 15
Munich, 81377
Germany

<http://www.tumorregister-muenchen.de/en>

Cancer statistics: Baseline statistics

C52: Vaginal cancer

Year of diagnosis	1998-2012
Patients	253
Diseases	253
Creation date	03/20/2014
Export date	02/12/2014
Population (females)	2.3 m



http://www.tumorregister-muenchen.de/en/facts/base/base_C52__E.pdf

**Global Statements about the statistics on the Internet –
Baseline Statistics** (grey button ) , **Survival** (red button )

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.5 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases^{###} are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR. The time-delayed acquisition of data and the occasionally high DCO-rates indicate optimizing reserves, among others, because of current financial and legal conditions that hinder the analyses.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, March 2014

- [#] Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007). Death certificates from 2013 are incorporated into these analyses.
- ^{##} Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- ^{###} DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate. A high proportion of DCO cases ($\geq 5\%$) in particular cancer types indicate insufficient participation of specific cancer specializations.

INCIDENCE

Table 1

Patient cohorts by year of diagnosis including DCO cases and multiple primaries, and with proportion of deaths and active follow-up

Year of diagnosis	Cases n	DCO cases n	Prop. DCO %	Prop. mult. primaries %	Prop. deaths %	Prop. actively followed %
1998	12			33.3	75.0	91.7
1999	7	1	14.3	28.6	85.7	100.0
2000	10			10.0	30.0	100.0
2001	12	1	8.3	41.7	75.0	100.0
2002	11	1	9.1	45.5	81.8	100.0 #
2003	20	1	5.0	25.0	80.0	100.0 #
2004	19	4	21.1	15.8	78.9	100.0 #
2005	15	1	6.7	20.0	46.7	100.0 #
2006	18	1	5.6	33.3	77.8	88.9 #
2007	28			39.3	50.0	89.3 # ##
2008	13	2	15.4	15.4	46.2	76.9
2009	23	3	13.0	52.2	52.2	69.6
2010	24	2	8.3	41.7	58.3	70.8
2011	25	1	4.0	20.0	48.0	88.0
2012	16	1	6.3	25.0	31.3	93.8 ###
1998-2012	253	19	7.5	30.8	59.7	89.3

The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

Since 2007 the percentage of actively followed patients sharply declined compared to the previous years. This is a consequence of ambiguous data protection rules that currently forbid cancer registries in Bavaria to obtain the essential life status informations from competent registration offices.

Please be aware that data of recent annual patient cohorts may not yet be fully processed. Therefore, the presented figures and tables are potentially related to different time periods as pointed out in the respective headlines or legends.

Table 2

Incidence measures by year of diagnosis and gender including DCO cases
(with respect to registry area expansion from 2.51 to 3.96 m as of 2002,
and from 3.96 to 4.52 m as of 2007, respectively)

Year of diagnosis	Cases n	Incidence raw	Incidence WS	Incidence ES	Incidence BRD-S
1998	12	1.0	0.5	0.7	0.9
1999	7	0.6	0.2	0.4	0.5
2000	10	0.8	0.5	0.6	0.8
2001	12	1.0	0.5	0.7	0.9
2002	11	0.6	0.2	0.4	0.5
2003	20	1.0	0.4	0.6	0.8
2004	19	1.0	0.6	0.7	0.8
2005	15	0.8	0.4	0.5	0.6
2006	18	0.9	0.3	0.5	0.7
2007	28	1.2	0.5	0.7	1.0
2008	13	0.6	0.2	0.3	0.4
2009	23	1.0	0.4	0.6	0.8
2010	24	1.0	0.5	0.7	0.8
2011	25	1.1	0.4	0.6	0.8
2012	16	0.7	0.3	0.4	0.5
1998-2012	253	0.9	0.4	0.6	0.7

The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

Table 3

Age distribution parameters by year of diagnosis
(incl. DCO)

Year of diagnosis	Cases n	Std.		Min.	Max.	10%	25%	Median		
		Mean	dev.					50%	75%	90%
1998	12	67.0	17.0	32.6	89.4	48.6	54.3	71.0	80.8	82.7
1999	7	71.3	15.8	42.4	89.2	42.4	59.1	75.0	82.3	89.2
2000	10	59.6	15.7	37.5	80.1	40.3	45.7	60.2	75.0	77.9
2001	12	70.2	20.9	14.7	93.2	55.1	61.7	75.6	83.7	84.9
2002	11	72.1	14.6	46.1	96.1	52.8	63.0	73.3	82.5	83.9
2003	20	73.1	11.0	54.0	85.8	57.7	61.3	78.9	82.4	84.5
2004	19	65.0	22.4	2.9	92.5	25.2	57.9	67.0	82.0	91.1
2005	15	70.2	22.2	17.6	89.4	25.4	59.5	75.5	85.1	88.7
2006	18	74.5	15.8	49.9	96.0	50.2	60.8	79.4	86.9	95.1
2007	28	72.4	14.7	26.6	92.7	52.6	65.8	76.9	82.4	88.1
2008	13	76.6	12.5	47.8	91.3	64.5	68.1	80.4	85.5	88.8
2009	23	70.0	16.0	33.3	95.5	45.9	59.7	71.9	82.0	87.9
2010	24	69.7	14.4	46.7	95.3	49.5	57.4	70.1	81.1	87.7
2011	25	72.3	13.3	42.2	90.3	49.5	67.4	74.7	82.0	88.0
2012	16	69.3	12.0	29.9	84.3	60.0	69.3	71.1	73.6	81.7
1998-2012	253	70.5	15.9	2.9	96.1	49.5	61.6	73.3	82.3	87.9

Table 4

Age distribution by 5-year age group for period 1998-2012
(incl. DCO)

Age at diagnosis Years	Cases		Cum.%
	n	%	
0-4	1	0.4	0.4
5-9	0	0.0	0.4
10-14	1	0.4	0.8
15-19	1	0.4	1.2
20-24	0	0.0	1.2
25-29	4	1.6	2.8
30-34	2	0.8	3.6
35-39	2	0.8	4.3
40-44	4	1.6	5.9
45-49	13	5.1	11.1
50-54	13	5.1	16.2
55-59	18	7.1	23.3
60-64	16	6.3	29.6
65-69	28	11.1	40.7
70-74	35	13.8	54.5
75-79	30	11.9	66.4
80-84	47	18.6	85.0
85+	38	15.0	100.0
All ages	253	100.0	

Included in the statistics are 45.1% multiple primaries.

Table 5

Age-specific incidence, DCO rate and proportion of all cancers
for period 1998-2012

Age at diagnosis Years	Cases n	Age-spec. incidence	DCO rate n=19 %	Prop. all cancers n=142297 %
0- 4	1	0.1		0.4
5- 9		0.0		
10-14	1	0.1		0.6
15-19	1	0.1		0.4
20-24		0.0		
25-29	4	0.2		0.4
30-34	2	0.1		0.1
35-39	2	0.1		0.1
40-44	4	0.2		0.1
45-49	13	0.6		0.2
50-54	13	0.7		0.1
55-59	18	1.0		0.1
60-64	16	0.9	6.3	0.1
65-69	28	1.7	3.6	0.2
70-74	35	2.5	5.7	0.2
75-79	30	2.7	3.3	0.2
80-84	47	5.4	10.6	0.3
85+	38	4.6	23.7	0.2
All ages	253		7.5	0.2
Incidence				
Raw		0.9		
WS		0.4		
ES		0.6		
BRD-S		0.7		

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

Table 6

Standardized incidence ratio (SIR, with 95% confidence limits),
excess absolute risk (EAR) and DCO rate of second primaries
for period 1998-2012

Diagnosis	Observed n	Expected n	SIR	LCL 95%	UCL 95%	EAR	DCO %
C18 Colon	3	0.7	4.4	0.9	13.0	44.8	33.3
C19-C20 Rectum	2	0.3	7.1	0.9	25.7	33.2	50.0
C53 Cervix uteri	2	0.1	24.7	3.0	89.2 #	37.0	
C54 Corpus uteri	4	0.3	13.2	3.6	33.9 #	71.4	25.0
Other primaries	4	2.0	2.0	0.6	5.2	39.5	
Not observed	0	2.7	0.0	0.0	1.4	-52.4	
All mult. primaries	15	6.0	2.5	1.4	4.1 #	173.5	20.0

Patients	178
Mean age at second malignancy (years)	77.0
Person-years	518
Mean observation time (years)	2.9
Median observation time (years)	1.4

The occurrence of second malignancy is statistically significant.

Observed second malignancies with count 1 are pooled in category "Other primaries".




Figure 7. Age distribution and age-specific incidence




Figure 7a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).

Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2012, based on the November 2011 submission. <http://www.seer.cancer.gov>.




Figure 8. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.

Average incidence (world standard population) 2003 - 2008




Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (0.4/100,000 WS N=110). Since cancer data are not available in some counties until 2007, the local incidence rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 1 women were identified with newly diagnosed vaginal cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.1/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 0.9/100,000.

Standardized incidence ratio (SIR) 2003 - 2008




Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (N=110). Since cancer data are not available in some counties until 2007, the local SIR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 1 women were identified with newly diagnosed vaginal cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.32. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 2.38, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts, and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

Year of diagnosis	Incident cases n	Prop. actively followed %	Prop. DCO %	Deaths n	Prop. deaths %	Prop. deaths with death certific. %
1998	12	91.7		9	75.0	100.0
1999	7	100.0	14.3	6	85.7	83.3
2000	10	100.0		3	30.0	66.7
2001	12	100.0	8.3	9	75.0	88.9
2002	11	100.0	9.1	9	81.8	88.9
2003	20	100.0	5.0	16	80.0	93.8
2004	19	100.0	21.1	15	78.9	100.0
2005	15	100.0	6.7	7	46.7	100.0
2006	18	88.9	5.6	14	77.8	92.9
2007	28	89.3		14	50.0	100.0
2008	13	76.9	15.4	6	46.2	100.0
2009	23	69.6	13.0	12	52.2	100.0
2010	24	70.8	8.3	14	58.3	100.0
2011	25	88.0	4.0	12	48.0	100.0
2012	16	93.8	6.3	5	31.3	100.0
1998-2012	253	89.3	7.5	151	59.7	96.0

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)
 (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

Year of diagnosis/ death	Incident cases n	Deaths n	Prop. deaths with death certific. %	Deaths in same year n	Prop. deaths in same year %
1998	12	9	100.0	2	16.7
1999	7	11	90.9	1	14.3
2000	10	10	90.0	1	10.0
2001	12	10	90.0	6	50.0
2002	11	12	83.3	2	18.2
2003	20	6	100.0	3	15.0
2004	19	22	95.5	6	31.6
2005	15	13	100.0	4	26.7
2006	18	12	100.0	5	27.8
2007	28	21	95.2	4	14.3
2008	13	12	100.0	2	15.4
2009	23	9	100.0	4	17.4
2010	24	13	100.0	5	20.8
2011	25	11	100.0	2	8.0
2012	16	21	100.0	3	18.8
1998-2012	253	192	96.4	50	19.8

Table 10c

Annual cohorts of deaths, proportion of cancer-related and not cancer-related deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

Year of death	Deaths n	Prop. cancer-related %	Prop. not cancer-related %	Prop. cancer recorded on death certificate %
1998	9	66.7	33.3	77.8
1999	11	72.7	27.3	80.0
2000	10	90.0	10.0	100.0
2001	10	70.0	30.0	88.9
2002	12	50.0	50.0	60.0
2003	6	66.7	33.3	83.3
2004	22	86.4	13.6	90.5
2005	13	92.3	7.7	92.3
2006	12	66.7	33.3	75.0
2007	21	71.4	28.6	80.0
2008	12	66.7	33.3	83.3
2009	9	33.3	66.7	44.4
2010	13	84.6	15.4	84.6
2011	11	90.9	9.1	90.9
2012	21	76.2	23.8	85.7
1998-2012	192	74.0	26.0	82.2

Table 11

Means of age at death according to the grouping in Table 10

Year of death	Deaths n	Age at death (all causes) Years	Age at death (cancer-related) Years	Age at death (not cancer-related) Years	Age at death (according to death certificate) Years
1998	9	70.5	64.3	82.9	67.0
1999	11	76.0	72.9	84.1	73.1
2000	10	73.5	72.8	80.2	73.6
2001	10	73.4	72.0	76.6	75.3
2002	12	79.5	74.6	84.4	81.2
2003	6	81.3	77.6	88.8	79.5
2004	22	73.7	73.3	76.5	73.3
2005	13	77.0	76.0	89.0	76.0
2006	12	81.2	79.2	85.1	80.2
2007	21	75.3	71.1	85.7	74.3
2008	12	76.4	74.4	80.3	76.0
2009	9	78.9	83.7	76.5	84.0
2010	13	80.0	77.8	91.8	77.8
2011	11	79.3	79.9	74.1	79.9
2012	21	76.0	73.8	83.2	74.4
1998-2012	192	76.5	74.5	82.4	75.7

By 2010, life expectancy for a newborn male in Germany is 77.5 years compared with 82.6 years for his female counterpart.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 12

Mortality measures (cancer-related death) and mortality-incidence-index
by year of death

Year of death	Deaths n	Mort. raw	MI-Index raw	Mort. WS	MI-Index WS	Mort. ES	MI-Index ES	Mort. BRD-S	MI-Index BRD-S
1998	6	0.5	0.50	0.3	0.51	0.4	0.51	0.4	0.42
1999	8	0.7	1.14	0.2	1.03	0.4	1.02	0.6	1.17
2000	9	0.7	0.90	0.3	0.65	0.5	0.71	0.6	0.79
2001	7	0.6	0.58	0.3	0.53	0.4	0.60	0.5	0.53
2002	6	0.3	0.55	0.1	0.53	0.2	0.54	0.3	0.56
2003	4	0.2	0.20	0.1	0.14	0.1	0.15	0.2	0.19
2004	19	1.0	1.00	0.4	0.64	0.6	0.81	0.7	0.90
2005	12	0.6	0.80	0.2	0.59	0.3	0.67	0.4	0.70
2006	8	0.4	0.44	0.1	0.34	0.2	0.36	0.3	0.44
2007	15	0.6	0.54	0.3	0.66	0.4	0.60	0.5	0.52
2008	8	0.3	0.62	0.1	0.65	0.2	0.70	0.3	0.68
2009	3	0.1	0.13	0.0	0.05	0.0	0.07	0.1	0.11
2010	11	0.5	0.46	0.1	0.30	0.2	0.33	0.3	0.39
2011	10	0.4	0.40	0.1	0.29	0.2	0.33	0.3	0.37
2012	16	0.7	1.00	0.2	0.77	0.4	0.88	0.5	0.96
1998-2012	142	0.5	0.56	0.2	0.48	0.3	0.51	0.4	0.53

Table 13

Age distribution of age at death (cancer-related) for period 1998-2012
(incl. multiple primaries)

Age at death Years	Cases		Cum. %
	n	%	
5-9	1	0.7	0.7
10-14	0	0.0	0.7
15-19	0	0.0	0.7
20-24	0	0.0	0.7
25-29	0	0.0	0.7
30-34	1	0.7	1.4
35-39	0	0.0	1.4
40-44	1	0.7	2.1
45-49	5	3.5	5.6
50-54	7	4.9	10.6
55-59	8	5.6	16.2
60-64	10	7.0	23.2
65-69	10	7.0	30.3
70-74	18	12.7	43.0
75-79	18	12.7	55.6
80-84	30	21.1	76.8
85+	33	23.2	100.0
All ages	142	100.0	

Included in the statistics are 45.1% multiple primaries.

Table 14

Age-specific mortality (cancer-related) and proportion of all cancers
for period 1998-2012
(incl. multiple primaries)

Age at death Years	Cases n	Age-spec. mortality	MI-index	Prop. all cancers %
0- 4		0.0		
5- 9	1	0.1	1.00	2.6
10-14		0.0		
15-19		0.0		
20-24		0.0		
25-29		0.0		
30-34	1	0.0	0.50	0.5
35-39		0.0		
40-44	1	0.0	0.25	0.1
45-49	5	0.2	0.38	0.3
50-54	7	0.4	0.54	0.2
55-59	8	0.4	0.44	0.2
60-64	10	0.6	0.63	0.2
65-69	10	0.6	0.36	0.1
70-74	18	1.3	0.51	0.2
75-79	18	1.6	0.60	0.2
80-84	30	3.5	0.64	0.3
85+	33	4.0	0.87	0.3
All ages	142			0.2
Mortality				
Raw		0.5	0.56	
WS		0.2	0.48	
ES		0.3	0.51	
BRD-S		0.4	0.53	
PYLL-70				
per 100,000		2.3		
ES		2.1		
AYLL-70		13.1		

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15

Multiple primaries in deaths in period 1998-2012

Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C03-C06 Oral cavity	1	1.3					1	100.0
C16 Stomach	1	1.3	1	100.0				
C17 Small intestine	1	1.3					1	100.0
C18 Colon	3	3.9	1	33.3	1	33.3	1	33.3
C19-C20 Rectum	5	6.5	2	40.0			3	60.0
C21 Anus/canal	1	1.3					1	100.0
C23-C24 Bile	1	1.3					1	100.0
C33-C34 Lung	5	6.5	1	20.0	1	20.0	3	60.0
C44 Skin others	2	2.6	1	50.0			1	50.0
C50 Breast	10	13.0	6	60.0			4	40.0
C51 Vulva	4	5.2	1	25.0	1	25.0	2	50.0
C52 Vagina	4	5.2					4	100.0
C53 Cervix uteri	15	19.5	11	73.3	3	20.0	1	6.7
C54 Corpus uteri	11	14.3	7	63.6	3	27.3	1	9.1
C55,C57 Fem. genitals un	2	2.6			1	50.0	1	50.0
C56 Ovary	1	1.3					1	100.0
C64 Kidney	1	1.3					1	100.0
C65 Renal pelvis	1	1.3	1	100.0				
C67 Bladder	3	3.9	2	66.7			1	33.3
C69 Eye carcinoma	1	1.3	1	100.0				
C70-C72 CNS cancer	1	1.3					1	100.0
C91-C96 Leukaemia	3	3.9					3	100.0
All mult. primaries	77	100.0	35	45.5	10	13.0	32	41.6

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers
for period 1998-2012
(Singular primaries only *)

Age at death Years	Cases n	Age-spec. mortality	MI-index	Prop. all cancers %
0- 4		0.0		
5- 9	1	0.1	1.00	2.8
10-14		0.0		
15-19		0.0		
20-24		0.0		
25-29		0.0		
30-34	1	0.0	1.00	0.5
35-39		0.0		
40-44		0.0		
45-49	5	0.2	0.42	0.3
50-54	6	0.3	0.60	0.2
55-59	6	0.3	0.46	0.2
60-64	8	0.5	0.50	0.2
65-69	8	0.5	0.36	0.1
70-74	13	0.9	0.72	0.2
75-79	11	1.0	0.42	0.1
80-84	20	2.3	0.59	0.2
85+	24	2.9	0.80	0.2
All ages	103			0.2
Mortality				
Raw		0.4	0.53	
WS		0.1	0.46	
ES		0.2	0.49	
BRD-S		0.3	0.50	
PYLL-70 per 100,000		1.9		
ES		1.8		
AYLL-70		13.5		

* See corresponding tables with multiple primaries.

Table 17

Age-specific mortality (cancer-related) and proportion of all cancers
for period 1998-2012
(**Single primaries only ***)

Age at death Years	Cases n	Age-spec. mortality	MI-index	Prop. all cancers %
0- 4		0.0		
5- 9	1	0.1	1.00	2.9
10-14		0.0		
15-19		0.0		
20-24		0.0		
25-29		0.0		
30-34	1	0.0	1.00	0.6
35-39		0.0		
40-44		0.0		
45-49	5	0.2	0.45	0.3
50-54	4	0.2	0.50	0.2
55-59	6	0.3	0.50	0.2
60-64	7	0.4	0.54	0.2
65-69	6	0.4	0.29	0.1
70-74	10	0.7	0.59	0.2
75-79	7	0.6	0.32	0.1
80-84	18	2.1	0.56	0.3
85+	21	2.6	0.78	0.2
All ages	86			0.2
Mortality				
Raw		0.3	0.49	
WS		0.1	0.43	
ES		0.2	0.45	
BRD-S		0.2	0.46	
PYLL-70				
per 100,000		1.7		
ES		1.6		
AYLL-70		14.2		

* See corresponding tables with multiple primaries.

C52: Malignant neoplasm of vagina

Age distribution and age-specific mortality 1998 - 2012 (n=142)




Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at vaginal cancer-related death (see Table 10) should be considered.

Average mortality (world standard population) 2003 - 2008




Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2003 to 2008. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (0.2/100,000 WS N=61). Since cancer data are not available in some counties until 2007, the local mortality rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 0 women died from vaginal cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.0/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.4/100,000.

Standardized mortality ratio (SMR) 2003 - 2008




Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (N=61). Since cancer data are not available in some counties until 2007, the local SMR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 0 women died from vaginal cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.00. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 3.08, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the tumor-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

AYLL-70	Average years of life lost prior to age 70 given a person dies before that age
BRD-S	German standard population
DCO	Death certificate only
EAR	Excess absolute risk = excess cancer cases (O - E) per 10,000 person-years
ES	European standard population (old)
FRG	Federal Republic of Germany
GEKID	Association of Population-based Cancer Registries in Germany (Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)
LCL	Lower confidence limit
MI-index	Ratio between mortality and incidence
MCR	Munich Cancer Registry (Tumorregister München)
PYLL-70	Potential years of life lost prior to age 70 given a person dies before that age
SEER	Surveillance, Epidemiology, and End Results (USA)
SIR	Standardized incidence ratio
SMR	Standardized mortality ratio
UCL	Upper confidence limit
WS	World standard population

Recommended Citation

Munich Cancer Registry. Baseline statistics C52: Vaginal cancer [Internet]. 2014 [updated 2014 Mar 20; cited 2014 May 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/base/base_C52__E.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

Fig./Tbl.		Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	3
2	Incidence by year of diagnosis	4
3	Age distribution parameters by year of diagnosis	5
4	Age distribution by 5-year age group	6
5	Age-specific incidence and DCO rate	7
6	Standardized incidence ratio of second primaries	8
7	Age distribution and age-specific incidence (chart)	9
7a	Age-specific incidence internationally (chart)	10
8	Cumulative follow-up years (chart)	11
9a	Map of cancer incidence (WS) by county (chart)	12
9b	Standardized incidence ratio (SIR) by county (chart)	13
10a	Pts incident cohorts and mortality / yr	14
10b	Incidence and mortality by year of diagnosis	15
10c	Cancer-related deaths, death certification available / yr	16
11	Means of age at death / yr	17
12	Mortality by year of death	18
13	Distribution of age at death	19
14	Age-specific mortality	20
15	Multiple primaries in deaths	21
16	Age-specific mortality (first primaries)	22
17	Age-specific mortality (single primaries)	23
18	Age distribution and age-specific mortality (chart)	24
19a	Map of cancer mortality (WS) by county (chart)	25
19b	Standardized mortality ratio (SMR) by county (chart)	26