Munich Cancer Registry



- Survival
- Selection Matrix
- ► Homepage

Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

Cancer statistics: Baseline statistics

C75: Other endocrine glands cancer

Year of diagnosis	1998-2012
Patients	111
Diseases	111
Creation date	03/20/2014
Export date	02/12/2014
Population	4.5 m



http://www.tumorregister-muenchen.de/en/facts/base/base_C75__E.pdf

Global Statements about the statistics on the Internet -

Baseline Statistics (grey button ___), Survival (red button ___)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.5 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases^{###} are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR. The time-delayed acquisition of data and the occasionally high DCO-rates indicate optimizing reserves, among others, because of current financial and legal conditions that hinder the analyses.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, March 2014

- [#] Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007). Death certificates from 2013 are incorporated into these analyses.
- ^{##} Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- ^{###} DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate. A high proportion of DCO cases (≥5%) in particular cancer types indicate insufficient participation of specific cancer specializations.

ICD-10 codes used for specifying cancer site

ICD-10	Description
C75	Malignant neoplasm of other endocrine glands and related structures
C75.0	Parathyroid gland
C75.1	Pituitary gland
C75.2	Craniopharyngeal duct
C75.3	Pineal gland
C75.4	Carotid body
C75.5	Aortic body and other paraganglia
C75.8	Pluriglandular involvement, unspecified
C75.9	Endocrine gland, unspecified

INCIDENCE

Table 1

Patient cohorts by year of diagnosis including DCO cases and multiple primaries, and with proportion of deaths and active follow-up

		DCO	Prop.	Prop. mult.	Prop.	Prop. actively
Year of	Cases	cases	DCO	primaries	deaths	followed
diagnosis	n	n	90	8	00	ତ
1998	2				100.0	100.0
1999	2				50.0	100.0
2000	2			50.0	50.0	50.0
2001	4			25.0	25.0	100.0
2002	6			16.7	33.3	100.0 #
2003	11	1	9.1		45.5	81.8 #
2004	9			22.2	44.4	100.0 #
2005	12	4	33.3	16.7	50.0	91.7 #
2006	5			40.0	40.0	60.0 #
2007	8	1	12.5	25.0	50.0	62.5 # ##
2008	13	7	53.8	30.8	61.5	61.5
2009	4	1	25.0	25.0	25.0	100.0
2010	11	3	27.3	36.4	27.3	54.5
2011	11	10	90.9	18.2	90.9	100.0
2012	11	5	45.5	45.5	72.7	90.9 ###
1998-2012	111	32	28.8	24.3	52.3	82.0

The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

- ## Since 2007 the percentage of actively followed patients sharply declined compared to the previous years. This is a consequence of ambiguous data protection rules that currently forbid cancer registries in Bavaria to obtain the essential life status informations from competent registration offices.
- ### Please be aware that data of recent annual patient cohorts may not yet be fully processed. Therefore, the presented figures and tables are potentially related to different time periods as pointed out in the respective headlines or legends.

Table 1a

Patient cohorts by year of diagnosis and gender including DCO cases

Year of	All	Males	Females	Prop. males %	
diagnosis	n	n	n	6	
1998	2	1	1	50.0	
1999	2	1	1	50.0	
2000	2 2		2		
2001	4	2	2	50.0	
2002	6	3	3	50.0	
2003	11	5	6	45.5	
2004	9	6	3 7 1	66.7	
2005	12	5 4	7	41.7	
2006	5	4		80.0	
2007 2008	13	5 7	36	62.5 53.8	
2008	13 4	2	2	50.0	
2010	11	4	7	36.4	
2011	11	7	4	63.6	
2012	11	6	5	54.5	
1998-2012	111	58	53	52.3	

Incidence measures by year of diagnosis and gender including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

Year of diagnosis	Males n	Females n	Males Inc. raw	Fem. Inc. raw	Males Inc. WS	Fem. Inc. WS	Males Inc. ES	Fem. Inc. ES	Males Inc. BRD-S	Fem. Inc. BRD-S
1998	1	1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1
1999	1	1	0.1	0.1	0.1	0.0	0.1	0.0	0.1	0.1
2000		2		0.2		0.1		0.1		0.1
2001	2	2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2002	3	3	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.1
2003	5	6	0.3	0.3	0.2	0.3	0.2	0.3	0.3	0.3
2004	6	3	0.3	0.2	0.3	0.1	0.3	0.1	0.3	0.1
2005	5	7 <	0.3	0.4	0.2	0.2	0.3	0.3	0.3	0.3
2006	4	1	0.2	0.0	0.2	0.0	0.2	0.0	0.2	0.0
2007	5	3	0.2	0.1	0.1	0.1	0.2	0.1	0.2	0.1
2008	7	б	0.3	0.3	0.1	0.1	0.2	0.2	0.3	0.2
2009	2	2	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.1
2010	4	7	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.3
2011	7	4	0.3	0.2	0.1	0.1	0.2	0.1	0.3	0.1
2012	6	5	0.3	0.2	0.1	0.1	0.2	0.2	0.2	0.2
1998-2012	58	53	0.2	0.2	0.2	0.1	0.2	0.1	0.2	0.2

The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	2	38.0	37.6	11.4	64.6	11.4	11.4	38.0	64.6	64.6
1999	2	65.3	19.0	51.8	78.7	51.8	51.8	65.3	78.7	78.7
2000	2	68.0	14.5	57,7	78.3	57.7	57.7	68.0	78.3	78.3
2001	4	36.1	24.6	18.5	72.0	18.5	20.2	26.9	52.0	72.0
2002	6	48.8	16.7	29.4	74.2	29.4	34.1	48.7	57.7	74.2
2003	11	47.2	20.0	14.0	79.7	30.1	32.6	46.1	59.5	74.3
2004	9	50.6	25.8	2.2	81.1	2.2	31.7	53.7	69.8	81.1
2005	12	53.7	24.8	13.0	92.8	23.1	36.1	51.1	72.2	84.6
2006	5	53.0	28.3	10.9	86.0	10.9	42.3	59.2	66.6	86.0
2007	8	55.3	21.6	14.9	81.7	14.9	41.2	62.6	68.9	81.7
2008	13	69.5	19.7	15.9	86.5	43.0	68.6	74.1	80.4	85.2
2009	4	52.6	22.2	25.1	71.5	25.1	34.6	56.8	70.6	71.5
2010	11	54.7	28.7	15.4	91.9	19.9	21.8	62.6	81.1	82.4
2011	11	77.1	8.1	63.3	87.8	68.2	68.5	80.9	83.1	84.8
2012	11	69.3	18.2	21.1	90.0	58.6	64.2	73.8	76.8	85.7
1998-2012	111	58.0	23.2	2.2	92.8	21.8	42.3	64.6	76.8	82.7

Age distribution parameters by year of diagnosis (All) (incl. DCO)

Table 3a

Age distribution parameters by year of diagnosis (MALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
diagnosis	_ <u> </u>	Mean	uev.		nax.	100	200	508	/ 5 0	20.8
1998	1	64.6		64.6	64.6	64.6	64.6	64.6	64.6	64.6
1999	1	51.8		51.8	51.8	51.8	51.8	51.8	51.8	51.8
2001	2	45.2	37.8	18.5	72.0	18.5	18.5	45.2	72.0	72.0
2002	3	42.3	12.6	29.4	54.7	29.4	29.4	42.6	54.7	54.7
2003	5	56.9	21.0	34.8	79.7	34.8	36.0	59.5	74.3	79.7
2004	6	48.1	28.5	2.2	81.1	2.2	29.7	52.8	69.8	81.1
2005	5	46.9	26.4	13.0	76.9	13.0	28.1	50.3	66.2	76.9
2006	4	44.8	24.7	10.9	66.6	10.9	26.6	50.8	62.9	66.6
2007	5	61.8	9.3	46.2	70.1	46.2	62.4	62.8	67.7	70.1
2008	7	72.3	14.3	43.0	86.5	43.0	68.6	74.1	82.2	86.5
2009	2	34.6	13.3	25.1	44.0	25.1	25.1	34.6	44.0	44.0
2010	4	43.1	25.8	19.9	68.1	19.9	20.9	42.2	65.4	68.1
2011	7	80.1	4.7	72.7	84.8	72.7	74.1	82.4	83.1	84.8
2012	6	76.4	9.7	64.2	90.0	64.2	70.4	74.2	85.7	90.0
1998-2012	58	58.5	22.4	2.2	90.0	21.8	43.0	65.4	74.3	82.7

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of diagnosis	Cases n	Mean	Std. dev.	Min.	Max.	10%	25%	Median 50%	75%	90%
1998	1	11.4		11.4	11.4	11.4	11.4	11.4	11.4	11.4
1999	1	78.7		78.7	78.7	78.7	78.7	78.7	78.7	78.7
2000	2	68.0	14.5	57.7	78.3	57.7	57.7	68.0	78.3	78.3
2001	2	26.9	7.2	21.8	32.0	21.8	21.8	26.9	32.0	32.0
2002	3	55.4	20.2	34.1	74.2	34.1	34.1	57.7	74.2	74.2
2003	6	39.1	16.6	14.0	59.4	14.0	30.1	39.4	52.1	59.4
2004	3	55.6	24.3	31.7	80.3	31.7	31.7	54.8	80.3	80.3
2005	7	58.6	24.5	23.1	92.8	23.1	44.0	51.9	84.6	92.8
2006	1	86.0		86.0	86.0	86.0	86.0	86.0	86.0	86.0
2007	3	44.3	34.1	14.9	81.7	14.9	14.9	36.3	81.7	81.7
2008	б	66.3	25.7	15.9	85.2	15.9	65.8	75.4	80.4	85.2
2009	2	70.6	1.3	69.7	71.5	69.7	69.7	70.6	71.5	71.5
2010	7	61.3	30.1	15.4	91.9	15.4	24.1	74.0	82.4	91.9
2011	4	72.0	10.8	63.3	87.8	63.3	65.8	68.4	78.1	87.8
2012	5	60.8	23.4	21.1	76.8	21.1	58.6	73.3	74.4	76.8
1998-2012	53	57.3	24.3	11.4	92.8	21.1	34.1	63.3	78.3	84.6

Age at									
diagnosis	Cases			Males			Females		
Years	n	00	Cum.%	n	olo	Cum.%	n	00	Cum.%
0-4	1	0.9	0.9	1	1.7	1.7			0.0
5-9	0	0.0	0.9			1.7			0.0
10-14	5	4.5	5.4	2	3.4	5.2	3	5.7	5.7
15-19	4	3.6	9.0	2	3.4	8.6	2	3.8	9.4
20-24	5	4.5	13.5	/ 1	1.7	10.3	4	7.5	17.0
25-29	4	3.6	17.1	4	6.9	17.2			17.0
30-34	6	5.4	22.5	1	1.7	19.0	5	9.4	26.4
35-39	2	1.8	24.3	1	1.7	20.7	1	1.9	28.3
40 - 44	5	4.5	28.8	4	6.9	27.6	1	1.9	30.2
45-49	3	2.7	31.5	1	1.7	29.3	2	3.8	34.0
50-54	8	7.2	38.7	5	8.6	37.9	3	5.7	39.6
55-59	б	5.4	44.1	2	3.4	41.4	4	7.5	47.2
60-64	7	6.3	50.5	5	8.6	50.0	2	3.8	50.9
65-69	11	9.9	60.4	б	10.3	60.3	5	9.4	60.4
70-74	16	14.4	74.8	10	17.2	77.6	6	11.3	71.7
75-79	7	6.3	81.1	3	5.2	82.8	4	7.5	79.2
80-84	13	11.7	92.8	7	12.1	94.8	6	11.3	90.6
85+	8		100.0	3	5.2	100.0	5	9.4	100.0
	-			-					
All ages	111	100.0		58	100.0		53	100.0	

Age distribution by 5-year age group and gender for period 1998-2012 (incl. DCO)

Table 4

Included in the statistics are 27.6% multiple primaries in males and 24.5% in females.

Males Females Males Females Males Females Prop.all Prop.all Age at DCO rate DCO rate cancers cancers Age- Agediagnosis Males Females n=21 n=11 n=146755 n=142297 spec. spec. Years incid. incid. % n n % % % 0- 4 0.1 0.0 0.3 1 5-9 0.0 0.0 10 - 142 0.1 0.2 1.9 3 1.4 15-19 2 2 0.1 0.1 0.6 0.7 20-24 1 4 0.1 0.2 0.2 0.8 25-29 4 0.2 0.0 25.0 0.5 30-34 1 5 0.0 0.2 0.1 0.3 35-39 1 1 0.0 0.0 0.0 0.0 40 - 444 1 0.2 0.0 25.0 0.1 0.0 45-49 1 2 0.0 0.1 0.0 0.0 50-54 5 3 0.3 0.2 0.0 0.1 55-59 2 4 0.2 0.0 0.1 0.0 60-64 5 2 0.1 20.0 50.0 0.0 0.3 0.0 65-69 6 5 33.3 40.0 0.0 0.4 0.3 0.0 70-74 10 б 0.9 0.4 70.0 0.0 0.0 75-79 4 0.4 0.4 0.0 3 25.0 0.0 7 85.7 50.0 80-84 6 1.5 0.7 0.0 0.1 3 5 1.0 80.0 0.0 85+ 0.6 100.0 0.0 36.2 53 20.8 0.0 0.0 All ages 58 Incidence 0.2 0.2 Raw 0.2 WS 0.1 ES 0.2 0.1 BRD-S 0.2 0.2

Age-specific incidence, DCO rate and proportion of all cancers for period 1998-2012

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

Table 6a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2012 MALES Observed Expected LCL UCL DCO Diagnosis SIR 95% 95% EAR % n n Other primaries 1 0.2 4.1 0.1 23.0 71.6 Not observed 0 0.7 0.0 0.0 5.5 -63.1 All mult. primaries 1 0.9 1.1 0.0 6.1 8.5 Patients 33 74.2 Mean age at second malignancy (years) 106 Person-years Mean observation time (years) 3.2 Median observation time (years) 2.5

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries".


Table 6b

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2012 FEMALES

Diagnosis	Observed n	Expected n	SIR	LCL 95%	UCL 95%	EAR	DCO %
Other primaries Not observed	1 0	0.0 0.9	45.1 0.0	1.1 0.0	251.1 # 4.0	73.7 -68.9	
All mult. primaries	1	0.9	1.1	0.0	6.0	4.9	
Patients Mean age at second ma Person-years Mean observation time Median observation ti	(years)	years)	36 58.6 133 3.7 2.8				

The occurrence of second malignancy is statistically significant.


Observed second malignancies with count 1 are pooled in category "Other primaries".



C75: Malignant neoplasm of other endocrine glands and related structures Age distribution and age-specific incidence 1998 - 2012 (Males: 58, Females: 53)

Figure 7. Age distribution and age-specific incidence









Figure 7a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).

Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2012, based on the November 2011 submission. http://www.seer.cancer.gov.




C75: Malignant neoplasm of other endocrine glands and related structures Cumulative follow-up years since diagnosis for period 1998 - 2012 (excl. DCO)

Figure 8. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.






Average incidence (world standard population) 2003 - 2008: Males

Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.2/100,000 WS N=31, females 0.2/100,000 WS N=25). Since cancer data are not available in some counties until 2007, the local incidence rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 1 women were identified with newly diagnosed other endocrine glands cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.4/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 3.1/100,000.



Standardized incidence ratio (SIR) 2003 - 2008: Males

Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=31, females N=25). Since cancer data are not available in some counties until 2007, the local SIR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 1 women were identified with newly diagnosed other endocrine glands cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 1.32. Though, the value of this parameter may vary with an underlying probability of 99% between 0.01 and 9.83, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts, and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

		Prop.				Prop. deaths
	Incident	actively	Prop.		Prop.	with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	90 90	olo	n	00	90
1998	2	100.0		2	100.0	100.0
1999	2	100.0		1	50.0	100.0
2000	2	50.0		1	50.0	100.0
2001	4	100.0		1	25.0	
2002	6	100.0		2	33.3	100.0
2003	11	81.8	9.1	5	45.5	100.0
2004	9	100.0		4	44.4	100.0
2005	12	91.7	33.3	6	50.0	100.0
2006	5	60.0		2	40.0	100.0
2007	8	62.5	12.5	4	50.0	100.0
2008	13	61.5	53.8	8	61.5	100.0
2009	4	100.0	25.0	1	25.0	100.0
2010	11	54.5	27.3	3	27.3	100.0
2011	11	100.0	90.9	10	90.9	100.0
2012	11	90.9	45.5	8	72.7	100.0
1998-2012	111	82.0	28.8	58	52.3	98.3

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

			Prop.		
			deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	n	o.	n	8
1998	2	2	100.0		
1999	2	1	100.0		
2000	2	1	100.0		
2001	4	2	100.0		
2002	б				
2003	11	5	80.0	3	27.3
2004	9	2	100.0	1	11.1
2005	12	5	100.0	4	33.3
2006	5	4	100.0	1	20.0
2007	8	5	100.0	4	50.0
2008	13	4	100.0	6	46.2
2009	4	4	100.0	1	25.0
2010	11	4	100.0	3	27.3
2011	11	4	100.0	9	81.8
2012	11	5	100.0	6	54.5
1998-2012	111	48	97.9	38	34.2

Table 10c

		Prop.	Prop.	Prop. cancer recorded	
		cancer-	not cancer-	on death	
Year of	Deaths	related	related	certificate	
death	n	%	8	8	
1998	2	50.0	50.0	100.0	
1999	1/	100.0		100.0	
2000	1	100.0		100.0	
2001	2	100.0		100.0	
2002					
2003	5	80.0	20.0	100.0	
2004	2	100.0		100.0	
2005	5	60.0	40.0	100.0	
2006	4	50.0	50.0	50.0	
2007	5	60.0	40.0	80.0	
2008	4	75.0	25.0	100.0	
2009	4	100.0		100.0	
2010	4	75.0	25.0	100.0	
2011	4	50.0	50.0	75.0	
2012	5	60.0	40.0	100.0	
1998-2012	48	70.8	29.2	91.5	



					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(not cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	1	84.5	84.5		84.5
1999	1	56.4	56.4		56.4
2000					
2001	1	67.1	67.1		67.1
2002					
2003	4	71.9	71.9		71.4
2004	1	30.4	30.4		30.4
2005	1	3.0	3.0		3.0
2006	2	56.5	56.5		56.5
2007	2	58.4	58.4		58.4
2008	3	39.9	25.5	68.6	39.9
2009	3	57.0	57.0		57.0
2010	2	75.2	75.2		75.2
2011	4	73.0	83.6	62.4	79.9
2012	4 2	82.0	73.9	90.0	82.0
1998-2012	27	61.3	59.7	70.9	61.2

Table 11a

Means of age at death according to the grouping in Table 10 $$\rm MALES$$

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(not cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	1	74.7		74.7	74.7
1999					
2000	1	10.5	10.5		10.5
2001	1	58.9	58.9		58.9
2002					
2003	1	83.4		83.4	83.4
2004	1	15.5	15.5		15.5
2005	4	72.8	69.5	76.1	72.8
2006	2	50.8		50.8	
2007	3	70.9	70.1	71.3	78.6
2008	1	80.4	80.4		80.4
2009	1	72.8	72.8		72.8
2010	2	86.5	91.9	81.1	86.5
2011					
2012	3	68.3	73.7	57.5	68.3
1998-2012	21	65.7	62.4	69.3	67.9

Table 11b

Means of age at death according to the grouping in Table 10 FEMALES

By 2010, life expectancy for a newborn male in Germany is 77.5 years compared with 82.6 years for his female counterpart.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 12a

Mortality measures (cancer-related death) and mortality-incidence-index by year of death MALES

Year of death	Deaths n	Mort. raw	MI-Index raw	Mort. WS	MI-Index WS	Mort. ES	MI-Index ES	Mort. BRD-S	MI-Index BRD-S
1998	1	0.1	1.00	0.0	0.77	0.1	1.22	0.2	2.54
1999	1	0.1	1.00	0.0	0.69	0.1	0.74	0.1	0.76
2000									
2001	1	0.1	0.50	0.1	0.26	0.1	0.36	0.1	0.35
2002									
2003	4	0.2	0.80	0.1	0.74	0.2	0.83	0.2	0.91
2004	1	0.1	0.17	0.0	0.13	0.0	0.15	0.0	0.14
2005	1	0.1	0.20	0.1	0.52	0.1	0.33	0.1	0.19
2006	2	0.1	0.50	0.1	0.41	0.1	0.57	0.1	0.62
2007	2	0.1	0.40	0.1	0.38	0.1	0.39	0.1	0.41
2008	2	0.1	0.29	0.1	0.83	0.1	0.48	0.1	0.35
2009	3	0.1	1.50	0.1	1.07	0.1	1.39	0.1	1.46
2010	2	0.1	0.50	0.0	0.19	0.1	0.31	0.1	0.43
2011	2	0.1	0.29	0.0	0.25	0.0	0.28	0.1	0.32
2012	1	0.0	0.17	0.0	0.14	0.0	0.14	0.0	0.14
1000 2012	22	0 1	0 10	0 1	0 20	0 1	0 20	0 1	0.40
1998-2012	23	0.1	0.40	0.1	0.38	0.1	0.39	0.1	0.40

Table 12b

Mortality measures (cancer-related death) and mortality-incidence-index by year of death FEMALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998									
1999									
2000	1	0.1	0.50	0.2	2.46	0.1	1.16	0.1	0.63
2001	1	0.1	0.50	0.1	0.30	0.1	0.47	0.1	0.40
2002									
2003									
2004	1	0.1	0.33	0.1	1.06	0.1	0.61	0.1	0.56
2005	2	0.1	0.29	0.1	0.23	0.1	0.25	0.1	0.26
2006									
2007	1	0.0	0.33	0.0	0.15	0.0	0.23	0.0	0.30
2008	1	0.0	0.17	0.0	0.05	0.0	0.08	0.0	0.15
2009	1	0.0	0.50	0.0	0.42	0.0	0.45	0.0	0.49
2010	1	0.0	0.14	0.0	0.03	0.0	0.06	0.0	0.06
2011									
2012	2	0.1	0.40	0.0	0.29	0.1	0.33	0.1	0.37
1998-2012	11	0.0	0.21	0.0	0.20	0.0	0.19	0.0	0.19

Age at									
death	Cases		1	Males			Females		
Years	n	% C	um.%	n	olo	Cum.%	n	00	Cum.%
0-4	1	2.9	2.9	1	4.3	4.3			0.0
5-9	0	0.0	2.9			4.3			0.0
10-14	1	2.9	5.9			4.3	1	9.1	9.1
15-19	2	5.9	11.8	1	4.3	8.7	1	9.1	18.2
20-24	0	0.0	11.8			8.7			18.2
25-29	0	0.0	11.8			8.7			18.2
30-34	2	5.9	17.6	2	8.7	17.4			18.2
35-39	1	2.9	20.6	1	4.3	21.7			18.2
40 - 44	0	0.0	20.6			21.7			18.2
45-49	2	5.9	26.5	1	4.3	26.1	1	9.1	27.3
50-54	1	2.9	29.4	1	4.3	30.4			27.3
55-59	4	11.8	41.2	3	13.0	43.5	1	9.1	36.4
60-64	2	5.9	47.1	1	4.3	47.8	1	9.1	45.5
65-69	2	5.9	52.9	2	8.7	56.5			45.5
70-74	7	20.6	73.5	5	21.7	78.3	2	18.2	63.6
75-79	1	2.9	76.5	1	4.3	82.6			63.6
80-84	6	17.6	94.1	4	17.4	100.0	2	18.2	81.8
85+	2	5.9 1	00.0			100.0	2	18.2	100.0
All ages	34	100.0		23	100.0		11	100.0	

Age distribution of age at death (cancer-related) for period 1998-2012 (incl. multiple primaries)

Included in the statistics are 27.6% multiple primaries in males and 24.5% in females.

Age at death	Males	Females	÷ /		Females Age- spec.		cancers	Females Prop.all cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	00	0/0
0- 4 5- 9	1		0.1	1.00	0.0		3.2	
10-14 15-19	1	1 1	0.0 0.1	0.50	0.1 0.1	0.33 0.50	2.4	3.6 2.9
20-24 25-29 30-34	2		0.0	2.00	0.0 0.0 0.0		1.1	
30-34 35-39 40-44	1		0.1 0.0 0.0	1.00	0.0		0.3	
45-49 50-54	1 1	1	0.0	1.00	0.0	0.50	0.1	0.1
55-59 60-64 65-69	3 1 2	1	0.2 0.1 0.1	0.20	0.1 0.1 0.0	0.25 0.50	0.1 0.0 0.0	0.0 0.0
70-74 75-79	5 1	2	0.4 0.1	0.50 0.33	0.1 0.0	0.33	0.0 0.0	0.0
80-84 85+	4	2 2	0.9 0.0	0.57	0.2	0.33 0.40	0.0	0.0 0.0
All ages	23	11					0.0	0.0
Mortality Raw			0.1	0.40	0.0	0.21		
WS ES			0.1 0.1	0.38 0.39	0.0	0.20 0.19		
BRD-S			0.1	0.40	0.0	0.19		
PYLL-70 per 100,000			1.3		0.6			
ES AYLL-70			1.4 24.4		0.8 30.5			

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2012 (incl. multiple primaries)

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

	Multiple	primar		death MALES		riod 19	98-2012		
Diagnos	is	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n		Post n	Post ←%
C25 C44 C61 C76-C79	Pancreas Skin others Prostate CUP	1 1 4 1	14.3 14.3 57.1 14.3	1 3 1	100.0 75.0 100.0	1	100.0	1	25.0
All mul	t. primaries	7	100.0	5	71.4	1	14.3	1	14.3

Table 15a

Multiple primaries in deaths in period 1998-2012

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 15b

Multiple primaries in deaths in period 1998-2012 FEMALES

					Syn- chron	Syn- chron		
Diagnosis	Total n	Total %↓	Pre n	Pre ←%	±30d n	±30d ←%	Post n	Post ←%
C50 Breast C67 Bladder C76-C79 CUP C91-C96 Leukaemia	1 1 1	25.0 25.0 25.0 25.0	1 1 1	100.0 100.0 100.0	1	100.0		

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

		(Singular	primarie	s only *)		
		Males		Females		Males	Females
Age at		Age-		Age-			Prop.all
death	Males Female			spec.		cancers	cancers
Years	n n	- /	MI-index		MI-index		90
0- 4	1	0.1	1.00	0.0		3.8	
5-9		0.0		0.0			
10-14	1	0.0		0.1	0.33		3.8
15-19	1 1	0.1		0.1	0.50	2.6	3.2
20-24		0.0		0.0			
25-29		0.0		0.0			
30-34	2	0.1	2.00	0.0		1.2	
35-39	1	0.0	1.00	0.0		0.3	
40-44		0.0		0.0			
45-49	1 1	0.0	1.00	0.0	0.50	0.1	0.1
50-54	1	0.1	0.25	0.0		0.0	
55-59	3 1	0.2	1.50	0.1	0.33	0.1	0.0
60-64	1	0.0		0.1	0.50		0.0
65-69	1	0.1	0.25	0.0		0.0	
70-74	3 1	0.3	0.60	0.1	0.17	0.0	0.0
75-79	1/ /	0.1	0.50	0.0		0.0	
80-84	3 1	0.7	0.60	0.1	0.33	0.0	0.0
85+		0.0		0.0			
All ages	18 7					0.0	0.0
Mortality							
Raw		0.1		0.0	0.16		
WS		0.0	0.37	0.0	0.19		
ES		0.1	0.39	0.0	0.18		
BRD-S		0.1	0.41	0.0	0.17		
PYLL-70							
per 100,000		1.2		0.6			
ES		1.4		0.8			
AYLL-70		28.0		30.5			


Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2012 (Singular primaries only *)

* See corresponding tables with multiple primaries.

			(Single P	primaries	only ^)			
			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males F	Temales	- /		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	010	010
0-4	1		0.1	1.05	0.0		4.0	
5-9			0.0		0.0			
10-14		1	0.0		0.1	0.35		4.2
15-19	1	1	0.1	0.52	0.1	0.52	2.6	3.8
20-24			0.0		0.0			
25-29			0.0		0.0			
30-34	2		0.1	2.12	0.0		1.2	
35-39	1		0.0	1.05	0.0		0.3	
40-44			0.0		0.0			
45-49	1	1	0.0	1.04	0.0	0.52	0.1	0.1
50-54	1	_	0.1	0.26	0.0		0.0	
55-59	3	1	0.2	1.58	0.1	0.35	0.1	0.0
60-64	-	1	0.0	0 50	0.1	0.52		0.0
65-69	1		0.1	0.52	0.0	0 01	0.0	
70-74	3	1	0.3	0.62	0.1	0.21	0.0	0.0
75-79	1 2	-	0.1	0.52	0.0	0.50	0.0	0 0
80-84 85+	Z	1	0.5	0.41	0.1 0.0	0.52	0.0	0.0
+60			0.0		0.0			
All ages	17	7					0.0	0.0
Mortality								
Raw			0.1	0.41	0.0	0.18		
WS			0.0	0.39	0.0	0.20		
ES			0.1		0.0	0.19		
BRD-S			0.1	0.41	0.0	0.18		
PYLL-70								
per 100,000			1.3		0.6			
ES			1.3		0.0			
AYLL-70			28.0		30.5			
			20.0		50.5			

Age-specific mortality (cancer-related) and proportion of all cancers for period 1999-2012 (Single primaries only *)


* See corresponding tables with multiple primaries.



C75: Malignant neoplasm of other endocrine glands and related structures Age distribution and age-specific mortality 1999 - 2012 (Males: 23, Females: 11)

Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).


The difference between age at diagnosis (Table 3) and age at other endocrine glands cancerrelated death (see Table 10) should be considered.



Average mortality (world standard population) 2003 - 2008: Males

Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2003 to 2008. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.1/100,000 WS N=11, females 0.0/100,000 WS N=4). Since cancer data are not available in some counties until 2007, the local mortality rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 0 women died from other endocrine glands cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.0/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.4/100,000.



Standardized mortality ratio (SMR) 2003 - 2008: Males

Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=11, females N=4). Since cancer data are not available in some counties until 2007, the local SMR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 0 women died from other endocrine glands cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.00. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 44.60, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the tumor-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

AYLL-70 BRD-S	Average years of life lost prior to age 70 given a person dies before that age German standard population
DCO	Death certificate only
EAR	Excess absolute risk
	= excess cancer cases (O - E) per 10,000 person-years
ES	European standard population (old)
FRG	Federal Republic of Germany
GEKID	Association of Population-based Cancer Registries in Germany
	(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)
LCL	Lower confidence limit
MI-index	Ratio between mortality and incidence
MCR	Munich Cancer Registry (Tumorregister München)
PYLL-70	Potential years of life lost prior to age 70 given a person dies before that age
SEER	Surveillance, Epidemiology, and End Results (USA)
SIR	Standardized incidence ratio
SMR	Standardized mortality ratio
UCL	Upper confidence limit
WS	World standard population

Recommended Citation

Munich Cancer Registry. Baseline statistics C75: Other endocrine glands cancer [Internet]. 2014 [updated 2014 Mar 20; cited 2014 May 1]. Available from: http://www.tumorregistermuenchen.de/en/facts/base/base_C75__E.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Munich Cancer Registry

Index of figures and tables

Fig./Tb	l.	Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	3
1a	Gender distribution by year of diagnosis	4
2	Incidence by year of diagnosis	5
3	Age distribution parameters by year of diagnosis	6
4	Age distribution by 5-year age group and gender	8
5	Age-specific incidence and DCO rate	9
6	Standardized incidence ratio of second primaries	10
7	Age distribution and age-specific incidence (chart)	12
7a	Age-specific incidence internationally (chart)	13
8	Cumulative follow-up years (chart)	14
9a	Map of cancer incidence (WS) by county (chart)	15
9b	Standardized incidence ratio (SIR) by county (chart)	16
10a	Pts incident cohorts and mortality / yr	17
10b	Incidence and mortality by year of diagnosis	18
10c	Cancer-related deaths, death certification available / yr	19
11	Means of age at death / yr	20
12	Mortality by year of death	22
13	Distribution of age at death	23
14	Age-specific mortality	24
15	Multiple primaries in deaths	25
16	Age-specific mortality (first primaries)	27
17	Age-specific mortality (single primaries)	28
18	Age distribution and age-specific mortality (chart)	29
19a	Map of cancer mortality (WS) by county (chart)	30
19b	Standardized mortality ratio (SMR) by county (chart)	31