# **Munich Cancer Registry**



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage

Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

# **Cancer statistics: Baseline statistics**

# **GEP-NET:** Gastr.ent.pancr. neuroend. tumor

Year of diagnosis	1998-2012
Patients	1,945
Diseases	1,965
Creation date	03/20/2014
Export date	02/12/2014
Population	4.5 m



http://www.tumorregister-muenchen.de/en/facts/base/base\_hDNETE.pdf

# Global Statements about the statistics on the Internet – Baseline Statistics (grey button —), Survival (red button —)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut<sup>#</sup>, with a total of 4.5 million inhabitants, account for the frequency of cancer diseases<sup>##</sup> and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases\*\*\*\* are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR. The time-delayed acquisition of data and the occasionally high DCO-rates indicate optimizing reserves, among others, because of current financial and legal conditions that hinder the analyses.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, March 2014

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007). Death certificates from 2013 are incorporated into these analyses.
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate. A high proportion of DCO cases (≥5%) in particular cancer types indicate insufficient participation of specific cancer specializations.



## ICD-10 codes used for specifying cancer site

Code	Description
C15	Oesophagus
C16	Stomach
C17	Small intestine
C18	Colon
C19	Rectosigmoid junction
C20	Rectum
C21	Anus and anal canal
C22	Liver and intrahepatic bile ducts
C23	Gallbladder
C24	Other and unspecified parts of biliary tract
C25	Pancreas
C26	Other and ill-defined digestive organs

## ICD-O-3 codes (morphology) used for specifying cancer site

Code	Description
8013/3	Large cell neuroendocrine carcinoma
8041/3	Small cell carcinoma, NOS
8150/0	Islet cell adenoma
8150/3	Islet cell carcinoma
8151/3	Insulinoma, malignant
8152/1	Glucagonoma, NOS
8152/3	Glucagonoma, malignant
8153/3	Gastrinoma, malignant
8155/3	Vipoma, malignant
8156/3	Somatostatinoma, malignant
8240/3	Carcinoid tumor, NOS
8241/3	Enterochromaffin cell carcinoid
8243/3	Goblet cell carcinoid
8244/3	Composite carcinoid
8245/1	Tubular carcinoid
8246/3	Neuroendocrine carcinoma, NOS
8249/3	Atypical carcinoid tumor
8683/0	Gangliocytic paraganglioma

#### Reference:

Bosman FT, Carneiro F, Hruban RH, Theise ND, editors. WHO Classification of Tumours of the Digestive System 4th edition, IARC, Lyon (2010).

#### **INCIDENCE**

Table 1

Patient cohorts by year of diagnosis including DCO cases and multiple primaries, and with proportion of deaths and active follow-up

				Prop.		Prop.
		DCO	Prop.	mult.	Prop.	actively
Year of	Cases	cases	DCO	primaries	deaths	followed
diagnosis	n	'n	%	્ર	%	%
1998	55			23.6	61.8	100.0
1999	62			22.6	50.0	91.9
2000	48			22.9	43.8	97.9
2001	54			38.9	48.1	100.0
2002	93			20.4	46.2	96.8 #
2003	94			29.8	51.1	94.7 #
2004	122			28.7	42.6	94.3 #
2005	118			33.1	45.8	94.1 #
2006	152			28.9	44.1	91.4 #
2007	183			29.5	40.4	79.8 # ##
2008	182			25.3	35.7	65.4
2009	183			25.1	38.8	64.5
2010	205			28.8	28.3	64.4
2011	215			23.3	22.8	69.8
2012	199			26.6	18.6	98.0 ###
1998-2012	1965			27.1	37.2	82.3

<sup>#</sup> The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

<sup>##</sup> Since 2007 the percentage of actively followed patients sharply declined compared to the previous years. This is a consequence of ambiguous data protection rules that currently forbid cancer registries in Bavaria to obtain the essential life status informations from competent registration offices.

<sup>###</sup> Please be aware that data of recent annual patient cohorts may not yet be fully processed. Therefore, the presented figures and tables are potentially related to different time periods as pointed out in the respective headlines or legends.

Table 1a

Patient cohorts by year of diagnosis and gender including DCO cases

Year of	All	Males	Females	Prop. males	
diagnosis	n	n	n	ર્જ	
1998	55 /	30	25	54.5	
1999	62/	32	30	51.6	
2000	48	23	25	47.9	
2001	54	30	24	55.6	
2002	93	52	41	55.9	
2003	94	49	45	52.1	
2004	122	69	53	56.6	
2005	/118	67	51	56.8	
2006	152	93	59	61.2	
2007	183	116	67	63.4	
2008	182	92	90	50.5	
2009	183	99	84	54.1	
2010	205	102	103	49.8	
2011	215	118	97	54.9	
2012	199	96	103	48.2	
1998-2012	1965	1068	897	54.4	

Table 2

Incidence measures by year of diagnosis and gender including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

			Males	Fem.	Males	Fem.	Males	Fem.	Males	Fem.
Year of	Males	Females	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	30	25	2.7	2.1	1.7	1.3	2.3	1.7	2.8	1.9
1999	32	30	2.9	2.5	2.0	1.6	2.5	2.0	2.9	2.2
2000	23	25	2.0	2.1	1.3	1.2	1.8	1.5	2.2	1.8
2001	30	24	2.6	2.0	1.6	1.2	2.3	1.6	2.6	1.8
2002	52	41	2.8	2.1	1.8	1.3	2.4	1.6	2.7	1.9
2003	49	45	2.6	2.3	1.5	1.4	2.2	1.9	2.6	2.1
2004	69	53	3.7	2.7	2.2	1.6	3.1	2.1	3.7	2.4
2005	67	51 <	3.5	2.6	2.1	1.3	2.9	1.8	3.6	2.2
2006	93	59	4.9	2.9	2.7	1.6	3.8	2.2	4.7	2.5
2007	116	67	5.2	2.9	3.3	1.7	4.4	2.2	5.1	2.6
2008	92	90	4.1	3.9	2.4	2.2	3.4	2.8	4.0	3.4
2009	99	84	4.4	3.6	2.5	2.1	3.5	2.8	4.2	3.3
2010	102	103	4.5	4.4	2.6	2.6	3.6	3.5	4.3	3.9
2011	118	97	5.2	4.1	2.8	2.5	4.0	3.1	4.9	3.6
2012	96	103	4.2	4.4	2.4	2.8	3.3	3.4	4.0	3.9
1998-2012	1068	897	3.9	3.1	2.3	1.9	3.2	2.4	3.8	2.8



The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

Table 3 Age distribution parameters by year of diagnosis (All) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	55	62.6	14.9	13.2	87.9	46.6	55.9	63.4	71.5	81.7
1999	62	58.7	15.4	24.9	87.9	33.5	52.5	60.7	67.7	73.7
2000	48	61.4	16.4	24,7	92.1	39.0	49.5	60.5	74.6	80.6
2001	54	61.8	12.6	29.6	88.5	47.7	54.0	62.4	69.1	77.8
2002	93	61.2	15.8	17.7	90.9	37.0	51.2	62.8	72.0	80.3
2003	94	62.7	13.0	23.5	87.5	48.8	56.2	63.4	72.3	78.9
2004	122	61.6	14.6	13.8	93.3	40.8	52.8	63.3	72.3	77.8
2005	118	64.8	14.5	16.1	90.8	45.4	56.9	66.8	74.9	82.5
2006	152	64.4	13.2	16.8	91.9	45.6	56.6	66.7	73.9	79.2
2007	183	61.9	15.0	15.8	91.2	41.0	54.8	64.2	71.2	79.4
2008	182	62.7	15.1	18.9	93.9	42.4	53.3	65.8	73.1	79.9
2009	183	63.7	16.2	12.4	92.6	38.8	55.5	66.4	75.3	84.0
2010	205	62.1	14.1	14.9	92.4	43.9	53.8	62.5	72.5	79.0
2011	215	63.1	15.5	17.1	92.9	42.3	52.0	65.1	74.4	81.4
2012	199	61.6	18.1	13.7	101	30.9	52.7	65.6	74.8	80.7
1998-2012	1965	62.5	15.2	12.4	101	41.7	54.2	64.4	73.2	80.0

Table 3a Age distribution parameters by year of diagnosis (MALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	30	62.9	10.7	32.1	81.7	51.2	58.6	63.5	66.8	77.2
1999	32	57.5	15.9	24.9	85.3	29.8	52.3	59.8	67.7	73.5
2000	23	61.8	14.2	37.4	92.1	43.9	50.0	60.1	73.9	76.6
2001	30	62.6	11.6	34.3	88.5	51.5	54.9	61.2	69.1	78.8
2002	52	60.0	12.5	27.1	88.3	42.3	52.7	61.0	66.6	74.9
2003	49	64.6	9.9	32.4	85.0	52.9	58.8	64.1	70.9	76.9
2004	69	62.4	11.8	27.8	78.9	47.7	54.1	63.4	71.4	76.5
2005	67	64.7	13.6	19.0	87.6	47.7	57.8	66.3	74.8	79.4
2006	93	64.7	12.3	16.8	85.7	48.1	59.1	67.2	73.5	77.0
2007	116	61.3	14.2	15.8	91.2	40.4	55.0	63.3	69.7	78.7
2008	92	62.7	13.0	19.3	85.6	48.2	54.7	64.2	71.7	79.1
2009	99	65.5	13.2	12.4	89.0	49.6	58.1	67.1	74.6	81.8
2010	102	63.9	12.4	26.3	92.4	49.0	56.4	63.5	72.5	79.0
2011	118	65.3	12.4	32.0	89.3	46.5	56.4	67.2	75.4	81.1
2012	96	63.7	15.0	25.8	89.0	42.5	53.8	66.0	75.0	80.4
1998-2012	1068	63.4	13.1	12.4	92.4	46.4	56.1	64.6	72.4	78.8

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	25	62.2	19.0	13.2	87.9	32.4	54.3	62.1	77.0	84.2
1999	30	59.9	14.9	26.9	87.9	38.6	52.5	61.6	72.1	76.0
2000	25	61.0	18.4	24,7	92.0	33.1	46.0	63.0	75.5	81.0
2001	24	60.9	13.9	29.6	84.4	38.6	53.7	63.0	70.0	75.7
2002	41	62.9	19.2	17.7	90.9	32.9	49.4	66.3	76.4	84.7
2003	45	60.7	15.5	23.5	87.5	33.3	53.0	61.7	72.5	79.4
2004	53	60.6	17.7	13.8	93.3	35.4	51.7	62.6	74.4	78.9
2005	51	65.0	15.7	16.1	90.8	42.1	55.6	67.8	77.4	82.5
2006	59	64.0	14.5	29.9	91.9	43.5	52.4	64.7	74.2	83.1
2007	67	62.8	16.4	17.8	88.4	41.0	53.5	65.2	74.3	81.7
2008	90	62.8	17.1	18.9	93.9	38.6	50.4	66.8	73.5	82.0
2009	84	61.6	19.1	15.9	92.6	35.0	49.9	63.4	76.5	84.9
2010	103	60.3	15.5	14.9	89.6	39.8	50.8	60.7	72.5	78.8
2011	97	60.5	18.4	17.1	92.9	34.6	47.7	63.2	74.4	82.6
2012	103	59.7	20.4	13.7	101	25.4	46.9	64.1	74.7	81.5
1998-2012	897	61.5	17.3	13.2	101	36.9	50.8	64.0	74.3	81.7



Table 4

Age distribution by 5-year age group and gender for period 1998-2012 (incl. DCO)

Age at									
diagnosis	Cases			Males			Females	S	
Years	n	%	Cum.%	n	96	Cum.%	n	%	Cum.%
10-14	5	0.3	0.3	1	0.1	0.1	4	0.4	0.4
15-19	20	1.0	1.3	4	0.4	0.5	16	1.8	2.2
20-24	18	0.9	2.2	2	0.2	0.7	16	1.8	4.0
25-29	34	1.7	3.9	16	1.5	2.2	18	2.0	6.0
30-34	43	2.2	6.1	17	1.6	3.7	26	2.9	8.9
35-39	53	2.7	8.8	18	1.7	5.4	35	3.9	12.8
40-44	73	3.7	12.5	36	3.4	8.8	37	4.1	16.9
45-49	120	6.1	18.6	59	5.5	14.3	61	6.8	23.7
50-54	154	7.8	26.5	87	8.1	22.5	67	7.5	31.2
55-59	210	10.7	37.2	128	12.0	34.5	82	9.1	40.4
60-64	281	14.3	51.5	177	16.6	51.0	104	11.6	52.0
65-69	292	14.9	66.3	183	17.1	68.2	109	12.2	64.1
70-74	248	12.6	78.9	134	12.5	80.7	114	12.7	76.8
75-79	218	11.1	90.0	125	11.7	92.4	93	10.4	87.2
80-84	121	6.2	96.2	57	5.3	97.8	64	7.1	94.3
85+	75	3.8	100.0	24	2.2	100.0	51	5.7	100.0
All ages	1965	100.0		1068	100.0		897	100.0	
=									

Included in the statistics are 39.3% multiple primaries in males and 27.3% in females.

Table 5 Age-specific incidence, DCO rate and proportion of all cancers for period 1998-2012

Age at diagnosis Years	Males n	Females n	Age- spec. incid.	incid.		Females DCO rate n=0 %	cancers	Females Prop.all cancers n=142297
0- 4 5- 9			0.0	0.0				
10-14	1	4	0.1	0.3			0.7	2.5
15-19	4	16	0.3	1.2			1.3	6.0
20-24	2	16	0.1	1.0			0.4	3.3
25-29	16	18	0.9	1.0			1.8	1.8
30-34	17	26	0.8	1.3			1.2	1.4
35-39	18	35	0.8	1.6			0.9	1.0
40-44	36	37	1.5	1.6			1.2	0.6
45-49	59	61	2.7	2.9			1.2	0.8
50-54	87	67	4.7	3.5			1.1	0.7
55-59 60-64	128 172	82 103	7.5 10.4	4.6 5.9			0.9	0.6
65-69	181	109	12.3	6.8			0.8	7 0.6
70-74	133	114	11.5	8.3			0.5	0.7
75-79	124	90	16.5	8.2			0.7	0.6
80-84	56	64	12.3	7.4			0.4	0.4
85+	24	51	7.7	6.2			0.3	0.3
All ages	1058	893			0.0	0.0	0.7	0.6
- ' 1								
Incidence Raw			3.9	3.1				
Kaw WS			2.3					
ES			3.2	2.4				
BRD-S			3.8	2.8				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

Table 6a Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2012 MALES

	Observed	Expected		LCL	UCL		DCO
Diagnosis	'n	n	SIR	95%	95%	EAR	%
C16 Stomach	/ 5 /	1.4	3.5	1.1	8.2 #	13.9	
C17 Small intestine	/ 2 /	0.2	10.9	1.3	39.3 #	7.1	
C18 Colon	16/	3.4	4.7	2.7	7.6 #	48.9	
C19-C20 Rectum	12	2.1	5.7	3.0	10.0 #	38.6	
C22 Liver	6	1.0	5.8	2.1	12.7 #	19.3	16.7
C23-C24 Bile	/ 2	0.3	6.0	0.7	21.5	6.5	
C25 Pancreas	7	1.3	5.6	2.2	11.5 #	22.3	
C33-C34 Lung	8	4.4	1.8	0.8	3.6	14.1	25.0
C43 Malign. melanoma	5	1.5	3.3	1.1	7.6 #	13.5	
C61 Prostate	21	10.9	1.9	1.2	3.0 #	39.5	
C64 Kidney	7	1.3	5.3	2.1	10.8 #	22.0	
C65 Renal pelvis	2	0.1	14.4	1.7	51.9 #	7.2	
C67 Bladder	2	1.5	1.4	0.2	5.0	2.1	
C73 Thyroid	2	0.3	7.2	0.9	26.1	6.7	
C76-C79 CUP	2	0.6	3.4	0.4	12.2	5.5	
C82-C85 NHL	5	1.4	3.6	_1.2	8.3 #	14.0	
C91-C96 Leukaemia	2	0.6	3.6/	0.4	13.1	5.6	100.0
Other primaries	7	2.2	3.2	1.3	6.5 #	18.7	
Not observed	0	2.4	0.0	0.0	1.5	-9.4	
All mult. primaries	113	36.9	3.1	2.5	3.7 #	296.1	4.4

Patients	721
Mean age at second malignancy (years)	68.6
Person-years	2571
Mean observation time (years)	3.6
Median observation time (years)	2.5

# The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries".

Table 6b

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2012 FEMALES

	Observed Ex	rpected		LCL UCL		DCO
Diagnosis	ń	n	SIR	95% 95%	EAR	%
C16 Stomach	5	0.8	6.6	2.1 15.3	# 19.6	40.0
C18 Colon	/ 11 /	2.1	5.2	2.6 9.4	# 41.1	
C19-C20 Rectum	6	0.9	6.5	2.4 14.0	# 23.4	16.7
C25 Pancreas	/ 7/	0.9	7.8	3.1 16.1	# 28.2	
C33-C34 Lung	5	1.5	3.3	1.1 7.8	# 16.2	20.0
C43 Malign. melanoma	2	0.8	2.6	0.3 9.2	5.6	
C50 Breast	10	6.7	1.5	0.7 2.7	15.3	
C54 Corpus uteri	2	1.2	1.7	0.2 6.0	3.7	
C56 Ovary	3	0.9	3.3	0.7 9.8	9.7	33.3
C64 Kidney	3	0.5	5.7	1.2 16.7	# 11.4	33.3
C82-C85 NHL	5	0.8	6.2	2.0 14.5	# 19.4	
Other primaries	5	1.2	4.1	1.3 9.5	# 17.4	
Not observed	0	3.3	0.0	0.0 1.1	-15.1	
All mult. primaries	64	21.6	3.0	2.3 3.8	# 195.8	9.4

Patients	562
Mean age at second malignancy (years)	70.5
Person-years	2166
Mean observation time (years)	3.9
Median observation time (years)	2.8

# The occurrence of second malignancy is statistically significant.

Observed second malignancies with count 1 are pooled in category "Other primaries".

## GEP-NET: Gastroenteropancreatic neuroendocrine tumor






Figure 7. Age distribution and age-specific incidence







**Figure 7a.** Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).



#### Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER\*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2012, based on the November 2011 submission. http://www.seer.cancer.gov.







Figure 8. Cumulative follow-up years depending on time since diagnosis


The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.



#### Average incidence (world standard population) 2003 - 2008: Males




#### Average incidence (world standard population) 2003 - 2008: Females



**Figure 9a.** Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 2.4/100,000 WS N=463, females 1.6/100,000 WS N=352). Since cancer data are not available in some counties until 2007, the local incidence rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 11 women were identified with newly diagnosed gastr.ent.pancr. neuroend. tumor. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 1.4/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.5 and 3.3/100,000.

#### Standardized incidence ratio (SIR) 2003 - 2008: Males



**Figure 9b.** Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=463, females N=352). Since cancer data are not available in some counties until 2007, the local SIR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 11 women were identified with newly diagnosed gastr.ent.pancr. neuroend. tumor. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 1.03. Though, the value of this parameter may vary with an underlying probability of 99% between 0.41 and 2.14, and is therefore not statistically striking.

#### **MORTALITY**

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts, and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

		Prop.				Prop. deaths
	Incident	actively	Prop.		Prop.	with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	96	%	n	%	%
1998	55	100.0		34	61.8	91.2
1999	62	91.9		31	50.0	100.0
2000	48	97.9		21	43.8	90.5
2001	54	100.0		26	48.1	92.3
2002	93	96.8		43	46.2	97.7
2003	94	94.7		48	51.1	97.9
2004	122	94.3		52	42.6	98.1
2005	118	94.1		54	45.8	98.1
2006	152	91.4		67	44.1	98.5
2007	183	79.8		74	40.4	94.6
2008	182	65.4		65	35.7	96.9
2009	183	64.5		71	38.8	98.6
2010	205	64.4		58	28.3	94.8
2011	215	69.8		49	22.8	95.9
2012	199	98.0		37	18.6	94.6
1998-2012	1965	82.3		730	37.2	96.4

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

(with respect to registry area expansion from  $2.51\ \mathrm{to}\ 3.96\ \mathrm{m}$  as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

			Prop.		
			deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	/ n /	%	n	%
1998	55	/ 1/1	81.8	5	9.1
1999	62	16	100.0	7	11.3
2000	48	23	95.7	6	12.5
2001	54	15	100.0	3	5.6
2002	93	29	100.0	/ 11	11.8
2003	94	41	97.6	14	14.9
2004	122	44	100.0	11	9.0
2005	118	54	94.4	20	16.9
2006	152	51	96.1	15	9.9
2007	183	72	97.2	21	11.5
2008	182	83	96.4	23	12.6
2009	183	81	98.8	28	15.3
2010	205	84	97.6	23	11.2
2011	215	80	97.5	23	10.7
2012	199	97	97.9	29	14.6
1998-2012	1965	781	97.3	239	12.2

Table 10c

Annual cohorts of deaths, proportion of cancer-related and not cancerrelated deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from  $2.51\ \mathrm{to}\ 3.96\ \mathrm{m}$  as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

				Prop. cancer	
		Prop.	Prop.	recorded	
		/ =		on death	
		cancer-	not cancer-		
Year of	Deaths	related	related	certificate	
death	n	%	8	%	
1998	11/	81.8	18.2	66.7	
1999	16	87.5	12.5	93.8	
2000	23	69.6	30.4	77.3	
2001	15	80.0	20.0	86.7	
2002	29	86.2	13.8	89.7	
2003	41	70.7	29.3	82.5	
2004	44	75.0	25.0	79.5	
2005	54	70.4	29.6	80.4	
2006	51	86.3	13.7	89.8	
2007	72	83.3	16.7	84.3	
2008	83	86.7	13.3	93.8	
2009	81	70.4	29.6	78.8	
2010	84	76.2	23.8	78.0	
2011	80	80.0	20.0	88.5	
2012	97	77.3	22.7	81.1	
1998-2012	781	78.4	21.6	83.8	

Table 11a Means of age at death according to the grouping in Table 10 MALES

Year of	Deaths	Age at death (all causes)	Age at death (cancer-related)	Age at death (not cancer- related)	Age at death (according to death certificate)
death	n	Years	Years	Years	Years
1998	5	69.0	71.4	59.3	68.5
1999	12	67.8	65.6	78.7	66.3
2000	16	68.4	65.4	81.2	65.4
2001	12	70.1	67.1	84.8	68.5
2002	21	67.2	65.9	92.4	67.1
2003	20	71.5	69.6	75.1	70.6
2004	19	70.2	67.9	73.5	68.5
2005	33	72.6	72.2	74.0	72.8
2006	29	72.2	71.3	77.8	71.3
2007	37	70.0	70.2	69.0	70.2
2008	56	67.9	66.9	76.1	67.2
2009	50	69.4	68.2	71.9	68.1
2010	51/	71.3	70.0	76.1	69.8
2011	46	69.8	68.5	77.2	68.8
2012	49	74.1	72.3	79.7	72.3
1998-2012	456	70.4	69.1	75.5	69.3




Table 11b Means of age at death according to the grouping in Table 10 FEMALES

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(not cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	6	82.4	81.2	88.1	82.2
1999	4	78.9	78.9		78.9
2000	7	71.1	66.0	75.0	71.6
2001	3	77.4	69.9	92.2	69.9
2002	8	74.1	73.3	75.4	71.6
2003	21	69.8	67.0	78.6	67.7
2004	25	71.3	70.8	74.9	72.0
2005	21	76.7	72.5	83.5	74.0
2006	22	75.5	75.4	75.6	75.7
2007	35	68.3	67.0	73.3	68.3
2008	27	74.4	71.7	86.4	74.2
2009	31	72.7	69.2	82.6	71.1
2010	33/	70.7	66.1	82.9	68.9
2011	34	71.1	67.5	81.3	69.8
2012	48	75.9	72.3	89.6	74.0
1998-2012	325	72.9	70.3	81.6	71.8

By 2010, life expectancy for a newborn male in Germany is 77.5 years compared with 82.6 years for his female counterpart.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 12a Mortality measures (cancer-related death) and mortality-incidence-index by year of death MALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	4	0.4	0.13	0.2	0.13	0.3	0.15	0.5	0.17
1999	10	0.9	0.31	0.6	0.29	0.8	0.31	1.0	0.33
2000	13	1.1	0.57	0.7	0.58	1.1	0.59	1.2	0.57
2001	10	0.9	0.34	0.5	0.32	0.7	0.33	1.0	0.38
2002	20	1.1	0.38	0.6	0.35	0.9	0.37	1.1	0.39
2003	13	0.7	0.27	0.4	0.25	0.6	0.27	0.8	0.30
2004	11	0.6	0.16	0.3	0.14	0.5	0.16	0.6	0.17
2005	25	1.3	0.38	0.7	0.31	1.0	0.35	1.4	0.40
2006	25	1.3	0.27	0.6	0.23	1.0	0.26	1.4	0.29
2007	32	1.4	0.28	0.7	0.23	1,1	0.25	1.4	0.29
2008	50	2.2	0.54	1.3	0.51	1.8	0.53	2.2	0.55
2009	34	1.5	0.35	0.8	0.31	1.2	0.33	1.4	0.35
2010	40	1.8	0.40	0.9	0.36	1.3	0.38	1.8	0.42
2011	39	1.7	0.33	0.9	0.31	1.3	0.32	1.6	0.32
2012	37	1.6	0.39	0.7	0.31	1.1	0.35	1.6	0.41
1998-2012	363	1.3	0.34	0.7	0.30	1.0	0.33	1.4	0.36

Table 12b Mortality measures (cancer-related death) and mortality-incidence-index by year of death FEMALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	5	0.4	0.20	0.1	0.08	0.2	0.12	0.4	0.21
1999	4	0.3	0.13	0.1	0.08	0.2	0.09	0.2	0.10
2000	3	0.2	0.12	0.1	0.11	0.2	0.12	0.2	0.10
2001	2	0.2	0.08	0.1	0.08	0.1	0.08	0.2	0.08
2002	5	0.3	0.12	0.1	0.09	0.2	0.10	0.2	0.10
2003	16	0.8	0.36	0.4	0.30	0.6	0.31	0.7	0.33
2004	22	1.1	0.42	0.5	0.29	0.7	0.34	0.9	0.38
2005	13	0.7	0.25	0.3	0.19	0.4	0.21	0.5	0.24
2006	19	0.9	0.33	0.3	0.21	0.5	0.25	0.7	0.30
2007	28	1.2	0.42	0.6	0.37	0.8	0.39	1.0	0.40
2008	22	0.9	0.24	0.4	0.17	0.5	0.19	0.8	0.22
2009	23	1.0	0.28	0.4	0.21	0.6	0.23	0.8	0.23
2010	24	1.0	0.23	0.5	0.20	0.7	0.21	0.9	0.23
2011	25	1.1	0.26	0.5	0.20	0.7	0.22	0.9	0.24
2012	38	1.6	0.37	0.7	0.23	1.0	0.29	1.2	0.31
1998-2012	249	0.9	0.28	0.4	0.21	0.6	0.23	0.7	0.25

Table 13 Age distribution of age at death (cancer-related) for period 1998-2012 (incl. multiple primaries)

Age at								
death	Cases		Males			Females		
Years	n	% Cum.%	n	%	Cum.%	n	%	Cum.%
15-19	1	0.2 0.2	1	0.3	0.3			0.0
20-24	0	0.0 0.2			0.3			0.0
25-29	3	0.5 0.6	/ 1	0.3	0.5	2	0.8	0.8
30-34	4	0.6 1.3			0.5	4	1.6	2.4
35-39	7	1.1 2.4	_ 2	0.5	1.1	5	2.0	4.4
40 - 44	10	1.6 4.1	4	1.1	2.2	6	2.4	6.8
45-49	12	1.9 6.0	4	1.1	3.3	8	3.2	10.0
50-54	38	6.2 12.2	24	6.6	9.8/	14	5.6	15.6
55-59	52	8.4 20.6	33	9.0	18.9	19	7.6	23.2
60-64	71	11.5 32.1	49	13.4	32.2	22	8.8	32.0
65-69	103	16.7 48.9	74	20.2	52.5	29	11.6	43.6
70-74	91	14.8 63.6	57	15.6	68.0	34	13.6	57.2
75-79	93	15.1 78.7	60	16.4	84.4	33	13.2	70.4
80-84	80	13.0 91.7	40	10.9	95.4	40	16.0	86.4
85+	51	8.3 100.0	17	4.6	100.0	34	13.6	100.0
All ages	616	100.0	366	100.0		250	100.0	

Included in the statistics are 39.3% multiple primaries in males and 27.3% in females.

Table 14 Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2012 (incl. multiple primaries)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males	Females	spec.		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0- 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19	1		0.1	0.25	0.0		2.4	
20-24			0.0		0.0			
25-29	1	2 /	0.1	0.06	0.1	0.11	1.0	1.8
30-34		4	0.0		0.2			1.9
35-39	2	5	0.1	0.11	0.2		0.5	1.0
40-44	4	6	0.2		0.3	0.16	0.5	0.6
45-49	4	8	0.2	0.07	0.4		0.2	0.4
50-54	24	14	1.3	0.28	0.7	0.21	0.8	0.5
55-59	33	19	1.9		1.1		0.6	0.4
60-64	49	22	3.0		1.3	0.21	0.6	0.4
65-69	74	29	5.0		1.8	0.27	0.7	0.4
70-74	57	34	4.9		2.5		0.5	0.4
75-79	60	33	8.0	0.48	3.0	0.35	0.5	0.3
80-84	40	40	8.8		4.6		0.4	0.4
85+	17	34	5.5	0.71	4.2	0.67	0.2	0.3
All ages	366	250					0.5	0.4
Mortality								
Raw			1.3		0.9	0.28		
WS			0.7		0.4			
ES			1.0		0.6	0.23		
BRD-S			1.4	0.36	0.7	0.25		
PYLL-70								
per 100,000			7.0		6.0			
ES 100,000			6.3		5.2			
AYLL-70			9.1		13.4			
YITT - / 0			9.1		13.4			

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15a

Multiple primaries in deaths in period 1998-2012

MALES

	_	_			Syn- chron	Syn- chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n /	%↓	n	<b>←%</b>	n	<b>←</b> %	n	<b>←%</b>
G00 G10 0 1	_	0.6	1	100 0				
C09-C10 Oropharynx	1	0.6	1	100.0			-	100 0
C12-C13 Hypopharynx	/1	0.6					1	100.0
C15 Oesophagus	/ 1	0.6					1	100.0
C16 Stomach	4	2.5			_		4	100.0
C17 Small intestine	16	9.9			7	43.8	9	56.3
C18 Colon	/ 17 /	10.6			15	88.2	2	11.8
C19-C20 Rectum	/ 7 \	4.3			4	57.1	3	42.9
C21 Anus/canal	1	0.6					1	100.0
C22 Liver	4	2.5			1	25.0	3	75.0
C23-C24 Bile	2	1.2			1	50.0	1	50.0
C25 Pancreas	10	6.2			4	40.0	6	60.0
C32 Larynx	2	1.2	2	100.0				
C33-C34 Lung	11	6.8	3	27.3	2	18.2	6	54.5
C38,C45 Mesothelioma	1	0.6					1	100.0
C43 Malign. melanoma	4	2.5	2	50.0			2	50.0
C44 Skin others	6	3.7	3	50.0			3	50.0
C46,C49 Soft tissue	2	1.2	1	50.0			1	50.0
C50 Breast	1	0.6	1	100.0				
C61 Prostate	35	21.7	26	74.3	3	8.6	6	17.1
C62 Testis	2	1.2	2	100.0				
C64 Kidney	6	3.7	3	50.0	2	33.3	1	16.7
C65 Renal pelvis	2	1.2					2	100.0
C66 Ureter	1	0.6					1	100.0
C67 Bladder	10	6.2	7	70.0			3	30.0
C70-C72 CNS cancer	3	1.9	•		1/	33.3	2	66.7
C76-C79 CUP	3	1.9	2	66.7	1	33.3	_	00.7
C82-C85 NHL	5	3.1	2	40.0	1	20.0	2	40.0
C91-C96 Leukaemia	3	1.9		10.0	_	20.0	3	100.0
COT COO LICUNACIIITA	J	1.5					5	100.0
All mult. primaries	161	100.0	55	34.2	42	26.1	64	39.8
AII maic. Piimaiies	101	100.0	23	J <b>T.</b> 4	74	∠0.1	υ <del>ı</del>	39.0

Multiple primaries with number of cases n<1 are pooled in category "Other primaries".

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 15b Multiple primaries in deaths in period 1998-2012 FEMALES

						Syn- chron	Syn- chron		
		Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnos	is	n /	왕↓	n	<b>←</b> %	n	<b>←</b> %	n	<b>←</b> %
C15	Oesophagus	1	1.1					1	100.0
C16	Stomach	/7	7.4			2	28.6	5	71.4
C17	Small intestine	4	4.2			3	75.0	1	25.0
C18	Colon	10	10.5			5	50.0	5	50.0
C19-C20	Rectum	4	4.2			2	50.0	2	50.0
C25	Pancreas	3	3.2			1	33.3	2	66.7
C33-C34	Lung	10	10.5	4	40.0			6	60.0
C43	Malign. melanoma	2	2.1	2	100.0				
C48	Peritoneal	1	1.1	1	100.0				
C50	Breast	21	22.1	15	71.4	2	9.5	4	19.0
C51	Vulva	2	2.1	2	100.0				
C53	Cervix uteri	1	1.1			1	100.0		
C54	Corpus uteri	5	5.3	3	60.0	2	40.0		
C56	Ovary	8	8.4	2	25.0	5	62.5	1	12.5
C64	Kidney	3	3.2	1	33.3	1	33.3	1	33.3
C67	Bladder	3	3.2	3	100.0				
C70-C72	CNS cancer	2	2.1					2	100.0
C73	Thyroid	1	1.1	1	100.0				
C76-C79	_	1	1.1					1	100.0
C81	Hodgkin lymphoma	1	1.1	1	100.0				
C82-C85		4	4.2	2	50.0			2	50.0
C91-C96	Leukaemia	1	1.1					1	100.0
All mul	t. primaries	95	100.0	37	38.9	24	25.3	34	35.8

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 16 Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2012 (Singular primaries only \*)

Acc at			Males Age-		Females Age-		Males	Females Prop.all
Age at death	Males	Females			spec.		cancers	cancers
Years	n	n		MI-index		MT-index		%
10012				/	01 00.11	\	ŭ	v
0- 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19	1		0.1	0.25	0.0		2.6	
20-24			0.0		0.0			
25-29	1	2 /	0.1	0.06	0.1	0.12	1.1	1.9
30-34		2 <	0.0		0.1	0.09		1.1
35-39	2	3	0.1		0.1	0.09	0.6	0.7
40-44	3	6	0.1		0.3	0.17	0.4	0.6
45-49	4	7	0.2		0.3	0.12	0.3	0.4
50-54	23	11	1.2		0.6	0.19	0.9	0.5
55-59	23	15	1.4		0.8	0.22	0.5	0.4
60-64	40	21	2.4		1.2	0.24	0.6	0.4
65-69	59	20	4.0		1.2	0.25	0.6	0.3
70-74	41	25	3.5	0.42	1.8	0.29	0.4	0.3
75-79	38	22	5.0		2.0	0.33	0.4	0.3
80-84	22	28	4.8		3.2	0.57	0.3	0.3
85+	12	26	3.9	0.86	3.2	0.65	0.2	0.3
	\	_ \ _					\	
All ages	269	188					0.5	0.3
Mortality			1.0	0 22	0 7	0.06		
Raw			1.0	0.33	0.7	0.26		
WS			0.5	0.29	0.3	0.19		
ES			0.8	0.32	0.4	0.21		
BRD-S			1.0	0.34	0.5	0.23		
PYLL-70								
per 100,000			5.9		4.8			
ES			5.4		4.1			
AYLL-70			9.4		13.4			


<sup>\*</sup> See corresponding tables with multiple primaries.

Table 17 Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2012 (Single primaries only \*)

Age at death	Males	Females	Males Age- spec.		Females Age- spec.		Males Prop.all cancers	Females Prop.all cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	8	%
0 - 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19	1		0.1	0.25	0.0		2.6	
20-24			0.0		0.0			
25-29	1	2 /	0.1	0.07	0.1	0.12	1.2	2.1
30-34		2	0.0		0.1			1.2
35-39	2	3	0.1		0.1	0.10	0.6	0.7
40-44	3	6	0.1		0.3	0.17	0.4	0.7
45-49	4	6	0.2	0.08	0.3	0.11	0.3	0.4
50-54	23	11	1.2	0.34	0.6	0.21	0.9	0.5
55-59	23	12	1.4	0.23	0.7	0.19	0.5	0.4
60-64	39	20	2.4	0.33	1.1	0.25	0.6	0.5
65-69	52	16	3.5	0.43	1.0	0.22	0.7	0.3
70-74	31	24	2.7	0.36	1.7	0.29	0.4	0.4
75-79	31	21	4.1	0.46	1.9	0.33	0.4	0.3
80-84	22	25	4.8	0.85	2.9	0.56	0.4	0.4
85+	9	23	2.9	0.69	2.8	0.59	0.2	0.3
All ages	241	171					0.5	0.4
Mortality								
Raw			0.9	0.33	0.6	0.25		
WS			0.5	0.29	0.3	0.18		
ES			0.7	0.31	0.4	0.20		
BRD-S			0.9	0.34	0.5	0.22		
PYLL-70								
per 100,000			5.8		4.4			
ES			5.3		3.8			
AYLL-70			9.8		14.0			

<sup>\*</sup> See corresponding tables with multiple primaries.

## GEP-NET: Gastroenteropancreatic neuroendocrine tumor




**Figure 18.** Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).


The difference between age at diagnosis (Table 3) and age at gastr.ent.pancr. neuroend. tumor-related death (see Table 10) should be considered.



#### Average mortality (world standard population) 2003 - 2008: Males




#### Average mortality (world standard population) 2003 - 2008: Females





**Figure 19a.** Map of cancer mortality (world standard population) by county averaged for period 2003 to 2008. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.7/100,000 WS N=147, females 0.4/100,000 WS N=114). Since cancer data are not available in some counties until 2007, the local mortality rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 3 women died from gastr.ent.pancr. neuroend. tumor. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.3/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.5/100,000.

### Standardized mortality ratio (SMR) 2003 - 2008: Males







**Figure 19b.** Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=147, females N=114). Since cancer data are not available in some counties until 2007, the local SMR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 3 women died from gastr.ent.pancr. neuroend. tumor. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.91. Though, the value of this parameter may vary with an underlying probability of 99% between 0.10 and 3.32, and is therefore not statistically striking.

#### **Statistical Notes**

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

#### 1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the tumor-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the age at death, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, MI-index) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and agespecific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (PYLL, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (AYLL, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

#### **Shortcuts**

AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

BRD-S German standard population

DCO Death certificate only EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

ES European standard population (old)
FRG Federal Republic of Germany

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

LCL Lower confidence limit

MI-index Ratio between mortality and incidence

MCR Munich Cancer Registry (Tumorregister München)

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age

SEER Surveillance, Epidemiology, and End Results (USA)

SIR Standardized incidence ratio
SMR Standardized mortality ratio
UCL Upper confidence limit
WS World standard population

#### **Recommended Citation**

Munich Cancer Registry. Baseline statistics GEP-NET: Gastr.ent.pancr. neuroend. tumor [Internet]. 2014 [updated 2014 Mar 20; cited 2014 May 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/base/base\_hDNETE.pdf

#### Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

#### **Disclaimer**

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

# Index of figures and tables

I.	Page
Pts cohorts, DCO, mult. prim., follow-up / yr	4
Gender distribution by year of diagnosis	5
Incidence by year of diagnosis	6
Age distribution parameters by year of diagnosis	7
Age distribution by 5-year age group and gender	9
Age-specific incidence and DCO rate	10
Standardized incidence ratio of second primaries	11
Age distribution and age-specific incidence (chart)	13
Age-specific incidence internationally (chart)	14
Cumulative follow-up years (chart)	15
Map of cancer incidence (WS) by county (chart)	16
Standardized incidence ratio (SIR) by county (chart)	17
Pts incident cohorts and mortality / yr	18
Incidence and mortality by year of diagnosis	19
Cancer-related deaths, death certification available / yr	20
Means of age at death / yr	21
Mortality by year of death	23
Distribution of age at death	24
Age-specific mortality	25
Multiple primaries in deaths	26
Age-specific mortality (first primaries)	28
Age-specific mortality (single primaries)	29
Age distribution and age-specific mortality (chart)	30
Map of cancer mortality (WS) by county (chart)	31
Standardized mortality ratio (SMR) by county (chart)	32
	Pts cohorts, DCO, mult. prim., follow-up / yr Gender distribution by year of diagnosis Incidence by year of diagnosis Age distribution parameters by year of diagnosis Age distribution by 5-year age group and gender Age-specific incidence and DCO rate Standardized incidence ratio of second primaries Age distribution and age-specific incidence (chart) Age-specific incidence internationally (chart) Cumulative follow-up years (chart) Map of cancer incidence (WS) by county (chart) Standardized incidence ratio (SIR) by county (chart) Pts incident cohorts and mortality / yr Incidence and mortality by year of diagnosis Cancer-related deaths, death certification available / yr Means of age at death / yr Mortality by year of death Distribution of age at death Age-specific mortality Multiple primaries in deaths Age-specific mortality (first primaries) Age-specific mortality (single primaries) Age distribution and age-specific mortality (chart) Map of cancer mortality (WS) by county (chart)