Munich Cancer Registry



- Survival
- Selection Matrix
- ► Homepage

Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

Cancer statistics: Baseline statistics

Year of diagnosis	1998-2012
Patients	955
Diseases	957
Creation date	03/20/2014
Export date	02/12/2014
Population	4.5 m



http://www.tumorregister-muenchen.de/en/facts/base/base_hGISTE.pdf

GIST: Gastroint. stromal tumor

Global Statements about the statistics on the Internet -

Baseline Statistics (grey button ___), Survival (red button ___)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.5 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases^{###} are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR. The time-delayed acquisition of data and the occasionally high DCO-rates indicate optimizing reserves, among others, because of current financial and legal conditions that hinder the analyses.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, March 2014

- [#] Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007). Death certificates from 2013 are incorporated into these analyses.
- ^{##} Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- ^{###} DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate. A high proportion of DCO cases (≥5%) in particular cancer types indicate insufficient participation of specific cancer specializations.

ICD-O-3 codes (morphology) used for specifying cancer site

Description	
Gastrointestinal stromal tumor, benign Gastrointestinal stromal tumor, NOS Gastrointestinal stromal sarcoma	
	Gastrointestinal stromal tumor, benign Gastrointestinal stromal tumor, NOS

INCIDENCE

Table 1

Patient cohorts by year of diagnosis including DCO cases and multiple primaries, and with proportion of deaths and active follow-up

		DCO	Prop.	Prop. mult.	Prop.	Prop. actively
Year of	Cases	cases	DCO	primaries	deaths	followed
diagnosis	n	n	00	90	00	ଌ
1998	4			25.0	100.0	100.0
1999	8				75.0	100.0
2000	18			38.9	66.7	94.4
2001	28			35.7	60.7	100.0
2002	48			29.2	58.3	97.9 #
2003	50			34.0	36.0	96.0 #
2004	73			28.8	46.6	97.3 #
2005	67			35.8	37.3	92.5 #
2006	62			24.2	19.4	88.7 #
2007	91			27.5	26.4	75.8 # ##
2008	92			30.4	22.8	55.4
2009	99			36.4	20.2	53.5
2010	103			35.9	17.5	53.4
2011	100			38.0	10.0	57.0
2012	114			38.6	9.6	95.6 ###
1998-2012	957			33.1	27.2	76.7

The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

- ## Since 2007 the percentage of actively followed patients sharply declined compared to the previous years. This is a consequence of ambiguous data protection rules that currently forbid cancer registries in Bavaria to obtain the essential life status informations from competent registration offices.
- ### Please be aware that data of recent annual patient cohorts may not yet be fully processed. Therefore, the presented figures and tables are potentially related to different time periods as pointed out in the respective headlines or legends.

Table 1a

Patient cohorts by year of diagnosis and gender including DCO cases

Year of	All	Males	Females	Prop. males	
diagnosis	n	n	n	00	
1998	4	1	3	25.0	
1999	8	5	3	62.5	
2000	18	9	9	50.0	
2001	28	17	11	60.7	
2002	48	24	24	50.0	
2003	50	23	27	46.0	
2004	73	36	37	49.3	
2005	67	33	34	49.3	
2006	62	27	35	43.5	
2007	91	49	42	53.8	
2008	92	45	47	48.9	
2009	99	50	49	50.5	
2010	103	59	44	57.3	
2011	100	60	40	60.0	
2012	114	59	55	51.8	
1998-2012	957	497	460	51.9	

Incidence measures by year of diagnosis and gender including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

Year of diagnosis	Males n	Females n	Males Inc. raw	Fem. Inc. raw	Males Inc. WS	Fem. Inc. WS	Males Inc. ES	Fem. Inc. ES	Males Inc. BRD-S	Fem. Inc. BRD-S
1998	1	3	0.1	0.3	0.0	0.1	0.1	0.2	0.1	0.2
1999	5	3	0.4	0.3	0.3	0.2	0.5	0.2	0.5	0.2
2000	9	9	0.8	0.7	0.4	0.4	0.7	0.6	0.8	0.7
2001	17	11	1.5	0.9	0.8	0.5	1.3	0.7	1.7	0.8
2002	24	24	1.3	1.2	0.7	0.8	1.1	1.0	1.4	1.1
2003	23	27	1.2	1.4	0.8	0.7	1.0	1.0	1.2	1.2
2004	36	37	1.9	1.9	1.1	1.0	1.5	1.4	2.0	1.7
2005	33	34	1.7	1.7	1.0	0.8	1.4	1.1	1.7	1.4
2006	27	35	1.4	1.7	0.8	0.9	1.2	1.3	1.3	1.5
2007	49	42	2.2	1.8	1.3	0.9	1.9	1.3	2.2	1.6
2008	45	47	2.0	2.0	1.1	1.0	1.6	1.4	1.9	1.6
2009	50	49	2.2	2.1	1.2	0.9	1.8	1.4	2.1	1.8
2010	59	44	2.6	1.9	1.4	1.0	2.0	1.4	2.5	1.6
2011	60	40	2.6	1.7	1.3	0.8	1.9	1.1	2.4	1.4
2012	59	55	2.6	2.3	1.4	1.0	1.9	1.5	2.3	1.9
1998-2012	497	460	1.8	1.6	1.0	0.8	1.5	1.1	1.8	1.4

The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	4	80.5	9.8	69.7	90.0	69.7	72.2	81.1	88.8	90.0
1999	8	61.9	12.9	46.4	88.3	46.4	52.2	62.5	65.7	88.3
2000	18	61.2	14.3	36.6	79.1	37.6	55.1	62.0	73.2	77.9
2001	28	65.6	11,9	31.9	80.5	52.7	57.9	67.9	73.9	80.3
2002	48	65.5	13.3	19.3	89.0	51.5	57.4	65.8	75.8	81.6
2003	50	64.4	13.7	17.9	92.1	49.7	58.2	65.5	72.7	80.7
2004	73	64.8	14.5	16.7	87.8	47.1	56.3	65.1	76.2	80.9
2005	67	67.5	12.9	34.3	89.4	48.8	60.4	69.0	77.6	82.3
2006	62	65.0	12.2	35.9	90.5	50.9	57.2	65.3	72.7	80.0
2007	91	66.7	9.9	46.5	88.3	52.4	59.4	66.8	74.0	79.6
2008	92	66.6	13.9	26.5	97.0	45.9	59.5	68.9	74.3	82.7
2009	99	66.8	12.4	32.6	93.5	49.8	56.8	68.7	77.0	81.4
2010	103	66.9	11.2	30.6	88.5	52.7	59.2	67.6	75.2	81.4
2011	100	66.9	14.4	30.0	90.6	44.6	54.8	71.6	77.7	83.6
2012	114	69.4	10.7	34.9	91.8	54.8	62.7	69.9	77.4	82.2
1998-2012	957	66.6	12.6	16.7	97.0	50.2	58.5	67.7	75.7	81.7

Age distribution parameters by year of diagnosis (All) (incl. DCO)

Table 3a

Age distribution parameters by year of diagnosis (MALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	1	87.5		87.5	87.5	87.5	87.5	87.5	87.5	87.5
1999	5	61.2	17.0	46.4	88.3	46.4	51.4	53.0	67.1	88.3
2000	9	60.7	15.8	36.6	79.1	36.6	55.1	64.0	72.5	79.1
2001	17	66.5	12.8	31.9	80.5	53.6	58.2	70.2	76.7	80.4
2002	24	69.1	10.9	44.8	89.0	54.6	61.3	68.4	79.0	81.6
2003	23	62.0	15.4	17.9	92.1	39.2	57.1	64.7	69.5	77.9
2004	36	64.1	14.9	16.7	84.6	41.0	57.2	65.3	76.1	80.5
2005	33	64.9	13.1	34.3	89.4	44.2	60.4	67.2	74.6	77.7
2006	27	62.3	12.8	35.9	89.2	46.3	53.7	62.0	68.2	80.0
2007	49	65.2	9.3	47.4	83.2	52.0	58.5	64.3	72.3	78.8
2008	45	65.5	12.2	26.5	88.0	46.3	60.6	68.8	73.2	78.4
2009	50	64.6	12.7	40.8	93.5	48.1	53.5	64.9	74.9	79.5
2010	59	67.0	12.4	30.6	88.5	51.3	57.6	68.5	75.5	82.5
2011	60	68.1	14.6	30.0	90.6	46.5	55.7	72.3	78.9	85.1
2012	59	68.0	10.4	34.9	91.0	54.1	63.0	68.5	73.7	80.8
1998-2012	497	65.8	12.7	16.7	93.5	49.7	58.1	67.2	74.7	80.8

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of diagnosis	Cases n	Mean	Std. dev.	Min.	Max.	10%	25%	Median 50%	75%	90%
1998	3	78.2	10.6	69.7	90.0	69.7	69.7	74.8	90.0	90.0
1999	3	63.1	1.3	61.7	64.3	61.7	61.7	63.2	64.3	64.3
2000	9	61.6	13.5	41,1	77.9	41.1	55.2	60.0	73.2	77.9
2001	11	64.1	10.7	40.4	79.2	52.7	57.0	66.9	69.9	73.7
2002	24	61.9	14.6	19.3	85.9	46.8	52.6	62.4	73.6	79.1
2003	27	66.5	12.1	27.1	83.9	56.0	59.6	66.9	74.2	81.8
2004	37	65.5	14.2	22.1	87.8	47.9	56.2	64.6	76.9	81.0
2005	34	70.1	12.3	43.8	85.7	51.2	64.3	73.0	80.8	84.0
2006	35	67.2	11.4	39.5	90.5	55.6	58.4	65.7	76.2	84.7
2007	42	68.5	10.4	46.5	88.3	56.9	60.6	68.2	77.0	81.5
2008	47	67.6	15.4	27.2	97.0	44.3	57.2	70.7	78.4	86.3
2009	49	69.2	11.8	32.6	89.4	52.8	61.3	70.0	78.4	84.6
2010	44	66.8	9.4	47.1	85.0	53.9	60.1	67.1	72.8	79.8
2011	40	65.1	14.2	34.4	86.6	44.3	50.5	69.5	74.9	81.8
2012	55	70.9	10.9	47.2	91.8	55.0	62.4	72.3	78.8	82.8
1998-2012	460	67.4	12.5	19.3	97.0	51.5	59.5	68.2	76.9	82.7

Age at diagnosis	Cases		Males			Females		
Years	n	% Cum.		olo	Cum.%	n	00	Cum.%
IEals	11		5 11	6	Culli. o	11	6	Culli. 8
1 - 1 0	2			0.4		-	0 0	
15-19	3	0.3 0.3		0.4	0.4	1	0.2	0.2
20-24	1	0.1 0.4	1		0.4	1	0.2	0.4
25-29	3	0.3 0.7	7 / 1	0.2	0.6	2	0.4	0.9
30-34	9	0.9 1.7	7 7	1.4	2.0	2	0.4	1.3
35-39	14	1.5 3.3	L 10	2.0	4.0	4	0.9	2.2
40 - 44	25	2.6 5.7	7 13	2.6	6.6	12	2.6	4.8
45-49	39	4.1 9.8	3 18	3.6	10.3	21	4.6	9.3
50-54	73	7.6 17.5	5 49	9.9	20.1	24	5.2	14.6
55-59	102	10.7 28.2	L 46	9.3	29.4	56	12.2	26.7
60-64	125	13.1 41.2	2 64	12.9	42.3	61	13.3	40.0
65-69	160	16.7 57.9	88	17.7	60.0	72	15.7	55.7
70-74	148	15.5 73.4	1 79	15.9	75.9	69	15.0	70.7
75-79	123	12.9 86.2	2 59	11.9	87.7	64	13.9	84.6
80-84	88	9.2 95.4	42	8.5	96.2	46	10.0	94.6
85+	44	4.6 100.0) 19	3.8	100.0	25	5.4	100.0
All ages	957	100.0	497	100.0		460	100.0	

Age distribution by 5-year age group and gender for period 1998-2012 (incl. DCO)

Table 4

Included in the statistics are 47.8% multiple primaries in males and 36.8% in females.

Males Females Males Females Males Females Prop.all Prop.all Age at DCO rate DCO rate cancers cancers Age- Agediagnosis Males Females n=0n=0 n=146755 n=142297 spec. spec. Years incid. incid. % n n % % % 0- 4 0.0 0.0 5-9 0.0 0.0 10 - 140.0 0.0 15-19 2 0.1 0.1 0.6 0.4 1 20-24 0.0 0.1 0.2 1 25-29 1 2 0.1 0.1 0.1 0.2 30-34 7 2 0.3 0.1 0.5 0.1 35-39 10 4 0.4 0.2 0.5 0.1 40 - 4413 0.5 0.5 0.4 0.2 11 45-49 0.8 1.0 0.4 0.3 18 21 50-54 49 24 0.2 2.7 1.3 0.6 55-59 46 2.7 0.4 56 3.1 0.3 60-64 3.5 0.4 64 61 3.9 0.3 65-69 4.5 0.4 88 72 6.0 0.3 70-74 78 0.4 6.7 5.0 69 0.3 75-79 59 64 7.8 5.9 0.4 0.3 80-84 42 46 9.2 5.3 0.3 0.3 6.1 3.1 0.2 85+ 19 25 0.2 459 0.0 0.0 0.3 0.3 All ages 496 Incidence Raw 1.8 1.6 WS 1.0 0.8 ES 1.5 1.1 BRD-S 1.8 1.4

Age-specific incidence, DCO rate and proportion of all cancers for period 1998-2012

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

Table 6a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2012 MALES

		Observed	Expected		LCL	UCL		DCO
Diagnos	is	n	n	SIR	95%	95%	EAR	00
C15	Oesophagus	2	0.3	6.8	0.8	24.4	15.9	
C17	Small intestine	4	0.1	48.7	13.3	124.7 ‡	\$ 36.7	
C18	Colon	9	1.6	5.7	2.6	10.8	69.4	
C32	Larynx	2	0.2	12.1	1.5	43.6	\$ 17.2	
C33-C34	Lung	2	1.9	1.0	0.1	3.8	0.8	
C43	Malign. melanoma	2	0.7	3.0	0.4	11.0	12.6	
C61	Prostate	9	4.7	1.9	0.9	3.7	40.4	
C64	Kidney	4	0.6	7.1	1.9	18.1 ‡	\$ 32.1	
C76-C79	CUP	3	0.3	10.7	2.2	31.2 ‡	\$ 25.4	
C82-C85	NHL	4	0.6	6.3	1.7	16.2 ‡	\$ 31.5	
C90	Mult. myeloma	2	0.2	9.9	1.2	35.7 ‡	\$ 16.8	
Other p	rimaries	б	2.4	2.6	0.9	5.6	34.1	
Not obs	erved	0	2.9	0.0	0.0	1.3	-27.0	
All mul	t. primaries	49	16.3	3.0	2.2	4.0 ‡	\$ 306.0	

Patients	308	
Mean age at second malignancy (years)	70.2	
Person-years	1069	
Mean observation time (years)	3.5	
Median observation time (years)	3.0	

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries".

Table 6b

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2012 FEMALES

	Observed Ex	pected		LCL	UCL		DCO
Diagnosis	n	n	SIR	95%	95%	EAR	00
Cl6 Stomach	2	0.4	4.5	0.5	16.1	14.5	
C18 Colon	3	1.3	2.4	0.5	7.0	16.4	
C19-C20 Rectum	2	0.6	3.6	0.4	13.1	13.6	
C25 Pancreas	4	0.6	7.1	1.9	18.1 #	32.2	
C50 Breast	5	3.8	1.3	0.4	3.1	11.5	
C54 Corpus uteri	2	0.7	2.8	0.3	10.2	12.1	
C64 Kidney	3	0.3	9.3	1.9	27.2 #	25.1	
C70-C72 CNS cancer	2	0.2	11.2	1.4	40.4 #	17.1	
C82-C85 NHL	2	0.5	4.1	0.5	14.9	14.2	
Other primaries	8	2.5	3.1	1.4	6.2 #	51.1	12.5
Not observed	0	1.8	0.0	0.0	2.1	-16.8	
All mult. primaries	33	12.6	2.6	1.8	3.7 #	191.0	3.0
-							

Patients	289
Mean age at second malignancy (years)	71.5
Person-years	1067
Mean observation time (years)	3.7
Median observation time (years)	3.3

The occurrence of second malignancy is statistically significant.

Observed second malignancies with count 1 are pooled in category "Other primaries".





Figure 7. Age distribution and age-specific incidence






GIST: Gastrointestinal stromal tumor

Figure 8. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.






Average incidence (world standard population) 2003 - 2008: Males

Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 1.0/100,000 WS N=199, females 0.9/100,000 WS N=211). Since cancer data are not available in some counties until 2007, the local incidence rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 6 women were identified with newly diagnosed gastroint. stromal tumor. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.7/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.2 and 2.2/100,000.






Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=199, females N=211). Since cancer data are not available in some counties until 2007, the local SIR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 6 women were identified with newly diagnosed gastroint. stromal tumor. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.96. Though, the value of this parameter may vary with an underlying probability of 99% between 0.25 and 2.50, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts, and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

		Prop.				Prop. deaths
	Incident	actively	Prop.		Prop.	with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	%	010	n	00	90
1998	4	100.0		4	100.0	100.0
1999	8	100.0		6	75.0	66.7
2000	18	94.4		12	66.7	100.0
2001	28	100.0		17	60.7	82.4
2002	48	97.9		28	58.3	96.4
2003	50	96.0		18	36.0	100.0
2004	73	97.3		34	46.6	94.1
2005	67	92.5		25	37.3	96.0
2006	62	88.7		12	19.4	100.0
2007	91	75.8		24	26.4	95.8
2008	92	55.4		21	22.8	95.2
2009	99	53.5		20	20.2	100.0
2010	103	53.4		18	17.5	100.0
2011	100	57.0		10	10.0	90.0
2012	114	95.6		11	9.6	90.9
1998-2012	957	76.7		260	27.2	95.0

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

			Prop.		
			deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	n	90	n	8
1998	4	1	100.0	1	25.0
1999	8	/ 1	100.0		
2000	18	3	66.7	2	11.1
2001	28	5	100.0	2	7.1
2002	48	12	100.0	6	12.5
2003	50	13	92.3	2	4.0
2004	73	23	82.6	10	13.7
2005	67	21	95.2	5	7.5
2006	62	18	94.4	2	3.2
2007	91	22	95.5	5	5.5
2008	92	28	100.0	4	4.3
2009	99	23	100.0	6	6.1
2010	103	32	93.8	6	5.8
2011	100	27	96.3	6	6.0
2012	114	35	100.0	9	7.9
1998-2012	957	264	95.5	66	6.9

Table 10c

Annual cohorts of deaths, proportion of cancer-related and not cancerrelated deaths, and cancer recorded on death certificates (incl. DCO) (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.52 m as of 2007, respectively)

				Prop.	
				cancer	
		Prop.	Prop.	recorded	
		cancer-	not cancer-	on death	
Year of	Deaths	related	related	certificate	
death	n	00	00	00	
1998	1/		100.0	100.0	
1999	1	100.0		100.0	
2000	3	33.3	66.7	100.0	
2001	5	100.0		100.0	
2002	12	58.3	41.7	66.7	
2003	13	76.9	23.1	91.7	
2004	23	65.2	34.8	89.5	
2005	21	81.0	19.0	80.0	
2006	18	61.1	38.9	70.6	
2007	22	77.3	22.7	81.0	
2008	28	85.7	14.3	85.7	
2009	23	65.2	34.8	73.9	
2010	32	62.5	37.5	76.7	
2011	27	66.7	33.3	73.1	
2012	35	65.7	34.3	77.1	
1998-2012	264	69.7	30.3	79.4	

Munich Cancer Registry

Year of death	Deaths n	Age at death (all causes) Years	Age at death (cancer- related) Years	Age at death (not cancer- related) Years	Age at death (according to death certificate) Years
1998	1	87.6		87.6	87.6
1999					
2000	2	84.4	79.2	89.5	79.2
2001	3	71.1	71.1		71.1
2002	6	79.3	80.3	78.2	80.3
2003	9	69.5	73.2	62.2	71.7
2004	11	71.0	68.9	80.5	71.0
2005	9	71.1	70.6	72.8	72.9
2006	9	75.2	73.5	77.2	73.5
2007	13	67.6	66.2	75.1	66.2
2008	9	75.2	72.1	85.8	72.1
2009	11	75.2	73.0	81.2	73.0
2010	21	73.7	71.9	76.6	72.3
2011	13	77.6	78.0	77.0	77.1
2012	14	75.5	70.6	84.3	70.6
1998-2012	131	73.8	71.8	78.0	72.2

Table 11a

Means of age at death according to the grouping in Table 10 $$\rm MALES$$

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Year of	Deaths	Age at death (all causes)	Age at death (cancer- related)	Age at death (not cancer- related)	Age at death (according to death certificate)
death		/ •			
ueatli	n	Years	Years	Years	Years
1998					
1999	1	77.8	77.8		77.8
2000	1	78.5		78.5	78.5
2001	2	60.2	60.2		60.2
2002	б	71.8	69.5	76.3	72.0
2003	4	64.5	64.5		64.5
2004	12	78.1	75.6	80.7	78.3
2005	12	69.4	69.6	68.3	69.6
2006	9	78.1	78.5	77.3	78.0
2007	9	76.5	71.4	86.5	72.6
2008	19	75.0	74.1	81.9	74.1
2009	12	79.7	74.4	87.1	76.0
2010	11	74.3	70.8	80.6	71.9
2011	14	71.6	69.9	77.7	69.4
2012	21	76.1	72.3	83.8	74.4
1998-2012	133	74.6	72.0	81.2	72.9

Table 11b

Means of age at death according to the grouping in Table 10 FEMALES

By 2010, life expectancy for a newborn male in Germany is 77.5 years compared with 82.6 years for his female counterpart.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 12a

Mortality measures (cancer-related death) and mortality-incidence-index by year of death MALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1999									
2000	1	0.1	0.11	0.0	0.09	0.1	0.12	0.1	0.18
2001	3	0.3	0.18	0.2	0.19	0.2	0.18	0.3	0.16
2002	3	0.2	0.13	0.1	0.08	0.1	0.11	0.2	0.17
2003	6	0.3	0.26	0.2	0.19	0.2	0.24	0.4	0.32
2004	9	0.5	0.25	0.2	0.23	0.3	0.23	0.5	0.23
2005	7	0.4	0.22	0.2	0.19	0.3	0.23	0.4	0.27
2006	5	0.3	0.19	0.1	0.15	0.2	0.18	0.3	0.20
2007	11	0.5	0.22	0.3	0.20	0.4	0.21	0.5	0.21
2008	7	0.3	0.16	0.2	0.14	0.2	0.15	0.3	0.16
2009	8	0.4	0.16	0.2	0.13	0.2	0.14	0.4	0.16
2010	13	0.6	0.22	0.3	0.19	0.4	0.19	0.5	0.21
2011	7	0.3	0.12	0.1	0.09	0.2	0.10	0.3	0.12
2012	9	0.4	0.15	0.2	0.15	0.3	0.16	0.4	0.17
1999-2012	89	0.3	0.19	0.2	0.16	0.3	0.17	0.4	0.20

Table 12b

Mortality measures (cancer-related death) and mortality-incidence-index by year of death FEMALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1999	1	0.1	0.33	0.0	0.13	0.0	0.21	0.1	0.35
2000									
2001	2	0.2	0.18	0.1	0.18	0.1	0.18	0.1	0.17
2002	4	0.2	0.17	0.1	0.12	0.1	0.14	0.2	0.16
2003	4	0.2	0.15	0.1	0.16	0.2	0.16	0.2	0.14
2004	б	0.3	0.16	0.1	0.09	0.2	0.11	0.2	0.14
2005	10	0.5	0.29	0.2	0.30	0.3	0.31	0.4	0.31
2006	б	0.3	0.17	0.1	0.11	0.2	0.12	0.2	0.15
2007	б	0.3	0.14	0.1	0.13	0.2	0.13	0.2	0.13
2008	17	0.7	0.36	0.3	0.29	0.4	0.32	0.5	0.33
2009	7	0.3	0.14	0.1	0.12	0.2	0.13	0.3	0.14
2010	7	0.3	0.16	0.1	0.13	0.2	0.13	0.3	0.15
2011	11	0.5	0.28	0.2	0.26	0.3	0.25	0.4	0.27
2012	14	0.6	0.25	0.2	0.24	0.4	0.24	0.5	0.26
1999-2012	95	0.3	0.22	0.1	0.18	0.2	0.19	0.3	0.20

Age at									
death	Cases			Males			Females		
Years	n	010	Cum.%	n	010	Cum.%	n	00	Cum.%
25-29	1	0.5	0.5			0.0	1	1.1	1.1
30-34	0	0.0	0.5			0.0			1.1
35-39	1	0.5	1.1	/ 1	1.1	1.1			1.1
40 - 44	2	1.1	2.2			1.1	2	2.1	3.2
45-49	5	2.7	4.9	2	2.2	3.3	3	3.2	6.3
50-54	6	3.2	8.1	3	3.3	6.7	3	3.2	9.5
55-59	12	6.5	14.6	6	6.7	13.3	6	6.3	15.8
60-64	19	10.3	24.9	9	10.0	23.3	10	10.5	26.3
65-69	27	14.6	39.5	16	17.8	41.1	11	11.6	37.9
70-74	31	16.8	56.2	17	18.9	60.0	14	14.7	52.6
75-79	28	15.1	71.4	11	12.2	72.2	17	17.9	70.5
80-84	33	17.8	89.2	18	20.0	92.2	15	15.8	86.3
85+	20	10.8	100.0	7	7.8	100.0	13	13.7	100.0
All ages	185	100.0		90	100.0		95	100.0	

Age distribution of age at death (cancer-related) for period 1999-2012 (incl. multiple primaries)

Included in the statistics are 47.8% multiple primaries in males and 36.8% in females.

Age-spe	ecific mortal		riod 199	9-2012	ortion o	t all can	cers
		Males		Females		Males	Females
Age at		Age-		Age-		Prop.all	Prop.all
death	Males Fema			spec.		cancers	cancers
Years	n n	mortal.	MI-inde	x mortal.	MI-index	olo	olo
0- 4		0.0		0.0			
5-9		0.0		0.0			
10-14		0.0		0.0			
15-19		0.0		0.0			
20-24		0.0		0.0			
25-29		1 0.0		0.1	0.52		0.9
30-34		0.0		0.0			
35-39	1	0.0	0.10	0.0		0.3	
40 - 44		2 0.0		0.1	0.17		0.2
45-49	2	3 0.1	0.12	0.1	0.15	0.1	0.2
50-54	3	3 0.2	0.06	0.2	0.13	0.1	0.1
55-59	б	6 0.4	0.14	0.4	0.11	0.1	0.1
60-64	9 1	0 0.6	0.15	0.6	0.17	0.1	0.2
65-69	16 1	1 1.1	0.19	0.7	0.16	0.1	0.1
70-74	17 1	4 1.5	0.22	1.1	0.21	0.1	0.2
75-79	11 1	7 1.5	0.19	1.6	0.28	0.1	0.2
80-84	18 1	5 4.1	0.44	1.8	0.34	0.2	0.1
85+	7 1	.3 2.3	0.38	1.7	0.54	0.1	0.1
All ages	90 9	5				0.1	0.1
Mortality							
Raw		0.3	0.19	0.3	0.22		
WS		0.2	0.16	0.1	0.18		
ES		0.3	0.18	0.2	0.19		
BRD-S		0.4	0.20	0.3	0.20		
PYLL-70							
per 100,00	00	1.3		1.7			
ES		1.1		1.4			
AYLL-70		8.4		11.0			

Age-specific mortality (cancer-related) and proportion of all cancers

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15a

Multiple primaries in deaths in period 1999-2012 $$\rm MALES$$

					Syn- chron	Syn- chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosia	/	/		+16 €		⊥30a ←%		POSL ~%
Diagnosis	n	° ↓	n	6→	n	₽	n	\$→
	-	1 0	1	100 0				
C03-C06 Oral cavity	1	1.6	T	100.0			0	22.2
Cl6 Stomach	6	9.5			4	66.7	2	33.3
C17 Small intestine	5	7.9			3	60.0	2	40.0
C18 Colon	8	12.7			5	62.5	3	37.5
C19-C20 Rectum	5	7.9			2	40.0	3	60.0
C21 Anus/canal	/ 1 /	1.6					1	100.0
C22 Liver	1 -	1.6			1	100.0		
C25 Pancreas	5	7.9	1	20.0	4	80.0		
C32 Larynx	2	3.2	2	100.0				
C33-C34 Lung	4	6.3	2	50.0	2	50.0		
C38,C45 Mesothelioma	1	1.6					1	100.0
C43 Malign. melanoma	2	3.2	2	100.0				
C44 Skin others	1	1.6					1	100.0
C61 Prostate	8	12.7	7	87.5			1	12.5
C64 Kidney	3	4.8	1	33.3	2	66.7		
C67 Bladder	1	1.6	1	100.0				
C76-C79 CUP	1	1.6			1	100.0		
C82-C85 NHL	3	4.8	1	33.3			2	66.7
C90 Mult. myeloma	1	1.6					1	100.0
C91-C96 Leukaemia	4	6.3	2	50.0			2	50.0
			_					
All mult. primaries	63	100.0	20	31.7	24	38.1	19	30.2

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 15b

Multiple primaries in deaths in period 1999-2012 FEMALES

					Syn-	Syn-		
					chron	chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n	% ↓	n	60	n	se →	n	¢
C16 Stomach	7	13.5			6	85.7	1	14.3
C17 Small intestine	4	7.7			3	75.0	1	25.0
C18 Colon	3	5.8			2	66.7	1	33.3
C22 Liver	1	1.9			1	100.0		
C25 Pancreas	7	13.5			1	14.3	6	85.7
C33-C34 Lung	4	7.7	2	50.0	1	25.0	1	25.0
C43 Malign. melanoma	/ 1 4	1.9					1	100.0
C46,C49 Soft tissue	2	3.8			1	50.0	1	50.0
C48 Peritoneal	1	1.9			1	100.0		
C50 Breast	11	21.2	8	72.7			3	27.3
C53 Cervix uteri	2	3.8	2	100.0				
C56 Ovary	5	9.6			3	60.0	2	40.0
C67 Bladder	1	1.9	1	100.0				
C70-C72 CNS cancer	2	3.8	2	100.0				
C74-C80 Cancer others	1	1.9					1	100.0
All mult. primaries	52	100.0	15	28.8	19	36.5	18	34.6
- / /								

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Age-specific mortality (cancer-related) and proportion of all cancers for period 2000-2012 (Singular primaries only *)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	MI-index	Females Age- spec. mortal.	MI-index	cancers	Females Prop.all cancers %
0- 4			0.0		0.0			
5-9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29			0.0		0.0			
30-34 35-39	1		0.0	0 10	0.0		0.3	
40-44	1	2	0.0	0.12	0.0	0.24	0.3	0.2
45-49		3	0.0		0.1	0.24		0.2
50-54	2	2	0.0	0.05	0.2	0.17	0.1	0.2
55-59	4	5	0.1		0.1	0.12	0.1	0.1
60-64	6	6	0.4		0.4	0.12	0.1	0.1
65-69	11	6	0.8		0.4	0.12	0.1	0.1
70-74	8	7	0.7	0.21	0.5	0.16	0.1	0.1
75-79	5	12	0.7		1.2	0.29	0.1	0.2
80-84	10	10	2.3		1.2	0.34	0.1	0.1
85+	3	9	1.0	0.36	1.2	0.49	0.0	0.1
All ages	50	62					0.1	0.1
Mortality								
Raw			0.2		0.2	0.19		
WS			0.1	0.13	0.1	0.16		
ES			0.1		0.1	0.17		
BRD-S			0.2	0.16	0.2	0.18		
PYLL-70								
per 100,000			0.8		1.2			
ES			0.7		1.1			
AYLL-70			7.9		11.7			

* See corresponding tables with multiple primaries.



Age-specific mortality (cancer-related) and proportion of all cancers for period 2000-2012 (Single primaries only *)

Age at death Years	Males n	Females n	- /	MI-index	Females Age- spec. mortal.	MI-index	cancers	Females Prop.all cancers %
0- 4			0.0		0.0			
5-9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29			0.0		0.0			
30-34			0.0		0.0			
35-39	1		0.0	0.12	0.0		0.3	
40 - 44		1	0.0		0.0	0.13		0.1
45-49		2	0.0		0.1	0.11		0.1
50-54	2	2	0.1	0.05	0.1	0.11	0.1	0.1
55-59	4	5	0.3		0.3	0.13	0.1	0.1
60-64	б	4	0.4	0.16	0.3	0.10	0.1	0.1
65-69	9	6	0.7		0.4	0.14	0.1	0.1
70-74	7	6	0.6	0.22	0.5	0.15	0.1	0.1
75-79	5	11	0.7		1.1	0.27	0.1	0.2
80-84	7	8	1.6	0.33	1.0	0.28	0.1	0.1
85+	2	9	0.7	0.27	1.2	0.49	0.0	0.1
All ages	43	54					0.1	0.1
Mortality								
Raw			0.2	0.15	0.2	0.18		
WS			0.1	0.13	0.1	0.14		
ES			0.1		0.1	0.15		
BRD-S			0.2	0.15	0.2	0.17		
PYLL-70								
per 100,000			0.8		1.0			
ES NULL RO			0.7		0.8			
AYLL-70			8.4		10.8			

* See corresponding tables with multiple primaries.





Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at gastroint. stromal tumor-related death (see Table 10) should be considered.







Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2003 to 2008. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.2/100,000 WS N=45, females 0.2/100,000 WS N=46). Since cancer data are not available in some counties until 2007, the local mortality rates were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 0 women died from gastroint. stromal tumor. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.0/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.4/100,000.






Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2003 to 2008. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=45, females N=46). Since cancer data are not available in some counties until 2007, the local SMR values were not calculated, and the map tiles show as shaded.

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,131 female residents (averaged) in the period from 2003 to 2008 a total of 0 women died from gastroint. stromal tumor. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.00. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 3.99, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the tumor-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

AYLL-70 BRD-S DCO	Average years of life lost prior to age 70 given a person dies before that age German standard population Death certificate only
EAR	Excess absolute risk
	= excess cancer cases (O - E) per 10,000 person-years
ES	European standard population (old)
FRG	Federal Republic of Germany
GEKID	Association of Population-based Cancer Registries in Germany
	(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)
LCL	Lower confidence limit
MI-index	Ratio between mortality and incidence
MCR	Munich Cancer Registry (Tumorregister München)
PYLL-70	Potential years of life lost prior to age 70 given a person dies before that age
SEER	Surveillance, Epidemiology, and End Results (USA)
SIR	Standardized incidence ratio
SMR	Standardized mortality ratio
UCL	Upper confidence limit
WS	World standard population

Recommended Citation

Munich Cancer Registry. Baseline statistics GIST: Gastroint. stromal tumor [Internet]. 2014 [updated 2014 Mar 20; cited 2014 May 1]. Available from: http://www.tumorregistermuenchen.de/en/facts/base/base_hGISTE.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Munich Cancer Registry

Index of figures and tables

Fig./Tb	l.	Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	3
1a	Gender distribution by year of diagnosis	4
2	Incidence by year of diagnosis	5
3	Age distribution parameters by year of diagnosis	6
4	Age distribution by 5-year age group and gender	8
5	Age-specific incidence and DCO rate	9
6	Standardized incidence ratio of second primaries	10
7	Age distribution and age-specific incidence (chart)	12
8	Cumulative follow-up years (chart)	13
9a	Map of cancer incidence (WS) by county (chart)	14
9b	Standardized incidence ratio (SIR) by county (chart)	15
10a	Pts incident cohorts and mortality / yr	16
10b	Incidence and mortality by year of diagnosis	17
10c	Cancer-related deaths, death certification available / yr	18
11	Means of age at death / yr	19
12	Mortality by year of death	21
13	Distribution of age at death	22
14	Age-specific mortality	23
15	Multiple primaries in deaths	24
16	Age-specific mortality (first primaries)	26
17	Age-specific mortality (single primaries)	27
18	Age distribution and age-specific mortality (chart)	28
19a	Map of cancer mortality (WS) by county (chart)	29
19b	Standardized mortality ratio (SMR) by county (chart)	30