Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ Deutsch

Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

Cancer statistics: Baseline statistics

C01: Base of tongue cancer

Year of diagnosis	1998-2013
Patients	645
Diseases	646
Creation date	05/19/2015
Export date	12/30/2014
Population	4.64 m



http://www.tumorregister-muenchen.de/en/facts/base/base_C01__E.pdf

Global Statements about the statistics on the Internet – Baseline Statistics (grey button ——), Survival (red button ——)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.64 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases**** are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR. The time-delayed acquisition of data and the occasionally high DCO-rates indicate optimizing reserves, among others, because of current financial and legal conditions that hinder the analyses.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, May 2015

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007). Death certificates from 2014 are incorporated into these analyses.
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate. A high proportion of DCO cases (≥5%) in particular cancer types indicate insufficient participation of specific cancer specializations.

Some remarks regarding this cancer type

As a general rule, these few results from the TRM form the basis of sophisticated analyses. For head and neck tumors this is not the case. Therefore the results for head and neck tumors should be interpreted with caution. In part this is due to problems of classification because of limited specific details of locality. Additionally, with advanced tumors in a close topographic location it is often not possible to determine the exact ICD localization of a tumor.

Topography codes (ICD-O-3 2000) used for specifying cancer site

Code	Description	
C01.9	Base of tongue, NOS	

INCIDENCE

Table 1

Patient cohorts by year of diagnosis including DCO cases and multiple primaries, and with proportion of deaths and active follow-up

				Prop.		Prop.
		DCO	Prop.	mult.	Prop.	actively
Year of	Cases	cases	DCO	primaries	deaths	followed
diagnosis	n	n	%	8	%	%
1998	14			21.4	78.6	100.0
1999	22			36.4	90.9	100.0
2000	13	1	7.7	38.5	76.9	100.0
2001	19	2	10.5	31.6	84.2	100.0
2002	33			27.3	72.7	97.0 #
2003	42	5	11.9	57.1	85.7	100.0
2004	49	4	8.2	22.4	73.5	93.9
2005	53	3	5.7	35.8	58.5	94.3
2006	51	/ 1	2.0	29.4	66.7	98.0
2007	53	1	1.9	32.1	64.2	86.8 # ##
2008	52	4	7.7	23.1	63.5	82.7
2009	46			32.6	50.0	93.5
2010	56	1	1.8	26.8	53.6	78.6
2011	65	2	3.1	30.8	40.0	75.4
2012	55	1	1.8	38.2	27.3	78.2
2013	23	1	4.3	21.7	30.4	100.0 ###
1998-2013	646	26	4.0	31.7	59.8	89.6

[#] The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

^{##} Since 2007 the percentage of actively followed patients sharply declined compared to the previous years. This is a consequence of ambiguous data protection rules that currently forbid cancer registries in Bavaria to obtain the essential life status informations from competent registration offices.

^{###} Please be aware that data of recent annual patient cohorts may not yet be fully processed. Therefore, the presented figures and tables are potentially related to different time periods as pointed out in the respective headlines or legends.

Table 1a

Patient cohorts by year of diagnosis and gender including DCO cases

Year of	All	Males	Females	Prop. males
diagnosis	n /	'n	n	%
1998	14	10	4	71.4
1999	22	18	4	81.8
2000	/13	9	4	69.2
2001	19 /	12	7/	63.2
2002	33	26	7	78.8
2003	42	37	5 /	88.1
2004	49	33	16	67.3
2005	53	40	13	75.5
2006	51	38	13	74.5
2007	53	43	10	81.1
2008	52	42	10	80.8
2009	46	31	15	67.4
2010	56	40	16	71.4
2011	65	50	15	76.9
2012	55	43	12	78.2
2013	23	20	3	87.0
1998-2013	646	492	154	76.2

Table 2

Incidence measures by year of diagnosis and gender including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

			Males	Fem.	Males	Fem.	Males	Fem.	Males	Fem.
Year of	Males	Females	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	10	4	0.9/	0.3	0.6	0.2	0.8	0.2	0.9	0.3
1999	18	4	1.6	0.3	1.1	0.2	1.4	0.3	1.5	0.3
2000	9	4	0.8	0.3	0.5	0.2	0.7	0.3	0.9	0.3
2001	12	7 /	1.0	0.6	0.7	0.3	1.0	0.4	1.1	0.5
2002	26	7 🛴	1.4	0.4	0.9	0.2	1.3	0.3	1.4	0.3
2003	37	5	2.0	0.3	1.2	0.1	1.7	0.2	2.0	0.2
2004	33	16	1.8	0.8	1.0	0.4	1.4	0.6	1.6	0.7
2005	40	13	2.1	0.7	1.4	0.3	1.9	0.5	2.1	0.6
2006	38	13	2.0	0.6	1.2	0.4	1.7	0.5	1.9	0.6
2007	43	10	1.9	0.4	1.3	0.3	1.7	0.4	1.8	0.4
2008	42	10	1.9	0.4	1.2	0.2	1.6	0.3	1.9	0.4
2009	31	15	1.4	0.6	0.8	0.3	1.2	0.5	1.3	0.5
2010	40	16	1.8	0.7	1.0	0.3	1.5	0.5	1.7	0.6
2011	50	1,5	2.2	0.6	1.3	0.3	1.8	0.5	2.0	0.5
2012	43	12	1.9	0.5	1.1	0.3	1.5	0.4	1.7	0.4
2013	20	3	0.9	0.1	0.5	0.1	0.7	0.1	0.8	0.1
1998-2013	492	154	1.7	0.5	1.0	0.3	1.4	0.4	1.6	0.4




Table 3

Age distribution parameters by year of diagnosis (All) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	14	57.7	13.2	31.1	78.2	44.9	50.3	58.0	67.5	77.8
1999	22	58.6	9.0	40.4	74.9	48.5	52.2	59.7	64.3	70.5
2000	13	60.3	14.0	35.9	78.3	42.5	47.6	65.5	68.7	78.2
2001	19	63.2	13.6	48.6	92.5	49.0	50.6	61.3	71.2	85.1
2002	33	60.5	9.0	43.8	81.5	50.3	54.8	58.2	65.1	72.1
2003	42	63.1	9.1	45.1	83.3	52.1	56.0	62.7	69.4	76.9
2004	49	63.6	10.0	38.6	93.3	52.0	58.2	62.5	69.5	75.4
2005	53	61.3	13.4	4.1	87.2	50.1	54.8	61.1	65.9	78.5
2006	51	60.5	12.0	19.0	84.8	46.0	52.6	61.3	68.0	76.2
2007	53	57.6	9.7	35.2	76.9	45.0	52.3	57.9	63.3	69.5
2008	52	63.9	10.3	38.3	87.7	50.1	57.6	63.2	71.1	77.4
2009	46	63.5	10.9	36.7	87.3	48.0	57.0	62.9	72.6	76.2
2010	56	64.5	11.3	38.0	92.1	50.0	57.2	65.6	72.7	77.9
2011	65	64.3	11.1	40.0	93.8	49.9	56.5	64.0	71.8	77.5
2012	55	62.5	10.9	39.8	87.9	49.3	53.6	62.0	70.6	76.3
2013	23	62.1	9.9	48.9	85.2	50.6	54.7	59.7	67.9	75.2
1998-2013	646	62.1	11.1	4.1	93.8	48.8	54.9	61.6	69.5	76.5

Table 3a

Age distribution parameters by year of diagnosis (MALES)

(incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	10	57.3	10.8	44.9	78.2	46.1	50.3	54.9	60.4	75.2
1999	18	56.9	8.8	40.4	72.5	43.1	50.5	58.4	63.4	70.5
2000	9	59.7	15.8	35.9	78.3	35.9	47.6	59.2	73.3	78.3
2001	12	61.1	12.1	48.6	85.1	49.0	50.4	59.2	68.2	78.7
2002	26	59.7	8.4	43.8	80.2	48.0	54.6	58.2	64.2	70.8
2003	37	62.3	8.9	45.1	83.3	50.6	56.0	62.4	66.7	76.9
2004	33	62.9	8.8	38.6	80.3	54.8	58.9	62.5	68.3	72.5
2005	40	59.5	14.1	4.1	87.1	46.3	52.9	59.8	65.6	78.2
2006	38	61.3	10.5	38.7	84.8	46.7	54.4	60.7	66.8	77.4
2007	43	58.4	9.6	37.1	76.9	45.6	54.7	58.0	64.5	69.5
2008	42	63.7	9.6	38.3	85.9	52.5	58.7	62.7	71.1	76.3
2009	31	62.2	11.0	36.7	81.1	48.0	56.1	61.9	72.1	76.2
2010	40	63.7	12.2	38.0	92.1	49.7	55.3	63.9	71.7	80.0
2011	50	63.8	10.7	40.0	86.5	49.8	55.0	64.4	71.0	77.0
2012	43	62.7	11.0	39.8	87.9	49.3	53.6	62.0	70.6	76.3
2013	20	61.1	9.0	48.9	76.7	50.1	53.0	59.5	66.9	74.9
1998-2013	492	61.5	10.7	4.1	92.1	48.8	54.7	61.2	68.7	76.2

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of	Cases	S	td.				Median		
diagnosis	n	Mean d	lev. Min.	Max.	10%	25%	50%	75%	90%
_									
1998	4	58.8 2	0.0 31.1	77.8	31.1	45.0	63.2	72.6	77.8
1999	4	66.1	6.6 59.0	74.9	59.0	61.6	65.3	70.6	74.9
2000	4	61.8 1	0.6 46.0	68.2	46.0	55.7	66.5	67.9	68.2
2001	7	66.8 1	6.2 50.2	92.5	50.2	50.6	63.0	83.0	92.5
2002	7	63.3 1	1.4 / 50.3	81.5	50.3	55.2	58.3	73.9	81.5
2003	5	68.8	9.6 53.1	77.3	53,1	68.5	69.4	75.7	77.3
2004	16	65.3 1	2.2 48.0	93.3	50.3	56.4	62.1	74.7	81.4
2005	13	66.9	9.3 57.1	87.2	57.2	60.9	64.4	72.9	79.3
2006	13	58.1 1	5.9 19.0	76.4	45.9	48.1	65.4	70.1	71.3
2007	10	53.9 1	0.1 35.2	70.5	40.1	47.5	55.1	61.3	66.3
2008	10	64.9 1	3.5 45.6	87.7	47.6	51.5	65.3	76.3	82.6
2009	15	66.2 1	0.6 45.8	87.3	54.2	57.8	66.8	73.7	75.6
2010	16	66.5	8.7 47.1	77.9	53.4	60.6	67.8	73.2	77.8
2011	15	65.9 1	2.5 48.6	93.8	54.5	56.5	60.8	72.9	84.0
2012	12 /	61.8 1	1.2 45.5	83.5	50.0	53.3	60.7	69.8	74.0
2013	3	69.4 1	5.1 55.0	85.2	55.0	55.0	67.9	85.2	85.2
1998-2013	154	64.0 1	2.1 19.0	93.8	49.5	56.5	64.4	72.0	77.8

Table 4

Age distribution by 5-year age group and gender for period 1998-2013 (incl. DCO)

Age at									
diagnosis	Cases			Males			Females		
Years	n	%	Cum.%	'n	%	Cum.%	n	%	Cum.%
				/ .					
0 - 4	1	0.2	0.2	/ 1	0.2	0.2			0.0
5-9	0	0.0	0.2			0.2			0.0
10-14	0	0.0	0.2			0.2			0.0
15-19	1	0.2	0.3			0.2	1	0.6	0.6
20-24	0	0.0	0.3			0.2			0.6
25-29	0	0.0	0.3			0.2			0.6
30-34	1	0.2	0.5			0.2	1	0.6	1.3
35-39	12	1.9	2.3	11	2.2	2.4	1	0.6	1.9
40-44	16	2.5	4.8	16	3.3	5.7			1.9
45-49	49	7.6	12.4	33	6.7	12.4	16	10.4	12.3
50-54	82	12.7	25.1	69	14.0	26.4	13	8.4	20.8
55-59	116	18.0	43.0	89	18.1	44.5	27	17.5	38.3
60-64	116	18.0	61.0	97	19.7	64.2	19	12.3	50.6
65-69	95	14.7	75.7	68	13.8	78.0	27	17.5	68.2
70-74	71	11.0	86.7	50	10.2	88.2	21	13.6	81.8
75-79	55	8.5	95.2	39	7.9	96.1	16	10.4	92.2
80-84	17	2.6	97.8	12	2.4	98.6	5	3.2	95.5
85+	14	2.2	100.0	7	1.4	100.0	7	4.5	100.0
All ages	646	100.0		492	100.0		154	100.0	

Included in the statistics are 43.0% multiple primaries in males and 36.4% in females.

Table 5

Age-specific incidence, DCO rate and proportion of all cancers for period 1998-2013

							Males	Females
			Males	Females	Males	Females	Prop.all	Prop.all
Age at			Age-	Age-	DCO rate	DCO rate	cancers	cancers
diagnosis	Males	Females		spec.	n=19	n=7	n=158258	n=153136
Years	n	n	incid.		8	%	%	%
0- 4	1		0.1	0.0			0.3	
5- 9			0.0	0.0				
10-14			0.0	0.0				
15-19		1 4	0.0	0.1				0.3
20-24			0.0	0.0				
25-29			0.0	0.0				
30-34		1	0.0	0.0				0.0
35-39	11	1	0.4	0.0			0.5	0.0
40-44	16		0.6	0.0			0.5	
45-49	33	16	1.4	0.7			0.6	0.2
50-54	69	13	3.4	0.6	1.4		0.8	0.1
55-59	89	27	4.9	1.4	3.4	_3.7	0.6	0.2
60-64	97	19	5.5	1.0	2.1		0.4	0.1
65-69	67	27	4.2	1.6	6.0		0.2	0.1
70-74	50	21	3.9	1.4	4.0	9.5	0.2	0.1
75-79	39	16	4.7	1.3	15.4	12.5	0.2	0.1
80-84	12	5	2.4	0.5			0.1	0.0
85+	7	7	2.1	0.8	14.3	28.6	0.1	0.0
All ages	491	154			3.9	4.5	0.3	0.1
Incidence								
Raw			1.7	0.5				
WS			1.0	0.3				
ES			1.4	0.4				
BRD-S			1.6	0.4				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

Table 6a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2013

MALES

	Observed	Expected		LCL	UCL		DCO
Diagnosis	n /	n	SIR	95%	95%	EAR	%
C03-C06 Oral cavity	5/	0.2	32.9	10.7	76.7 #	49.3	
C09-C10 Oropharynx	3	0.2	15.0	3.1	43.9 #	28.5	
C12-C13 Hypopharynx	/3	0.1	27.5	5.7	80.4 #	29.4	
C15 Oesophagus	/ 5	0.3	18.4	6.0	43.0 #	48.1	20.0
C18 Colon	2	1.2	1./7	0.2	6.1	8.3	
C19-C20 Rectum	2	0.8	2.6	0.3	9.5	12.6	
C22 Liver	2	0.4	5.5	0.7	19.8	16.6	50.0
C32 Larynx	3	0.2	18.6	3.8	54.4 #	28.9	66.7
C33-C34 Lung	12	1.6	7.6	3.9	13.3 #	106.0	8.3
C61 Prostate	5	3.7	1.3	0.4	3.1	12.8	
C64 Kidney	3	0.5	6.2	1.3	18.2 #	25.6	
C91-C96 Leukaemia	3	0.2	15.9	3.3	46.4 #	28.6	33.3
Other primaries	6	1.6	3.7	1.4	8.1 #	44.6	16.7
Not observed	0	2.4	0.0	< 0.0	1.6	-24.2	
All mult. primaries	54	13.2	4.1	3.1	5.3 #	415.1	13.0

Patients	338
Median age at second malignancy (years)	63.8
Person-years	984
Mean observation time (years)	2.9
Median observation time (years)	1.7

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries".

Table 6b

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2013

FEMALES

		Observed	Expected		LCL	UCL		DCO
Diagnos	is	n	n	SIR	95%	95%	EAR	%
C15	Oesophagus	2	0.0	76.5	9.3	276.5 #	47.5	
C19-C20	Rectum	2	0.2	10.6	1.3	38.4 #	43.6	
C33-C34	Lung	/5	0.3	15.4	5.0	35.9 #	112.5	40.0
C50	Breast	/ 3	1.4	2.1	0.4	6.3	38.6	33.3
C53	Cervix uteri	2	0.1	33.1	4.0	119.5 #	46.7	
Other pr	rimaries	4	0.2	18.3	5.0	46.9 #	91.0	
Not obse		0	2.3	0.0	0.0	1.6	-54.6	
All mult	t. primaries	18	4.5	4.0	2.4	6.3 #	325.4	16.7

Patients	108
Median age at second malignancy (years)	69.2
Person-years	415
Mean observation time (years)	3.8
Median observation time (years)	2.7

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries".




Figure 7. Age distribution and age-specific incidence






Figure 7a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).



Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2014, based on the November 2013 submission. http://www.seer.cancer.gov.





Figure 8. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.



Average incidence (world standard population) 2007 - 2013: Males



Average incidence (world standard population) 2007 - 2013: Females





Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2007 to 2013. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 1.0/100,000 WS N=269, females 0.3/100,000 WS N=81).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,928 female residents (averaged) in the period from 2007 to 2013 a total of 4 women were identified with newly diagnosed base of tongue cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.4/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.1 and 1.6/100,000.

Standardized incidence ratio (SIR) 2007 - 2013: Males



Standardized incidence ratio (SIR) 2007 - 2013: Females




Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2013. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=269, females N=81).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,642 female residents (averaged) in the period from 2007 to 2013 a total of 4 women were identified with newly diagnosed base of tongue cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 1.80. Though, the value of this parameter may vary with an underlying probability of 99% between 0.30 and 5.66, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts, and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

	Incident	Prop.	Prop.		Prop.	Prop. deaths with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	90	olo	n	%	%
1998	14	100.0		11	78.6	100.0
1999	22	100.0		20	90.9	90.0
2000	13	100.0	7.7	10	76.9	100.0
2001	19	100.0	10.5	16	84.2	93.8
2002	33	97.0		24	72.7	95.8
2003	42	100.0	11.9	36	85.7	94.4
2004	49	93.9	8.2	36	73.5	100.0
2005	53	94.3	5.7	31	58.5	96.8
2006	51	98.0	2.0	34	66.7	97.1
2007	53	86.8	1.9	34	64.2	100.0
2008	52	82.7	7.7	33	63.5	100.0
2009	46	93.5		23	50.0	100.0
2010	56	78.6	1.8	30	53.6	100.0
2011	65	75.4	3.1	26	40.0	96.2
2012	55	78.2	1.8	15	27.3	86.7
2013	23	100.0	4.3	7	30.4	100.0
1998-2013	646	89.6	4.0	386	59.8	97.2

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

			Prop. deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n /	n	8	n	૾ૢ
1998	14	12	91.7		
1999	22	11	90.9	/ 1	4.5
2000	13	18	100.0	3	23.1
2001	19	16	93.8	4	21.1
2002	33	20	100.0		
2003	42	24	91.7	10	23.8
2004	49	29	100.0	10	20.4
2005	53	26	100.0	7	13.2
2006	/ 51	47	97.9	12	23.5
2007	53 /	33	97.0	7	13.2
2008	52	30	96.7	12	23.1
2009	46	34	100.0	4	8.7
2010	56	38	100.0	10	17.9
2011	65	46	100.0	6	9.2
2012	55	39	97.4	4	7.3
2013	23	33	100.0	5	21.7
1998-2013	646	456	98.0	95	14.7

Table 10c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

				Prop.
				cancer
		Prop.	Prop.	recorded
		cancer-	non-cancer-	on death
Year of	Deaths	related	related	certificate
death	n/	્રું	8	%
1998	12	75.0	25.0	90.9
1999	11	72.7	27.3	90.0
2000	18	66.7	33.3	72.2
2001	16	75.0	25.0	93.3
2002	20	95.0	5.0	100.0
2003	24	62.5	37.5	90.9
2004	29	93.1	6.9	96.6
2005	26	96.2	3.8	96.2
2006	47	89.4	10.6	95.7
2007	33	78.8	21.2	90.6
2008	30	83.3	16.7	96.6
2009	34	85.3	14.7	91.2
2010	38	89.5	10.5	94.7
2011	46	82.6	17.4	89.1
2012	39	79.5	20.5	92.1
2013	33	84.8	15.2	90.9
1998-2013	456	83.3	16.7	92.4

Table 11a $\begin{tabular}{ll} Medians of age at death according to the grouping in Table 10 \\ \hline MALES \end{tabular}$

		Age at death (all	Age at death (cancer-	Age at death (non-cancer-	Age at death (according to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	8	63.2	61.8	65.7	62.1
1999	8	62.6	59.6	64.2	59.6
2000	16	65.7	61.8	70.4	61.5
2001	12	61.0	58.6	76.6	60.1
2002	15	60.5	56.3	67.3	60.5
2003	21	62.6	63.2	56.5	63.2
2004	23	63.1	62.7	86.1	62.7
2005	19	64.0	63.5	70.1	63.5
2006	33	66.2	66.1	72.2	66.1
2007	26	61.4	61.0	62.7	63.2
2008	25	61.0	61.2	59.2	60.7
2009	28	63.8	62.5	66.4	63.8
2010	27	65.3	65.6	61.7	65.3
2011	38	66.5	64.4	75.8	65.0
2012	31	70.1	68.2	70.7	66.3
2013	23	66.2	65.1	69.4	65.9
1998-2013	353	64.5	63.9	68.1	64.0

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 11b $\begin{tabular}{ll} \begin{tabular}{ll} \begin{tabula$

Year of death	Deaths n	Age at death (all causes)	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate)
1998	4	63.2	52.8	85.2	63.2
1999	3	58.4	55.9	70.0	58.4
2000	2	71.4	71.4		71.4
2001	4	74.4	59.3	84.7	71.9
2002	5	71.9	71.9		71.9
2003	3	52.1	85.4	51.8	68.4
2004	6	65.2	61.3	81.6	65.2
2005	7	62.1	62.1		62.1
2006	14	65.0	65.0		65.0
2007	7/	74.2	72.1	74.2	69.3
2008	5	66.7	58.3	71.9	62.5
2009	6	60.7	61.4	60.0	61.4
2010	11	72.7	67.7	76.0	70.6
2011	8	72.4	72.4	68.5	70.7
2012	8	67.0	65.0	72.3	65.3
2013	10	78.1	75.0	92.1	75.0
1998-2013	103	68.5	65.6	76.0	66.2

By 2010, life expectancy for a newborn male in Germany is 77.5 years compared with 82.6 years for his female counterpart.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 12a

Mortality measures (cancer-related death) and mortality-incidence-index by year of death

MALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	6	0.5	0.60	0.4	0.60	0.4	0.54	0.5	0.53
1999	6	0.5	0.33	0.3	0.32	0.5	0.33	0.5	0.33
2000	10	0.9	1,11	0.5	1.02	0.8	1.08	1.1	1.28
2001	10	0.9	0.83	0.6	0.83	0.8	0.79	0.8	0.72
2002	14	0.8	0.54	0.5	0.55	0.7	0.56	0.9	0.65
2003	14	0.7	0.39	0.5	0.38	0.6	0.38	0.7	0.36
2004	22	1.2	0.67	0.7	0.68	1,0	0.69	1.2	0.76
2005	18	1.0	0.45	0.6	0.39	0.8	0.42	0.9	0.45
2006	28	1.5	0.74	0.8	0.68	1.2	0.68	1.4	0.71
2007	22	1.0	0.51	0.6	0.48	0.9	0.50	0.9	0.52
2008	22	1.0	0.52	0.6	0.50	0.8	0.50	0.9	0.50
2009	24	1.1	0.77	0.6	0.76	0.9	0.76	1.0	0.74
2010	26	1.2	0.65	0.6	0.64	0.9	0.64	1.1	0.66
2011	32	1.4	0.64	0.8	0.65	1.2	0.67	1.4	0.68
2012	26	1.1	0.60	0.6	0.53	0.9	0.57	1.0	0.59
2013	20	0.9	1.00	0.5	0.92	0.7	0.91	0.9	1.06
1998-2013	300	1.0	0.61	0.6	0.58	0.8	0.60	1.0	0.62

Table 12b

Mortality measures (cancer-related death) and mortality-incidence-index by year of death

FEMALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	3	0.3	0.75	0.2	1.00	0.2	0.94	0.3	0.90
1999	2	0.2	0.50	0.1	0.56	0.2	0.58	0.2	0.51
2000	2	0.2	0.50	0.1	0.27	0.1	0.31	0.1	0.42
2001	2	0.2	0.29	0.1	0.40	0.2	0.35	0.2	0.37
2002	5	0.3	0.71	0.1	0.60	0.2	0.59	0.2	0.68
2003	1	0.1	0.20	0.0	0.09	0.0	0.11	0.0	0.11
2004	5	0.3	0.31	0.2	0.35	0.2	0.33	0.2	0.33
2005	7	0.4	0.54	0.2	0.64	0.3	0.64	0.3	0.61
2006	14	0.7	1.08	0.4	0.83	0.5	0.95	0.6	0.94
2007	4	0.2	0.40	0.1	0.26	0.1	0.28	0.1	0.30
2008	3	0.1	0.30	0.1	0.33	0.1	0.33	0.1	0.31
2009	5	0.2	0.33	0.1	0.39	0.2	0.36	0.2	0.37
2010	8	0.3	0.50	0.2	0.51	0.3	0.51	0.3	0.50
2011	6	0.3	0.40	0.1	0.29	0.1	0.30	0.2	0.30
2012	5	0.2	0.42	0.1	0.41	0.2	0.39	0.2	0.39
2013	8	0.3	2.67	0.1	2.22	0.2	2.31	0.3	3.04
1998-2013	80	0.3	0.52	0.1	0.48	0.2	0.49	0.2	0.50

Table 13

Age distribution of age at death (cancer-related) for period 1998-2013

(incl. multiple primaries)

Age at									
death	Cases			Males			Females		
Years	n	%	Cum.%	n	%	Cum.%	n	%	Cum.%
35-39	2	0.5	0.5	2	0.7	0.7			0.0
40-44	8	2.1	2.6	8	2.7	3.3			0.0
45-49	26	6.8	9.4	20	6.6	10.0	6	7.5	7.5
50-54	40	10.5	19.9	33	11.0	20.9	7	8.8	16.3
55-59	50	13.1	33.1	43	14.3	35.2	7	8.8	25.0
60-64	80	21.0	54.1	62	20.6	55.8	18	22.5	47.5
65-69	56	14.7	68.8	47	15.6	71.4	9	11.3	58.8
70-74	45	11.8	80.6	30	10.0	81.4	15	18.8	77.5
75-79	42	11.0	91.6	34	11.3	92.7	8	10.0	87.5
80-84	17	4.5	96.1	13	4.3	97.0	4	5.0	92.5
85+	15	3.9	100.0	9	3.0	100.0	6	7.5	100.0
All ages	381	100.0		301	100.0		80	100.0	

Included in the statistics are 43.0% multiple primaries in males and 36.4% in females.

Table 14

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2013 (incl. multiple primaries)

Age at death	Males Femal	/ - /		Females Age- spec.		cancers	Females Prop.all cancers
Years	n n	mortal.	MI-index	mortal.	MI-index	%	%
0- 4		0.0		0.0			
5- 9		0.0		0.0			
10-14		0.0		0.0			
15-19		0.0		0.0			
20-24		0.0		0.0			
25-29		0.0		0.0			
30-34		0.0		0.0			
35-39	2	0.1	0.18	0.0		0.5	
40-44	8	0.3	0.50	0.0		0.9	
45-49	20 6			0.3	0.38	1.1	0.3
50-54	33 / 7			0.3	0.54	1.0	0.2
55-59	43 / 7			0.4	0.26	0.7	0.1
60-64	62 18			1.0	0.95	0.7	0.3
65-69	47 9			0.5	0.33	0.4	0.1
70-74	30 15			1.0	0.71	0.2	0.2
75-79	34 8			0.7		0.3	0.1
80-84	13 4			0.4		0.1	0.0
85+	9 6	2.6	1.29	0.7	0.86	0.1	0.0
All ages	301 80					0.4	0.1
AII ages	201 00					0.4	0.1
Mortality							
Raw		1.0	0.61	0.3	0.52		
WS		0.6		0.1	0.48		
ES		0.8		0.2	0.49		
BRD-S		1.0		0.2	0.50		
PYLL-70							
per 100,000		9.1		1.9			
ES		8.0		1.6			
AYLL-70		11.3		10.7			

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15a

Multiple primaries in deaths in period 1998-2013

MALES

Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C03-C06 Oral cavity	5	3.2			1	20.0	4	80.0
C09-C10 Oropharynx	13	8.3			2	15.4	11	84.6
C12-C13 Hypopharynx	8	5.1	3	37.5	3	37.5	2	25.0
C15 Oesophagus	14	8.9	5	35.7	1	7.1	8	57.1
C16 Stomach	4	2.5	2	50.0		/ • 1	2	50.0
C18 Colon	5	3.2	1	20.0	1	20.0	3	60.0
C22 Liver	2	1.3	_	20.0		20.0	2	100.0
C25 Pancreas	5	3.2	2	40.0			3	60.0
C32 Larynx	13	8.3	8	61.5	1	7.7	3 4	30.8
C33-C34 Lung	24	15.3	3	12.5	1	4.2	20	83.3
_	3	15.3	3			4.2	20	03.3
C43 Malign. melanoma C44 Skin others	13	8.3	_	100.0	1	7.7	7	53.8
			5		Τ.	7.1		
C46,C49 Soft tissue	3	1.9	2	66.7				33.3
C61 Prostate	6	3.8	3	50.0			3	50.0
C62 Testis	2	1.3	2	100.0	\	\	_	
C64 Kidney	6	3.8	2	33.3	1	16.7	3	50.0
C67 Bladder	6	3.8	4	66.7			2	33.3
C70-C72 CNS cancer	2	1.3	1	50.0	1	50.0		
C73 Thyroid	2	1.3	1	50.0			1	50.0
C76-C79 CUP	9	5.7	7	77.8	2	22.2		
C82-C85 NHL	2	1.3			1	50.0	1	50.0
C91-C96 Leukaemia	2	1.3					2	100.0
Other primaries	8	5.1	5	62.5			3	37.5
All mult. primaries	157	100.0	59	37.6	16	10.2	82	52.2

Multiple primaries with number of cases 1 are pooled in category "Other primaries".

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 15b

Multiple primaries in deaths in period 1998-2013
FEMALES

Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C03-C06 Oral cavity	1	2.6					1	100.0
C09-C10 Oropharynx	$\sqrt{7}$	17.9			2	28.6	5	71.4
C12-C13 Hypopharynx	2	5.1			1	50.0	1	50.0
C15 Oesophagus	4	10.3	1	25.0			3	75.0
C16 Stomach	1	2.6			/1	100.0		
C19-C20 Rectum	1	2.6	1	100.0				
C25 Pancreas	1	2.6					1	100.0
C32 Larynx	4	10.3	1	25.0	2	50.0	1	25.0
C33-C34 Lung	7	17.9			2	28.6	5	71.4
C44 Skin others	2	5.1	2	100.0				
C50 Breast	4	10.3	2	50.0			2	50.0
C53 Cervix uteri	1	2.6					1 /	100.0
C54 Corpus uteri	1	2.6	1	100.0				
C76-C79 CUP	/ 3	7.7	3	100.0				
All mult. primaries	39	100.0	11	28.2	8	20.5	20	51.3




Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2013

(Singular primaries only *)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males	Females	spec.		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0 - 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29			0.0		0.0			
30-34			0.0		0.0			
35-39	1		0.0	0.11	0.0		0.3	
40-44	6		0.2		0.0		0.8	
45-49	14	6	0.6	0.50	0.3	0.43	0.9	0.3
50-54	30	5	1.5		0.2	0.56	1.0	0.2
55-59	34	6	1.9		0.3	0.27	0.7	0.1
60-64	48	13	2.7		0.7	0.81	0.6	0.2
65-69	39	6	2.5	0.68	0.3	0.29	0.4	0.1
70-74	23	12	1.8	0.66	0.8	0.71	0.2	0.2
75-79	21	6	2.5		0.5	0.43	0.2	0.1
80-84	9	\3	1.8		0.3	0.75	0.1	0.0
85+	6	4	1.8	1.50	0.4	1.33	0.1	0.0
All ages	231	61					0.4	0.1
Mortality								
Raw			0.8	0.61	0.2	0.50		
WS			0.5	0.57	0.1	0.46		
ES			0.7	0.59	0.1	0.47		
BRD-S			0.8	0.62	0.2	0.48		
PYLL-70								
per 100,000			7.2		1.5			
ES ES			6.4		1.3			
AYLL-70			11.2		11.4			
עוחח //			11.2		11.4			

^{*} See corresponding tables with multiple primaries.

Table 17

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2013

(Single primaries only *)

Age at death Years	Males n	Females	Males Age- spec. mortal.	MI-index	Females Age- spec. mortal.	MI-index	cancers	Females Prop.all cancers
			/ ./.					
0 - 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29 30-34			0.0		0.0			
35-39	1		0.0	0.11	0.0		0.3	
40-44	5		0.0		0.0		0.3	
45-49	10	4	0.4	0.45	0.2	0.31	0.7	0.3
50-54	25	5	1.2		0.2		1.0	0.2
55-59	27	5	1.5	0.42	0.3		0.6	0.1
60-64	41	10	2.3	0.65	0.5		0.6	0.2
65-69	29	5	1.8	0.58	0.3		0.3	0.1
70-74	17	8	1.3	0.61	0.5		0.2	0.1
75-79	17	5	2.1	0.89	0.4		0.2	0.1
80-84	6	\3	1.2	0.86	0.3		0.1	0.0
85+	6	3	1.8	1.50	0.3	1.00	0.1	0.0
All ages	184	48					0.3	0.1
Mortality								
Raw			0.6	0.55	0.2			
WS			0.4	0.53	0.1	0.41		
ES			0.5	0.54	0.1	0.42		
BRD-S			0.6	0.57	0.1	0.43		
PYLL-70								
per 100,000			5.8		1.2			
ES ES			5.0		1.1			
AYLL-70			11.2		11.3			
, 0			2		11.3			

^{*} See corresponding tables with multiple primaries.





Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at base of tongue cancer-related death (see Table 10) should be considered.



Average mortality (world standard population) 2007 - 2013: Males



Average mortality (world standard population) 2007 - 2013: Females





Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2007 to 2013. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.6/100,000 WS N=171, females 0.1/100,000 WS N=39).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,928 female residents (averaged) in the period from 2007 to 2013 a total of 1 women died from base of tongue cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.1/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 0.5/100,000.

Standardized mortality ratio (SMR) 2007 - 2013: Males



Standardized mortality ratio (SMR) 2007 - 2013: Females




Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2013. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=171, females N=39).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,642 female residents (averaged) in the period from 2007 to 2013 a total of 1 women died from base of tongue cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.95. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 7.09, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

FRG Federal Republic of Germany

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

MCR Munich Cancer Registry (Tumorregister München)
SEER Surveillance, Epidemiology, and End Results (USA)

AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

BRD-S German standard population

DCO Death certificate only EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

ES European standard population (old)

LCL Lower confidence limit

MI-index Ratio between mortality and incidence

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age

SIR Standardized incidence ratio SMR Standardized mortality ratio UCL Upper confidence limit WS World standard population

Recommended Citation

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

Fig./Tbl		Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	3
1a	Gender distribution by year of diagnosis	4
2	Incidence by year of diagnosis	5
3	Age distribution parameters by year of diagnosis	6
4	Age distribution by 5-year age group and gender	8
5	Age-specific incidence and DCO rate	9
6	Standardized incidence ratio of second primaries	10
7	Age distribution and age-specific incidence (chart)	12
7a	Age-specific incidence internationally (chart)	13
8	Cumulative follow-up years (chart)	14
9a	Map of cancer incidence (WS) by county (chart)	15
9b	Standardized incidence ratio (SIR) by county (chart)	16
10a	Pts incident cohorts and mortality / yr	17
10b	Incidence and mortality by year of diagnosis	18
10c	Cancer-related deaths, death certification available / yr	19
11	Medians of age at death / yr	20
12	Mortality by year of death	22
13	Distribution of age at death	23
14	Age-specific mortality	24
15	Multiple primaries in deaths	25
16	Age-specific mortality (first primaries)	27
17	Age-specific mortality (single primaries)	28
18	Age distribution and age-specific mortality (chart)	29
19a	Map of cancer mortality (WS) by county (chart)	30
19b	Standardized mortality ratio (SMR) by county (chart)	31