# **Munich Cancer Registry**



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ Deutsch

Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

# **Cancer statistics: Baseline statistics**

C51: Vulva cancer

Year of diagnosis	1998-2013
Patients	1,329
Diseases	1,332
Creation date	05/19/2015
Export date	12/30/2014
Population (females)	2.36 m



http://www.tumorregister-muenchen.de/en/facts/base/base\_C51\_\_E.pdf

# Global Statements about the statistics on the Internet – Baseline Statistics (grey button ——), Survival (red button ——)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut<sup>#</sup>, with a total of 4.64 million inhabitants, account for the frequency of cancer diseases<sup>##</sup> and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases\*\*\*\* are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR. The time-delayed acquisition of data and the occasionally high DCO-rates indicate optimizing reserves, among others, because of current financial and legal conditions that hinder the analyses.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, May 2015

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007). Death certificates from 2014 are incorporated into these analyses.
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate. A high proportion of DCO cases (≥5%) in particular cancer types indicate insufficient participation of specific cancer specializations.

### ICD-10 codes (ICD-10 2015) used for specifying cancer site

Code	Description
C51	Malignant neoplasm of vulva
C51.0	Labium majus
C51.1	Labium minus
C51.2	Clitoris
C51.8	Overlapping lesion of vulva
C51.9	Vulva, unspecified

Page 3 of 29

## **INCIDENCE**

Table 1

Patient cohorts by year of diagnosis including DCO cases and multiple primaries, and with proportion of deaths and active follow-up

				Prop.		Prop.
		DCO	Prop.	mult.	Prop.	actively
Year of	Cases	cases	DCO	primaries	deaths	followed
diagnosis	n	n	%	8	%	%
1998	42	2	4.8	26.2	81.0	97.6
1999	37	3	8.1	10.8	78.4	100.0
2000	39	5	12.8	23.1	71.8	100.0
2001	37	4	10.8	21.6	70.3	97.3
2002	70	5	7.1	17.1	65.7	98.6 #
2003	78	5	6.4	34.6	73.1	98.7
2004	/75	5	6.7	30.7	65.3	94.7
2005	91	6	6.6	29.7	59.3	97.8
2006	88	/ 1	1.1	25.0	51.1	95.5
2007	106	4	3.8	22.6	50.0	84.0 # ##
2008	110	4	3.6	26.4	49.1	67.3
2009	103	3	2.9	22.3	50.5	72.8
2010	137	4	2.9	25.5	43.1	70.8
2011	113	6	5.3	23.0	39.8	61.9
2012	119	3	2.5	20.2	27.7	61.3
2013	87	4	4.6	17.2	9.2	98.9 ###
1998-2013	1332	64	4.8	23.9	50.5	83.1

<sup>#</sup> The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

<sup>##</sup> Since 2007 the percentage of actively followed patients sharply declined compared to the previous years. This is a consequence of ambiguous data protection rules that currently forbid cancer registries in Bavaria to obtain the essential life status informations from competent registration offices.

<sup>###</sup> Please be aware that data of recent annual patient cohorts may not yet be fully processed. Therefore, the presented figures and tables are potentially related to different time periods as pointed out in the respective headlines or legends.

Table 2

Incidence measures by year of diagnosis and gender including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

Year of	Cases	Incidence	Incidence	Incidence	Incidence
diagnosis	n	raw	WS	ES	BRD-S
1998	42	3.6	1.5	2.3	3.1
1999	37	/ 3.1/	1.2	1.9	2.5
2000	39	3.2	1.5	2.2	2.7
2001	37	3.0	1.4	2.0	2.6
2002	70	3.6	1.5	2.2	2.9
2003	78	4.0	1.5	2.3	3.1
2004	75	3.8	1.6	2.3	3.0
2005	91	4.6	1.7	2.6	3.6
2006	88	4.4	1.8	2.6	3.4
2007	106	4.6	1.9	2.8	3.7
2008	110	4.7	1.9	2.8	3.6
2009	103	4.4	1.7	2.6	3.4
2010	137	5.9	2.4	3.5	4.4
2011	113	4.8	1.8	2.7	3.5
2012	/119	5.0	2.2	3.1	4.0
2013	87	3.7	1.4	2.1	2.7
1998-2013	1332	4.3	1.7	2.6	3.4

The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

Table 3

Age distribution parameters by year of diagnosis (incl. DCO)

	~		~. ]							
Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	42	72.1	13,6	36.3	93.3	52.1	67.5	74.2	80.6	87.2
1999	37	71.8	15.5	34.3	94.0	48.7	62.4	75.6	84.8	88.0
2000	39	69.2	16.4	34.5	92.3	42.6	59.3	74.3	83.0	88.7
2001	37	69.8	13.3	33.3	89.1	49.6	61.0	70.0	80.0	86.2
2002	70	71.2	16.1	26.9	96.1	43.0	65.7	75.1	81.7	89.6
2003	78	74.5	14.8	34.3	94.8	50,2	63.8	79.5	85.7	91.0
2004	75	72.0	14.8	32.7	95.5	50.7	61.1	76.3	83.9	88.6
2005	91	74.5	13.6	34.9	96.9	53.8	67.4	78.0	83.6	89.4
2006	88	73.0	14.2	34.6	97.2	47.7	65.3	74.0	84.0	88.7
2007	106	71.5	14.4	34.1	96.7	48.6	62.8	74.8	82.8	86.8
2008	110	73.2	14.0	28.7	96.9	52.8	65.9	74.9	83.5	89.1
2009	103	72.7	15.2	24.9	100	50.5	64.4	76.7	84.6	88.3
2010	137	71.8	13.6	36.5	95.1	48.6	65.0	72.6	82.1	88.7
2011	113	72.9	14.8	30.3	98.5	49.7	66.9	75.7	83.8	89.1
2012	119/	70.4	15.3	27.3	101	46.9	61.5	73.4	82.3	87.8
2013	87	73.4	13.2	33.1	98.6	53.0	66.4	75.9	82.4	88.3
1998-2013	1332	72.3	14.5	24.9	101	49.7	64.1	75.3	83.0	88.4

Table 4

Age distribution by 5-year age group for period 1998-2013 (incl. DCO)

Age at			
diagnosis	Cases		
Years	'n	%	Cum.%
20-24	/ 1	0.1	0.1
25-29	/ 4	0.3	0.4
30-34	13	1.0	1.4
35-39	22	1.7/	3.0
40-44	34	2.6	5.6
45-49	63	4.7	10.3
50-54	50	3.8	14.0
55-59	77	5.8	19.8
60-64	96	7.2	27.0
65-69	128	9.6	36.6
70-74	166	12.5	49.1
75-79	206	15.5	64.6
80-84	207	15.5	80.1
85+	265	19.9	100.0
All ages	1332	100.0	

Included in the statistics are 31.3% multiple primaries.

Table 5

Age-specific incidence, DCO rate and proportion of all cancers for period 1998-2013

				Prop. all	
Age at			DCO rate	cancers	
diagnosis	Cases	Age-spec.	n=64	n=153136	
Years	n /	incidence	8	8	
0- 4		0.0			
5- 9		0.0			
10-14		0.0			
15-19		0.0			
20-24	1	0.1		0.2	
25-29	4	0.2		0.4	
30-34	13	0.6		0.6	
35-39	22	0.9		0.6	
40-44	34	1.4		0.5	
45-49	63	2.7		0.7	
50-54	50	2.4		0.5	
55-59	77	4.0	2.6	0.6	
60-64	96	5.1		0.6	
65-69	128	7.4	0.8	0.7	
70-74	166	10.9	2.4	0.9	
75-79	206	17.3	3.4	1.2	
80-84	207	22.2	4.8	1.3	
85+	265	29.6	15.1	1.5	
All ages	1332		4.8	0.9	
Incidence					
Raw		4.3			
WS		1.7			
ES		2.6			
BRD-S		3.4			

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

Table 6

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2013

	Observed	Expected		LCL	UCL		DCO
Diagnosis	n	n	SIR	95%	95%	EAR	%
C18 Colon	/ 10 /	4.5	2.2	1.1	4.1 #	17.5	30.0
C19-C20 Rectum	2 /	1.8	1.1	0.1	4.0	0.6	
C21 Anus/canal	8 /	0.2	38.7	16.7	76.2 #	24.8	
C33-C34 Lung	/ 7/	2.6	2.7	1.1	5.6 #	14.0	
C43 Malign. melanoma	5	1.3	3.8	1.2	9.0 #	11.8	40.0
C50 Breast	18	10.9	1.7	/1.0	2.6	22.7	11.1
C51 Vulva	3	0.4	6.7	1.4	19.6 #	8.1	
C52 Vagina	2	0.1	24.3	2.9	87.8 #	6.1	
C53 Cervix uteri	3	0.5	6.0	1.2	17.6 #	8.0	
C54 Corpus uteri	3	2.0	1.5	0.3	4.3	3.1	
C56 Ovary	4	1.6	2.5	0.7	6.4	7.6	25.0
C67 Bladder	2	0.9	2.3	0.3	8.2	3.6	
C68 Urethra	2	0.0	169.6	20.5	612.8 #	6.3	
C70-C72 CNS cancer	2	0.5	3.8	0.5	13.8	4.7	50.0
C76-C79 CUP	2	0.8	2.4	0.3	8.6	3.7	
C82-C85 NHL	5	1.6	3.2	1.0	7.4 #	10.9	
C91-C96 Leukaemia	2	0.7	2.9	0.4	10.4	4.2	100.0
Other primaries	9	6.4	1.4	0.6	2.7	8.2	
Not observed	0	2.8	0.0	0.0	1.3	-8.9	
All mult. primaries	89	39.7	2.2	1.8	2.8 #	157.0	12.4

Patients	844
Median age at second malignancy (years)	73.3
Person-years	3140
Mean observation time (years)	3.7
Median observation time (years)	2.5

# The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries".








Figure 7. Age distribution and age-specific incidence



## C51: Malignant neoplasm of vulva Age-specific incidence in international comparison Average 35 Region Period population MCR SEER 2007-2013 2007-2011 2.3 m 32.5 m 30 5 2 **FEMALES** Age at diagnosis (years)

**Figure 7a.** Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).



Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER\*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2014, based on the November 2013 submission. http://www.seer.cancer.gov.





Figure 8. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.



### Average incidence (world standard population) 2007 - 2013




**Figure 9a.** Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2007 to 2013. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (1.9/100,000 WS N=775).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,928 female residents (averaged) in the period from 2007 to 2013 a total of 18 women were identified with newly diagnosed vulva cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 1.4/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.6 and 2.8/100,000.



### Standardized incidence ratio (SIR) 2007 - 2013



**Figure 9b.** Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2013. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (N=775).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,642 female residents (averaged) in the period from 2007 to 2013 a total of 18 women were identified with newly diagnosed vulva cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.89. Though, the value of this parameter may vary with an underlying probability of 99% between 0.44 and 1.58, and is therefore not statistically striking.



## **MORTALITY**

#### Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts, and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

Year of	Incident cases	Prop. actively followed	Prop. DCO	Deaths	Prop. deaths	Prop. deaths with death certific.
diagnosis	n	%	00	n	%	%
1998	42	97.6	4.8	34	81.0	88.2
1999	37	100.0	8.1	29	78.4	89.7
2000	39	100.0	12.8	28	71.8	85.7
2001	37	97.3	10.8	26	70.3	100.0
2002	70	98.6	7.1	46	65.7	95.7
2003	78	98.7	6.4	57	73.1	98.2
2004	75	94.7	6.7	49	65.3	98.0
2005	91	97.8	6.6	54	59.3	96.3
2006	88	95.5	1.1	45	51.1	97.8
2007	106	84.0	3.8	53	50.0	100.0
2008	110	67.3	3.6	54	49.1	96.3
2009	103	72.8	2.9	52	50.5	100.0
2010	137	70.8	2.9	59	43.1	98.3
2011	113	61.9	5.3	45	39.8	93.3
2012	119	61.3	2.5	33	27.7	90.9
2013	87	98.9	4.6	8	9.2	75.0
1998-2013	1332	83.1	4.8	672	50.5	95.7

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

			Prop. deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	n	%	n	96
1998	42	24	75.0	4	9.5
1999	37	27	81.5	6	16.2
2000	39	31	96.8	4	10.3
2001	37	29	89.7	4	10.8
2002	70	40	95.0	12	17.1
2003	78	44	97.7	11	14.1
2004	75	53	100.0	11	14.7
2005	91	51	100.0	12	13.2
2006	88	49	95.9	_ 5	5.7
2007	106	63	95.2	14	13.2
2008	110	67	100.0	17	15.5
2009	103	59	98.3	7	6.8
2010	137	68	100.0	17	12.4
2011	113	83	98.8	16	14.2
2012	119	86	97.7	17	14.3
2013	87	63	96.8	5	5.7
1998-2013	1332	837	96.5	162	12.2

#### Table 10c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

				Prop.
				cancer
		Prop.	Prop.	recorded
		cancer-	non-cancer-	on death
Year of	Deaths	related	related	certificate
death	n	્રે	8	%
1998	24	70.8	29.2	83.3
1999	27	63.0	37.0	95.5
2000	31	61.3	38.7	76.7
2001	29	62.1	37.9	80.8
2002	40	72.5	27.5	73.7
2003	44	75.0	25.0	79.1
2004	53	73.6	26.4	84.9
2005	51	64.7	35.3	74.5
2006	49	59.2	40.8	74.5
2007	63	69.8	30.2	81.7
2008	67	67.2	32.8	71.6
2009	59	62.7	37.3	77.6
2010	\ 68	63.2	36.8	73.5
2011	83	67.5	32.5	76.8
2012	86	60.5	39.5	73.8
2013	63	52.4	47.6	70.5
1998-2013	837	65.0	35.0	76.7

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(non-cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	24	82.7	77.8	87.0	83.0
1999	27	79.5	75.6	84.6	76.9
2000	31	84.7	85.4	84.6	85.4
2001	29	79.3	78.6	80.5	79.0
2002	40	81.9	81.9	83.7	81.9
2003	44	82.3	78.2	88.9	77.9
2004	53	83.4	83.2	84.5	83.2
2005	51	83.2	81.4	85.1	82.9
2006	49	80.2	77.7	84.7	78.3
2007	63	82.1	80.5	87.4	80.8
2008	67	85.5	81.8	86.5	82.5
2009	59	82.9	81.1	84.9	82.6
2010	68	82.4	81.5	84.9	82.0
2011	83	83.9	79.1	86.0	79.4
2012	86	84.9	81.2	88.2	83.4
2013	63	84.3	78.7	88.3	81.3
1998-2013	837	83.1	81.1	85.9	81.5

By 2010, life expectancy for a newborn male in Germany is 77.5 years compared with 82.6 years for his female counterpart.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 12

Mortality measures (cancer-related death) and mortality-incidence-index by year of death

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	17	1.4	0.40	0.5	0.34	0.8	0.35	1.1	0.35
1999	17	1.4	0.46	0.6	0.47	0.9	0.47	1.1	0.46
2000	19	1.6	0.49	0.4	0.27	0.7	0.34	1.1	0.39
2001	18	1.5	0.49	0.5	0.38	0.9	0.42	1.2	0.46
2002	29	1.5	0.41	0.4	0.28	0.7	0.32	1.1	0.36
2003	33	1.7	0.42	0.6	0.40	0.9	0.41	1.3	0.42
2004	39	2.0	0.52	0.6	0.36	0.9	0.40	1.3	0.43
2005	33	1.7	0.36	0.5	0.31	0.8	0.32	1.2	0.32
2006	29	1.4	0.33	0.5	0.26	0.8	0.29	1.2	0.35
2007	44	1.9	0.42	0.5	0.29	0.9	0.33	1.4	0.37
2008	45	1.9	0.41	0.6	0.31	0.9	0.33	1.3	0.35
2009	37	1.6	0.36	0.5	0.27	0.8	0.29	1.1	0.32
2010	43	1.8	0.31	0.5	0.23	0.9	0.26	1.3	0.29
2011	56	2.4	0.50	0.8	0.45	1.2	0.45	1.6	0.45
2012	52	2.2	0.44	0.6	0.26	1.0	0.32	1.5	0.39
2013	33	1.4	0.38	0.5	0.35	0.8	0.37	1.0	0.37
1998-2013	544	1.8	0.41	0.5	0.31	0.9	0.34	1.2	0.37

Table 13

Age distribution of age at death (cancer-related) for period 1998-2013

(incl. multiple primaries)

Age at				
death	Cases			
Years	'n	%	Cum.%	
30-34	/ 3	0.6	0.6	
35-39	/ 2	0.4	0.9	
40-44	4	0.7	1.7	
45-49	3	0.6	2.2	
50-54	8	1.5	3.7	
55-59	19	3.5	7.2	
60-64	27	5.0	12.1	
65-69	54	9.9	22.1	
70-74	53	9.7	31.8	
75-79	81	14.9	46.7	
80-84	115	21.1	67.8	
85+	175	32.2	100.0	
All ages	544	100.0		

Included in the statistics are 31.3% multiple primaries.

Table 14

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2013

(incl. multiple primaries)

Age at				Prop. all
death	Cases	Age-spec.		cancers
Years	n	mortality	MI-index	96
0- 4		0.0		
5- 9		0.0		
10-14		0.0		
15-19		0.0		
20-24		0.0		
25-29		0.0		
30-34	3	0.1	0.23	1.3
35-39	2	0.1	0.09	0.4
40-44	4	0.2	0.12	0.4
45-49	3	0.1	0.05	0.1
50-54	8	0.4	0.16	0.3
55-59	19	1.0	0.25	0.4
60-64	27	1.4	0.28	0.4
65-69	54	3.1	0.42	0.6
70-74	53	3.5	0.32	0.5
75-79	81	6.8	0.39	0.8
80-84	115	12.3	0.56	1.0
85+	175	19.6	0.66	1.3
All ages	544			0.8
Mortality			/	
Raw		1.8	0.41	
WS		0.5	0.31	
ES		0.9	0.34	
BRD-S		1.2	0.37	
PYLL-70				
per 100,000		4.0		
ES ES		3.4		
		8.9		

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15

Multiple primaries in deaths in period 1998-2013

						Syn- chron	Syn- chron		
		Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	5	n/	% ↓	n	<b>←</b> %	n	<b>~</b> %	n	<b>~</b> %
_									
C16 S	Stomach	6	2.5	1	16.7			5	83.3
C18 C	Colon	/17	7.2	6	35.3	1	5.9	10	58.8
C19-C20 R	Rectum	10	4.2	6	60.0	1	10.0	3	30.0
C21 A	Anus/canal	6 /	2.5	4	66.7			2	33.3
C22 I	Liver	4	1.7	1	25.0			3	75.0
C25 P	Pancreas	3	1.3	1	33.3			2	66.7
C33-C34 L	Lung	19	8.0	3	15.8	2	10.5	14	73.7
C43 M	Malign. melanoma	10	4.2	2	20.0	1	10.0	7	70.0
C44 S	Skin others	12	5.1	2	16.7	4	33.3	6	50.0
C46,C49 S	Soft tissue	4	1.7	1	25.0	1	25.0	2	50.0
C50 E	Breast	35	14.8	21	60.0	3	8.6	11	31.4
C51 V	/ulva	6	2.5			1	16.7	5	83.3
C52 V	<i>l</i> agina	4	1.7	1	25.0	1	25.0	2	50.0
C53 C	Cervix uteri	19	8.0	14	73.7	_ 3	15.8	2	10.5
C54 C	Corpus uteri	17	7.2	16	94.1	1	5.9		
C56 C	Ovary	12	5.1	7	58.3	2	16.7	3	25.0
C64 K	Kidney	3	1.3	2	66.7			1	33.3
C67 E	Bladder	11	4.6	3	27.3	1	9.1	7	63.6
C68 U	Jrethra \	3	1.3	1	33.3			2	66.7
C70-C72 C	CNS cancer	4	1.7	3	75.0			1	25.0
C76-C79 C	CUP	3	1.3			2	66.7	1	33.3
C82-C85 N	NHL	9	3.8	3	33.3			6	66.7
C91-C96 I	Leukaemia	6	2.5	1	16.7			5	83.3
Other pri	imaries	14	5.9	7	50.0			7	50.0
All mult.	. primaries	237	100.0	106	44.7	24	10.1	107	45.1

Multiple primaries with number of cases 1 to 2 are pooled in category "Other primaries".

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2013

(Singular primaries only \*)

Age at				Prop. all	
death	Cases	Age-spec.		cancers	
Years	n	mortality	MI-index	%	
0- 4		0.0			
5- 9		0.0			
10-14		0.0			
15-19		0.0			
20-24		0.0			
25-29		0.0			
30-34	2	0.1	0.17	1.0	
35-39	2	0.1	0.10	0.4	
40-44	2	0.1	0.07	0.2	
45-49	2	0.1	0.04	0.1	
50-54	7	0.3	0.16	0.3	
55-59	15	0.8	0.23	0.4	
60-64	20	1.1	0.25	0.4	
65-69	41	2.4	0.41	0.6	
70-74	45	3.0	0.33	0.6	
75-79	62	5.2	0.37	0.7	
80-84	88	9.4	0.53	1.0	
85+	146	16.3	0.66	1.3	
All ages	432			0.7	
Mortality					
Raw		1.4	0.39		
WS		0.4	0.29		
ES		0.7	0.32		
BRD-S		1.0	0.35		
DVI I 70					
PYLL-70		2.0			
per 100,000		3.0			
ES 70		2.5			
AYLL-70		8.8			

<sup>\*</sup> See corresponding tables with multiple primaries.


Table 17

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2013

(Single primaries only \*)

1 2 2 6 12	Age-spec. mortality  0.0 0.0 0.0 0.0 0.0 0.0 0.1 0.1 0.1 0.	0.08 0.10 0.07 0.04 0.14 0.20	Prop. all cancers %  0.5 0.5 0.2 0.1 0.3
1 2 2 2 6	0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.1 0.1 0.1	0.08 0.10 0.07 0.04 0.14	0.5 0.5 0.2 0.1
2 2 2 6	0.0 0.0 0.0 0.0 0.0 0.1 0.1 0.1	0.10 0.07 0.04 0.14	0.5 0.2 0.1 0.3
2 2 2 6	0.0 0.0 0.0 0.0 0.0 0.1 0.1 0.1	0.10 0.07 0.04 0.14	0.5 0.2 0.1 0.3
2 2 2 6	0.0 0.0 0.0 0.0 0.1 0.1 0.1 0.3	0.10 0.07 0.04 0.14	0.5 0.2 0.1 0.3
2 2 2 6	0.0 0.0 0.0 0.0 0.1 0.1 0.1 0.3	0.10 0.07 0.04 0.14	0.5 0.2 0.1 0.3
2 2 2 6	0.0 0.0 0.0 0.1 0.1 0.1 0.3	0.10 0.07 0.04 0.14	0.5 0.2 0.1 0.3
2 2 2 6	0.0 0.0 0.1 0.1 0.1 0.3	0.10 0.07 0.04 0.14	0.5 0.2 0.1 0.3
2 2 2 6	0.0 0.1 0.1 0.1 0.3	0.10 0.07 0.04 0.14	0.5 0.2 0.1 0.3
2 2 2 6	0.1 0.1 0.1 0.3	0.10 0.07 0.04 0.14	0.5 0.2 0.1 0.3
2 2 6	0.1 0.1 0.3	0.07 0.04 0.14	0.2 0.1 0.3
2 6	0.1 0.3	0.04 0.14	0.1
6	0.3	0.14	0.3
12	0.6	በ ኃስ	
			0.3
			0.4
			0.5
			0.6
_			0.6
			0.9
130	14.5	0.60	1.4
354			0.7
33 <sup>4</sup>			0.7
	1.1		
	0.3	0.26	
	0.6	0.29	
	0.8	0.31	
	2.5		
	2.1		
	9.5		
	18 26 40 45 70 130	26	26

<sup>\*</sup> See corresponding tables with multiple primaries.




**Figure 18.** Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at vulva cancer-related death (see Table 10) should be considered.




### Average mortality (world standard population) 2007 - 2013



**Figure 19a.** Map of cancer mortality (world standard population) by county averaged for period 2007 to 2013. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (0.6/100,000 WS N=306).


The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,928 female residents (averaged) in the period from 2007 to 2013 a total of 6 women died from vulva cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.4/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.1 and 1.2/100,000.

## Standardized mortality ratio (SMR) 2007 - 2013



**Figure 19b.** Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2013. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (N=306).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,642 female residents (averaged) in the period from 2007 to 2013 a total of 6 women died from vulva cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.78. Though, the value of this parameter may vary with an underlying probability of 99% between 0.20 and 2.03, and is therefore not statistically striking.



#### Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

#### 1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

#### 2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

## **3. Single primary** (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

#### **Shortcuts**

FRG Federal Republic of Germany

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

MCR Munich Cancer Registry (Tumorregister München)
SEER Surveillance, Epidemiology, and End Results (USA)

AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

BRD-S German standard population

DCO Death certificate only EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

ES European standard population (old)

LCL Lower confidence limit

MI-index Ratio between mortality and incidence

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age

SIR Standardized incidence ratio
SMR Standardized mortality ratio
UCL Upper confidence limit
WS World standard population

#### **Recommended Citation**

Munich Cancer Registry. Baseline statistics C51: Vulva cancer [Internet]. 2015 [updated 2015 May 19; cited 2015 Jul 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/base/base C51 E.pdf

#### Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

#### Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

## Index of figures and tables

Fig./Tbl		Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	3
2	Incidence by year of diagnosis	4
3	Age distribution parameters by year of diagnosis	5
4	Age distribution by 5-year age group	6
5	Age-specific incidence and DCO rate	7
6	Standardized incidence ratio of second primaries	8
7	Age distribution and age-specific incidence (chart)	9
7a	Age-specific incidence internationally (chart)	10
8	Cumulative follow-up years (chart)	11
9a	Map of cancer incidence (WS) by county (chart)	12
9b	Standardized incidence ratio (SIR) by county (chart)	13
10a	Pts incident cohorts and mortality / yr	14
10b	Incidence and mortality by year of diagnosis	15
10c	Cancer-related deaths, death certification available / yr	16
11	Medians of age at death / yr	17
12	Mortality by year of death	18
13	Distribution of age at death	19
14	Age-specific mortality	20
15	Multiple primaries in deaths	21
16	Age-specific mortality (first primaries)	22
17	Age-specific mortality (single primaries)	23
18	Age distribution and age-specific mortality (chart)	24
19a	Map of cancer mortality (WS) by county (chart)	25
19b	Standardized mortality ratio (SMR) by county (chart)	26