# **Munich Cancer Registry**



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ Deutsch

Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

# **Cancer statistics: Baseline statistics**

# C53: Cervical cancer

Year of diagnosis	1998-2013
Patients	3,474
Diseases	3,477
Creation date	05/19/2015
Export date	12/30/2014
Population (females)	2.36 m



http://www.tumorregister-muenchen.de/en/facts/base/base\_C53\_\_E.pdf

# Global Statements about the statistics on the Internet – Baseline Statistics (grey button ——), Survival (red button ——)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut<sup>#</sup>, with a total of 4.64 million inhabitants, account for the frequency of cancer diseases<sup>##</sup> and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases\*\*\*\* are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR. The time-delayed acquisition of data and the occasionally high DCO-rates indicate optimizing reserves, among others, because of current financial and legal conditions that hinder the analyses.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, May 2015

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007). Death certificates from 2014 are incorporated into these analyses.
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate. A high proportion of DCO cases (≥5%) in particular cancer types indicate insufficient participation of specific cancer specializations.

## ICD-10 codes (ICD-10 2015) used for specifying cancer site

cervix uteri
ervix uteri
=

## **INCIDENCE**

Table 1

Patient cohorts by year of diagnosis including DCO cases and multiple primaries, and with proportion of deaths and active follow-up

				Prop.		Prop.
		DCO	Prop.	mult.	Prop.	actively
Year of	Cases	cases	DCO	primaries	deaths	followed
diagnosis	n	n	%	8	%	%
1998	144	8	5.6	16.0	46.5	91.7
1999	161	2	1.2	18.0	40.4	93.8
2000	148	8	5.4	24.3	39.9	95.9
2001	151	6	4.0	20.5	42.4	93.4
2002	238	23	9.7	18.5	45.4	96.2 #
2003	218	10	4.6	18.3	50.5	93.6
2004	214	14	6.5	19.2	47.2	96.3
2005	235	12	5.1	14.9	41.3	93.2
2006	242	8	3.3	13.6	38.8	86.8
2007	243	8	3.3	13.2	38.3	79.0 # ##
2008	271	6	2.2	16.2	36.2	59.4
2009	267	10	3.7	13.9	38.2	57.3
2010	251	12	4.8	21.5	32.7	61.4
2011	245	8	3.3	11.4	28.2	58.8
2012	243	19	7.8	20.6	31.7	66.3
2013	206	11	5.3	14.6	17.0	97.6 ###
1998-2013	3477	165	4.7	16.9	38.0	80.5

<sup>#</sup> The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

<sup>##</sup> Since 2007 the percentage of actively followed patients sharply declined compared to the previous years. This is a consequence of ambiguous data protection rules that currently forbid cancer registries in Bavaria to obtain the essential life status informations from competent registration offices.

<sup>###</sup> Please be aware that data of recent annual patient cohorts may not yet be fully processed. Therefore, the presented figures and tables are potentially related to different time periods as pointed out in the respective headlines or legends.

Table 2

Incidence measures by year of diagnosis and gender including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

Year of diagnosis	Cases n	Incidence raw	Incidence WS	Incidence ES	Incidence BRD-S
1998	144	12.2	8.1	10.3	11.2
1999	161	13.6	9.2	11.5	12.4
2000	148	12.3	8.2	10.3	11.2
2001	151	12.4	8.0	10.2	11.0
2002	238	12.2	7.7	9.8	10.8
2003	218	11.1	7.0	9.0	9.8
2004	214	10.8	6.8	8.8	9.5
2005	235	11.8	7.5	9.5	10.3
2006	242	12.0	7.8	9.9	10.6
2007	243	10.5	6.9	8.7	9.2
2008	271	11.7	7.7	9.7	10.3
2009	267	11.5	7.5	9.6	10.3
2010	251	10.7	7.0	8.9	9.5
2011	245	10.4	7.1	8.8	9.3
2012	243	10.3	6.4	8.3	9.0
2013	206	8.7	6.0	7.5	8.0
1998-2013	3477	11.2	7.3	9.3	10.0




Table 3

Age distribution parameters by year of diagnosis (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	144	53.4	16.0	23.8	89.4	34.5	41.1	49.8	64.5	79.7
1999	161	51.3	16.5	24.4	90.0	33.8	37.7	47.9	63.8	77.7
2000	148	53.1	16.8	23.9	90.7	33.6	39.2	49.3	64.0	79.9
2001	151	54.3	17.0	28.8	96.0	34.4	40.0	48.9	63.6	80.9
2002	238	55.3	18.0	25.9	96.1	34.9	39.4	51.6	70.3	81.7
2003	218	56.2	16.9	27.3	93.4	36.0	42.7	53.4	67.5	82.0
2004	214	55.6	17.1	21.0	95.2	36.0	42.1	53.0	67.2	82.7
2005	235	55.5	17.4	24.0	100	35.1	40.3	54.5	68.7	80.8
2006	242	54.6	16.8	22.9	99.4	35.6	41.5	50.7	65.2	81.3
2007	243	53.3	17.1	22.0	96.6	34.4	40.7	48.8	66.8	79.9
2008	271	54.0	15.6	24.0	92.8	36.5	42.4	50.7	66.9	75.3
2009	267	55.1	17.1	23.1	95.1	35.9	41.0	52.2	66.3	81.9
2010	251	55.1	16.4	25.1	93.2	35.5	42.3	52.5	67.5	80.7
2011	245	53.0	16.2	25.7	95.6	32.9	41.5	50.2	63.6	77.8
2012	243	57.0	17.0	25.4	95.7	36.2	43.1	56.2	70.4	80.7
2013	206	53.1	15.7	22.1	96.0	33.8	42.8	50.0	63.6	77.2
1998-2013	3477	54.5	16.8	21.0	100	34.9	41.2	51.5	66.8	80.5

Table 4

Age distribution by 5-year age group for period 1998-2013 (incl. DCO)

Age at			
diagnosis	Cases		
Years	'n	%	Cum.%
20-24	13	0.4	0.4
25-29	100	2.9	3.2
30-34	244	7.0	10.3
35-39	401	11.5	21.8
40-44	493	14.2	36.0
45-49	392	11.3	47.3
50-54	324	9.3	56.6
55-59	308	8.9	65.4
60-64	259	7.4	72.9
65-69	226	6.5	79.4
70-74	198	5.7	85.1
75-79	158	4.5	89.6
80-84	173	5.0	94.6
85+	188	5.4	100.0
All ages	3477	100.0	

Included in the statistics are 19.5% multiple primaries.

Table 5

Age-specific incidence, DCO rate and proportion of all cancers for period 1998-2013

				Prop. all	
Age at			DCO rate	cancers	
diagnosis	Cases	Age-spec.	n=165	n=153136	
Years	n /	incidence	%	%	
0 - 4		0.0			
5- 9		0.0			
10-14		0.0			
15-19		0.0			
20-24	13	0.7		2.5	
25-29	99	4.9		8.9	
30-34	244	11.0		11.9	
35-39	401	16.9	0.2	10.7	
40-44	492	19.8	0.4	7.9	
45-49	392	17.0	1.8	4.5	
50-54	324	15.8	1.5	2.9	
55-59	308	16.0	1.6	2.2	
60-64	259	13.8	1.5	1.5	
65-69	226	13.1	4.0	1.2	
70-74	198	13.0	7.1	1.1	
75-79	158	13.3	10.1	0.9	
80-84	173	18.5	21.4	\1.1	
85+	188	21.0	34.6	1.1	
All ages	3475		4.7	2.3	
Incidence					
Raw		11.2			
WS		7.3			
ES		9.3			
BRD-S		10.0			

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

Table 6

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2013

	Observed E	Expected		LCL	UCL		DCO
Diagnosis	/n /	n	SIR	95%	95%	EAR	%
C03-C06 Oral cavity	/ 2 /	0.4	4.7	0.6	17.0	1.6	
C09-C10 Oropharynx	3 /	0.3	8.8	1.8	25.8	# 2.6	
C16 Stomach	2	1.9	1./1	0.1	3.8	0.1	
C18 Colon	12	5.2	2.3	1.2	4.1	# 6.8	8.3
C19-C20 Rectum	12	2.5	4.9	2.5	8.5	# 9.4	16.7
C21 Anus/canal	6	0.4	16.4	6.0	35.7	# 5.6	16.7
C25 Pancreas	8	2.1	3.8	1.7	7.5	<sup>#</sup> 5.8	37.5
C33-C34 Lung	33	4.3	7.7	5.3	10.8	# 28.4	3.0
C43 Malign. melanoma	5	3.0	1.6	0.5	3.8	1.9	
C46,C49 Soft tissue	3	0.4	7.8	1.6	22.9	# 2.6	
C50 Breast	39	23.3	1.7	1.2	2.3	# 15.5	2.6
C51 Vulva	6	0.5	11.4	4.2	24.9	\$ 5.4	
C52 Vagina	5	0.1	46.8	15.2	109.3	# 4.8	
C54 Corpus uteri	20	3.3	6.0	3.7	9.3	# 16.5	25.0
C56 Ovary	28	2.6	10.6	7.1	15.4	<sup>‡</sup> 25.1	35.7
C64 Kidney	2	1.4	1.5	0.2	5.3	0.6	
C67 Bladder	10	0.9	11.0	5.3	20.3	# 9.0	10.0
C73 Thyroid	6	2.0	2.9	1.1	6.4	# 3.9	
C76-C79 CUP	2	0.9	2.2	0.3	7.9	1.1	
C82-C85 NHL	4	2.2	1.8	0.5	4.7	1.8	
C91-C96 Leukaemia	4	0.9	4.4	1.2	11.2	3.1	50.0
Other primaries	8	2.8	2.9	1.2	5.6	<sup>#</sup> 5.1	
Not observed	0	4.2	0.0	0.0	0.9/	+ -4.1	
All mult. primaries	220	65.7	3.3	2.9	3.8	# 152.6	12.3

Patients	2279
Median age at second malignancy (years)	64.6
Person-years	10112
Mean observation time (years)	4.4
Median observation time (years)	3.1

# The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries".

# C53: Malignant neoplasm of cervix uteri Age distribution and age-specific incidence 1998 - 2013 (n=3475)





Figure 7. Age distribution and age-specific incidence





**Figure 7a.** Age-specific incidence in MCR registry areas compared to Germany (FRG, GEKID extrapolation) and SEER (Surveillance, Epidemiology, and End Results, USA).



#### Reference:

Extrapolated age-specific patient population of Germany, data status middle of 2010. Association of Population-based Cancer Registries in Germany (GEKID e.V.). Berlin, 2014. http://www.gekid.de. Last access: 02/11/2015

Surveillance, Epidemiology, and End Results (SEER) Program SEER\*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2014, based on the November 2013 submission. http://www.seer.cancer.gov.





Figure 8. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.




## Average incidence (world standard population) 2007 - 2013




**Figure 9a.** Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2007 to 2013. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (6.9/100,000 WS N=1,724).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,928 female residents (averaged) in the period from 2007 to 2013 a total of 56 women were identified with newly diagnosed cervical cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 7.7/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 5.2 and 11.0/100,000.



## Standardized incidence ratio (SIR) 2007 - 2013



**Figure 9b.** Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2013. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (N=1,724).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,642 female residents (averaged) in the period from 2007 to 2013 a total of 56 women were identified with newly diagnosed cervical cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 1.17. Though, the value of this parameter may vary with an underlying probability of 99% between 0.81 and 1.64, and is therefore not statistically striking.



## **MORTALITY**

#### Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts, and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

	Incident	Prop.	Prop.		Prop.	Prop. deaths with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	96	%	n	%	%
1998	144	91.7	5.6	67	46.5	86.6
1999	161	93.8	1.2	65	40.4	90.8
2000	148	95.9	5.4	59	39.9	98.3
2001	151	93.4	4.0	64	42.4	89.1
2002	238	96.2	9.7	108	45.4	95.4
2003	218	93.6	4.6	110	50.5	96.4
2004	214	96.3	6.5	101	47.2	97.0
2005	235	93.2	5.1	97	41.3	94.8
2006	242	86.8	3.3	94	38.8	100.0
2007	243	79.0	3.3	93	38.3	95.7
2008	271	59.4	2.2	98	36.2	98.0
2009	267	57.3	3.7	102	38.2	96.1
2010	251	61.4	4.8	82	32.7	98.8
2011	245	58.8	3.3	69	28.2	94.2
2012	243	66.3	7.8	77	31.7	100.0
2013	206	97.6	5.3	35	17.0	88.6
1998-2013	3477	80.5	4.7	1321	38.0	95.5

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

			Prop. deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	n	%	n	%
1998	144	92	85.9	14	9.7
1999	161	93	87.1	14	8.7
2000	148	91	92.3	15	10.1
2001	151	68	89.7	13	8.6
2002	238	132	93.9	36	15.1
2003	218	152	94.7	27	12.4
2004	214	153	96.7	26	12.1
2005	235	151	96.0	25	10.6
2006	242	143	95.1	19	7.9
2007	243	146	94.5	28	11.5
2008	271	165	98.8	24	8.9
2009	267	169	98.8	26	9.7
2010	251	174	98.9	29	11.6
2011	245	173	98.8	28	11.4
2012	243	150	96.7	40	16.5
2013	206	168	98.2	25	12.1
1998-2013	3477	2220	95.6	389	11.2

Table 10c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

				Prop.
				cancer
		Prop.	Prop.	recorded
		cancer-	non-cancer-	on death
Year of	Deaths	related	related	certificate
death	n	%	8	%
1998	92	58.7	41.3	83.5
1999	93	68.8	31.2	82.7
2000	91	67.0	33.0	83.3
2001	68	64.7	35.3	86.9
2002	132	71.2	28.8	83.1
2003	152	75.0	25.0	85.4
2004	153	67.3	32.7	76.4
2005	151	72.8	27.2	83.4
2006	143	64.3	35.7	79.4
2007	146	74.0	26.0	78.3
2008	165	70.3	29.7	76.1
2009	169	65.7	34.3	76.0
2010	174	74.1	25.9	81.4
2011	173	69.9	30.1	74.9
2012	150	67.3	32.7	80.0
2013	168	70.2	29.8	78.2
1998-2013	2220	69.4	30.6	79.9

		Age at death (all	Age at death (cancer-	Age at death (non-cancer-	Age at death (according to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	92	74.8	70.4	80.8	74.6
1999	93	74.8	71.5	79.6	75.1
2000	91	78.4	74.9	83.8	77.0
2001	68	74.6	69.7	81.4	71.5
2002	132	74.6	65.2	87.6	69.4
2003	152	73.1	66.6	81.9	72.3
2004	153	74.9	64.6	85.0	69.9
2005	151	76.9	68.4	84.0	70.0
2006	143	74.6	68.9	84.0	72.6
2007	146	71.8	68.5	83.4	69.5
2008	165	70.2	65.8	85.7	67.0
2009	169	75.1	66.0	86.0	68.0
2010	174	69.6	63.5	80.9	65.7
2011	173	74.1	68.4	84.9	69.6
2012	150	73.3	69.0	85.1	69.3
2013	168	70.6	62.2	82.6	64.5
1998-2013	2220	73.1	67.4	84.0	69.9

By 2010, life expectancy for a newborn male in Germany is 77.5 years compared with 82.6 years for his female counterpart.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 12

Mortality measures (cancer-related death) and mortality-incidence-index by year of death

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	54	4.6	0.38	2.2	0.28	3.2	0.31	4.0	0.36
1999	64	5.4	0.40	2.5	0.28	3.7	0.32	4.7	0.38
2000	61	5.1	0.41	2.3	0.28	3.3	0.32	4.2	0.37
2001	44	3.6	0.29	1.7	0.22	2.4	0.24	3.0	0.27
2002	94	4.8	0.39	2.5	0.32	3.5	0.36	4.2	0.39
2003	114	5.8	0.52	2.9	0.42	4.1	0.45	4.9	0.50
2004	103	5.2	0.48	2.7	0.39	3.8	0.43	4.6	0.48
2005	110	5.5	0.47	2.6	0.34	3.7	0.39	4.5	0.43
2006	92	4.6	0.38	2.1	0.27	3.0	0.31	3.7	0.35
2007	108	4.7	0.44	2.3	0.34	3.2	0.37	3.8	0.42
2008	116	5.0	0.43	2.5	0.33	3.5	0.36	4.0	0.39
2009	111	4.8	0.42	2.4	0.32	3.3	0.35	3.9	0.38
2010	129	5.5	0.51	2.9	0.42	4.0	0.45	4.5	0.47
2011	121	5.1	0.49	2.4	0.35	3.5	0.39	4.1	0.44
2012	101	4.3	0.42	2.0	0.32	2.9	0.35	3.5	0.39
2013	118	5.0	0.57	2.7	0.46	3.7	0.50	4.3	0.54
1998-2013	1540	5.0	0.44	2.5	0.34	3.5	0.37	4.1	0.41

Table 13

Age distribution of age at death (cancer-related) for period 1998-2013 (incl. multiple primaries)

Age at				
death	Cases			
Years	'n	%	Cum.%	
20-24	/ 1	0.1	0.1	
25-29	/ 5	0.3	0.4	
30-34	17	1.1	1.5	
35-39	51	3.3	4.8	
40-44	93	6.0	10.8	
45-49	112	7.3	18.1	
50-54	119	7.7	25.8	
55-59	165	10.7	36.5	
60-64	140	9.1	45.6	
65-69	162	10.5	56.1	
70-74	156	10.1	66.2	
75-79	149	9.7	75.9	
80-84	191	12.4	88.3	
85+	181	11.7	100.0	
All ages	1542	100.0		

Included in the statistics are 19.5% multiple primaries.

Table 14

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2013 (incl. multiple primaries)

Age at				Prop. all
death	Cases	Age-spec.	\ . \	cancers
Years	n	mortality	MI-index	%
0 - 4		0.0		
5- 9		0.0		
10-14		0.0		
15-19		0.0		
20-24	1	0.1	0.08	2.0
25-29	5	0.2	0.05	4.3
30-34	17	0.8	0.07	7.5
35-39	51	2.2	0.13	9.9
40-44	93	3.7	0.19	8.2
45-49	112	4.8	0.29	5.6
50-54	119	5.8	0.37	3.8
55-59	165	8.6	0.54	3.5
60-64	140	7.5	0.54	2.1
65-69	162	9.4	0.72	1.9
70-74	156	10.3	0.79	1.6
75-79	149	12.5	0.94	1.4
80-84	191	20.5	1.10	1.7
85+	181	20.3	0.96	1.3
All ages	1542			2.1
Mortality			/	
Raw		5.0	0.44	
WS		2.5	0.34	
ES		3.5	0.37	
BRD-S		4.1	0.41	
PYLL-70				
per 100,000		49.9		
ES		42.9		
AYLL-70		15.3		

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15

Multiple primaries in deaths in period 1998-2013

					Syn- chron	Syn- chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n	% ↓	n	-%	n	<b>←</b> %	n	-%
C03-C06 Oral cavity	/ 9	1.2	1	11.1			8	88.9
C16 Stomach	/16	2.2	1	6.3	1	6.3	14	87.5
C18 Colon	53	7.3	9	17.0	5	9.4	39	73.6
C19-C20 Rectum	43	5.9	8	18.6	1	2.3	34	79.1
C21 Anus/canal	16	2.2	2	12.5			14	87.5
C25 Pancreas	21	2.9	1	4.8	/ 1	4.8	19	90.5
C33-C34 Lung	91	12.6	9	9.9	7	7.7	75	82.4
C43 Malign. melanoma	20	2.8	7	35.0	1	5.0	12	60.0
C44 Skin others	9	1.2	2	22.2			7	77.8
C50 Breast	117	16.1	33	28.2	10	8.5	74	63.2
C51 Vulva	19	2.6	4	21.1	5	26.3	10	52.6
C52 Vagina	18	2.5			5	27.8	13	72.2
C53 Cervix uteri	10	1.4			1	10.0	9	90.0
C54 Corpus uteri	33	4.6	6	18.2	_ 5	15.2	22	66.7
C56 Ovary	61	8.4	6	9.8	12	19.7	43	70.5
C64 Kidney	14	1.9	5	35.7	1	7.1	8	57.1
C65 Renal pelvis	8	1.1	2	25.0			6	75.0
C67 Bladder	54	7.4	5	9.3	5	9.3	44	81.5
C70-C72 CNS cancer	12	1.7	2	16.7	1	8.3	9	75.0
C76-C79 CUP	12	1.7	2	16.7	1 \	8.3	9	75.0
C82-C85 NHL	15	2.1	4	26.7	1	6.7	10	66.7
C91-C96 Leukaemia	12	1.7			2	16.7	10	83.3
Other primaries	62	8.6	13	21.0			49	79.0
All mult. primaries	725	100.0	122	16.8	65	9.0	538	74.2

Multiple primaries with number of cases 1 to 7 are pooled in category "Other primaries".

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2013

(Singular primaries only \*)

Age at				Prop. all	
death	Cases	Age-spec.		cancers	
Years	n/	mortality	MI-index	%	
0 - 4		0.0			
5- 9		0.0			
10-14		0.0			
15-19		0.0			
20-24	1	0.1	0.08	2.1	
25-29	5	0.2	0.05	4.6	
30-34	15	0.7	0.06	7.5	
35-39	49	2.1	0.12	10.5	
40-44	89	3.6	0.19	9.0	
45-49	98	4.2	0.28	5.7	
50-54	107	5.2	0.36	4.1	
55-59	149	7.7	0.55	3.7	
60-64	129	6.9	0.56	2.4	
65-69	129	7.5	0.71	1.9	
70-74	132	8.7	0.86	1.7	
75-79	132	11.1	1.02	1.5	
80-84	171	18.3	1.22	1.9	
85+	156	17.5	1.07	1.4	
All ages	1362			2.3	
Mortality					
Raw		4.4	0.44		
WS		2.2	0.33		
ES		3.1	0.37		
BRD-S		3.6	0.41		
PYLL-70					
per 100,000		45.6			
ES		39.2			
AYLL-70		15.7			
-					

<sup>\*</sup> See corresponding tables with multiple primaries.

Table 17


Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2013

(Single primaries only \*)

Age at				Prop. all
death	Cases	Age-spec.		cancers
Years	n	mortality	MI-index	%
iears	11/	mortality	MI-Index	***
0- 4		0.0		
5- 9		0.0		
10-14		0.0		
15-19		0.0		
20-24	1	0.1	0.08	2.3
25-29	5	0.2	0.05	4.9
30-34	15	0.7	0.06	8.2
35-39	46	1.9	0.12	10.8
40-44	82	3.3	0.18	9.0
45-49	85	3.7	0.26	5.5
50-54	86	4.2	0.32	3.7
55-59	116	6.0	0.48	3.3
60-64	92	4.9	0.45	2.0
65-69	82	4.8	0.49	1.4
70-74	72	4.7	0.55	1.1
75-79	72	6.1	0.61	1.0
80-84	100	10.7	0.77	1.3
85+	106	11.9	0.79	1.1
All ages	960			1.9
Mortality			/	
Raw		3.1	0.33	
WS		1.7	0.27	
ES		2.3	0.29	
BRD-S		2.6	0.31	
PYLL-70				
per 100,000		39.0		
ES		33.6		
AYLL-70		16.9		
		10.7		

<sup>\*</sup> See corresponding tables with multiple primaries.

base\_C53\_\_E.pdf




**Figure 18.** Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at cervical cancer-related death (see Table 10) should be considered.




## Average mortality (world standard population) 2007 - 2013



**Figure 19a.** Map of cancer mortality (world standard population) by county averaged for period 2007 to 2013. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (2.5/100,000 WS N=795).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,928 female residents (averaged) in the period from 2007 to 2013 a total of 26 women died from cervical cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 3.0/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 1.7 and 5.2/100,000.

## Standardized mortality ratio (SMR) 2007 - 2013



**Figure 19b.** Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2013. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (N=795).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,642 female residents (averaged) in the period from 2007 to 2013 a total of 26 women died from cervical cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 1.21. Though, the value of this parameter may vary with an underlying probability of 99% between 0.68 and 1.96, and is therefore not statistically striking.

#### Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

## 1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

#### 2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

# 3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

#### **Shortcuts**

FRG Federal Republic of Germany

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

MCR Munich Cancer Registry (Tumorregister München)
SEER Surveillance, Epidemiology, and End Results (USA)

AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

BRD-S German standard population

DCO Death certificate only EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

ES European standard population (old)

LCL Lower confidence limit

MI-index Ratio between mortality and incidence

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age

SIR Standardized incidence ratio
SMR Standardized mortality ratio
UCL Upper confidence limit
WS World standard population

#### **Recommended Citation**

Munich Cancer Registry. Baseline statistics C53: Cervical cancer [Internet]. 2015 [updated 2015 May 19; cited 2015 Jul 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/base/base C53 E.pdf

#### Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

#### Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

# Index of figures and tables

Fig./Tbl	l.	Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	3
2	Incidence by year of diagnosis	4
3	Age distribution parameters by year of diagnosis	5
4	Age distribution by 5-year age group	6
5	Age-specific incidence and DCO rate	7
6	Standardized incidence ratio of second primaries	8
7	Age distribution and age-specific incidence (chart)	9
7a	Age-specific incidence internationally (chart)	10
8	Cumulative follow-up years (chart)	11
9a	Map of cancer incidence (WS) by county (chart)	12
9b	Standardized incidence ratio (SIR) by county (chart)	13
10a	Pts incident cohorts and mortality / yr	14
10b	Incidence and mortality by year of diagnosis	15
10c	Cancer-related deaths, death certification available / yr	16
11	Medians of age at death / yr	17
12	Mortality by year of death	18
13	Distribution of age at death	19
14	Age-specific mortality	20
15	Multiple primaries in deaths	21
16	Age-specific mortality (first primaries)	22
17	Age-specific mortality (single primaries)	23
18	Age distribution and age-specific mortality (chart)	24
19a	Map of cancer mortality (WS) by county (chart)	25
19b	Standardized mortality ratio (SMR) by county (chart)	26