Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ Deutsch

Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

Cancer statistics: Baseline statistics

C70-C72: Brain/nerves cancer

Year of diagnosis	1998-2013
Patients	5,017
Diseases	5,023
Creation date	05/19/2015
Export date	12/30/2014
Population	4.64 m



http://www.tumorregister-muenchen.de/en/facts/base/base_C7072E.pdf

Global Statements about the statistics on the Internet – Baseline Statistics (grey button ——), Survival (red button ——)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.64 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases**** are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR. The time-delayed acquisition of data and the occasionally high DCO-rates indicate optimizing reserves, among others, because of current financial and legal conditions that hinder the analyses.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, May 2015

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007). Death certificates from 2014 are incorporated into these analyses.
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- """ DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate. A high proportion of DCO cases (≥5%) in particular cancer types indicate insufficient participation of specific cancer specializations.

base_C7072E.pdf

ICD-10 codes (ICD-10 2015) used for specifying cancer site

Code	Description
C70 C70.0 C70.1 C70.9	Malignant neoplasm of meninges Cerebral meninges Spinal meninges Meninges, unspecified
C71 C71.0 C71.1 C71.2 C71.3 C71.4 C71.5 C71.6 C71.7 C71.8	Malignant neoplasm of brain Cerebrum, except lobes and ventricles Frontal lobe Temporal lobe Parietal lobe Occipital lobe Cerebral ventricle Cerebellum Brain stem Overlapping lesion of brain Brain, unspecified
C72	Malignant neoplasm of spinal cord, cranial nerves and other parts of central
C72.0 C72.1 C72.2 C72.3 C72.4 C72.5 C72.8 C72.9	nervous system Spinal cord Cauda equina Olfactory nerve Optic nerve Acoustic nerve Other and unspecified cranial nerves Overlapping lesion of brain and other parts of central nervous system Central nervous system, unspecified

INCIDENCE

Table 1

Patient cohorts by year of diagnosis including DCO cases and multiple primaries, and with proportion of deaths and active follow-up

				Prop.		Prop.
		DCO	Prop.	mult.	Prop.	actively
Year of	Cases	cases	DCO	primaries	deaths	followed
diagnosis	n	n	%	8	%	%
1998	188	48	25.5	10.1	86.7	98.9
1999	176	50	28.4	10.2	89.2	98.3
2000	198	50	25.3	10.6	81.8	98.0
2001	235	55	23.4	9.4	84.3	96.6
2002	340	63	18.5	12.1	82.4	98.8 #
2003	378	72	19.0	11.4	79.9	99.2
2004	337	73	21.7	13.1	80.4	97.6
2005	374	62	16.6	13.9	79.7	97.1
2006	299	40	13.4	12.4	80.9	96.0
2007	348	48	13.8	14.1	72.7	90.5 # ##
2008	403	54	13.4	13.9	75.2	83.9
2009	439	53	12.1	12.5	78.6	87.7
2010	378	51	13.5	15.3	80.7	89.7
2011	396	47	11.9	15.2	68.9	83.8
2012	372	44	11.8	15.1	58.9	82.8
2013	162	37	22.8	20.4	61.7	100.0 ###
1998-2013	5023	847	16.9	13.2	77.1	92.6

[#] The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

Please be aware that data of recent annual patient cohorts may not yet be fully processed. Therefore, the presented figures and tables are potentially related to different time periods as pointed out in the respective headlines or legends.

^{##} Since 2007 the percentage of actively followed patients sharply declined compared to the previous years. This is a consequence of ambiguous data protection rules that currently forbid cancer registries in Bavaria to obtain the essential life status informations from competent registration offices.

Table 1a

Patient cohorts by year of diagnosis and gender including DCO cases

Year of	All	Males	Females	Prop. males
diagnosis	n /	'n	n	%
1998	188	97	91	51.6
1999	176	88	88	50.0
2000	198	103	95	52.0
2001	235	116	119	49.4
2002	340	175	165	51.5
2003	378	197	181	52.1
2004	337	168	169	49.9
2005	374	198	176	52.9
2006	299	170	129	56.9
2007	348	186	162	53.4
2008	403	228	175	56.6
2009	439	253	186	57.6
2010	378	212	166	56.1
2011	396	210	186	53.0
2012	372	198	174	53.2
2013	162	99	63	61.1
1998-2013	5023	2698	2325	53.7

Table 2

Incidence measures by year of diagnosis and gender including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

			Males	Fem.	Males	Fem.	Males	Fem.	Males	Fem.
Year of	Males	Females	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	97	91	8.8	7.7	6.2	5.3	7.9	6.4	8.9	7.3
1999	88	88	7.9	7.4	5.4	4.5	7.2	5.8	8.4	6.6
2000	103	95	9.0	7.9	6.5	4.8	8.3	6.1	9.8	7.0
2001	116	119	10.0	9.8	6.8	5.8	9.1	7.5	10.6	8.7
2002	175	165	9.4	8.4	6.7	4.8	8.3	6.3	9.4	7.6
2003	197	181	10.5	9.2	7.3	5.7	9.3	7.2	10.8	8.3
2004	168	169	8.9	8.5	5.7	5.6	7.7	6.6	9.0	7.4
2005	198	176	10.5	8.8	7.0	5.4	8.8	6.7	10.2	7.6
2006	170	129	8.9	6.4	5.9	3.7	7.5	4.7	8.5	5.5
2007	186	162	8.4	7.0	5.8	4.3	7.2	5.4	8.0	6.3
2008	228	175	10.2	7.5	6.8	4.8	8.7	5.8	10.2	6.5
2009	253	186	11.3	8.0	7.0	4.8	9.3	6.1	11.0	6.9
2010	212	166	9.4	7.1	6.0	4.1	7.7	5.1	9.0	5.9
2011	210	186	9.2	7.9	5.8	4.3	7.5	5.7	8.7	6.8
2012	198	174	8.7	7.4	5.6	4.5	7.0	5.6	8.2	6.3
2013	99	63	4.3	2.7	2.6	1.4	3.5	1.8	4.1	2.2
1998-2013	2698	2325	9.1	7.5	6.0	4.5	7.7	5.7	8.9	6.5




Table 3

Age distribution parameters by year of diagnosis (All) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	188	57.0	18,7	1.8	90.6	32.4	46.7	57.8	71.5	79.8
1999	176	59.9	17.8	1.6	93.4	36.5	50.6	61.6	72.6	79.4
2000	198	59.2	18.5	2.9	93.8	33.7	48.5	62.3	71.7	79.7
2001	235	60.3	18.0	1.0	92.0	37.0	50.8	62.1	73.3	80.4
2002	340	59.9	18.4	0.8	91.2	32.6	50.7	63.0	72.9	80.9
2003	378	58.5	18.4	0.6	95.4	31.7	45.9	62.0	72.9	79.7
2004	337	60.0	18.9	0.0	92.8	34.9	49.9	63.1	73.7	81.9
2005	374	59.6	19.6	0.8	94.3	32.8	47.5	64.3	73.2	81.3
2006	299	60.5	18.6	3.0	97.0	34.7	48.3	64.2	73.2	81.6
2007	348	58.6	19.6	0.8	93.5	31.1	44.2	62.7	73.3	80.9
2008	403	59.5	19.9	0.1	94.1	31.0	48.0	64.2	73.8	80.8
2009	439	61.7	18.0	0.2	94.2	36.5	52.4	65.1	74.7	82.5
2010	378	61.7	19.5	0.1	91.6	37.0	51.7	67.1	75.1	82.6
2011	396	60.5	18.4	1.7	94.0	37.1	47.9	63.3	75.5	81.7
2012	372	60.8	19.9	0.0	96.0	35.0	49.8	65.4	74.1	82.9
2013	162	64.3	18.5	2.0	93.9	38.4	56.3	69.1	76.2	84.3
1998-2013	5023	60.1	18.9	0.0	97.0	34.2	49.3	63.8	73.8	81.3

Table 3a

Age distribution parameters by year of diagnosis (MALES)

(incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	97	54.7	15.7	3.4	83.3	32.6	46.3	55.8	65.5	74.6
1999	88	59.0	16.3	1.6	85.5	38.9	51.1	60.4	71.2	77.6
2000	103	56.4	18.2	4.1	88.2	30.0	46.6	59.4	68.6	77.6
2001	116	58.6	17.3	1.0	91.2	37.0	50.7	60.2	71.5	77.4
2002	175	56.2	18.2	0.8	87.7	30.9	45.4	61.7	68.8	75.9
2003	197	56.8	18.8	6.2	89.4	28.3	44.3	60.8	71.8	78.1
2004	168	59.6	16.2	0.1	90.1	37.4	48.6	61.0	70.4	81.9
2005	198	58.6	19.5	0.8	94.3	32.7	47.5	63.6	72.1	80.3
2006	170	58.1	17.3	3.0	90.4	34.5	46.7	62.4	69.7	77.2
2007	186	57.2	18.9	1.5	92.6	32.5	44.5	60.6	69.9	79.2
2008	228	58.6	18.5	1.2	94.1	31.9	47.6	63.4	72.9	78.7
2009	253	61.0	17.8	5.0	90.3	35.5	51.0	64.8	74.2	81.9
2010	212	60.2	19.1	0.1	90.8	34.4	49.2	64.2	74.0	81.0
2011	210	58.5	18.4	1.7	91.9	35.9	44.5	59.5	73.7	80.5
2012	198	60.4	20.2	0.3	96.0	33.5	48.6	66.0	73.9	81.3
2013	99	62.9	18.3	2.0	93.9	37.6	54.6	68.6	75.5	79.3
1998-2013	2698	58.7	18.3	0.1	96.0	33.5	47.7	62.0	71.9	79.3

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	91	59.5	21,3	1.8	90.6	30.9	47.5	62.9	77.2	83.4
1999	88	60.7	19.3	4.7	93.4	33.0	49.6	64.1	75.9	84.0
2000	95	62.1	18.4	2.9	93.8	35.8	55.5	67.6	74.2	80.6
2001	119	62.0	18.5	2.3	92.0	36.1	51.0	64.5	76.0	81.8
2002	165	63.9	17.8	2.6	91.2	37.8	54.3	67.2	78.1	83.0
2003	181	60.3	17.9	0.6	95.4	36.6	49.6	62.9	74.1	80.3
2004	169	60.3	21.2	0.0	92.8	30.5	51.8	65.1	76.4	82.6
2005	176	60.6	19.6	2.7	91.7	32.8	47.6	65.0	75.3	83.4
2006	129	63.6	19.7	7.3	97.0	35.0	54.1	67.4	77.7	86.1
2007	162	60.3	20.3	0.8	93.5	30.7	44.1	65.5	77.7	82.3
2008	175	60.6	21.6	0.1	92.9	29.1	48.7	65.9	77.1	85.1
2009	186	62.7	18.1	0.2	94.2	38.0	53.3	65.6	75.8	83.6
2010	166	63.7	20.0	0.6	91.6	41.8	56.6	68.9	76.6	83.5
2011	186	62.9	18.0	11.1	94.0	37.7	50.0	66.8	77.0	82.3
2012	174	61.3	19.5	0.0	90.4	35.0	50.4	65.0	74.3	83.2
2013	63	66.6	18.9	2.7	92.8	38.4	60.3	69.6	77.4	87.6
1998-2013	2325	61.8	19.4	0.0	97.0	34.8	51.3	65.9	76.3	83.2

Table 4

Age distribution by 5-year age group and gender for period 1998-2013 (incl. DCO)

Age at									
diagnosis	Cases			Males			Females		
Years	n	%	Cum.%	n	%	Cum.%	n	%	Cum.%
0-4	71	1.4	1.4	33	1.2	1.2	38	1.6	1.6
5-9	50	1.0	2.4	29	1.1	2.3	21	0.9	2.5
10-14	40	0.8	3.2	19	0.7	3.0	21	0.9	3.4
15-19	41	0.8	4.0	24	0.9	3.9	17	0.7	4.2
20-24	65	1.3	5.3	40	1.5	5.4	25	1.1	5.2
25-29	108	2.2	7.5	61	2.3	7.6	47	2.0	7.3
30-34	163	3.2	10.7	91	3.4	11.0	72	3.1	10.4
35-39	202	4.0	14.7	116	4.3	15.3	86	3.7	14.1
40 - 44	263	5.2	20.0	172	6.4	21.7	91	3.9	18.0
45-49	298	5.9	25.9	172	6.4	28.1	126	5.4	23.4
50-54	363	7.2	33.1	209	7.7	35.8	154	6.6	30.0
55-59	452	9.0	42.1	265	9.8	45.6	187	8.0	38.1
60-64	525	10.5	52.6	291	10.8	56.4	234	10.1	48.1
65-69	663	13.2	65.8	377	14.0	70.4	286	12.3	60.4
70-74	571	11.4	77.1	297	11.0	81.4	274	11.8	72.2
75-79	522	10.4	87.5	262	9.7	91.1	260	11.2	83.4
80-84	381	7.6	95.1	159	5.9	97.0	222	9.5	92.9
85+	245	4.9	100.0	81	3.0	100.0	164	7.1	100.0
All ages	5023	100.0		2698	100.0		2325	100.0	

Included in the statistics are 15.4% multiple primaries in males and 14.9% in females.

Table 5

Age-specific incidence, DCO rate and proportion of all cancers for period 1998-2013

							Males	Females
			Males	Females		Females	_	Prop.all
Age at			Age-	Age-	DCO rate	DCO rate	cancers	cancers
diagnosis	Males	Females	spec.	spec.	n=372	n=475	n=158258	n=153136
Years	n	n	incid.	incid.	%	%	%	%
0- 4	33	38	2.2	2.7		13.2	10.2	15.6
5- 9	29	21	1.9	1.5	6.9		16.5	16.8
10-14	19	21	1.2	1.4	10.5	4.8	11.4	12.4
15-19	24	17	1.6	1.2	4.2	5.9	6.8	5.8
20-24	40	25	2.3	1.4	2.5		6.5	4.7
25-29	61	47	3.0	2.3	3.3		6.3	4.2
30-34	91	72	4.0	3.2	3.3	4.2	6.1	3.5
35-39	116	86	4.7	3.6	7.8	5.8	5.2	2.3
40-44	172	91	6.6	3.7	4.1	2.2	5.4	1.5
45-49	172	126	7.3	5.5	2.9	2.4	3.2	1.4
50-54	209	154	10.4	7.5	8.6	7.1	2.4	1.4
55-59	265	187	14.4	9.7	10.9	9.1	1.8	1.4
60-64	290	234	16.4	12.5	5.5	11.1	1.3	1.4
65-69	377	286	23.9	16.6	10.9	10.5	1.4	1.5
70-74	296	273	23.1	18.0	16.6	18.7	1.1	1.5
75-79	262	260	31.7	21.9	29.8	30.4	1.3	1.5
80-84	159	222	31.8	23.8	37.1	55.4	1.2	1.4
85+	81	164	23.8	18.3	61.7	72.0	0.8	1.0
All ages	2696	2324			13.8	20.4	1.7	1.5
Incidence								
Raw			9.1	7.5				
WS			6.0	4.5				
ES			7.7	5.7				
BRD-S			8.9	6.5				
21.2 2			3.5					


The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

Table 6a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2013

ΙVΙ		

Diagnosis	Observed h	Expected n	SIR	LCL 95%	UCL 95%	EAR	DCO %
C16 Stomach	2	1.0	1.9	0.2	6.9	2.7	
C18 Colon	2 /	2.5	0.8	0.1	2.9	-1.3	
C25 Pancreas	/ 2/	0.9	2.2	0.3	7.9	3.0	
C30-C31 Sinuses	2	0.1	35.7	4.3	128.9 #	5.4	
C33-C34 Lung	3	3.3	0.9	0.2	2.6	-0.9	
C43 Malign. melanoma	4	1.3	3.0	0.8	7.7	7.4	25.0
C61 Prostate	10	8.0	1.2	0.6	2.3	5.5	20.0
C62 Testis	3	0.4	8.2	1.7	23.9 #	7.3	
C64 Kidney	3	1.1	2.8	0.6	8.1	5.3	
C70-C72 CNS cancer	3	0.5	6.4	1.3	18.8 #	7.0	
C76-C79 CUP	2	0.4	4.5	0.5	16.3	4.3	
Other primaries	6	2.6	2.3	0.8	5.0	9.4	33.3
Not observed	0	6.5	0.0	0.0	0.6 #	-18.0	
All mult. primaries	42	28.6	1.5	1.1	2.0 #	36.9	11.9
_ \ \ \ \					\ "		



The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries".

12.3 12.5

-9.6

Other primaries

Not observed

Table 6b

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2013

FEMALES

		Observ	ed Expected		LCL	UCL		DCO
Diagnos	is	n	n	SIR	95%	95%	EAR	%
C18	Colon	5	1.7	2.9	0.9	6.8	10.3	20.0
C19-C20	Rectum	3	0.8	3.6	0.7	10.6	6.8	33.3
C43	Malign. melanoma	2	0.9	2.2	0.3	7.9	3.4	
C50	Breast	12	7.3	/1.7	0.9	2.9	15.0	8.3
C54	Corpus uteri	3	1.2	2.6	0.5	7.5	5.8	
C56	Ovary	2	0.9	2.3	0.3	8.3	3.6	
C82-C85	NHL	3	0.7	4.1	0.8	12.0	7.2	33.3
C90	Mult. myeloma	2	0.2	9.1	1.1	32.8 ‡	‡ 5.6	
C91-C96	Leukaemia	2.	0.3	6.6	0.8	23.9	5.4	

4.1


3.0

1.9

0.0

0.8 3.8

0.0 1.2



8

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries".

C70-C72: Malignant neoplasm of brain and nerves Age distribution and age-specific incidence 1998 - 2013 (Males: 2696, Females: 2324)




Figure 7. Age distribution and age-specific incidence



C70-C72: Malignant neoplasm of brain and nerves




Figure 7a. Age-specific incidence in MCR registry areas compared to Germany (FRG, GEKID extrapolation) and SEER (Surveillance, Epidemiology, and End Results, USA).



Reference:

Extrapolated age-specific patient population of Germany, data status middle of 2010. Association of Population-based Cancer Registries in Germany (GEKID e.V.). Berlin, 2014. http://www.gekid.de. Last access: 02/11/2015

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2014, based on the November 2013 submission. http://www.seer.cancer.gov.





Figure 8. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.



Average incidence (world standard population) 2007 - 2013: Males



Average incidence (world standard population) 2007 - 2013: Females





Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2007 to 2013. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 5.7/100,000 WS N=1,384, females 4.1/100,000 WS N=1,111).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,928 female residents (averaged) in the period from 2007 to 2013 a total of 39 women were identified with newly diagnosed brain/nerves cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 5.4/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 3.1 and 9.6/100,000.

Standardized incidence ratio (SIR) 2007 - 2013: Males



Standardized incidence ratio (SIR) 2007 - 2013: Females




Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2013. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=1,384, females N=1,111).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,642 female residents (averaged) in the period from 2007 to 2013 a total of 39 women were identified with newly diagnosed brain/nerves cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 1.30. Though, the value of this parameter may vary with an underlying probability of 99% between 0.83 and 1.94, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts, and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

	Incident	Prop. actively	Prop.		Prop.	Prop. deaths with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	%	olo	n	%	%
1998	188	98.9	25.5	163	86.7	96.3
1999	176	98.3	28.4	157	89.2	94.9
2000	198	98.0	25.3	162	81.8	93.8
2001	235	96.6	23.4	198	84.3	92.9
2002	340	98.8	18.5	280	82.4	98.6
2003	378	99.2	19.0	302	79.9	95.7
2004	337	97.6	21.7	271	80.4	97.8
2005	374	97.1	16.6	298	79.7	97.3
2006	299	96.0	13.4	242	80.9	97.9
2007	348	90.5	13.8	253	72.7	97.6
2008	403	83.9	13.4	303	75.2	97.7
2009	439	87.7	12.1	345	78.6	98.0
2010	378	89.7	13.5	305	80.7	98.4
2011	396	83.8	11.9	273	68.9	97.4
2012	372	82.8	11.8	219	58.9	96.8
2013	162	100.0	22.8	100	61.7	94.0
1998-2013	5023	92.6	16.9	3871	77.1	96.9

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

			Prop.		D
			deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	'n	8	n	96
1998	188	136	95.6	72	38.3
1999	176	172	97.1	83	47.2
2000	198	156	94.9	75	37.9
2001	235	192	92.2	103	43.8
2002	340	233	97.0	124	36.5
2003	378	253	95.7	127	33.6
2004	337	273	97.4	122	36.2
2005	374	256	97.3	138	36.9
2006	299	267	96.3	108	36.1
2007	348	256	98.0	121	34.8
2008	403	265	97.4	127	31.5
2009	439	310	98.4	142	32.3
2010	378	357	98.9	147	38.9
2011	396	319	98.7	123	31.1
2012	372	300	97.3	126	33.9
2013	162	250	98.8	78	48.1
1998-2013	5023	3995	97.2	1816	36.2

Table 10c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

				Prop.
				cancer
		Prop.	Prop.	recorded
		cancer-	non-cancer-	on death
Year of	Deaths	related	related	certificate
death	n	%	8	%
1998	136	71.3	28.7	100.0
1999	172	79.1	20.9	97.6
2000	156	78.8	21.2	98.0
2001	192	83.3	16.7	98.9
2002	233	91.4	8.6	99.6
2003	253	93.3	6.7	98.8
2004	273	93.0	7.0	98.9
2005	256	91.0	9.0	98.0
2006	267	91.0	9.0	96.9
2007	256	93.4	6.6	98.0
2008	265	94.7	5.3	98.4
2009	310	91.3	8.7	96.7
2010	357	93.6	6.4	97.5
2011	319	93.1	6.9	96.8
2012	300	92.7	7.3	98.3
2013	250	94.0	6.0	98.4
1998-2013	3995	90.4	9.6	98.0

Table 11a $\begin{tabular}{ll} Medians of age at death according to the grouping in Table 10 \\ \hline MALES \end{tabular}$

Year of death	Deaths n	Age at death (all causes)	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate) Years
1998	66	57.8	57.6	61.6	57.8
1999	91	62.1	61.4	63.5	62.1
2000	69	63.5	63.4	71.2	65.9
2001	104	64.1	63.4	67.8	64.9
2002	131	63.0	63.5	57.9	63.3
2003	123	66.3	66.6	56.0	66.8
2004	142	65.8	65.5	69.0	65.5
2005	147	65.3	64.0	73.3	64.2
2006	136	64.1	64.2	63.7	64.2
2007	137	66.3	66.3	65.0	66.4
2008	146	64.8	64.3	72.1	64.6
2009	185	68.7	66.6	71.6	66.4
2010	223	68.6	68.4	69.6	68.5
2011	169	67.4	67.1	72.6	67.1
2012	163	67.4	67.4	68.7	67.4
2013	140	66.5	66.8	61.8	66.5
1998-2013	2172	65.7	65.6	68.0	65.7

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(non-cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	70	72.2	72.7	70.7	72.5
1999	81	68.3	64.9	78.6	69.1
2000	87	69.2	66.4	73.6	69.3
2001	88	70.3	67.8	78.7	70.3
2002	102	70.3	70.4	70.3	70.4
2003	130	67.2	66.6	75.4	67.6
2004	131	66.5	66.2	68.6	66.5
2005	109	67.8	67.7	68.4	67.8
2006	131	68.0	67.9	69.1	68.5
2007	119	69.9	69.0	80.7	69.6
2008	119	68.0	67.9	76.8	68.2
2009	125	69.2	68.9	77.4	69.2
2010	134	69.2	68.6	78.5	69.2
2011	150	70.4	69.6	71.7	70.5
2012	137	68.4	67.9	78.7	69.1
2013	110	64.2	63.2	73.4	63.2
1998-2013	1823	68.7	68.2	75.0	69.0

By 2010, life expectancy for a newborn male in Germany is 77.5 years compared with 82.6 years for his female counterpart.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 12a

Mortality measures (cancer-related death) and mortality-incidence-index by year of death

MALES

Year of	Deaths	Mort.	MI-Index	Mort. N	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	46	4.2	0.47	3.0	0.49	3.8	0.48	4.3	0.48
1999	75	6.7	0.85	4.4	0.80	6.0	0.83	7.2	0.86
2000	52	4.6	0.50	3.0	0.46	4.1	0.50	4.7	0.48
2001	86	7.4	0.74	4.4	0.65	6.5	0.72	8.2	0.78
2002	121	6.5	0.69	4.1	0.61	5.6	0.67	6.6	0.70
2003	115	6.1	0.58	3.6	0.49	5.1	0.55	6.4	0.59
2004	132	7.0	0.79	4.3	0.76	6.0	0.77	7.2	0.80
2005	133	7.0	0.67	4.3	0.62	5.8	0.66	6.9	0.68
2006	127	6.6	0.75	4.4	0.75	5.6	0.75	6.4	0.75
2007	131	5.9	0.70	3.4	0.58	4.7	0.65	5.7	0.71
2008	137	6.2	0.60	3.8	0.55	5.0	0.58	5.9	0.58
2009	163	7.3	0.64	4.2	0.60	5.8	0.62	7.1	0.65
2010	213	9.5	1.00	5.4	0.90	7.4	0.95	8.9	0.99
2011	159	7.0	0.76	3.9	0.67	5.4	0.72	6.6	0.75
2012	154	6.7	0.79	4.0	0.71	5.2	0.75	6.3	0.78
2013	131	5.7	1.32	3.4	1.31	4.5	1.31	5.3	1.30
1998-2013	1975	6.6	0.73	4.0	0.67	5.4	0.71	6.5	0.73

Table 12b

Mortality measures (cancer-related death) and mortality-incidence-index by year of death

FEMALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	51	4.3	0.56	2.1	0.39	3.0	0.47	4.0	0.54
1999	61	5.1	0.69	3.3	0.73	4.1	0.71	4.6	0.69
2000	71	5.9	0.75	3.5	0.72	4.5	0.74	5.2	0.73
2001	74	6.1	0.62	3.5	0.60	4.5	0.60	5.4	0.62
2002	92	4.7	0.56	2.4	0.50	3.3	0.52	4.1	0.54
2003	121	6.1	0.67	3.5	0.62	4.7	0.65	5.5	0.66
2004	122	6.2	0.72	3.7	0.67	4.7	0.70	5.4	0.73
2005	100	5.0	0.57	2.7	0.51	3.6	0.53	4.3	0.56
2006	116	5.8	0.90	3.1	0.84	4.1	0.87	4.9	0.89
2007	108	4.7	0.67	2.3	0.54	3.2	0.60	4.0	0.63
2008	114	4.9	0.65	2.9	0.59	3.6	0.63	4.2	0.64
2009	120	5.2	0.65	2.6	0.55	3.5	0.57	4.1	0.60
2010	121	5.2	0.73	2.7	0.65	3.6	0.70	4.4	0.74
2011	138	5.8	0.74	3.0	0.70	4.0	0.71	4.8	0.72
2012	124	5.3	0.72	2.9	0.64	3.7	0.67	4.4	0.71
2013	104	4.4	1.65	2.5	1.71	3.3	1.81	3.8	1.71
1998-2013	1637	5.3	0.70	2.9	0.64	3.8	0.67	4.5	0.69

Table 13

Age distribution of age at death (cancer-related) for period 1998-2013

(incl. multiple primaries)

Age at								
death	Cases		Males			Females		
Years	n	% Cum. %	k n	%	Cum.%	n	%	Cum.%
0 - 4	17	0.5 0.5	5 / 8	0.4	0.4	9	0.5	0.5
5-9	33	0.9 /1.4	12	0.6	1.0	21	1.3	1.8/
10-14	23	0.6 2.0) 12	0.6	1.6	11	0.7	2.5
15-19	20	0.6 / 2.6	5 9	0.5	2.1	11	0.7	3.2
20-24	20	0.6 / 3.1	12	0.6	2.7/	8	0.5	3.7
25-29	34	0.9 4.1	20	1.0	3.7	14	0.9	4.5
30-34	51	1.4 5.5	32	1.6	5,3	19	1.2	5.7
35-39	85	2.3 7.8	50	2.5	7.8	35	2.1	7.8
40-44	144	4.0 11.8	91	4.6	12.4	53	3.2	11.0
45-49	208	5.7 17.6	127	6.4	18.9	81	4.9	16.0
50-54	231	6.4 23.9	136	6.9	25.7	95	5.8	21.8
55-59	346	9.6 33.5	207	10.5	36.2	139	8.5	30.2
60-64	430	11.9 45.4	242	12.2	48.5	188	11.5	41.7
65-69	547	15.1 60.5	316	16.0	64.4	231	14.1	55.8
70-74	526	14.5 75.0	286	14.5	78.9	240	14.6	70.4
75-79	429	11.9 86.9	216	10.9	89.8	213	13.0	83.4
80-84	295	8.2 95.1	138	7.0	96.8	157	9.6	92.9
85+	179	4.9 100.0	63	3.2	100.0	116	7.1	100.0
All ages	3618	100.0	1977	100.0		1641	100.0	

Included in the statistics are 15.4% multiple primaries in males and 14.9% in females.

Table 14

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2013 (incl. multiple primaries)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males	Females	spec.		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0 - 4	8	9	0.5	0.24	0.6	0.24	24.2	34.6
5- 9	12	21 /	0.8	0.41	/1.5	1.00	31.6	52.5
10-14	12	11/	0.8	0.63	0.8	0.52	34.3	35.5
15-19	9	11	0.6	0.38	0.7	0.65	20.0	29.7
20-24	12	8	0.7	0.30	0.4	0.32	13.3	15.7
25-29	20	14	1.0		0.7	0.30	18.5	12.2
30-34	32	19	1.4		0.9		17.2	8.4
35-39	50	35	2.0	0.43	1.5	0.41	12.5	6.8
40-44	91	53	3.5	0.53	2.1	0.58	10.6	4.7
45-49	127	81	5.4	0.74	3.5	0.64	7.0	4.0
50-54	136	95	6.7		4.6	0.62	4.1	3.1
55-59	207	139	11.3		7.2	0.74	3.5	2.9
60-64	242	188	13.7	0.83	10.0	0.80	2.7	2.9
65-69	316	231	20.0	0.84	13.4	0.81	2.6	2.8
70-74	286	240	22.3	0.96	15.8	0.88	2.1	2.4
75-79	216	213	26.1	0.82	17.9	0.82	1.6	2.0
80-84	138	157	27.6	0.87	16.8	0.71	1.3	1.4
85+	63	116	18.5	0.78	13.0	0.71	0.7	0.8
All ages	1977	1641					2.5	2.3
Mortality								
Raw			6.6	0.73	5.3	0.71		
WS			4.0	0.67	2.9	0.64		
ES			5.5	0.71	3.8	0.67		
BRD-S			6.6	0.73	4.5	0.69		
PYLL-70								
per 100,000			73.3		55.1			
ES			69.0		54.5			
AYLL-70			15.4		16.0			

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15a

Multiple primaries in deaths in period 1998-2013

MALES

Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C16 Stomach	6	1.7	5	83.3			1	16.7
C18 Colon	24	6.9	20	83.3	3	12.5	1	4.2
C19-C20 Rectum	21	6.1	19	90.5	1	4.8	1	4.8
C25 Pancreas	/ 4	1.2	1	25.0	1	25.0	2	50.0
C33-C34 Lung	17	4.9	6	35.3	4	23.5	7	41.2
C43 Malign. melanoma	24	6.9	17	70.8			7	29.2
C44 Skin others	16	4.6	8	50.0	3	18.8	5	31.3
C46,C49 Soft tissue	4	1.2	2	50.0			2	50.0
C61 Prostate	101	29.2	90	89.1	5	5.0	6	5.9
C62 Testis	7	2.0	5	71.4	_ 1	14.3	1	14.3
C64 Kidney	13	3.8	7	53.8	1	7.7	5	38.5
C65 Renal pelvis	4	1.2	1	25.0			3	75.0
C67 Bladder	23	6.6	21	91.3			2	8.7
C70-C72 CNS cancer	17	4.9			2	11.8	15	88.2
C76-C79 CUP	6	1.7	2	33.3			4	66.7
C82-C85 NHL	14	4.0	12	85.7	2	14.3		
C90 Mult. myeloma	7	2.0	3	42.9	1	14.3	3	42.9
C91-C96 Leukaemia	6	1.7	2	33.3	2	33.3	2	33.3
Other primaries	32	9.2	19	59.4	3	9.4	10	31.3
All mult. primaries	346	100.0	240	69.4	29	8.4	77	22.3

Multiple primaries with number of cases 1 to 3 are pooled in category "Other primaries".

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 15b

Multiple primaries in deaths in period 1998-2013
FEMALES

						Syn- chron	Syn- chron		
		Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnos	is	n	% ↓	n	← %	n	←%	n	← %
C18	Colon	14	5.9	9	64.3	1	7.1	4	28.6
C19-C20	Rectum	6	2.5	2	33.3	2	33.3	2	33.3
C33-C34	Lung	/ 4 /	1.7	4	100.0				
C43	Malign. melanoma	17	7.1	14	82.4			3	17.6
C44	Skin others	5	2.1	1	20.0	/ 1	20.0	3	60.0
C46,C49	Soft tissue	4	1.7	2	50.0	1	25.0	1	25.0
C50	Breast	72	30.3	57	79.2	7	9.7	8	11.1
C53	Cervix uteri	8	3.4	7	87.5			1	12.5
C54	Corpus uteri	15	6.3	13	86.7			2	13.3
C56	Ovary	8	3.4	6	75.0	1	12.5	1	12.5
C64	Kidney	8	3.4	6	75.0	2	25.0		
C67	Bladder	7	2.9	4	57.1			3	42.9
C69	Eye melanoma	3	1.3	3	100.0				
C70-C72	CNS cancer	24	10.1			6	25.0	18	75.0
C73	Thyroid	6	2.5	6	100.0				
C82-C85		8	3.4	4	50.0			4	50.0
C91-C96	Leukaemia	8	3.4	4	50.0			4	50.0
Other p	rimaries	21	8.8	12	57.1	1	4.8	8	38.1
All mul	t. primaries	238	100.0	154	64.7	22	9.2	62	26.1

Multiple primaries with number of cases 1 to 2 are pooled in category "Other primaries".

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2013

(Singular primaries only *)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males	Females	spec.		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	ે	8
0 - 4	7	9	0.5	0.22	0.6	0.24	25.0	37.5
5- 9	11	20 /	0.7		1.4	0.95	30.6	54.1
10-14	12	10	0.8		0.7	0.50	34.3	34.5
15-19	8	10	0.5	0.33	0.7	0.67	19.0	30.3
20-24	12	6	0.7		0.3	0.25	14.3	12.8
25-29	19	13	0.9		0.6	0.30	19.2	11.9
30-34	31	18	1.4		0.8		17.1	9.0
35-39	50	33	2.0	0.43	1.4	0.40	13.4	7.1
40-44	86	50	3.3	0.53	2.0	0.60	10.9	5.0
45-49	121	80	5.1	0.74	3.5	0.68	7.4	4.6
50-54	132	90	6.5		4.4	0.62	4.6	3.5
55-59	192	124	10.5	0.78	6.4	0.73	3.8	3.1
60-64	223	173	12.6	0.85	9.2	0.80	3.0	3.2
65-69	274	202	17.4	0.86	11.7	0.81	2.8	3.0
70-74	234	211	18.3	0.97	13.9	0.95	2.2	2.7
75-79	165	179	20.0	0.84	15.1	0.82	1.7	2.1
80-84	111	141	22.2	0.85	15.1	0.74	1.4	1.6
85+	41	100	12.0	0.76	11.2	0.70	0.6	0.9
All ages	1729	1469					2.7	2.5
Mortality								
Raw			5.8	0.73	4.7			
WS			3.6	0.66	2.6	0.64		
ES			4.8	0.70	3.5	0.67		
BRD-S			5.7	0.72	4.1	0.69		
PYLL-70								
per 100,000			69.4		51.4			
ES			65.3		50.9			
AYLL-70			15.8		16.3			

^{*} See corresponding tables with multiple primaries.

Table 17

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2013

(Single primaries only *)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males	Females	spec.		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	8
0 - 4	6	9	0.4	0.19	0.6	0.24	22.2	37.5
5- 9	11	20 /	0.7		1.4	0.95	31.4	55.6
10-14	12	10	0.8	0.67	0.7	0.53	34.3	37.0
15-19	8	8	0.5	0.33	0.5	0.53	19.0	28.6
20-24	12	6	0.7	0.31	0.3	0.26	15.2	13.6
25-29	18	10	0.9	0.31	0.5	0.24	19.6	9.7
30-34	31	18	1.4	0.36	0.8	0.25	17.7	9.9
35-39	48	32	1.9	0.43	1.4	0.41	13.4	7.5
40-44	82	50	3.1	0.51	2.0	0.61	11.0	5.5
45-49	119	77	5.0	0.73	3.3	0.69	7.8	5.0
50-54	127	87	6.3	0.64	4.2	0.62	4.9	3.7
55-59	188	119	10.2	0.77	6.2	0.72	4.1	3.3
60-64	218	169	12.3	0.87	9.0	0.80	3.3	3.6
65-69	268	199	17.0	0.85	11.5	0.81	3.2	3.5
70-74	226	208	17.6	0.97	13.7	0.97	2.5	3.2
75-79	161	172	19.5	0.83	14.5	0.80	2.0	2.4
80-84	110	140	22.0	0.85	15.0	0.74	1.7	1.9
85+	40	98	11.7	0.74	11.0	0.70	0.7	1.0
All ages	1685	1432					3.1	2.9
Mortality								
Raw			5.7	0.72	4.6	0.71		
WS			3.5	0.65	2.6	0.64		
ES			4.7	0.69	3.4	0.67		
BRD-S			5.5	0.72	3.9	0.69		
PYLL-70								
per 100,000			67.5		49.6			
ES			63.4		49.1			
AYLL-70			15.7		16.2			

^{*} See corresponding tables with multiple primaries.





Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at brain/nerves cancer-related death (see Table 10) should be considered.



Average mortality (world standard population) 2007 - 2013: Males



Average mortality (world standard population) 2007 - 2013: Females





Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2007 to 2013. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 4.0/100,000 WS N=1,079, females 2.7/100,000 WS N=820).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,928 female residents (averaged) in the period from 2007 to 2013 a total of 28 women died from brain/nerves cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 3.2/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 1.6 and 6.2/100,000.

Standardized mortality ratio (SMR) 2007 - 2013: Males



Standardized mortality ratio (SMR) 2007 - 2013: Females




Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2013. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=1,079, females N=820).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,642 female residents (averaged) in the period from 2007 to 2013 a total of 28 women died from brain/nerves cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 1.26. Though, the value of this parameter may vary with an underlying probability of 99% between 0.73 and 2.01, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

FRG Federal Republic of Germany

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

MCR Munich Cancer Registry (Tumorregister München)
SEER Surveillance, Epidemiology, and End Results (USA)

AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

BRD-S German standard population

DCO Death certificate only EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

ES European standard population (old)

LCL Lower confidence limit

MI-index Ratio between mortality and incidence

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age

SIR Standardized incidence ratio
SMR Standardized mortality ratio
UCL Upper confidence limit
WS World standard population

Recommended Citation

Munich Cancer Registry. Baseline statistics C70-C72: Brain/nerves cancer [Internet]. 2015 [updated 2015 May 19; cited 2015 Jul 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/base/base_C7072E.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

Fig./Tbl		Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	4
1a	Gender distribution by year of diagnosis	5
2	Incidence by year of diagnosis	6
3	Age distribution parameters by year of diagnosis	7
4	Age distribution by 5-year age group and gender	9
5	Age-specific incidence and DCO rate	10
6	Standardized incidence ratio of second primaries	11
7	Age distribution and age-specific incidence (chart)	13
7a	Age-specific incidence internationally (chart)	14
8	Cumulative follow-up years (chart)	15
9a	Map of cancer incidence (WS) by county (chart)	16
9b	Standardized incidence ratio (SIR) by county (chart)	17
10a	Pts incident cohorts and mortality / yr	18
10b	Incidence and mortality by year of diagnosis	19
10c	Cancer-related deaths, death certification available / yr	20
11	Medians of age at death / yr	21
12	Mortality by year of death	23
13	Distribution of age at death	24
14	Age-specific mortality	25
15	Multiple primaries in deaths	26
16	Age-specific mortality (first primaries)	28
17	Age-specific mortality (single primaries)	29
18	Age distribution and age-specific mortality (chart)	30
19a	Map of cancer mortality (WS) by county (chart)	31
19b	Standardized mortality ratio (SMR) by county (chart)	32