Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ Deutsch

Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

Cancer statistics: Baseline statistics

CLL: Chronic lymph. leukaemia

Year of diagnosis	1998-2013
Patients	2,933
Diseases	2,937
Creation date	05/19/2015
Export date	12/30/2014
Population	4.64 m



http://www.tumorregister-muenchen.de/en/facts/base/base_C911_E.pdf

Global Statements about the statistics on the Internet – Baseline Statistics (grey button ——), Survival (red button ——)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.64 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases**** are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR. The time-delayed acquisition of data and the occasionally high DCO-rates indicate optimizing reserves, among others, because of current financial and legal conditions that hinder the analyses.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, May 2015

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007). Death certificates from 2014 are incorporated into these analyses.
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate. A high proportion of DCO cases (≥5%) in particular cancer types indicate insufficient participation of specific cancer specializations.

Some remarks regarding this cancer type

The results for leukemias should be interpreted with caution. As with other primarily non-surgically or non-radiologically treated cancer diseases, the MCR hardly manages to obtain even the simplest information on this cancer. The proportion of DCO cases indicates a situation that is far away from a satisfying cooperation. In the group of institutions that potentially participate in reporting are a few hospitals that refuse any contribution to MCR.

ICD-10 codes (ICD-10 2015) used for specifying cancer site

Code	Description
C91.1	Chronic lymphocytic leukaemia of B-cell type

INCIDENCE

Table 1

Patient cohorts by year of diagnosis including DCO cases and multiple primaries, and with proportion of deaths and active follow-up

				Prop.		Prop.
		DCO	Prop.	mult.	Prop.	actively
Year of	Cases	cases	DCO	primaries	deaths	followed
diagnosis	n	n	%	8	%	%
1998	99	8	8.1	23.2	72.7	97.0
1999	93	7	7.5	37.6	66.7	98.9
2000	96	16	16.7	36.5	76.0	96.9
2001	139	36	25.9	28.1	73.4	99.3
2002	251	74	29.5	30.3	75.7	96.8 #
2003	210	64	30.5	28.1	68.6	97.6
2004	231	51	22.1	32.9	59.7	93.1
2005	204	54	26.5	38.2	69.6	95.6
2006	215	36	16.7	42.8	60.9	93.5
2007	252	55	21.8	31.0	57.9	84.5 # ##
2008	221	44	19.9	43.0	49.8	72.9
2009	228	48	21.1	32.5	45.2	67.1
2010	209	53	25.4	36.4	46.4	76.6
2011	196	49	25.0	37.8	41.3	73.5
2012	179	47	26.3	31.3	40.8	67.6
2013	114	45	39.5	40.4	46.5	93.9 ###
1998-2013	2937	687	23.4	34.5	58.5	86.4

[#] The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

^{##} Since 2007 the percentage of actively followed patients sharply declined compared to the previous years. This is a consequence of ambiguous data protection rules that currently forbid cancer registries in Bavaria to obtain the essential life status informations from competent registration offices.

^{###} Please be aware that data of recent annual patient cohorts may not yet be fully processed. Therefore, the presented figures and tables are potentially related to different time periods as pointed out in the respective headlines or legends.

Table 1a

Patient cohorts by year of diagnosis and gender including DCO cases

Year of	All	Males	Females	Prop. males
diagnosis	n /	'n	n	%
1998	99	52	47	52.5
1999	93	48	45	51.6
2000	96	60	36	62.5
2001	139	74	65	53.2
2002	251	138	113	55.0
2003	210	131	79	62.4
2004	231	129	102	55.8
2005	204	128	76	62.7
2006	215	135	80	62.8
2007	252	146	106	57.9
2008	221	127	94	57.5
2009	228	129	99	56.6
2010	209	120	89	57.4
2011	196	112	84	57.1
2012	179	107	72	59.8
2013	114	72	42	63.2
1998-2013	2937	1708	1229	58.2

Table 2

Incidence measures by year of diagnosis and gender including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

			Males	Fem.	Males	Fem.	Males	Fem.	Males	Fem.
Year of	Males	Females	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	52	47	4.7	4.0	2.8	1.8	4.2	2.7	5.5	3.4
1999	48	45	4.3	3.8	2.5	1.9	3.8	2.7	5.0	3.3
2000	60	36 /	5.3	3.0	3.2	1.3	4.7	2.0	5.6	2.5
2001	74	65	6.4	5.3	3.8	2.2	5.6	3.4	7.3	4.3
2002	138	113	7.4	5.8	4.2	2.2	6.2	3.4	8.0	4.5
2003	131	79	7.0	4.0	3.8	1.6	5.7	2.4	7.6	3.2
2004	129	102	6.9	5.2	3.7	2.2	5.6	3.2	7.3	4.1
2005	128	76	6.8	3.8	3.4	1.4	5.3	2.2	7.2	3.0
2006	135	80	7.0	4.0	3.6	1.5	5.6	2.3	7.3	3.0
2007	146	106	6.6	4.6	3.3	1.8	5.1	2.7	6.9	3.5
2008	127	94	5.7	4.1	2.6	1.5	4.1	2.3	5.6	3.0
2009	129	99	5.8	4.3	2.9	1.5	4.4	2.3	5.6	3.0
2010	120	89	5.3	3.8	2.4	1.2	3.8	1.9	5.2	2.7
2011	112	84	4.9	3.6	2.3	1.2	3.5	1.9	4.6	2.4
2012	107	72	4.7	3.1	2.0	1.0	3.1	1.6	4.5	2.1
2013	72	42	3.2	1.8	1.3	0.6	2.2	0.9	3.0	1.3
1998-2013	1708	1229	5.7	4.0	2.9	1.5	4.5	2.3	5.9	3.0




Table 3

Age distribution parameters by year of diagnosis (All) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	99	68.0	12.8	33.9	95.8	54.0	59.2	67.4	77.8	84.2
1999	93	66.4	12.1	31.9	89.4	52.0	58.5	66.1	76.8	80.4
2000	96	68.8	11.1	46.5	91.2	54.7	60.5	67.6	77.1	86.5
2001	139	70.9	11.8	40.5	94.0	56.9	63.3	69.3	78.7	87.7
2002	251	71.4	12.3	28.6	95.0	55.9	63.0	71.1	80.1	88.3
2003	210	71.6	12.1	35.6	98.9	55.8	63.5	72.3	80.7	87.7
2004	231	70.5	12.4	29.8	98.6	55.4	63.3	70.7	79.8	85.7
2005	204	73.0	11.2	34.4	97.1	57.6	65.8	74.7	80.1	86.1
2006	215	72.0	11.5	30.0	95.4	56.9	64.9	72.8	80.6	86.2
2007	252	72.0	12.7	37.6	99.8	53.3	63.4	73.4	81.4	87.0
2008	221	73.9	10.5	42.6	97.4	61.0	67.2	73.1	81.7	87.3
2009	228	72.5	12.6	28.2	98.6	56.0	64.2	72.7	82.4	88.1
2010	209	75.0	12.1	42.8	101	55.4	68.3	76.4	83.7	89.3
2011	196	73.0	12.7	41.8	101	54.1	65.9	73.0	82.1	89.4
2012	179	75.2	11.3	28.2	96.9	60.7	68.9	75.8	84.0	88.7
2013	114	76.6	10.9	44.9	100	63.3	69.1	77.0	85.5	90.5
1998-2013	2937	72.2	12.1	28.2	101	56.2	64.2	72.7	81.1	87.7

Table 3a

Age distribution parameters by year of diagnosis (MALES)

(incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	52	66.2	13.2	33.9	95.8	51.8	57.7	64.6	75.8	82.5
1999	48	66.0	12.9	31.9	89.4	51.8	56.2	64.3	78.3	82.6
2000	60	67.4	10.4	46.5	91.1	54.3	60.7	67.1	72.2	82.6
2001	74	68.0	11.1	40.5	90.7	56.2	60.7	67.5	76.2	83.2
2002	138	68.5	11.7	28.6	90.9	54.1	61.9	68.1	76.8	84.1
2003	131	69.7	11.0	35.6	90.7	56.0	62.8	69.9	77.4	83.8
2004	129	69.7	11.8	29.8	95.2	56.4	62.9	70.2	77.3	82.5
2005	128	71.1	11.1	34.4	91.3	55.2	65.5	71.8	78.2	85.0
2006	135	70.9	11.3	30.0	95.4	55.9	64.3	71.5	78.1	85.2
2007	146	70.2	12.4	37.6	97.8	51.2	61.1	71.7	80.1	85.0
2008	127	72.8	8.9	49.5	93.7	61.2	67.4	72.1	78.7	84.7
2009	129	70.7	11.5	42.1	97.0	54.6	63.1	71.1	79.1	86.5
2010	120	73.5	12.0	45.3	101	54.7	66.5	75.1	81.7	88.1
2011	112	70.8	12.1	41.8	101	53.2	63.6	71.4	79.8	85.4
2012	107	73.7	10.3	39.0	95.2	60.7	68.7	73.7	80.8	86.5
2013	72	76.6	11.0	46.3	100	63.0	69.5	76.9	85.4	88.3
1998-2013	1708	70.7	11.6	28.6	101	55.0	63.3	71.2	78.9	85.3

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	47	70.1	12,2	39.8	90.3	55.6	61.4	71.4	78.7	86.7
1999	45	66.9	11.3	39.1	88.4	52.3	60.4	68.0	75.1	78.2
2000	36	71.2	12.0	49.3	91.2	54.7	60.5	71.7	81.3	88.6
2001	65	74.3	11.8	51.2	94.0	59.4	65.0	73.7	84.7	92.5
2002	113	74.9	12.0	40.4	95.0	58.5	66.0	75.6	84.4	90.1
2003	79	74.8	13.2	47.3	98.9	53.7	65.0	77.5	83.5	92.1
2004	102	71.5	13.1	40.4	98.6	52.1	64.0	72.5	81.7	87.3
2005	76	76.2	10.7	43.7	97.1	62.4	69.2	77.5	84.1	90.3
2006	80	74.0	11.7	32.7	93.9	57.9	67.3	74.9	82.8	87.8
2007	106	74.5	12.7	39.5	99.8	56.7	64.8	77.0	85.0	89.0
2008	94	75.4	12.3	42.6	97.4	60.7	66.7	78.0	84.5	90.8
2009	99	74.7	13.6	28.2	98.6	57.5	69.1	75.9	85.3	88.7
2010	89	77.0	12.2	42.8	97.5	60.4	70.6	79.1	85.1	91.2
2011	84	75.9	13.0	44.4	96.7	56.6	69.0	76.5	86.9	90.9
2012	72	77.4	12.4	28.2	96.9	61.1	69.5	80.0	87.1	90.0
2013	42	76.7	10.8	44.9	91.6	63.3	67.6	77.0	86.2	90.6
1998-2013	1229	74.4	12.5	28.2	99.8	58.1	65.8	75.7	84.2	89.7

Table 4

Age distribution by 5-year age group and gender for period 1998-2013 (incl. DCO)

diagnosis Cases Males Females Years n % Cum.% n % Cum.% n % C	Cum.%
Years n % Cum.% n % Cum.% n % (um.%
25-29 5 0.2 0.2 3 0.2 0.2 2 0.2	0.2
30-34 5 0.2 0.3 4 0.2 0.4 1 0.1	0.2
35-39 11 0.4 0.7 8 0.5 0.9 3 0.2	0.5
40-44 39 1.3 2.0 19 1.1 2.0 20 1.6	2.1
45-49 63 2.1 4.2 41 2.4 4.4 22 1.8	3.9
50-54 134 4.6 8.8 95 5.6 10.0 39 3.2	7.1
55-59 199 6.8 15.5 131 7.7 17.6 68 5.5	12.6
60-64 328 11.2 26.7 198 11.6 29.2 130 10.6	23.2
65-69 430 14.6 41.3 279 16.3 45.6 151 12.3	35.5
70-74 451 15.4 56.7 293 17.2 62.7 158 12.9	48.3
75-79 439 14.9 71.6 257 15.0 77.8 182 14.8	63.1
80-84 373 12.7 84.3 200 11.7 89.5 173 14.1	77.2
85+ 460 15.7 100.0 180 10.5 100.0 280 22.8 1	00.0
All ages 2937 100.0 1708 100.0 1229 100.0	

Included in the statistics are 53.9% multiple primaries in males and 36.3% in females.

Table 5

Age-specific incidence, DCO rate and proportion of all cancers for period 1998-2013

							Males	Females
			Males	Females	Males	Females		Prop.all
Age at			Age-	Age-		DCO rate	_	cancers
diagnosis	Males	Females	~	spec.	n=325	n=361		n=153136
Years	n	n	incid.	/ -	%	%	%	%
10012				/			v	/
0- 4			0.0	0.0				
5- 9			0.0	0.0				
10-14			0.0	0.0				
15-19			0.0	0.0				
20-24			0.0	0.0				
25-29	3	2	0.1	0.1		50.0	0.3	0.2
30-34	4	1	0.2	0.0			0.3	0.0
35-39	8	3	0.3	0.1			0.4	0.1
40-44	19	20	0.7	0.8	5.3		0.6	0.3
45-49	41	22	1.7	1.0			0.8	0.3
50-54	95	39	4.7	1.9	4.2		1.1	0.4
55-59	131	68	7.1	3.5	6.1	_1.5	0.9	0.5
60-64	198	130	11.2	6.9	7.6	6.9	0.9	0.8
65-69	279	151	17.7	8.8	9.0	6.0	1.0	0.8
70-74	293	158	22.9	10.4	11.9	14.6	1.1	0.9
75-79	257	181	31.1	15.2	23.3	24.9	1.2	1.0
80-84	200	173	40.0	18.5	33.0	51.4	1.5	1.1
85+	180	279	52.8	31.2	61.7	65.9	1.8	1.6
All ages	1708	1227			19.0	29.4	1.1	0.8
Incidence				4 0				
Raw			5.7	4.0				
WS			2.9	1.5				
ES			4.5	2.3				
BRD-S			5.9	3.0				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

Table 6a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2013

1\/	7	т	777	
I٧	м	ш	ır.	

	Observed	Expected		LCL	UCL		DCO
Diagnosis	n /	n	SIR	95%	95%	EAR	%
C07-C08 Salivary gland	5	0.2	24.8	8.1	58.0 #	9.7	
C15 Oesophagus	4	1.6	2.6	0.7	6.5	4.9	
C16 Stomach	9	3.7	2.4	1.1	4.6 #	10.6	
C18 Colon	15	9.0	1.7	0.9	2.8	12.2	6.7
C19-C20 Rectum	11	5.0	2.2	/ 1.1	4.0 #	12.1	
C22 Liver	7	2.5	2.8/	1.1	5.8 #	9.1	14.3
C25 Pancreas	3	3.2	0.9	0.2	2.7	-0.5	
C33-C34 Lung	29	10.6	2.7	1.8	3.9 #	37.1	
C43 Malign. melanoma	14	3.5	4.0	2.2	6.6 #	21.1	
C46,C49 Soft tissue	3	0.5	6.4	1.3	18.6 #	5.1	
C50 Breast	2	0.2	8.7	1.0	31.3 #		
C61 Prostate	58	26.9	2.2	1.6	2.8 #	62.7	3.4
C62 Testis	2	0.2	11.0	1.3	39.8 #	3.7	
C64 Kidney	5	3.2	1.6	0.5	3.7	3.7	
C65 Renal pelvis	2	0.4	5.3	0.6	19.2	3.3	
C67 Bladder	9	4.0	2.2	1.0	4.2 #	10.0	
C70-C72 CNS cancer	2	1.2	1.7	0.2	6.2	1.7	
C76-C79 CUP	3	1.5	2.0	0.4	5.8	3.0	
C81 Hodgkin lymphoma	5	0.2	28.4	9.2	66.2 #		
C82-C85 NHL	9	3.6	2.5	1.1	4.7 #	10.9	33.3
C90 Mult. myeloma	2	1.2	1.7	0.2	6.2	1.7	
C91-C96 Leukaemia	8	1.5	5.5	2.4	10.8 #	13.2	37.5
Other primaries	5	1.8	2.7	0.9	6.4	6.4	20.0
Not observed	0	5.4	0.0	0.0	0.7 #	-10.9	
All mult. primaries	212	91.0	2.3	2.0	2.7 #	244.2	5.2

Patients	1035
Median age at second malignancy (years)	72.6
Person-years	4955
Mean observation time (years)	4.8
Median observation time (years)	4.4

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries".

Table 6b

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2013

FEMALES

	Observed I	Expected		LCL	UCL		DCO
Diagnosis	/ n /	n	SIR	95%	95%	EAR	8
C09-C10 Oropharynx	/ 2/	0.1	13.7	1.7	49.3 #	5.8	
C16 Stomach	5 6	1.5	3.3	1.1	7.7 #	10.9	
C18 Colon	6	4.2	1.4	0.5	3.1	5.7	16.7
C19-C20 Rectum	2	1.8	/1.1	0.1	4.1	0.7	
C21 Anus/canal	2	0.2	9.8	1.2	35.3 #	5.6	
C25 Pancreas	3	1.8	/ 1.7/	0.3	4.8	3.7	
C33-C34 Lung	7	2.8	2.5	1.0	5.2 #	13.2	
C43 Malign. melanoma	6	1.3	4.5	1.6	9.8 #	14.6	
C50 Breast	29	11.6	2.5	1.7	3.6 #	54.5	
C54 Corpus uteri	4	2.3	1.8	0.5	4.5	5.5	
C56 Ovary	3	1.7	1.8	0.4	5.3	4.2	
C73 Thyroid	2	0.6	3.2	0.4	11.6	4.3	
C76-C79 CUP	2	0.7	2.8	0.3	10.0	4.0	
C82-C85 NHL	10	1.6	6.3	3.0	11.6 #	26.4	20.0
C90 Mult. myeloma	2	0.5	3.9	0.5	13.9	4.6	
C91-C96 Leukaemia	3	0.6	4.7	1.0	13.6	7.4	
Other primaries	5	1.7	3.0	1.0	7.0	10.4	
Not observed	0	4.8	0.0	0.0	0.8 #	-15.2	
All mult. primaries	93	39.9	2.3	1.9	2.9 #	166.5	3.2
<u>-</u>							

Patients	672
Median age at second malignancy (years)	72.9
Person-years	3189
Mean observation time (years)	4.7
Median observation time (years)	4.3

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries".

CLL: Chronic lymphocytic leukaemia of B-cell type Age distribution and age-specific incidence 1998 - 2013 (Males: 1708, Females: 1227) 25




Figure 7. Age distribution and age-specific incidence






Figure 7a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).



Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2014, based on the November 2013 submission. http://www.seer.cancer.gov.





Figure 8. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.



Average incidence (world standard population) 2007 - 2013: Males



Average incidence (world standard population) 2007 - 2013: Females





Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2007 to 2013. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 2.4/100,000 WS N=813, females 1.3/100,000 WS N=586).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,928 female residents (averaged) in the period from 2007 to 2013 a total of 9 women were identified with newly diagnosed chronic lymph. leukaemia. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.6/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.2 and 1.4/100,000.

Standardized incidence ratio (SIR) 2007 - 2013: Males



Standardized incidence ratio (SIR) 2007 - 2013: Females




Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2013. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=813, females N=586).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,642 female residents (averaged) in the period from 2007 to 2013 a total of 9 women were identified with newly diagnosed chronic lymph. leukaemia. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.60. Though, the value of this parameter may vary with an underlying probability of 99% between 0.21 and 1.33, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts, and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

	Incident	Prop. actively	Prop.		Prop.	Prop. deaths with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	%	olo	n	%	%
1998	99	97.0	8.1	72	72.7	98.6
1999	93	98.9	7.5	62	66.7	95.2
2000	96	96.9	16.7	73	76.0	98.6
2001	139	99.3	25.9	102	73.4	97.1
2002	251	96.8	29.5	190	75.7	98.4
2003	210	97.6	30.5	144	68.6	98.6
2004	231	93.1	22.1	138	59.7	100.0
2005	204	95.6	26.5	142	69.6	98.6
2006	215	93.5	16.7	131	60.9	96.9
2007	252	84.5	21.8	146	57.9	99.3
2008	221	72.9	19.9	110	49.8	96.4
2009	228	67.1	21.1	103	45.2	97.1
2010	209	76.6	25.4	97	46.4	96.9
2011	196	73.5	25.0	81	41.3	100.0
2012	179	67.6	26.3	73	40.8	98.6
2013	114	93.9	39.5	53	46.5	100.0
1998-2013	2937	86.4	23.4	1717	58.5	98.2

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

			Prop. deaths		Dwon
Voor of	Incident		with death	Deaths in	Prop. deaths in
Year of		//			
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	ń	%	n	96
1998	99	40	97.5	/ 7	7.1
1999	93	50	92.0	5	5.4
2000	96	50	92.0	14	14.6
2001	139	89	96.6	40	28.8
2002	251	132	98.5	83	33.1
2003	210	121	98.3	75	35.7
2004	231	119	100.0	54	23.4
2005	204	149	100.0	63	30.9
2006	215	140	97.9	49	22.8
2007	252	154	100.0	66	26.2
2008	221	154	98.7	53	24.0
2009	228	133	100.0	52	22.8
2010	209	158	98.7	62	29.7
2011	196	154	99.4	55	28.1
2012	179	169	98.2	56	31.3
2013	114	159	99.4	51	44.7
1998-2013	2937	1971	98.6	785	26.7

Table 10c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

				Prop.
				cancer
		Prop.	Prop.	recorded
		cancer-	non-cancer-	on death
Year of	Deaths	related	related	certificate
death	n	%	8	%
1998	40	62.5	37.5	89.7
1999	50	62.0	38.0	87.0
2000	50	62.0	38.0	97.8
2001	89	59.6	40.4	93.0
2002	132	76.5	23.5	96.2
2003	121	80.2	19.8	94.1
2004	119	87.4	12.6	95.8
2005	149	79.9	20.1	96.6
2006	140	77.9	22.1	92.0
2007	154	74.7	25.3	89.6
2008	154	81.8	18.2	89.5
2009	133	82.7	17.3	94.0
2010	158	80.4	19.6	93.6
2011	154	75.3	24.7	90.2
2012	169	78.1	21.9	89.2
2013	159	73.0	27.0	86.1
1998-2013	1971	76.7	23.3	92.0

Table 11a $\begin{tabular}{ll} Medians of age at death according to the grouping in Table 10 \\ \hline MALES \end{tabular}$

Year of death	Deaths n	Age at death (all causes)	Age at death (cancer-related)	Age at death (non-cancer-related)	Age at death (according to death certificate)
1998	23	73.9	68.2	84.2	74.3
1999	28	76.0	68.3	82.9	73.5
2000	34	73.5	71.3	80.2	73.5
2001	42	74.2	73.7	80.8	74.0
2002	76	76.5	75.9	77.1	77.1
2003	67	74.2	73.5	75.5	74.2
2004	76	74.5	73.7	82.4	74.5
2005	90	77.2	75.9	79.4	77.1
2006	83	76.0	73.7	81.9	75.0
2007	82	78.1	77.1	82.1	78.0
2008	96	75.7	75.1	80.8	75.6
2009	78	78.9	76.7	84.0	78.9
2010	94	77.9	77.2	81.6	77.9
2011	98	77.4	76.3	79.3	77.4
2012	107	77.8	77.6	82.6	78.4
2013	104	77.9	74.8	83.6	77.9
1998-2013	1178	76.7	75.5	81.0	76.5

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 11b $\begin{tabular}{ll} \begin{tabular}{ll} \begin{tabula$

Year of death	Deaths n	Age at death (all causes)	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate)
1998	17	79.9	76.7	82.2	78.6
1999	22	79.8	76.5	83.2	76.8
2000	16	85.5	81.3	88.7	85.5
2001	47	78.8	77.3	81.3	78.8
2002	56	83.1	76.1	88.7	82.9
2003	54	79.4	77.9	84.5	79.1
2004	43	79.7	77.3	84.5	78.9
2005	59	81.6	78.2	89.3	81.4
2006	57	79.9	78.4	81.8	78.8
2007	72	81.8	79.3	87.5	81.8
2008	58	82.8	81.7	91.3	81.9
2009	55	81.8	79.4	84.5	80.9
2010	64	82.3	82.2	89.6	82.3
2011	56	82.4	80.7	85.4	81.6
2012	62	81.7	80.7	84.2	80.3
2013	55	82.7	82.4	85.5	83.1
1998-2013	793	81.7	79.3	86.2	80.9

By 2010, life expectancy for a newborn male in Germany is 77.5 years compared with 82.6 years for his female counterpart.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 12a

Mortality measures (cancer-related death) and mortality-incidence-index by year of death

MALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	17	1.5	0.33	0.9	0.34	1.4	0.34	1.7	0.32
1999	16	1.4	0.33	0.8	0.34	1.3	0.34	1.7	0.34
2000	23	2.0	0.38	1.2	0.38	1.8	0.39	2.3	0.41
2001	29	2.5	0.39	1.4	0.37	2.2	0.39	3.1	0.42
2002	63	3.4	0.46	1.7	0.41	2.8	0.45	4.1	0.51
2003	54	2.9	0.41	1.4	0.38	2.4	0.41	3.3	0.44
2004	68	3.6	0.53	1.8	0.49	2.9	0.52	4.0	0.55
2005	70	3.7	0.55	1.7	0.50	2.8	0.53	4.1	0.57
2006	65	3.4	0.48	1.6	0.43	2.5	0.45	3.6	0.49
2007	63	2.8	0.43	1.2	0.38	2.1	0.41	3.1	0.45
2008	77	3.5	0.61	1.5	0.57	2.5	0.61	3.6	0.64
2009	66	3.0	0.51	1.3	0.46	2.1	0.49	3.0	0.54
2010	76	3.4	0.63	1.3	0.56	2.2	0.59	3.4	0.65
2011	78	3.4	0.70	1.4	0.59	2.3	0.64	3.3	0.72
2012	80	3.5	0.75	1.4	0.70	2.3	0.74	3.5	0.78
2013	79	3.5	1.10	1.4	1.07	2.3	1.04	3.3	1.07
1998-2013	3 924	3.1	0.54	1.4	0.49	2.3	0.52	3.3	0.57

Table 12b

Mortality measures (cancer-related death) and mortality-incidence-index by year of death

FEMALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	8	0.7	0.17	0.3	0.14	0.4	0.15	0.5	0.16
1999	15	1.3	0.33	0.5	0.24	0.7	0.27	1.0	0.31
2000	8	0.7	0.22	0.2	0.18	0.4	0.20	0.5	0.20
2001	24	2.0	0.38	0.7	0.32	1.1	0.33	1.6	0.38
2002	38	1.9	0.34	0.7	0.30	1.1	0.32	1.5	0.34
2003	43	2.2	0.54	0.8	0.48	1.2	0.50	1.7	0.55
2004	36	1.8	0.35	0.6	0.28	1.0	0.30	1.4	0.34
2005	49	2.5	0.64	0.8	0.58	1.3	0.59	1.8	0.61
2006	44	2.2	0.55	0.7	0.44	1.1	0.48	1.7	0.55
2007	52	2.3	0.49	0.7	0.42	1.2	0.44	1.7	0.47
2008	49	2.1	0.52	0.6	0.41	1.0	0.45	1.5	0.50
2009	44	1.9	0.44	0.6	0.40	1.0	0.43	1.4	0.46
2010	51	2.2	0.57	0.6	0.47	1.0	0.51	1.5	0.55
2011	38	1.6	0.45	0.4	0.37	0.7	0.39	1.1	0.44
2012	52	2.2	0.72	0.5	0.53	0.9	0.60	1.5	0.72
2013	37	1.6	0.88	0.4	0.67	0.7	0.74	1.1	0.84
1998-2013	588	1.9	0.48	0.6	0.39	1.0	0.42	1.4	0.46

Table 13

Age distribution of age at death (cancer-related) for period 1998-2013

(incl. multiple primaries)

Age at									
death	Cases			Males			Females		
Years	n	%	Cum.%	'n	%	Cum.%	n	%	Cum.%
30-34	1	0.1	0.1	/ 1	0.1	0.1			0.0
35-39	0	0.0	0.1			0.1			0.0
40-44	5	0.3	0.4	4	0.4	0.5	1	0.2	0.2
45-49	12	0.8	1.2	8	0.9	1.4	4	0.7	0.8
50-54	23	1.5	2.7	14	1.5	2.9/	9	1.5	2.4
55-59	54	3.6	6.3	39	4.2	7.1	15	2.5	4.9
60-64	108	7.1	13.4	73	7.9	15.0	35	5.9	10.8
65-69	185	12.2	25.6	137	14.8	29.8	48	8.1	19.0
70-74	261	17.2	42.8	175	18.9	48.7	86	14.6	33.6
75-79	298	19.7	62.5	187	20.2	68.9	111	18.8	52.4
80-84	285	18.8	81.3	157	17.0	85.9	128	21.7	74.1
85+	284	18.7	100.0	131	14.1	100.0	153	25.9	100.0
All ages	1516	100.0		926	100.0		590	100.0	

Included in the statistics are 53.9% multiple primaries in males and 36.3% in females.

Table 14

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2013 (incl. multiple primaries)

Age at death	Males Females	Males Age- s spec.		Females Age- spec.		Males Prop.all cancers	Females Prop.all cancers
Years	n n	/ - /	MI-index		MT-index		%
10015		mor car.	111 1114611	mor car.	III IIIGEI	·	,
0 - 4		0.0		0.0			
5- 9		0.0		0.0			
10-14		0.0		0.0			
15-19		0.0		0.0			
20-24		0.0		0.0			
25-29		0.0		0.0			
30-34	1	0.0	0.25	0.0		0.5	
35-39		0.0		0.0			
40-44	4 1	0.2	0.21	0.0	0.05	0.5	0.1
45-49	8 4	0.3	0.20	0.2	0.18	0.4	0.2
50-54	14 9	0.7	0.15	0.4	0.23	0.4	0.3
55-59	39 15	2.1	0.30	0.8	0.22	0.7	0.3
60-64	73 35	4.1	0.37	1.9	0.27	0.8	0.5
65-69	137 48	8.7	0.49	2.8	0.32	1.1	0.6
70-74	175 86	13.7	0.60	5.7	0.54	1.3	0.9
75-79	187 111	22.6	0.73	9.3	0.61	1.4	1.0
80-84	157 128	31.4	0.79	13.7	0.74	1.4	1.1
85+	131 153	38.4	0.73	17.1	0.55	1.5	1.1
All ages	926 590					1.2	0.8
Mortality							
Raw		3.1	0.54	1.9	0.48		
WS		1.4		0.6	0.39		
ES		2.3	0.52	1.0	0.42		
BRD-S		3.4	0.57	1.4	0.46		
PYLL-70							
per 100,000		7.3		3.2			
ES		6.2		2.6			
AYLL-70		7.1		7.5			

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15a

Multiple primaries in deaths in period 1998-2013

MALES

	Total	Total	Pre	Pre	Syn- chron ±30d	Syn- chron ±30d	Post	Post
Diagnosis	n	%↓	n	←%	n	←%	n	←%
C16 Stomach	10	1.7	2	20.0			8	80.0
C18 Colon	43	7.2	17	39.5	6	14.0	20	46.5
C19-C20 Rectum	24 /	4.0	6	25.0	6	25.0	12	50.0
C25 Pancreas	7 -	1.2			1	14.3	6	85.7
C33-C34 Lung	59	9.9	7	11.9	14	23.7	38	64.4
C43 Malign. melanoma	32	5.4	14	43.8	3	9.4	15	46.9
C44 Skin others	159	26.6	16	10.1	9	5.7	134	84.3
C46,C49 Soft tissue	9	1.5	3	33.3			6	66.7
C61 Prostate	89	14.9	39	43.8	13	14.6	37	41.6
C64 Kidney	14	2.3	7	50.0	2	14.3	5	35.7
C67 Bladder	30	5.0	14	46.7	4	13.3	12	40.0
C70-C72 CNS cancer	6	1.0			1	16.7	/ 5	83.3
C76-C79 CUP	7	1.2					7	100.0
C81 Hodgkin lymphoma	11	1.8	3	27.3	2	18.2	6	54.5
C82-C85 NHL	29	4.9			7	24.1	22	75.9
C90 Mult. myeloma	9	1.5	2	22.2	4	44.4	3	33.3
C91-C96 Leukaemia	15	2.5					15	100.0
Other primaries	44	7.4	14	31.8	9	20.5	21	47.7
All mult. primaries	597	100.0	144	24.1	81	13.6	372	62.3

Multiple primaries with number of cases 1 to 5 are pooled in category "Other primaries".

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 15b

Multiple primaries in deaths in period 1998-2013
FEMALES

		Total	Total	Pre	Pre	Syn- chron ±30d	Syn- chron ±30d	Post	Post
Diagnos	is	n	10cai %↓	n	-%	n	±30α ←%	n	
Diagnos.		7	/ • •		\	\ 11	. 0		
C16	Stomach	7	2.8	1	14.3	4	57.1	2	28.6
C18	Colon	18	7.1	6	33.3	2	11.1	10	55.6
C19-C20	Rectum	6	2.4	3	50.0			3	50.0
C23-C24	Bile	3 <	1.2	1	33.3	1	33.3	1	33.3
C25	Pancreas	5	2.0					5	100.0
C33-C34	Lung	10	4.0			1	10.0	9	90.0
C43	Malign. melanoma	10	4.0	7	70.0			3	30.0
C44	Skin others	68	27.0	16	23.5	4	5.9	48	70.6
C50	Breast	42	16.7	22	52.4	5	11.9	15	35.7
C51	Vulva	4	1.6	3	75.0			1	25.0
C53	Cervix uteri	3	1.2	3	100.0				
C54	Corpus uteri	9	3.6	4	44.4	2	22.2	/ 3	33.3
C56	Ovary	9	3.6	2	22.2	_ 2	22.2	5	55.6
C64	Kidney	7	2.8	2	28.6	3	42.9	2	28.6
C67	Bladder	4	1.6	1	25.0	2	50.0	1	25.0
C70-C72	CNS cancer	6	2.4	2	33.3	2	33.3	2	33.3
C76-C79	CUP	3	1.2	1	33.3			2	66.7
C82-C85	NHL	13	5.2			1	7.7	12	92.3
C90	Mult. myeloma	5	2.0			2	40.0	3	60.0
C91-C96	Leukaemia	6	2.4	1	16.7	1	16.7	4	66.7
Other p	rimaries	14	5.6	4	28.6	2	14.3	8	57.1
All mul	t. primaries	252	100.0	79	31.3	34	13.5	139	55.2

Multiple primaries with number of cases 1 to 2 are pooled in category "Other primaries".

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2013

(Singular primaries only *)

Age at death	Males Female	Males Age- es spec.		Females Age- spec.		Males Prop.all cancers	Females Prop.all cancers
Years	n n	mortal.	MI-index	mortal.	MI-index	%	%
0 - 4		0.0		0.0			
5- 9		0.0		0.0			
10-14		0.0		0.0			
15-19		0.0		0.0			
20-24		0.0		0.0			
25-29	1	0.0		0.0		0 6	
30-34	1	0.0		0.0		0.6	
35-39	2 1	0.0		0.0	0 05	0 1	0 1
40-44 45-49	3 1	0.1		0.0		0.4	0.1
50-54	8 4 11 8	0.3		0.2		0.5	0.2 0.3
55-59	35 13	1.9		0.4		0.4	0.3
60-64	56 31	3.2		1.7		0.7	0.5
65-69	111 42	7.0		2.4		1.1	0.6
70-74	144 81	11.2		5.3		1.3	1.0
75-79	146 82	17.7		6.9		1.5	1.0
80-84	121 100	24.2		10.7		1.5	1.1
85+	96 121	28.2		13.5		1.4	1.1
051	JO 121	20.2	0.00	13.3	0.55	\	.
All ages	732 483					1.1	0.8
mii ageb	732 103					/	0.0
Mortality							
Raw		2.5	0.57	1.6	0.49		
WS		1.1		0.5			
ES		1.8		0.8			
BRD-S		2.6		1.1			
PYLL-70							
per 100,000		6.1		2.9			
ES		5.2		2.4			
AYLL-70		7.2		7.7			

^{*} See corresponding tables with multiple primaries.

Table 17

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2013

(Single primaries only *)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males	Females	/ - /		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0 4								
0 - 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29	1		0.0	0.00	0.0		0 6	
30-34	1		0.0	0.33	0.0		0.6	
35-39		-	0.0		0.0	2 25		
40-44	3	1	0.1		0.0	0.05	0.4	0.1
45-49	6	4	0.3		0.2	0.22	0.4	0.3
50-54	10	6	0.5		0.3	0.19	0.4	0.3
55-59	28	12	1.5	0.30	0.6	0.24	0.6	0.3
60-64	40	25	2.3		1.3	0.25	0.6	0.5
65-69	71	34	4.5	0.44	2.0	0.37	0.8	0.6
70-74	97	64	7.6	0.64	4.2	0.56	1.1	1.0
75-79	99	62	12.0	0.68	5.2	0.55	1.2	0.9
80-84	83	84	16.6	0.67	9.0	0.71	1.3	1.1
85+	75	100	22.0	0.65	11.2	0.46	1.4	1.1
All ages	513	392					0.9	0.8
Mortality								
Raw			1.7	0.49	1.3	0.44		
WS			0.8	0.44	0.4	0.37		
ES			1.3	0.47	0.7	0.39		
BRD-S			1.8	0.51	0.9	0.43		
PYLL-70								
per 100,000			4.7		2.4			
ES			4.1		2.0			
AYLL-70			7.9		7.9			

^{*} See corresponding tables with multiple primaries.





Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at chronic lymph. leukaemia-related death (see Table 10) should be considered.



Average mortality (world standard population) 2007 - 2013: Males



Average mortality (world standard population) 2007 - 2013: Females





Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2007 to 2013. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 1.4/100,000 WS N=516, females 0.6/100,000 WS N=322).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,928 female residents (averaged) in the period from 2007 to 2013 a total of 11 women died from chronic lymph. leukaemia. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.5/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.2 and 1.1/100,000.

Standardized mortality ratio (SMR) 2007 - 2013: Males



Standardized mortality ratio (SMR) 2007 - 2013: Females




Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2013. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=516, females N=322).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,642 female residents (averaged) in the period from 2007 to 2013 a total of 11 women died from chronic lymph. leukaemia. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 1.35. Though, the value of this parameter may vary with an underlying probability of 99% between 0.53 and 2.80, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

FRG Federal Republic of Germany

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

MCR Munich Cancer Registry (Tumorregister München)
SEER Surveillance, Epidemiology, and End Results (USA)

AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

BRD-S German standard population

DCO Death certificate only EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

ES European standard population (old)

LCL Lower confidence limit

MI-index Ratio between mortality and incidence

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age

SIR Standardized incidence ratio SMR Standardized mortality ratio UCL Upper confidence limit WS World standard population

Recommended Citation

Munich Cancer Registry. Baseline statistics CLL: Chronic lymph. leukaemia [Internet]. 2015 [updated 2015 May 19; cited 2015 Jul 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/base/base_C911_E.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

Fig./Tbl	l.	Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	3
1a	Gender distribution by year of diagnosis	4
2	Incidence by year of diagnosis	5
3	Age distribution parameters by year of diagnosis	6
4	Age distribution by 5-year age group and gender	8
5	Age-specific incidence and DCO rate	9
6	Standardized incidence ratio of second primaries	10
7	Age distribution and age-specific incidence (chart)	12
7a	Age-specific incidence internationally (chart)	13
8	Cumulative follow-up years (chart)	14
9a	Map of cancer incidence (WS) by county (chart)	15
9b	Standardized incidence ratio (SIR) by county (chart)	16
10a	Pts incident cohorts and mortality / yr	17
10b	Incidence and mortality by year of diagnosis	18
10c	Cancer-related deaths, death certification available / yr	19
11	Medians of age at death / yr	20
12	Mortality by year of death	22
13	Distribution of age at death	23
14	Age-specific mortality	24
15	Multiple primaries in deaths	25
16	Age-specific mortality (first primaries)	27
17	Age-specific mortality (single primaries)	28
18	Age distribution and age-specific mortality (chart)	29
19a	Map of cancer mortality (WS) by county (chart)	30
19b	Standardized mortality ratio (SMR) by county (chart)	31