Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ Deutsch

Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

Cancer statistics: Baseline statistics

BNET: Pulm. neuroend. tumor

Year of diagnosis	1998-2013
Patients	622
Diseases	623
Creation date	05/19/2015
Export date	12/30/2014
Population	4.64 m



http://www.tumorregister-muenchen.de/en/facts/base/base_hBNETE.pdf

Global Statements about the statistics on the Internet – Baseline Statistics (grey button ——), Survival (red button ——)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.64 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases**** are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR. The time-delayed acquisition of data and the occasionally high DCO-rates indicate optimizing reserves, among others, because of current financial and legal conditions that hinder the analyses.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, May 2015

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007). Death certificates from 2014 are incorporated into these analyses.
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- """ DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate. A high proportion of DCO cases (≥5%) in particular cancer types indicate insufficient participation of specific cancer specializations.

Topography codes (ICD-O-3 2000) used for specifying cancer site

Code	Description
C33	Malignant neoplasm of trachea
C34	Malignant neoplasm of bronchus and lung
C34.0	Main bronchus
C34.1	Upper lobe, bronchus or lung
C34.2	Middle lobe, bronchus or lung
C34.3	Lower lobe, bronchus or lung
C34.8	Overlapping lesion of bronchus and lung
C34.9	Bronchus or lung, unspecified

... if additionally existing any of ...

Morphology codes (ICD-O-3 2011) used for specifying cancer site

Code	Description	
8013/3 8240/3 8249/3	Large cell neuroendocrine carcinoma Carcinoid tumor, NOS Atypical carcinoid tumor	

Reference:

Travis WD, Brambilla E, Muller-Hermelink HK, Harris CC, editors. WHO Classification of Tumours. Pathology and Genetics of Tumours of the Lung, Pleura, Thymus and Heart. IARC, Lyon (2004).

INCIDENCE

Table 1

Patient cohorts by year of diagnosis including multiple primaries, and with proportion of deaths and active follow-up

		Prop.		Prop.
		mult.	Prop.	actively
Year of	Cases	primaries	deaths	followed
diagnosis	n	%	%	%
1998	12	16.7	50.0	83.3
1999	23	26.1	60.9	95.7
2000	21	33.3	52.4	100.0
2001	21	23.8	61.9	95.2
2002	25	28.0	40.0	96.0 #
2003	22	13.6	50.0	100.0
2004	29	17.2	37.9	93.1
2005	28	25.0	32.1	89.3
2006	29	13.8	27.6	89.7
2007	41	17.1	51.2	82.9 # ##
2008	61	34.4	44.3	72.1
2009	57	28.1	36.8	61.4
2010	50	32.0	46.0	64.0
2011	76	22.4	40.8	72.4
2012	67	28.4	40.3	76.1
2013	61	29.5	23.0	96.7 ###
1998-2013	623	25.7	41.3	81.4

Due to the pathohistological classification of the tumor and the lack of information on morphology on the death certificates, the proportion of DCO cases can not be determined.

- # The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.
- ## Since 2007 the percentage of actively followed patients sharply declined compared to the previous years. This is a consequence of ambiguous data protection rules that currently forbid cancer registries in Bavaria to obtain the essential life status informations from competent registration offices.
- ### Please be aware that data of recent annual patient cohorts may not yet be fully processed. Therefore, the presented figures and tables are potentially related to different time periods as pointed out in the respective headlines or legends.

Table 1a

Patient cohorts by year of diagnosis and gender including DCO cases

Year of	All	Males	Females	Prop. males
diagnosis	n /	'n	n	%
1998	12	6	6	50.0
1999	23	/ 11	12	47.8
2000	/21	8	13	38.1
2001	21	14	7/	66.7
2002	25	12	13	48.0
2003	22	9	1/3	40.9
2004	29	14	15	48.3
2005	28	16	12	57.1
2006	29	17	12	58.6
2007	41	24	17	58.5
2008	61	30	31	49.2
2009	57	26	31	45.6
2010	50	29	21	58.0
2011	76	40	36	52.6
2012	67	35	32	52.2
2013	61	32	29	52.5
1998-2013	623	323	300	51.8

Table 2

Incidence measures by year of diagnosis and gender including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

			Males	Fem.	Males	Fem.	Males	Fem.	Males	Fem.
Year of	Males	Females	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	6	6	0.5	0.5	0.4	0.3	0.5	0.4	0.6	0.5
1999	11	12	1.0	1.0	0.6	0.5	0.9	0.7	1.1	0.9
2000	8	13	0.7	1.1	0.4	0.8	0.6	1.0	0.7	1.1
2001	14	7 /	1.2	0.6	0.9	0.4	1.1	0.5	1.3	0.5
2002	12	13 <	0.6	0.7	0.4	0.4	0.6	0.6	0.7	0.6
2003	9	13	0.5	0.7	0.3	0.4	0.4	0.5	0.5	0.6
2004	14	15	0.7	0.8	0.4	0.5	0.6	0.6	0.7	0.7
2005	16	12	0.8	0.6	0.7	0.4	0.8	0.5	0.8	0.6
2006	17	12	0.9	0.6	0.5	0.4	0.7	0.5	0.8	0.6
2007	24	17	1.1	0.7	0.7	0.4	0.9	0.6	1.1	0.7
2008	30	31	1.3	1.3	0.8	0.8	1.1	1.1	1.2	1.2
2009	26	31	1.2	1.3	0.7	0.8	1.0	1.1	1.1	1.2
2010	29	21	1.3	0.9	0.7	0.5	1.0	0.7	1.2	0.8
2011	40	36	1.8	1.5	1.0	0.9	1.4	1.2	1.7	1.4
2012	35	32	1.5	1.4	0.9	0.7	1.2	1.0	1.4	1.1
2013	32	29	1.4	1.2	0.7	0.6	1.1	0.9	1.3	1.0
1998-2013	323	300	1.1	1.0	0.6	0.6	0.9	0.8	1.0	0.9

The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

Table 3

Age distribution parameters by year of diagnosis (All) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	12	60.1	8.0	49.4	70.9	50.9	51.9	60.3	68.3	69.6
1999	23	64.3	13.8	32.9	81.2	45.5	54.6	67.2	76.7	78.9
2000	21	59.2	15.8	15.8	80.5	44.9	52.5	63.2	71.3	74.9
2001	21	56.2	17.1	17.0	79.8	31.3	41.3	62.2	69.3	70.7
2002	25	63.1	13.4	31.0	82.8	48.2	53.9	67.4	74.9	75.6
2003	22	63.6	10.6	43.9	84.4	49.4	60.1	63.2	72.7	79.5
2004	29	58.8	15.2	27.6	84.1	37.8	47.8	62.6	69.2	78.6
2005	28	51.5	16.6	18.1	82.9	21.7	41.6	52.2	65.4	71.2
2006	29	58.9	16.1	27.5	82.6	30.4	47.3	60.4	70.6	79.5
2007	41	60.0	17.0	18.7	84.2	39.9	47.5	65.8	72.1	79.2
2008	61	63.9	12.2	29.4	88.0	43.3	59.6	65.0	71.7	77.0
2009	57	62.5	12.0	25.9	89.3	48.1	55.8	62.7	70.1	76.7
2010	50	63.1	12.5	15.6	82.0	46.0	56.0	65.7	71.0	75.4
2011	76	63.6	12.1	33.0	85.6	47.3	55.8	63.1	71.6	80.5
2012	67 /	66.3	10.5	39.5	89.1	53.8	58.5	68.3	73.7	78.5
2013	61	66.3	10.3	36.7	85.4	51.7	59.7	68.1	73.5	78.9
1998-2013	623	62.4	13.4	15.6	89.3	44.0	55.1	64.3	71.7	78.4

Table 3a

Age distribution parameters by year of diagnosis (MALES)

(incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	6	57.4	10.0	49.4	70.9	49.4	50.9	51.9	69.6	70.9
1999	11	63.4	13.2	38.6	80.4	45.5	53.5	66.2	74.2	78.9
2000	8	58.7	9.9	44.9	77.6	44.9	53.2	56.6	64.0	77.6
2001	14	56.5	19.5	17.0	79.8	29.8	38.4	65.1	70.7	72.6
2002	12	66.7	10.6	49.8	82.8	53.5	57.3	69.4	74.9	75.4
2003	9	63.5	12.3	49.4	84.4	49.4	53.2	62.7	65.5	84.4
2004	14	58.3	14.6	37.8	80.5	38.9	41.0	60.6	69.2	78.6
2005	16	50.4	16.2	18.1	71.2	21.7	41.0	52.2	66.0	70.1
2006	17	62.3	15.9	28.7	82.6	35.1	56.4	66.5	72.3	80.9
2007	24	61.3	15.9	18.7	80.7	44.2	53.7	65.9	72.1	79.2
2008	30	65.2	12.2	32.3	88.0	47.9	60.2	67.3	72.3	79.6
2009	26	61.4	9.5	30.8	76.7	52.2	55.8	62.6	69.0	70.6
2010	29	63.6	14.7	15.6	82.0	43.8	54.6	69.3	72.7	81.7
2011	40	65.1	11.8	34.4	83.0	50.0	57.1	66.4	74.6	80.7
2012	35	66.5	9.8	44.0	85.1	55.5	58.9	67.8	70.8	81.9
2013	32	66.8	10.1	36.7	82.5	53.7	61.1	68.2	73.8	79.2
1998-2013	323	62.8	13.3	15.6	88.0	44.5	55.5	65.0	71.5	79.2

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of	Cases	S	Std.					Median		
diagnosis	n	Mean o	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	6	62.8	4,7	56.6	68.7	56.6	59.9	61.8	67.8	68.7
1999	12	65.2	14.8	32.9	81.2	47.0	56.8	71.6	77.1	78.4
2000	13	59.5	19.0	15.8	80.5	29.3	52.5	67.2	72.2	74.9
2001	7	55.6	12.2	38.0	69.3	38.0	41.3	55.5	66.9	69.3
2002	13	59.8	15.3	31.0	76.2	33.6	51.4	65.9	72.8	75.6
2003	13	63.7	9.8	43.9	79.5	49.2	60.4	63.8	72.7	73.0
2004	15	59.3	16.2	27.6	84.1	29.9	47.8	64.9	69.4	73.2
2005	12	52.9	17.7	21.6	82.9	32.2	41.6	52.3	62.3	77.4
2006	12	54.0	15.8	27.5	78.6	30.4	43.8	56.6	63.9	74.3
2007	17	58.3	18.7	22.3	84.2	29.1	43.9	59.7	72.9	81.5
2008	31	62.6	12.3	29.4	79.9	43.3	56.3	64.3	71.7	75.8
2009	31	63.4	13.8	25.9	89.3	47.2	55.6	62.7	72.7	79.5
2010	21	62.3	8.9	39.9	75.8	51.2	60.0	65.3	66.4	71.8
2011	36	62.1	12.3	33.0	85.6	47.3	54.1	61.6	69.0	80.4
2012	32	66.2	11.5	39.5	89.1	51.2	56.6	69.0	74.3	76.9
2013	29	65.8	10.7	45.0	85.4	51.2	58.8	68.1	73.2	78.9
1998-2013	300	61.9	13.6	15.8	89.3	42.9	54.7	63.3	71.8	77.2

Table 4

Age distribution by 5-year age group and gender for period 1998-2013 (incl. DCO)

Age at									
diagnosis	Cases			Males			Females		
Years	n	왕	Cum.%	'n	%	Cum.%	n	%	Cum.%
15-19	5	0.8	0.8	4	1.2	1.2	1	0.3	0.3
20-24	3	0.5	1.3	/ 1	0.3	1.5	2	0.7	1.0
25-29	10	1.6	2.9	3	0.9	2.5	7	2.3	3.3
30-34	12	1.9	4.8	5	1.5	4.0	7	2.3	5.7
35-39	13	2.1	6.9	8	2.5	6.5	5	1.7	7.3
40 - 44	26	4.2	11.1	13	4.0	10.5	13	4.3	11.7
45-49	29	4.7	15.7	11	3.4	13.9	18	6.0	17.7
50-54	56	9.0	24.7	31	9.6	23.5	25	8.3	26.0
55-59	73	11.7	36.4	35	10.8	34.4	38	12.7	38.7
60-64	101	16.2	52.6	51	15.8	50.2	50	16.7	55.3
65-69	98	15.7	68.4	55	17.0	67.2	43	14.3	69.7
70-74	102	16.4	84.8	57	17.6	84.8	45	15.0	84.7
75-79	53	8.5	93.3	24	7.4	92.3	29	9.7	94.3
80-84	35	5.6	98.9	23	7.1	99.4	12	4.0	98.3
85+	7	/ 1.1	100.0	2	0.6	100.0	5	1.7	100.0
All ages	623	100.0		323	100.0		300	100.0	

Included in the statistics are 31.0% multiple primaries in males and 32.1% in females.



Table 5

Age-specific incidence, DCO rate and proportion of all cancers for period 1998-2013

							Males	Females
			Males	Females	Males	Females		Prop.all
Age at				Age-		DCO rate	_	cancers
diagnosis	Males	Females		spec.	n=0	n=0		n=153136
Years	n	n		incid.	%	%	%	%
icars	11	11	incia.	filera.	0	Ů	0	• /
0- 4			0.0	0.0				
5- 9			0.0	0.0				
10-14			0.0	0.0				
15-14	4	1	0.0	0.0			1.1	0.3
	1	2	0.3	0.1			0.2	
20-24	3	2 7						0.4
25-29		7	0.1	0.3			0.3	0.6
30-34	5		0.2	0.3			0.3	0.3
35-39	8	5	0.3	0.2			0.4	0.1
40-44	13	13	0.5	0.5			0.4	0.2
45-49	11	18	0.5	0.8			0.2	0.2
50-54	31	25	1.5	1.2			0.4	0.2
55-59	35	38	1.9	2.0			0.2	0.3
60-64	51	50	2.9	2.7			0.2	0.3
65-69	55	43	3.5	2.5			0.2	0.2
70-74	57	44	4.4	2.9			0.2	0.2
75-79	24	29	2.9	2.4			0.1	0.2
80-84	23	12	4.6	1.3			0.2	0.1
85+	2	5	0.6	0.6			0.0	0.0
All ages	323	299			0.0	0.0	0.2	0.2
Incidence								
Raw			1.1	1.0				
WS			0.6	0.6				
ES			0.9	0.8				
BRD-S			1.0	0.9				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

DCO

Table 6a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2013

MALES

Diagnosis	Observed :	Expected n	SIR	LCL UCL 95% 95%	EAR
C16 Stomach C33-C34 Lung C61 Prostate C67 Bladder	2 4 2 2	0.4 1.1 2.6 0.4	5.5 3.7 0.8 5.2	0.7 20.0 1.0 9.5 # 0.1 2.8 0.6 18.7	24.7 44.2 -9.3 24.4
Other primaries Not observed	6	2.2	2.8	1.0 6.0 # 0.0 1.5	57.9 -37.7
All mult. primaries Patients	16	9.1	212 71.9	1.0 2.9 #	104.1
Median age at second ma Person-years Mean observation time (663 3.1		

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries".


Mean observation time (years)
Median observation time (years)

Table 6b

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2013

FEMALES

	Observed	Expected		LCL	UCL		DCO
Diagnosis	n	n	SIR	95%	95%	EAR	ઇ
C33-C34 Lung	3	0.6	5.4	1.1	15.9 #	32.8	
C50 Breast	4	2.4	1.7	0.5	4.4	22.1	
Other primaries	/5	1.6	3.0	1.0	7.1	45.0	
Not observed	0	2.8	0.0	0.0	1.3	-37.0	
All mult. primaries	12	7.3	1.6	0.8	2.9	62.9	



The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries".






Figure 7. Age distribution and age-specific incidence



BNET: Bronchopulmonary neuroendocrine tumor (excl. SCLC) Age-specific incidence in international comparison




Figure 7a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).



Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2014, based on the November 2013 submission. http://www.seer.cancer.gov.





Figure 8. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.



Average incidence (world standard population) 2007 - 2013: Males



Average incidence (world standard population) 2007 - 2013: Females





Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2007 to 2013. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.8/100,000 WS N=216, females 0.7/100,000 WS N=197).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,928 female residents (averaged) in the period from 2007 to 2013 a total of 7 women were identified with newly diagnosed pulm. neuroend. tumor. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 1.2/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.3 and 3.1/100,000.

Standardized incidence ratio (SIR) 2007 - 2013: Males



Standardized incidence ratio (SIR) 2007 - 2013: Females




Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2013. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=216, females N=197).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,642 female residents (averaged) in the period from 2007 to 2013 a total of 7 women were identified with newly diagnosed pulm. neuroend. tumor. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 1.30. Though, the value of this parameter may vary with an underlying probability of 99% between 0.38 and 3.19, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts, and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

					Prop.
	Trondalone	Prop.		Decem	deaths
_	Incident	actively	_ /_	Prop.	with death
Year of	cases	followed	Deaths	deaths	certific.
diagnosis	n	%	n	%	%
1998	12	83.3	6	50.0	100.0
1999	23	95.7	14	60.9	100.0
2000	21	100.0	11	52.4	90.9
2001	21	95.2	13	61.9	100.0
2002	25	96.0	10	40.0	90.0
2003	22	100.0	11	50.0	72.7
2004	29	93.1	11	37.9	100.0
2005	28	89.3	9	32.1	100.0
2006	29	89.7	8	27.6	100.0
2007	41	82.9	21	51.2	100.0
2008	61	72.1	27	44.3	96.3
2009	57	61.4	21	36.8	90.5
2010	50	64.0	23	46.0	100.0
2011	76	72.4	31	40.8	96.8
2012	67	76.1	27	40.3	96.3
2013	61	96.7	14	23.0	71.4
1998-2013	623	81.4	257	41.3	94.6

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

			Prop. deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in /
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	n	%	n	%
1998	12	5	100.0	2	16.7
1999	23	8	87.5	/ 1	4.3
2000	21	9	100.0	2	9.5
2001	21	6	83.3	1	4.8
2002	25	12	100.0	5	20.0
2003	22	9	100.0	3	13.6
2004	29	10	90.0	2	6.9
2005	28	15	93.3	3	10.7
2006	29	16	100.0	2	6.9
2007	41 /	17	94.1	6	14.6
2008	61	25	100.0	10	16.4
2009	57	25	96.0	10	17.5
2010	50	28	96.4	7	14.0
2011	76	34	100.0	12	15.8
2012	67	46	97.8	15	22.4
2013	61	35	97.1	10	16.4
1998-2013	623	300	97.0	91	14.6

Table 10c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

				Prop.
				cancer
		Prop.	Prop.	recorded
		cancer-	non-cancer-	on death
Year of	Deaths	related	related	certificate
death	n	%	8	%
1998	5	100.0		80.0
1999	8	75.0	25.0	100.0
2000	9	77.8	22.2	88.9
2001	6	83.3	16.7	80.0
2002	12	83.3	16.7	91.7
2003	9	88.9	11.1	88.9
2004	10	80.0	20.0	88.9
2005	15	73.3	26.7	85.7
2006	/ 16	68.8	31.3	87.5
2007	17	70.6	29.4	81.3
2008	25	80.0	20.0	80.0
2009	25	92.0	8.0	95.8
2010	28	85.7	14.3	92.6
2011	34	85.3	14.7	85.3
2012	46	91.3	8.7	91.1
2013	35	91.4	8.6	94.1
1998-2013	300	84.3	15.7	89.0

Table 11a $\begin{tabular}{ll} Medians of age at death according to the grouping in Table 10 \\ \hline MALES \end{tabular}$

Year of death	Deaths n	Age at death (all causes)	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate)
1998	3	71.0	71.0		64.7
1999	4	69.7	69.7		69.7
2000	3	77.7	77.7		77.7
2001	4	60.7	56.1	65.3	62.3
2002	9	68.0	69.1	65.9	68.5
2003	7	71.3	70.0	71.3	70.0
2004	5	57.1	57.1		57.1
2005	10	70.7	71.0	70.7	71.0
2006	13	71.8	75.2	56.1	71.8
2007	11/	67.1	66.9	67.1	66.9
2008	18	71.7	70.6	74.2	70.6
2009	18	68.5	67.7	75.7	68.5
2010	18	66.3	67.3	65.3	67.7
2011	23	73.3	71.8	87.6	72.4
2012	29	68.7	68.0	83.5	68.3
2013	24	71.3	72.0	69.2	70.6
1998-2013	199	69.6	69.4	71.6	69.4

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 11b $\begin{tabular}{ll} \begin{tabular}{ll} \begin{tabula$

		Age at death (all	Age at death (cancer-	Age at death (non-cancer-	Age at death (according to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	2	74.6	74.6		74.6
1999	4	72.0	77.2	59.5	73.0
2000	6	78.6	76.6	78.7	79.0
2001	2	58.7	58.7		58.7
2002	3	72.9	72.9		72.9
2003	2	73.2	73.2		73.2
2004	5	74.9	73.8	75.5	76.2
2005	5 3	82.2	82.2	78.2	82.6
2006	3	80.4		80.4	80.4
2007	6/	71.6	64.2	76.1	67.4
2008	7	77.3	72.4	84.1	72.4
2009	/7	72.8	66.4	86.6	72.8
2010	10	76.2	73.2	80.6	73.2
2011	11	71.4	70.9	78.2	71.4
2012	17	70.3	69.7	87.3	69.7
2013	11	71.3	71.3	74.3	71.3
1998-2013	101	73.8	71.8	78.7	73.0

By 2010, life expectancy for a newborn male in Germany is 77.5 years compared with 82.6 years for his female counterpart.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

base hBNETE.pdf

Table 12a Mortality measures (cancer-related death) and mortality-incidence-index by year of death MALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	3	0.3	0.50	0.2	0.40	0.2	0.44	0.3	0.50
1999	4	0.4	0.36	0.2	0.32	0.3	0.36	0.4	0.36
2000	3	0.3	0.38	0.1	0.30	0.2	0.36	0.4	0.58
2001	3	0.3	0.21	0.2	0.19	0.2	0.22	0.3	0.21
2002	7	0.4	0.58	0.2	0.55	0.3	0.54	0.4	0.54
2003	6	0.3	0.67	0.2	0.62	0.3	0.69	0.4	0.73
2004	5	0.3	0.36	0.2	0.42	0.3	0.44	0.3	0.39
2005	8	0.4	0.50	0.2	0.33	0.3	0.41	0.5	0.54
2006	11	0.6	0.65	0.3	0.54	0.4	0.67	0.6	0.77
2007	8	0.4	0.33	0.2	0.29	0.3	0.30	0.3	0.30
2008	16	0.7	0.53	0.3	0.45	0.5	0.50	0.7	0.59
2009	17	0.8	0.65	0.4	0.55	0.6	0.56	0.7	0.63
2010	15	0.7	0.52	0.4	0.50	0.5	0.52	0.6	0.49
2011	19	0.8	0.48	0.4	0.42	0.6	0.43	0.8	0.49
2012	26	1.1	0.74	0.6	0.68	0.9	0.72	1.1	0.77
2013	23	1.0	0.72	0.5	0.64	0.7	0.67	0.9	0.73
1998-2013	174	0.6	0.54	0.3	0.47	0.5	0.51	0.6	0.56

Table 12b Mortality measures (cancer-related death) and mortality-incidence-index by year of death

FEMALES

Year of Deaths Mort. MI-Index Mort. MI-Index Mort. MI-Index n raw raw WS WS ES ES BRD-S BRD-S death 0.24 0.1 0.27 0.2 0.38 1998 2 0.2 0.33 0.1 0.18 0.1 2 0.1 0.15 0.2 0.19 1999 0.2 0.13 2000 4 0.1 0.15 0.3 0.28 0.3 0.31 0.2 0.20 2 2001 0.2 0.29 0.1 0.33 0.2 0.32 0.2 0.35 0.1 0.1 0.16 0.1 0.10 0.1 0.10 0.14 2002 3 0.2 0.23 0.1 0.23 2003 2 0.1 0.15 0.0 0.09 0.1 0.13 0.1 0.14 2004 3 0.2 0.20 0.1 0.11 0.1 0.17 0.2 0.25 2005 3 $0.0 \quad 0.12$ 0.1 0.14 0.1 0.19 0.24 0.1 0.1 2006 2007 4 0.2 0.21 0.1 0.22 0.1 0.22 2008 4 0.2 0.13 0.09 0.1 0.10 0.1 0.12 2009 6 0.3 0.19 0.1 0.17 0.2 0.17 0.2 0.17 2010 9 0.4 0.43 0.1 0.27 0.2 0.30 0.3 0.34 2011 10 0.4 0.28 0.2 0.19 0.2 0.21 0.3 0.25 2012 16 0.7 0.50 0.3 0.43 0.4 0.45 0.5 0.48 2013 9 0.4 0.31 0.2 0.25 0.2 0.27 0.3 0.29 1998-2013 79 0.3 0.26 0.1 0.19 0.2 0.21 0.2 0.24

Table 13

Age distribution of age at death (cancer-related) for period 1998-2013

(incl. multiple primaries)

Age at									
death	Cases			Males			Females		
Years	n	ે	Cum.%	n	્ર	Cum.%	n	ે	Cum.%
40-44	6	2.4	2.4	4	2.3	2.3	2	2.5	2.5
45-49	6	2.4	4.7	4	2.3	4.6	2	2.5	5.1
50-54	13	5.1	9.9	10	5.7	10.3	3	3.8	8.9
55-59	22	8.7	18.6	17	9.8	20.1	5	6.3	15.2
60-64	28	11.1	29.6	20	11.5	31.6	8	10.1	25.3
65-69	48	19.0	48.6	36	20.7	52.3	12	15.2	40.5
70-74	51	20.2	68.8	34	19.5	71.8	17	21.5	62.0
75-79	33	13.0	81.8	24	13.8	85.6	9	11.4	73.4
80-84	35	13.8	95.7	20	11.5	97.1	15	19.0	92.4
85+	11	4.3	100.0	5	2.9	100.0	6	7.6	100.0
All ages	253	100.0		174	100.0		79	100.0	

Included in the statistics are 31.0% multiple primaries in males and 32.1% in females.

Table 14

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2013 (incl. multiple primaries)

Age at			Males Age-		Females Age-		Males Prop.all	Females Prop.all
death	Males Fen		spec.		spec.		cancers	cancers
Years	n	n mo	ortal.	MI-index	mortal.	MI-index	%	%
0 - 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29 30-34			0.0		0.0			
35-39			0.0		0.0			
40-44	4	2	0.0 0.2	0.31	0.0	0.15	0.5	0.2
45-49	4	2	0.2	0.31	0.1	0.15	0.5	0.2
50-54	10	3	0.2		0.1		0.2	0.1
55-59	17	5	0.9		0.1	0.12	0.3	0.1
60-64	20	8	1.1	0.39	0.4		0.3	0.1
65-69	36	12	2.3	0.65	0.7		0.3	0.1
70-74	34	17	2.7		1.1	0.38	0.3	0.2
75-79	24	9	2.9		0.8	0.31	0.2	0.1
80-84	20	15	4.0	0.87	1.6		0.2	0.1
85+	5	6	1.5	2.50	0.7		0.1	0.0
All ages	174	79					0.2	0.1
Mortality								
Raw			0.6	0.54	0.3	0.26		
WS			0.3	0.47	0.1	0.19		
ES			0.5	0.51	0.2			
BRD-S			0.6	0.56	0.2	0.24		
DVII 70								
PYLL-70			3.1		1.1			
per 100,000 ES			2.7		1.1			
ES AYLL-70			9.1		9.5			
AITT-\0			۶.1		9.5			

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15a

Multiple primaries in deaths in period 1998-2013

MALES

					_	_		
					Syn-	Syn-		
	/_			`	chron	chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n	%↓	n	← %	n	← %	n	← %
	/ _	/						
C03-C06 Oral cavity	/ 2	3.8	2	100.0				
C09-C10 Oropharynx	2	3.8	1	50.0			1	50.0
C15 Oesophagus	/ 1 /	1.9	1	100.0				
C16 Stomach	3	5.7	1	33.3	1	33.3	1	33.3
C17 Small intestine	1	1.9			/ 1	100.0		
C18 Colon	2	3.8	1	50.0	1	50.0		
C19-C20 Rectum	2	3.8	2	100.0				
C22 Liver	2	3.8			1	50.0	1	50.0
C33-C34 Lung	10	18.9			4	40.0	6	60.0
C43 Malign. melanoma	3	5.7	3	100.0				
C44 Skin others	4	7.5	3	75.0			1	25.0
C48 Peritoneal	1	1.9					1	100.0
C50 Breast	1	1.9	1	100.0				
C61 Prostate	5	9.4	3	60.0			2	40.0
C64 Kidney	1	1.9	1	100.0				
C66 Ureter	1	1.9					1	100.0
C67 Bladder	10	18.9	6	60.0	1	10.0	3	30.0
C70-C72 CNS cancer	1	1.9					1	100.0
C81 Hodgkin lymphoma	1	1.9	1	100.0				
3 7 4								
All mult. primaries	53	100.0	26	49.1	9	17.0	18	34.0

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 15b

Multiple primaries in deaths in period 1998-2013

FEMALES

					Syn-	Syn-		
					chron	chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n	%↓	n	← %	n	~ %	n	← %
C07-C08 Salivary gland	/ 1	2.3	1	100.0				
C09-C10 Oropharynx	/ 1	2.3	1	100.0				
C16 Stomach	/ 1 /	2.3			1	100.0		
C18 Colon	4	9.1	1	25.0	1	25.0	2	50.0
C32 Larynx	1	2.3	1	100.0				
C33-C34 Lung	6	13.6			3	50.0	3	50.0
C43 Malign. melanoma	1	2.3	1	100.0				
C50 Breast	10	22.7	9	90.0			1	10.0
C51 Vulva	1	2.3					1	100.0
C53 Cervix uteri	2	4.5			2	100.0		
C54 Corpus uteri	1	2.3	1	100.0				
C56 Ovary	1	2.3	1	100.0				
C64 Kidney	3	6.8	2	66.7	_ 1	33.3		
C67 Bladder	1	2.3	1	100.0				
C70-C72 CNS cancer	4	9.1	1	25.0	1	25.0	2	50.0
C73 Thyroid	1	2.3			1	100.0		
C74-C80 Cancer others	1	2.3	1	100.0				
C90 Mult. myeloma	3	6.8	2	66.7			1	33.3
C91-C96 Leukaemia	1	2.3					1	100.0
All mult. primaries	44	100.0	23	52.3	10	22.7	11	25.0




Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2013

(Singular primaries only *)

			Males		Females		Males	Females
Age at			Age-		Age-		_	Prop.all
death	Males	Females	/ - /		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0 - 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29			0.0		0.0			
30-34			0.0		0.0			
35-39			0.0		0.0			
40-44	4	1	0.2	0.36	0.0	0.08	0.5	0.1
45-49	4	2	0.2	0.40	0.1	0.11	0.2	0.1
50-54	8	3	0.4	0.27	0.1	0.14	0.3	0.1
55-59	17	5	0.9	0.59	0.3	0.17	0.3	0.1
60-64	16	4	0.9	0.41	0.2	0.11	0.2	0.1
65-69	33	7	2.1	0.67	0.4	0.19	0.3	0.1
70-74	25	10	2.0	0.69	0.7		0.2	0.1
75-79	17	7	2.1		0.6		0.2	0.1
80-84	11	12	2.2		1.3		0.1	0.1
85+	3	3	0.9		0.3		0.0	0.0
All ages	138	54					0.2	0.1
5								
Mortality								
Raw			0.5	0.54	0.2	0.23		
WS			0.3		0.1			
ES			0.4		0.1			
BRD-S			0.5	0.56	0.1			
DKD 5			0.5	0.50	0.1	0.21		
PYLL-70								
per 100,000			2.8		0.9			
ES ES			2.5		0.9			
AYLL-70			9.2		10.7			
AILL-/U			9.4		10.7			

^{*} See corresponding tables with multiple primaries.

Table 17

Age-specific mortality (cancer-related) and proportion of all cancers for period 1998-2013

(Single primaries only *)

Age at death	Males	Females			Females Age- spec.		cancers	Females Prop.all cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0- 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29			0.0		0.0			
30-34			0.0		0.0			
35-39			0.0		0.0			
40-44	4	1	0.2	0.36	0.0	0.08	0.5	0.1
45-49	3	2	0.1	0.30	0.1	0.13	0.2	0.1
50-54	8	2	0.4	0.28	0.1	0.11	0.3	0.1
55-59	16	5	0.9	0.55	0.3	0.18	0.3	0.1
60-64	13	3	0.7	0.37	0.2	0.09	0.2	0.1
65-69	33	/ 5	2.1	0.73	0.3	0.16	0.4	0.1
70-74	23	9	1.8	0.68	0.6	0.32	0.3	0.1
75-79	15	7	1.8	0.88	0.6	0.39	0.2	0.1
80-84	10	11	2.0	1.00	1.2	1.00	0.2	0.1
85+	3	3	0.9	3.00	0.3	0.75	0.1	0.0
All ages	128	48					0.2	0.1
Mortality								
Raw			0.4		0.2	0.22		
WS			0.2		0.1	0.15		
ES			0.3	0.50	0.1	0.17		
BRD-S			0.4	0.54	0.1	0.20		
D								
PYLL-70					0 0			
per 100,000			2.6		0.8			
ES			2.3		0.7			
AYLL-70			9.1		11.4			

^{*} See corresponding tables with multiple primaries.





Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at pulm. neuroend. tumor-related death (see Table 10) should be considered.



Average mortality (world standard population) 2007 - 2013: Males



Average mortality (world standard population) 2007 - 2013: Females





Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2007 to 2013. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.4/100,000 WS N=123, females 0.2/100,000 WS N=57).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,928 female residents (averaged) in the period from 2007 to 2013 a total of 1 women died from pulm. neuroend. tumor. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.1/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 0.9/100,000.

Standardized mortality ratio (SMR) 2007 - 2013: Males



Standardized mortality ratio (SMR) 2007 - 2013: Females




Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2013. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=123, females N=57).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,642 female residents (averaged) in the period from 2007 to 2013 a total of 1 women died from pulm. neuroend. tumor. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.66. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 4.90, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

FRG Federal Republic of Germany

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

MCR Munich Cancer Registry (Tumorregister München)
SEER Surveillance, Epidemiology, and End Results (USA)

AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

BRD-S German standard population

DCO Death certificate only EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

ES European standard population (old)

LCL Lower confidence limit

MI-index Ratio between mortality and incidence

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age

SIR Standardized incidence ratio SMR Standardized mortality ratio UCL Upper confidence limit WS World standard population

Recommended Citation

Munich Cancer Registry. Baseline statistics BNET: Pulm. neuroend. tumor [Internet]. 2015 [updated 2015 May 19; cited 2015 Jul 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/base/base_hBNETE.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

Fig./Tbl	l.	Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	4
1a	Gender distribution by year of diagnosis	5
2	Incidence by year of diagnosis	6
3	Age distribution parameters by year of diagnosis	7
4	Age distribution by 5-year age group and gender	9
5	Age-specific incidence and DCO rate	10
6	Standardized incidence ratio of second primaries	11
7	Age distribution and age-specific incidence (chart)	13
7a	Age-specific incidence internationally (chart)	14
8	Cumulative follow-up years (chart)	15
9a	Map of cancer incidence (WS) by county (chart)	16
9b	Standardized incidence ratio (SIR) by county (chart)	17
10a	Pts incident cohorts and mortality / yr	18
10b	Incidence and mortality by year of diagnosis	19
10c	Cancer-related deaths, death certification available / yr	20
11	Medians of age at death / yr	21
12	Mortality by year of death	23
13	Distribution of age at death	24
14	Age-specific mortality	25
15	Multiple primaries in deaths	26
16	Age-specific mortality (first primaries)	28
17	Age-specific mortality (single primaries)	29
18	Age distribution and age-specific mortality (chart)	30
19a	Map of cancer mortality (WS) by county (chart)	31
19b	Standardized mortality ratio (SMR) by county (chart)	32