Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ Deutsch

ICD-10 C02-C06: Oral cavity cancer

Incidence and Mortality

Year of diagnosis	1998-2014
Patients	3,015
Diseases	3,038
Creation date	04/13/2016
Export date	12/23/2015
Population	4.64 m



Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

http://www.tumorregister-muenchen.de/en/facts/base/bC0206E-ICD-10-C02-C06-Oral-cavity-cancer-incidence-and-mortality.pdf

Global Statements about the statistics on the Internet – Baseline Statistics (grey button ——), Survival (red button ——)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.64 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases**** are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, April 2016

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007).
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate.

Some remarks regarding this cancer type

As a general rule, these few results from the TRM form the basis of sophisticated analyses. For head and neck tumors this is not the case. Therefore the results for head and neck tumors should be interpreted with caution. In part this is due to problems of classification because of limited specific details of locality. Additionally, with advanced tumors in a close topographic location it is often not possible to determine the exact ICD localization of a tumor.

ICD-10 codes (ICD-10 2015) used for specifying cancer site

Code	Description
C02	Other and unspecified parts of tongue Excl.: Topography code C02.4 Lingual tonsil
C03	Gum
C04	Floor of mouth
C05	Palate Excl.: Topography code C05.1 Soft palate, NOS C05.2 Uvula
C06	Other and unspecified parts of mouth

INCIDENCE

Table 1

All patients with invasive cancer by year of diagnosis, proportions of DCO, multiple primaries, deaths, and active follow-up (incl. DCO)

				Prop.		Prop.
		/DCO	Prop.	mult.	Prop.	actively
Year of	Cases	cases	DCO	primaries	deaths	followed
diagnosis	n	n	90	%	9	િ
1998	123	5	4.1	32.5	79.7	99.2
1999	129	2	1.6	41.9	79.8	96.1
2000	132	4	3.0	31.8	75.0	99.2
2001	136	6	4.4	33.8	77.2	97.8
2002	190	12	6.3	35.8	71.1	97.9 #
2003	191	9	4.7	35.6	69.1	97.4
2004	193	6	3.1	34.7	71.0	97.4
2005	161	6	3.7	29.8	68.3	96.9
2006	195	3	1.5	32.3	65.6	93.8
2007	223	9	4.0	30.5	63.2	81.2 #
2008	216	3	1.4	31.9	53.7	74.1
2009	232	4	1.7	29.3	53.0	78.4
2010	251	12	4.8	27.9	50.6	74.1
2011	182	5	2.7	29.1	41.8	76.4
2012	226	7	3.1	26.1	36.7	73.9
2013	215	6	2.8	24.2	33.0	100.0
2014	43	9	20.9	41.9	37.2	100.0 ##
1998-2014	3038	108	3.6	31.4	59.2	88.3

[#] The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

^{##} Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be found in the respective headings.

Table 1a

All patients with invasive cancer by year of diagnosis and gender (incl. DCO)

Year of	All	Males	Females	Prop. males	
diagnosis	n/	n	n	96	
1998	123	87	36	70.7	
1999	129	80	49	62.0	
2000	132	105	27	79.5	
2001	/ 136	93	43	68.4	
2002	190	126	64	66.3	
2003	191	135	56	70.7	
2004	193	138	55	71.5	
2005	161	104	57	64.6	
2006	195	129	66	66.2	
2007	223	150	73	67.3	
2008	216	142	74	65.7	
2009	232	154	78	66.4	
2010/	251	171	80	68.1	
2011	182	110	72	60.4	
2012	226	143	83	63.3	
2013	215	141	74	65.6	
2014	43	27	16	62.8	
1998-2014	3038	2035	1003	67.0	

Table 2

Incidence measures by year of diagnosis including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

			Males	Fem.	Males	Fem.	Males	Fem.	Males	Fem.
Year of	Males	Females	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	87	36	7.9	3.1	5.3	1.6	7.2	2.3	8.0	2.8
1999	80	49	7.1	4.1	4.7	2.4	6.5	3.3	7.1	3.7
2000	105	27/	9.2	2.2	6.2	/1.3	8.4	1.8	9.1	2.0
2001	93	43	8.0	3.5	5.1	1.9	7.1	2.7	8.0	3.0
2002	126	64	6.8	3.3	4.4	1.7	5.9	2.5	6.5	2.8
2003	135	56	7.2	2.8	4.8	1.6	6.5	2.2	7.1	2.5
2004	138	55	7.3	2.8	4.7	1.3	6.5	1.9	7.2	2.4
2005	104	57	5.5	2.9	3.4	1.6	4.6	2.2	5.2	2.5
2006	129	66	6.7	3.3	4.2	1.8	5.9	2.5	6.8	2.9
2007	150	73	6.8	3.2	4.3	1.7	5.9	2.3	6.5	2.8
2008	142	74	6.4	3.2	3.9	1.8	5.4	2.5	6.1	2.8
2009	154	78	6.9	3.4	4.2	1.8	5.8	2.4	6.5	2.9
2010	171	80	7.6	3.4	4.7	1.7	6.4	2.4	7.0	2.7
2011	110	72	4.8	3.1	2.9	1.5	4.0	2.2	4.5	2.5
2012	143	83	6.3	3.5	3.8	1.9	5.2	2.6	5.8	3.0
2013	141	74	6.2	3.1	3.7	1.5	5.0	2.1	5.7	2.5
2014	27	16	1.2	0.7	0.7	0.3	1.0	0.4	1.1	0.5
1998-2014	2035	1003	6.4	3.0	4.0	1.6	5.5	2.2	6.1	2.6




Table 3

Age distribution parameters by year of diagnosis (All patients) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	123	59.1	12.2	0.9	91.4	46.4	51.8	57.6	66.3	76.3
1999	129	60.8	12.7	25.6	91.9	47.1	52.8	59.1	66.9	77.7
2000	132	58.7	11.4	33.5	85.8	45.0	50.1	58.1	66.6	73.1
2001	136	61.9	12.2	33.7	96.4	45.7	53.6	60.7	69.3	78.0
2002	190	61.1	12.4	26.4	99.0	45,4	53.0	60.9	68.0	78.7
2003	191	60.1	12.8	10.7	98.2	45.8	52.2	59.3	66.7	78.7
2004	193	61.9	12.9	29.5	97.9	45.5	53.4	61.7	70.4	79.8
2005	161	61.0	12.9	22.8	98.7	45.5	52.3	60.8	67.7	80.8
2006	195	62.7	12.9	22.6	96.2	47.5	54.9	61.4	71.7	81.2
2007	223	61.9	12.8	26.0	101	46.0	53.7	61.7	70.4	78.1
2008	216	62.5	11.7	21.8	100	48.8	54.1	62.4	69.5	77.6
2009	232	62.7	12.4	29.6	98.4	47.6	54.0	62.8	71.0	79.9
2010	251	62.4	13.2	21.9	92.8	46.9	52.3	62.0	70.7	81.5
2011	182	62.4	13.8	27.0	96.9	43.5	53.1	63.0	72.0	78.8
2012	226	62.4	12.5	21.5	100	48.0	54.0	63.8	70.9	77.4
2013	215	64.3	12.6	28.1	95.5	48.3	55.7	64.6	72.5	81.1
2014	43	66.5	12.3	41.3	90.0	51.1	58.4	66.4	75.2	82.2
1998-2014	3038	61.9	12.7	0.9	101	46.5	53.3	61.3	70.0	78.9

Table 3a

Age distribution parameters by year of diagnosis (MALES)

(incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	87	57.0	11.5	0.9	81.3	45.5	51.1	56.2	62.3	72.0
1999	80	59.2	12.1	33.3	90.8	46.8	51.2	57.4	64.2	80.2
2000	105	58.0	10.1	35.8	85.5	45.1	50.1	58.1	65.0	72.0
2001	93	59.9	12.1	33.7	94.3	44.5	51.2	59.7	64.3	77.4
2002	126	59.0	10.7	26.4	92.2	45.2	52.3	60.1	64.5	72.1
2003	135	58.7	10.4	28.1	86.1	46.0	52.9	57.8	64.4	71.9
2004	138	59.4	11.6	29.7	88.7	44.9	51.9	59.7	65.4	75.0
2005	104	59.1	11.6	36.8	85.0	43.3	49.7	58.3	66.7	77.2
2006	129	61.5	12.2	23.9	92.0	46.7	53.8	59.6	69.4	78.2
2007	150	60.3	11.6	26.0	101	46.0	52.5	59.8	67.7	75.1
2008	142	61.6	11.2	21.8	100	48.5	53.5	61.9	69.0	75.1
2009	154	62.0	10.8	30.2	88.1	48.0	54.5	62.3	69.7	74.7
2010	171	60.3	12.3	24.5	92.8	45.4	51.8	59.7	69.0	75.3
2011	110	60.2	13.1	27.0	93.0	43.5	52.6	58.7	69.7	78.0
2012	143	61.3	11.3	21.6	85.9	48.0	52.7	62.3	69.6	75.1
2013	141	62.2	10.7	30.0	85.0	49.3	54.7	62.2	70.0	75.8
2014	27	63.7	12.2	41.3	90.0	43.9	55.3	63.5	72.4	80.1
1998-2014	2035	60.2	11.5	0.9	101	46.0	52.5	59.7	67.7	75.2

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	36	64.1	12.8	32.0	91.4	49.6	56.4	63.2	75.8	79.5
1999	49	63.3	13.4	25.6	91.9	47.1	56.1	65.4	72.2	77.7
2000	27	61.3	15.4	33.5	85.8	39.8	50.6	59.6	76.3	84.8
2001	43	66.4	11.3	44.0	96.4	53.2	59.8	63.7	71.3	84.0
2002	64	65.3	14.3	35.8	99.0	47,0	54.5	62.5	76.2	82.9
2003	56	63.4	16.8	10.7	98.2	44.8	51.1	62.3	78.2	83.7
2004	55	68.4	13.9	29.5	97.9	48.9	58.2	69.0	78.9	84.3
2005	57	64.5	14.4	22.8	98.7	50.2	54.9	62.3	76.0	83.7
2006	66	65.2	14.1	22.6	96.2	48.2	56.0	63.1	77.0	84.2
2007	73	65.1	14.6	31.0	98.2	46.1	55.3	65.0	75.2	83.6
2008	74	64.3	12.4	26.7	91.5	49.7	55.4	64.1	72.5	79.4
2009	78	64.1	15.1	29.6	98.4	43.1	53.7	65.0	75.2	82.8
2010	80	66.8	14.0	21.9	91.8	50.1	56.2	66.9	76.5	87.0
2011	72	65.7	14.2	31.2	96.9	47.7	57.7	67.3	75.1	84.1
2012	83	64.2	14.3	21.5	100	47.5	56.5	64.9	73.0	82.1
2013	74	68.3	14.9	28.1	95.5	46.8	58.7	67.9	79.9	88.9
2014	16	71.4	11.4	43.4	88.4	55.0	65.4	72.6	79.1	83.0
1998-2014	1003	65.3	14.2	10.7	100	47.5	55.8	65.0	75.4	83.7

Table 4

Age distribution by 5-year age group and gender for period 2007-2014 (incl. DCO)

Age at									
diagnosis	Cases			Males			Females		
Years	n	્રે	Cum.%	n	양	Cum.%	n	olo	Cum.%
20-24	5	0.3	0.3	3	0.3	0.3	2	0.4	0.4
25-29	10	0.6	0.9	6	0.6	0.9	4	0.7	1.1
30-34	19	1.2	2.1/	8	0.8	1.6	11	2.0	3.1
35-39	14	0.9	3.0	10	1.0	2.6	4	0.7	3.8
40 - 44	60	3.8	6.8	41	3.9	6.6	19	3.5	7.3
45-49	135	8.5	15.3	98	9.4	16.0	37	6.7	14.0
50-54	197	12.4	27.7	155	14.9	30.9	42	7.6	21.6
55-59	229	14.4	42.1	168	16.2	47.1	61	11.1	32.7
60-64	233	14.7	56.8	155	14.9	62.0	78	14.2	46.9
65-69	242	15.2	72.0	164	15.8	77.8	78	14.2	61.1
70 - 74	191	12.0	84.1	118	11.4	89.2	73	13.3	74.4
75-79	110	6.9	91.0	62	6.0	95.2	48	8.7	83.1
80-84	71	4.5	95.5	27	2.6	97.8	44	8.0	91.1
85+	72	4.5	100.0	23	2.2	100.0	49	8.9	100.0
All ages	1588	100.0		1038	100.0		550	100.0	

Included in the statistics are 41.0% multiple primaries in males and 29.4% in females.



Table 5

Age-specific incidence, DCO rate and proportion of all cancers for period_2007-2014

							Males	Females
			Males	Females	Males	Females	Prop.all	Prop.all
Age at			Age-	Age-	DCO rate	DCO rate	cancers	cancers
diagnosis	Males	Females	spec.	spec.	n=39	n=16	n=91183	n=89596
Years	n	n	incid.	incid.	90	90	%	%
0- 4			0.0	0.0				
5- 9			0.0	0.0				
10-14			0.0	0.0				
15-19			0.0	0.0				
20-24	3	2	0.3	0.2			0.8	0.6
25-29	6	4	0.5	0.3			1.1	0.6
30-34	8	11	0.6	0.9			1.0	1.0
35-39	10	4	0.8	0.3			0.9	0.2
40 - 44	41	19	2.5	1.2			2.2	0.5
45-49	98	37	6.2	2.4			3.0	0.7
50-54	154	42	11.9	3.3	0.6		3.2	0.6
55-59	167	61 /	15.7	5.4	3.0	6.6	2.3	0.8
60-64	155	78	15.8	7.4	3.2	1.3	1.4	0.8
65-69	164	78	17.0	7.5	4.9	1.3	1.0	0.7
70-74	117	72	12.9	6.9	7.7		0.7	0.6
75-79	62	48	11.3	6.7	6.5		0.5	0.5
80-84	27	43	7.7	7.7	3.7	2.3	0.3	0.5
85+	23	49	9.9	8.5	26.1	18.4	0.4	0.5
All ages	1035	548			3.8	2.9	1.1	0.6
Incidence								
Raw			5.7	2.9				
WS			3.5	1.5				
ES			4.8	2.1				
BRD-S			5.3	2.5				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).



ICD-10 C02-C06: Malign neoplasm of oral cavity




Figure 6. Age distribution and age-specific incidence



ICD-10 C02-C06: Malign neoplasm of oral cavity Age-specific incidence rates: international comparison Average 25 Region MCR Period population 2007-2014 4.6 m ······· SEER 2007-2011 64.6 m 1-year averaged incidence (per 100,000) 5 35-39 **MALES** Age at diagnosis (years)

Figure 6a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).



Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2014, based on the November 2013 submission. http://www.seer.cancer.gov.




Figure 7. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.



Table 8a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2014 MALES

		Observed	Expected		LCL	UCL		DCO
]	Diagnosis	/ n /	n	SIR	95%	95%	EAR	%
(C03-C06 Oral cavity	12	0.9	13.4	6.9	23.5	# 18.3	
	C09-C10 Oropharynx /	46	1.1	40.2	29.4	53.6	# 74.0	2.2
	C12-C13 Hypopharynx	39	0.6	62.4	44.4	85.3	# 63.3	12.8
(C15 Oesophagus	40	1.5	26.3	18.8	35.7	# 63.4	7.5
(C16 Stomach	7	2.6	2.7	1.1	5.5	# 7.2	14.3
(C17 Small intestine	2	0.4	5.2	0.6	18.6	2.7	50.0
(C18 Colon	20	6.3	3.2	1.9	4.9	# 22.6	10.0
(C19-C20 Rectum	14	4.2	3.4		5.6	# 16.2	
(C21 Anus/canal	3	0.2	16.9	3.5	49.3	# 4.7	
(C22 Liver	14	2.0	7.1		11.9		14.3
	C25 Pancreas	5	2.4	2.0	0.7	4.8	4.2	
	C30-C31 Sinuses	3	0.1	22.5	4.6	65.7	# 4.7	
	C32 Larynx	24	0.9	26.2			# 38.1	12.5
	C33-C34 Lung	97	8.6	11.2			# 145.7	13.4
	C43 Malign. melanoma	9	3.3	2.7	1.2	5.1		11.1
(C46,C49 Soft tissue	2	0.4	5.1	0.6	18.3	2.6	
	C61 Prostate	23	20.7	1.1	0.7		3.8	8.7
	C64 Kidney	11	2.7	4.1	2.0	7.3		
	C67 Bladder	7	2.7	2.6	1.0	5.3		14.3
	C73 Thyroid	3	0.7	4.5		13.2	3.8	
	C76-C79 CUP	6	1.2	5.1		11.2		
	C82-C85 NHL	10	2.8	3.6		6.7		20.0
	C91-C96 Leukaemia	4	1.0	3.9		10.0		
(Other primaries	13	3.6	3.6	1.9	6.2	# 15.5	7.7
	Not observed	0	1.6	0.0	0.0	2.3	-2.6	
-								
į	All mult. primaries	414	72.6	5.7	5.2	6.3	# 562.9	9.2
	1							
at:	ients		195	59				
ed.	ian age at second malign	ancy (year						
	son-years		606					
	n observation time (year	s)	3.	. 1				
,				_				

Patients			1959
Median age at second m	malignancy	(years)	63.9
Person-years			6065
Mean observation time	(years)		3.1
Median observation tim	me (vears)		1.7

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries"

Table 8b

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2014

FEMALES

	Observed	Expected		LCL	UCL			DCO
Diagnosis	/ n /	n	SIR	95%	95%		EAR	%
C03-C06 Oral cavity	8	0.2	36.5	15.8	71.9	#	23.7	
C07-C08 Salivary gland	2	0.1	36.6	4.4	132.4	#	5.9	
C09-C10 Oropharynx	16	0.2	104.0	59.4	168.8	#	48.4	
C12-C13 Hypopharynx	8	0.0	193.0	83.3	380.4	#	24.3	50.0
C15 Oesophagus	12	0.2	56.1	29.0	98.0	#	36.0	8.3
C16 Stomach	3	1.2	2.5	0.5	7.4		5.5	33.3
C18 Colon	4	3.3	1.2	0.3	3.1		2.0	
C19-C20 Rectum	2	1.5	1.4	0.2	5.0		1.7	
C22 Liver	6	0.4	15.1	5.6	33.0	#	17.1	16.7
C23-C24 Bile	3	0.5	6.3	1.3	18.5	#	7.7	
C25 Pancreas	3	1.5	2.0	0.4	5.9		4.6	33.3
C30-C31 Sinuses	4	0.0	95.0	25.9	243.2	#	12.1	25.0
C32 Larynx	4	0.1	57.8	15.7	148.0	#	12.0	25.0
C33-C34 Lung	35	2.6	13.6	9.5	18.9	#	99.0	11.4
C43 Malign. melanoma	3	1.3	2.3	0.5	6.7		5.2	33.3
C50 Breast	17	10.9	1.6	0.9	2.5		18.6	
C56 Ovary	2	1.4	1.4	0.2	5.0		1.7	
C67 Bladder	3	0.6	4.8	1.0	13.9		7.2	66.7
C70-C72 CNS cancer	2	0.5	4.1	0.5	14.9		4.6	50.0
C73 Thyroid	4	0.7	6.2	1.7	15.7	#	10.2	
C82-C85 NHL	4	1.3	3.0	0.8	7.8		8.2	
Other primaries	8	4.9	1.6	0.7	3.2		9.6	25.0
Not observed	0	1.8	0.0	0.0	2.0		-5.6	
All mult. primaries	153	35.1	4.4	3.7	5.1	#	359.7	13.1
Patients			959					
Median age at second malig	nancy (yea		3.5					
Person-years			276					
Mean observation time (year	rs)		3.4					


2.0

The occurrence of second malignancy is statistically significant.

Median observation time (years)

Observed second primaries with count 1 are pooled in category "Other primaries"

Average incidence (world standard population) 2007 - 2014: Males



Average incidence (world standard population) 2007 - 2014: Females





Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 3.5/100,000 WS N=1,035, females 1.5/100,000 WS N=548).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 65,347 female residents (averaged) in the period from 2007 to 2014 a total of 16 women were identified with newly diagnosed oral cavity cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 1.4/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.6 and 2.9/100,000.

Standardized incidence ratio (SIR) 2007 - 2014: Males



Standardized incidence ratio (SIR) 2007 - 2014: Females




Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=1,035, females N=548).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,924 female residents (averaged) in the period from 2007 to 2014 a total of 16 women were identified with newly diagnosed oral cavity cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 1.08. Though, the value of this parameter may vary with an underlying probability of 99% between 0.51 and 1.99, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

		_				Prop.
		Prop.				deaths
	Incident	actively	Prop.		Prop.	with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	용	olo	n	%	ଚ
1998	123	99.2	4.1	98	79.7	94.9
1999	129	96.1	1.6	103	79.8	85.4
2000	132	99.2	3.0	99	75.0	96.0
2001	136	97.8	4.4	105	77.2	92.4
2002	190	97.9	6.3	135	71.1	97.8
2003	191	97.4	4.7	132	69.1	98.5
2004	193	97.4	3.1	137	71.0	95.6
2005	161	96.9	3.7	110	68.3	100.0
2006	195	93.8	1.5	128	65.6	96.9
2007	223	81.2	4.0	141	63.2	100.0
2008	216	74.1	1.4	116	53.7	97.4
2009	232	78.4	1.7	123	53.0	97.6
2010	251	74.1	4.8	127	50.6	98.4
2011	182	76.4	2.7	76	41.8	97.4
2012	226	73.9	3.1	83	36.7	96.4
2013	215	100.0	2.8	71	33.0	95.8
2014	43	100.0	20.9	16	37.2	93.8
1998-2014	3038	88.3	3.6	1800	59.2	96.4

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

			Prop.		
			deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n /	n	%	n	용
1998	123	74	93.2	16	13.0
1999	129	70	87.1	12	9.3
2000	132	87	89.7	17	12.9
2001	136	117	88.9	23	16.9
2002	190	149	98.0	33	17.4
2003	191	151	98.0	28	14.7
2004	193	147	98.0	37	19.2
2005	/161	134	98.5	18	11.2
2006	195	150	94.7	24	12.3
2007	223	158	98.1	33	14.8
2008	216	146	97.9	21	9.7
2009	232	188	97.3	28	12.1
2010	251	172	99.4	32	12.7
2011	182	170	97.6	23	12.6
2012	226	178	98.3	31	13.7
2013	215	173	98.8	41	19.1
2014	43	127	95.3	16	37.2
				//	
1998-2014	3038	2391	96.6	433	14.3

Table 10c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

				Prop.
				cancer
		Prop.	Prop.	recorded
		cancer-	non-cancer-	on death
Year of	Deaths	related	related	certificate
death	n	%	- 0 - 0 - 0 - 0 - 0 - 0 - 0 - 0 - 0 - 0	%
000011		_	_ / • /	Ÿ
1998	74	71.6	28.4	88.4
1999	70	61.4	38.6	85.2
2000	87	64.4	35.6	88.5
2001	117	76.9	23.1	91.3
2002	149	75.2	24.8	89.7
2003	151	78.8	21.2	87.8
2004	147	76.2	23.8	88.9
2005	134	86.6	13.4	93.2
2006	150	71.3	28.7	85.2
2007	158	77.8	22.2	89.0
2007	146	78.8	21.2	90.2
2009	188	78.7	21.2	87.4
2019	172			92.4
		79.7	20.3	
2011	170	74.7	25.3	84.3
2012	178	78.7	21.3	89.7
2013	173	72.8	27.2	87.1
2014	127	70.9	29.1	83.5
1998-2014	2391	75.9	24.1	88.4

 $$\operatorname{\textsc{Table 11a}}$$ Medians of age at death according to the grouping in Table 10 $$\operatorname{\textsc{MALES}}$$

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(non-cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	56	60.5	60.5	60.6	61.0
1999	54	55.8	56.8	53.1	54.1
2000	69	62.4	61.8	63.9	65.4
2001	89	60.5	60.4	61.6	60.5
2002	107	62.1	62.0	63.6	61.7
2003	107	63.5	63.1	69.3	63.5
2004	104	63.2	61.1	65.8	62.1
2005	83	65.7	65.2	76.5	65.4
2006	109	62.9	62.6	63.9	62.6
2007	119	62.8	61.5	67.7	62.2
2008	103	64.8	64.6	69.1	64.2
2009	127	67.0	65.1	71.5	65.5
2010	124	66.0	64.5	69.1	65.4
2011	122	67.3	65.8	70.1	64.9
2012	130	65.4	65.2	67.2	65.3
2013	109	66.5	63.3	74.0	64.7
2014	81	68.3	66.9	72.5	68.1
1998-2014	1693	64.2	63.3	68.0	63.7

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 11b $\begin{tabular}{ll} \begin{tabular}{ll} \begin{tabula$

		Age at	Age at	Age at	Age at death
		death	death	death	(according
		(all	(cancer-	(non-cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	18	62.0	65.0	55.7	62.0
1999	16	76.3	65.8	82.3	65.8
2000	18	68.8	67.6	85.4	68.8
2001	28	69.6	69.6	67.4	69.6
2002	42	71.9	69.1	82.9	71.8
2003	44	71.0	70.3	76.5	70.9
2004	43	72.3	73.7	69.1	71.7
2005	51	69.2	67.5	87.2	68.5
2006	41	78.3	77.4	80.4	76.9
2007	3.9	75.1	75.1	77.7	72.0
2008	43	71.4	69.6	72.5	69.4
2009	61	70.4	68.9	84.0	69.8
2010	48	73.0	69.4	86.4	70.0
2011	48	73.3	71.9	73.4	71.8
2012	48	72.1	69.5	85.7	70.5
2013	64	74.3	72.8	81.1	73.7
2014	46	76.2	70.3	79.5	73.9
1998-2014	698	72.3	69.9	79.5	70.4

By 2010, life expectancy at birth was 77.5 years for boys and 82.6 years for girls.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Year of	Deaths	Mort.	MI-Inde	x Mort.	MI-Index	Mort.	MI-Index	Mort.	${\tt MI-Index}$
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	41	3.7	0.47	2.4	0.44	3.3	0.46	3.9	0.49
1999	35	3.1	0.44	2.0	0.43	2.8	0.44	3.1	0.44
2000	45	4.0	0.43	2.4	0.39	3.6	0.42	4.3	0.47
2001	68	5.9	0.74	3.8	0.74	5.3	0.76	6.1	0.78
2002	80	4.3	0.63	2.6	0.60	3.7	0.63	4.3	0.66
2003	87	4.6	0.64	2.8	0.59	4,0	0.61	4.6	0.65
2004	80	4.3	0.58	2.7	0.57	3.7	0.58	4.2	0.58
2005	71	3.7	0.68	2.1	0.62	3.1	0.67	3.8	0.72
2006	80	4.2	0.62	2.6	0.63	3.6	0.61	4.1	0.59
2007	94	4.2	0.63	2.6	0.61	3.6	0.62	4.1	0.63
2008	86	3.9	0.61	2.3	0.58	3.2	0.60	3.8	0.63
2009	103	4.6	0.67	2.7	0.64	3.7	0.65	4.3	0.67
2010	96	4.3	0.56	2.5	0.53	3.5	0.55	4.0	0.56
2011	90	3.9	0.82	2.2	0.75	3.1	0.78	3.7	0.83
2012	104	4.6	0.73	2.6	0.67	3.6	0.69	4.2	0.72
2013	80	3.5	0.57	2.0	0.55	2.8	0.56	3.3	0.58
2014	63	2.8	2.33	1.5	2.14	2.1	2.18	2.5	2.35
1998-2014	1303	4.1	0.64	2.4	0.61	3.4	0.62	3.9	0.65

Table 12b

Mortality measures (cancer-related death) and mortality-incidence-index by year of death FEMALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	12	1.0	0.33	0.5	0.33	0.8	0.33	0.9	0.33
1999	8	0.7	0.16	0.3	0.14	0.5	0.15	0.6	0.16
2000	11	0.9	0.41	0.5	0.40	0.7	0.41	0.8	0.39
2001	22	1.8	0.51	0.9	0.45	1.2	0.46	1.6	0.52
2002	32	1.6	0.50	0.8	0.45	1.2	0.47	1.4	0.48
2003	33	1.7	0.59	0.8	0.53	1.2	0.56	1.5	0.59
2004	32	1.6	0.58	0.7	0.51	1.0	0.54	1.3	0.55
2005	45	2.3	0.79	1.1	0.68	1.6	0.72	1.9	0.75
2006	27	1.3	0.41	0.5	0.27	0.8	0.30	1.0	0.35
2007	29	1.3	0.40	0.5	0.30	0.8	0.33	1.0	0.36
2008	29	1.2	0.39	0.6	0.34	0.9	0.35	1.0	0.37
2009	45	1.9	0.58	0.9	0.51	1.3	0.54	1.5	0.54
2010	41	1.8	0.52	0.8	0.47	1.1	0.49	1.4	0.51
2011	37	1.6	0.51	0.7	0.43	1.0	0.44	1.1	0.45
2012	37	1.6	0.45	0.7	0.37	1.0	0.39	1.2	0.40
2013	46	1.9	0.62	0.8	0.53	1.1	0.55	1.4	0.58
2014	27	1.1	1.69	0.5	1.91	0.7	1.76	0.9	1.66
1998-2014	513	1.5	0.51	0.7	0.44	1.0	0.46	1.2	0.48

Table 13

Age distribution of age at death (cancer-related) for period 2007-2014

(incl. multiple primaries)

Age at death	Cases		1	Males			Females		
Years	n	% Cı	ım.%	n	양	Cum.%	n	%	Cum.%
20-24	2	0.2	0.2	1	0.1	0.1	1	0.3	0.3
25-29	2	0.2	0.4	1	0.1	0.3	1	0.3	0.7
30-34	1	0.1	0.5			0.3	1	0.3	1.0
35-39	5	0.5	1.0	2	0.3	0.5	3	1.0	2.0
40 - 44	20	1.9	2.9	14	1.9	2.5	6	2.0	4.0
45-49	51	5.0	7.9	42	5.8	8.2	9	3.0	7.0
50-54	91	8.9 1	6.8	78	10.7	19.0	13	4.4	11.4
55-59	133	13.0 2	9.7	111	15.2	34.2	22	7.4	18.8
60-64	162	15.8	5.5	121	16.6	50.8	41	13.8	32.6
65-69	186	18.1	3.6	137	18.8	69.6	49	16.4	49.0
70 - 74	144	14.0	77.7	107	14.7	84.3	37	12.4	61.4
75-79	93	9.1 8	86.7	63	8.7	93.0	30	10.1	71.5
80-84	58	5.7	2.4	31	4.3	97.3	27	9.1	80.5
85+	78	7.6 10	0.0	20	2.7	100.0	58	19.5	100.0
All ages	1026	100.0		728	100.0		298	100.0	

Included in the statistics are 41.0% multiple primaries in males and 29.4% in females.



Table 14

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2014 (incl. multiple primaries)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males	Females	spec.		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	%	%
0 - 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24	1	1	0.1	0.33	0.1	0.50	2.1	3.6
25-29	1	1	0.1	0.17	0.1	0.25	1.6	1.6
30-34		1	0.0		0.1	0.09		0.9
35-39	2	3	0.2	0.20	0.2	0.75	1.1	1.2
40-44	14	6	0.9	0.34	0.4	0.32	3.0	0.9
45-49	42	9	2.7	0.43	0.6	0.24	4.1	0.7
50-54	78	13	6.0		1.0	0.31	4.2	0.7
55-59	111/	22	10.5	0.66	2.0	0.36	3.6	0.8
60-64	121	41	12.3	0.78	3.9	0.53	2.5	1.2
65-69	137	49	14.2	0.84	4.7	0.63	1.9	0.9
70-74	107	37	11.8		3.5	0.51	1.2	0.6
75-79	63	30	11.4		4.2	0.63	0.7	0.5
80-84	31	27	8.9		4.8	0.61	0.4	0.4
85+	20	58	8.6		10.0	1.18	0.3	0.7
	20	\ 33	3.0	0.07	10.0	1.10	0.0	0. 7
All ages	728	298					1.5	0.7
1111 0900	, 20						/ 1.0	• •
Mortality /								
Raw			4.0	0.70	1.6	0.54		
WS			2.3		0.7	0.46		
ES			3.3	0.68	1.0	0.48		
BRD-S			3.8	0.71	1.2	0.50		
DIE 0			3.0	0.71	1.2	0.00		
PYLL-70								
per 100,000			34.2		9.6			
ES			30.1		8.2			
AYLL-70			10.8		10.4			
111111 / 0			10.0		10.1			

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15a $\begin{tabular}{ll} Multiple primaries in deaths in period 1998-2014 \\ \hline MALES \end{tabular}$

					Syn-	Syn-		
					chron	chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n	/%↓	n	%	n	– %	n	_ 00 - 00 - 00
		/ *						
C03-C06 Oral cavity	58	7.6			3	5.2	55	94.8
C09-C10 Oropharynx	53	7.0			12	22.6	41	77.4
C12-C13 Hypopharynx	57	7.5	19	33.3	7	12.3	31	54.4
C15 Oesophagus	65	8.6	8	12.3	9	13.8	48	73.8
C16 Stomach	14	1.8	2	14.3		10.0	12	85.7
C18 Colon	25	3.3	7	28.0	2	8.0	16	64.0
C19-C20 Rectum	31	4.1	5	16.1	1	3.2	25	80.6
C22 Liver	24	3.2	2	8.3	2	8.3	20	83.3
C25 Pancreas	8	-1.1	1	12.5	۷	0.3	7	87.5
C30-C31 Sinuses	8	1.1	2	25.0	1	12.5	5	62.5
C32 Larynx	49	6.5	26	53.1	9	18.4	14	28.6
C32 Laryllx C33-C34 Lung	168	22.1	20 17	10.1	. 15	8.9	136	81.0
3 /			7	50.0		7.1		42.9
C43 Malign. melanoma	14	1.8			1 3		6	
C44 Skin others	34	4.5	14	41.2		8.8	17	50.0
C61 Prostate	40	5.3	21	52.5	2	5.0	17	42.5
C64 Kidney	11	1.4	2	18.2	1	9.1	8	72.7
C67 Bladder	26	3.4	15	57.7	1\	3.8	10	38.5
C76-C79 CUP	14	1.8	7	50.0			7	50.0
C82-C85 NHL	14	1.8	7	50.0	2	14.3	5	35.7
Other primaries	46	6.1	23	50.0	3	6.5	20	43.5
All mult. primaries	759	100.0	185	24.4	74	9.7	500	65.9

Multiple primaries with number of cases 1 to 7 are pooled in category "Other primaries"

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 15b

Multiple primaries in deaths in period 1998-2014
FEMALES

						Syn-	Syn-		
						chron	chron		
		Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosi	S	n	% ↓	n	← %	n	← %	n	← %
2									
C03-C06	Oral cavity	24	8.9			1	4.2	23	95.8
	Oropharynx	22 /	8.2			3	13.6	19	86.4
C12-C13	Hypopharynx	8	3.0					8	100.0
C14	ENT cancer	3	1.1			/ 1	33.3	2	66.7
C15	Oesophagus	16	5.9					16	100.0
C16	Stomach	6	2.2	1	16.7	1	16.7	4	66.7
C18	Colon	13	4.8	7	53.8			6	46.2
C21	Anus/canal	3	1.1					3	100.0
C22	Liver	5	1.9			_ 1	20.0	4	80.0
C23-C24	Bile	3	1.1					3	100.0
C25	Pancreas	4	1.5	1	25.0			/ 3	75.0
C30-C31	Sinuses	6	2.2	2	33.3			4	66.7
C32	Larynx	7	2.6	3	42.9			4	57.1
C33-C34	Lung	45	16.7	2	4.4	2	4.4	41	91.1
C43	Malign. melanoma	5	1.9	1	20.0			4	80.0
C44	Skin others	11	4.1	3	27.3	2	18.2	6	54.5
C50	Breast	37	13.8	24	64.9			13	35.1
C53	Cervix uteri	10	3.7	8	80.0			2	20.0
C54	Corpus uteri	5	1.9	4	80.0			1	20.0
C56	Ovary	5	1.9	3	60.0			2	40.0
C67	Bladder	5	1.9	3	60.0			2	40.0
C70-C72	CNS cancer	6	2.2	1	16.7			5	83.3
C82-C85	NHL	4	1.5	1	25.0	1	25.0	2	50.0
Other pr	rimaries	16	5.9	9	56.3			7	43.8
All mult	. primaries	269	100.0	73	27.1	12	4.5	184	68.4

Multiple primaries with number of cases 1 to 2 are pooled in category "Other primaries"

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2014

(First primaries only *)

Age at death Years	Males n	Females	Males Age- spec. mortal.	MI-index	Females Age- spec. mortal.	MI-index	cancers	Females Prop.all cancers %
0- 4 5- 9 10-14 15-19			0.0 0.0 0.0		0.0 0.0 0.0			
20-24 25-29 30-34	1 1	1 1 1	0.1 0.1 0.0	0.17	0.1 0.1 0.1	0.09	2.3	3.8 1.7 1.1
35-39 40-44 45-49 50-54	2 12 38 63	3 6 7 11	0.2 0.7 2.4 4.9	0.32 0.43	0.2 0.4 0.5 0.9	0.32	1.2 2.8 4.1 3.9	1.3 1.1 0.7 0.7
55-59 60-64 65-69 70-74	84 95 109 85	19 33 38 31	7.9 9.7 11.3 9.3	0.66 0.79 0.89	1.7 3.1 3.6 3.0	0.38 0.52 0.67	3.2 2.4 1.9 1.2	0.9 1.2 0.9 0.6
75-79 80-84 85+	46 21 17	26 22 47	8.4 6.0 7.3	1.02 1.24	3.6 3.9 8.1	0.65	0.7 0.4 0.4	0.5 0.4 0.7
All ages	574	246					1.5	0.7
Mortality Raw WS ES BRD-S			3.2 1.8 2.6 3.0	0.66	1.3 0.6 0.8 1.0	0.46		
PYLL-70 per 100,000 ES AYLL-70			27.9 24.5 11.1		8.3 7.1 11.0			

^{*} See corresponding tables with multiple primaries.

Table 17

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2014

(Single primaries only *)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	MI-index	Females Age- spec. mortal.	MI-index	cancers	Females Prop.all cancers
0- 4 5- 9 10-14 15-19 20-24	1	1	0.0 0.0 0.0 0.0 0.1	0.33	0.0 0.0 0.0 0.0 0.1	0.50	2.6	4.2
25-29 30-34 35-39 40-44	1 2 11	1 1 1 6	0.1 0.0 0.2 0.7	0.17	0.1 0.1 0.1 0.4	0.25 0.11 0.25	2.0 1.3 2.8	1.8 1.2 0.5
45-49 50-54 55-59 60-64	33 50 53 51	7 8 15 19	2.1 3.9 5.0 5.2	0.39 0.45 0.48	0.5 0.6 1.3	0.23	3.8 3.5 2.3 1.5	0.8 0.6 0.8
65-69 70-74 75-79 80-84	66 56 33 14	23 20 17 13	6.9 6.2 6.0 4.0	0.63 0.73 0.73	2.2 1.9 2.4 2.3	0.49 0.38 0.50	1.4 1.0 0.7 0.3	0.7 0.5 0.4 0.3
85+ All ages	14 385	38 170	6.0	1.00	6.6	0.95	1.2	0.7
Mortality Raw WS ES BRD-S			2.1 1.2 1.7 2.0	0.49	0.9 0.4 0.6 0.7			
PYLL-70 per 100,000 ES AYLL-70			20.5 17.8 12.3		6.4 5.5 12.3			

^{*} See corresponding tables with multiple primaries.





Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at oral cavity cancer-related death (see Table 10) should be considered.



Average mortality (world standard population) 2007 - 2014: Males



Average mortality (world standard population) 2007 - 2014: Females





Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2007 to 2014. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 2.2/100,000 WS N=710, females 0.7/100,000 WS N=290).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 65,347 female residents (averaged) in the period from 2007 to 2014 a total of 6 women died from oral cavity cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.5/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.1 and 1.5/100,000.

Standardized mortality ratio (SMR) 2007 - 2014: Males



Standardized mortality ratio (SMR) 2007 - 2014: Females




Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=710, females N=290).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,924 female residents (averaged) in the period from 2007 to 2014 a total of 6 women died from oral cavity cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.78. Though, the value of this parameter may vary with an underlying probability of 99% between 0.20 and 2.04, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases, where applicable. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

FRG Federal Republic of Germany

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

MCR Munich Cancer Registry (Tumorregister München)
SEER Surveillance, Epidemiology, and End Results (USA)

AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

BRD-S German standard population

DCO Death certificate only EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

ES European standard population (old)

LCL Lower confidence limit

MI-index Ratio between mortality and incidence

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age

SIR Standardized incidence ratio SMR Standardized mortality ratio UCL Upper confidence limit WS World standard population

Recommended Citation

Munich Cancer Registry. ICD-10 C02-C06: Oral cavity cancer - Incidence and Mortality [Internet]. 2016 [updated 2016 Apr 13; cited 2016 Jun 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/base/bC0206E-ICD-10-C02-C06-Oral-cavity-cancer-incidence-and-mortality.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

Fig./Tb	I.	Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	4
1a	Gender distribution by year of diagnosis	5
2	Incidence by year of diagnosis	6
3	Age distribution parameters by year of diagnosis	7
4	Age distribution by 5-year age group and gender	9
5	Age-specific incidence, DCO rate, proportion malignancies	10
6	Age distribution and age-specific incidence (chart)	11
6a	Age-specific incidence internationally (chart)	12
7	Cumulative follow-up years (chart)	13
8	Standardized incidence ratio of second primaries	14
9a	Map of cancer incidence (WS) by county (chart)	16
9b	Standardized incidence ratio (SIR) by county (chart)	17
10a	Pts incident cohorts and mortality / yr	18
10b	Incidence and mortality by year of diagnosis	19
10c	Cancer-related deaths, death certification available / yr	20
11	Medians of age at death / yr	21
12	Mortality by year of death	23
13	Distribution of age at death	24
14	Age-specific mortality	25
15	Multiple primaries in deaths	26
16	Age-specific mortality (first primaries)	28
17	Age-specific mortality (single primaries)	29
18	Age distribution and age-specific mortality (chart)	30
19a	Map of cancer mortality (WS) by county (chart)	31
19b	Standardized mortality ratio (SMR) by county (chart)	32