


Munich Cancer Registry



- ▶ [Survival](#)
- ▶ [Selection Matrix](#)
- ▶ [Homepage](#)
- ▶ [Deutsch](#)

ICD-10 C18-C20: Colorectal cancer

Incidence and Mortality

Year of diagnosis	1998-2014
Patients	45,104
Diseases	46,277
Creation date	04/13/2016
Export date	12/23/2015
Population	4.64 m



Munich Cancer Registry at Munich Cancer Center
Marchioninstr. 15
Munich, 81377
Germany

<http://www.tumorregister-muenchen.de/en>

<http://www.tumorregister-muenchen.de/en/facts/base/bC1820E-ICD-10-C18-C20-Colorectal-cancer-incidence-and-mortality.pdf>

**Global Statements about the statistics on the Internet –
Baseline Statistics** (grey button) , **Survival** (red button)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.64 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases^{###} are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, April 2016

[#] Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007).

^{##} Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.

^{###} DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate.

ICD-10 codes (ICD-10 2015) used for specifying cancer site

Code	Description
C18.-	Malignant neoplasm of colon
C18.0	Caecum
C18.1	Appendix
C18.2	Ascending colon
C18.3	Hepatic flexure
C18.4	Transverse colon
C18.5	Splenic flexure
C18.6	Descending colon
C18.7	Sigmoid colon
C18.8	Overlapping lesion of colon
C18.9	Colon, unspecified
C19	Malignant neoplasm of rectosigmoid junction
C20	Malignant neoplasm of rectum

INCIDENCE

Table 1

All patients with invasive cancer by year of diagnosis, proportions of DCO, multiple primaries, deaths, and active follow-up (incl. DCO)

Year of diagnosis	Cases n	DCO cases n	Prop. DCO %	Prop. mult. primaries %	Prop. deaths %	Prop. actively followed %
1998	1824	97	5.3	24.9	73.8	97.9
1999	1838	112	6.1	24.8	72.6	97.6
2000	1686	94	5.6	26.4	71.0	98.3
2001	1834	118	6.4	27.3	66.6	97.3
2002	3147	365	11.6	25.6	69.6	97.6 #
2003	3161	298	9.4	25.9	64.6	97.3
2004	3038	241	7.9	25.6	64.4	97.1
2005	2965	211	7.1	27.3	63.9	96.8
2006	3054	156	5.1	27.8	57.5	94.0
2007	3391	199	5.9	25.6	56.6	79.9 #
2008	3341	192	5.7	27.5	53.3	73.3
2009	3281	172	5.2	26.6	49.9	71.2
2010	3067	177	5.8	26.6	46.5	69.6
2011	2999	154	5.1	25.5	42.9	69.5
2012	2917	158	5.4	25.1	37.3	72.2
2013	2801	155	5.5	24.8	29.3	99.1
2014	1933	135	7.0	23.7	20.8	98.3 ##
1998–2014	46277	3034	6.6	26.0	54.7	87.2

The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be found in the respective headings.

Table 1a

All patients with invasive cancer
by year of diagnosis and gender
(incl. DCO)

Year of diagnosis	All n	Males n	Females n	Prop. males %
1998	1824	920	904	50.4
1999	1838	941	897	51.2
2000	1686	871	815	51.7
2001	1834	959	875	52.3
2002	3147	1671	1476	53.1
2003	3161	1684	1477	53.3
2004	3038	1623	1415	53.4
2005	2965	1574	1391	53.1
2006	3054	1665	1389	54.5
2007	3391	1874	1517	55.3
2008	3341	1852	1489	55.4
2009	3281	1844	1437	56.2
2010	3067	1726	1341	56.3
2011	2999	1647	1352	54.9
2012	2917	1612	1305	55.3
2013	2801	1608	1193	57.4
2014	1933	1071	862	55.4
1998–2014	46277	25142	21135	54.3

Table 2

Incidence measures by year of diagnosis including DCO cases
(with respect to registry area expansion from 2.51 to 3.96 m as of 2002,
and from 3.96 to 4.64 m as of 2007, respectively)

Year of diagnosis	Males n	Females n	Males Inc. raw	Fem. Inc. raw	Males Inc. WS	Fem. Inc. WS	Males Inc. ES	Fem. Inc. ES	Males Inc. BRD-S	Fem. Inc. BRD-S
1998	920	904	83.0	76.8	49.8	32.7	74.9	49.4	97.7	64.7
1999	941	897	84.1	75.6	49.8	32.1	75.3	48.4	98.5	63.2
2000	871	815	76.5	67.8	44.7	27.6	67.8	42.2	87.7	55.7
2001	959	875	82.7	71.9	48.5	30.6	72.7	46.1	92.9	60.2
2002	1671	1476	89.7	75.4	50.0	30.5	75.5	46.3	98.8	60.6
2003	1684	1477	89.8	75.0	49.3	30.6	74.3	46.2	96.8	59.9
2004	1623	1415	86.3	71.6	46.0	29.6	69.5	44.3	91.0	57.1
2005	1574	1391	83.1	69.9	44.1	27.3	66.2	41.4	85.9	54.5
2006	1665	1389	86.9	69.1	45.5	28.1	68.2	42.0	88.5	54.9
2007	1874	1517	84.6	65.7	44.0	26.3	65.6	39.4	85.3	51.2
2008	1852	1489	83.2	64.2	41.7	25.1	63.0	37.8	82.2	49.3
2009	1844	1437	82.6	61.8	40.8	23.9	61.2	36.0	80.2	47.4
2010	1726	1341	76.6	57.3	37.6	21.6	56.4	32.7	73.5	43.3
2011	1647	1352	72.1	57.3	35.1	22.4	52.4	33.4	68.1	42.9
2012	1612	1305	70.6	55.3	34.3	22.1	51.6	32.6	66.8	42.2
2013	1608	1193	70.4	50.6	33.5	20.4	50.7	30.1	66.7	38.8
2014	1071	862	46.9	36.5	23.1	14.7	34.5	21.8	44.8	28.1
1998-2014	25142	21135	78.5	63.2	40.7	25.3	61.1	38.0	79.4	49.5

The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

Table 3

Age distribution parameters by year of diagnosis (All patients)
(incl. DCO)

Year of diagnosis	Cases n	Mean	Std. dev.	Min.	Max.	10%	25%	Median 50%	75%	90%
1998	1824	70.0	12.4	13.2	102	54.1	61.0	71.0	78.7	86.1
1999	1838	70.3	12.4	24.9	102	54.3	61.7	71.1	79.3	86.4
2000	1686	70.6	12.1	24.7	103	55.1	61.8	71.6	79.3	86.7
2001	1834	70.1	12.4	28.3	103	54.5	61.7	70.4	79.3	86.5
2002	3147	71.0	12.1	17.7	104	55.5	62.7	71.9	80.1	86.7
2003	3161	71.0	11.8	8.4	101	56.1	63.1	71.5	79.9	86.0
2004	3038	70.7	12.3	13.8	101	55.2	63.0	71.2	79.9	85.5
2005	2965	71.4	12.2	15.1	99.9	55.6	63.8	71.8	80.4	86.1
2006	3054	70.6	12.1	17.9	102	54.8	63.4	71.0	79.7	85.3
2007	3391	70.7	12.4	15.8	103	54.2	63.9	71.3	80.2	85.7
2008	3341	71.5	12.3	18.9	105	55.3	64.2	72.1	80.4	86.6
2009	3281	71.3	12.3	12.4	102	54.8	64.1	72.1	80.3	86.0
2010	3067	71.5	12.6	14.9	101	54.4	63.8	72.6	81.0	86.4
2011	2999	71.3	12.8	15.5	101	53.4	63.4	72.4	80.8	87.0
2012	2917	71.2	12.9	9.7	101	54.3	63.4	72.7	80.4	86.4
2013	2801	70.9	13.1	15.7	105	53.1	63.2	72.8	80.2	86.2
2014	1933	70.8	13.3	20.3	103	52.7	62.7	72.7	80.1	86.5
1998-2014	46277	71.0	12.4	8.4	105	54.6	63.1	71.9	80.1	86.2

Table 3a

Age distribution parameters by year of diagnosis (MALES)
(incl. DCO)

Year of diagnosis	Cases n	Mean	Std. dev.	Min.	Max.	10%	25%	Median 50%	75%	90%
1998	920	67.9	11.7	31.4	98.1	53.7	59.5	68.3	75.9	83.9
1999	941	68.3	11.5	24.9	95.5	54.5	60.3	69.0	76.6	83.2
2000	871	68.3	11.0	34.4	95.9	54.3	60.3	68.1	76.6	83.0
2001	959	68.4	11.3	31.3	102	54.2	61.2	68.0	76.0	83.6
2002	1671	69.1	11.0	20.9	98.5	55.6	61.9	69.6	76.7	82.5
2003	1684	69.3	11.1	8.4	99.4	55.5	62.6	69.6	76.6	82.7
2004	1623	69.4	11.0	27.8	101	55.7	62.4	69.3	77.1	83.5
2005	1574	69.3	11.3	19.0	99.6	54.6	62.9	69.5	77.1	83.6
2006	1665	69.1	11.1	17.9	102	54.8	62.6	69.3	77.4	82.9
2007	1874	69.1	11.7	15.8	99.4	54.2	62.9	69.6	77.6	83.0
2008	1852	69.9	11.3	19.3	105	55.0	63.6	70.4	77.9	83.4
2009	1844	69.7	11.4	12.4	99.0	54.4	63.1	71.0	77.9	83.0
2010	1726	69.9	11.7	21.1	98.9	54.0	62.5	70.9	78.2	84.1
2011	1647	69.9	11.8	15.5	97.3	53.4	63.2	71.1	78.3	84.2
2012	1612	70.2	11.5	9.7	101	55.3	62.9	71.4	78.3	84.0
2013	1608	70.3	12.0	19.4	99.6	54.1	63.1	72.1	78.5	84.5
2014	1071	69.7	12.6	20.3	102	53.1	61.8	71.7	78.7	84.9
1998-2014	25142	69.4	11.5	8.4	105	54.5	62.3	70.1	77.5	83.5

Table 3b

Age distribution parameters by year of diagnosis (FEMALES)
(incl. DCO)

Year of diagnosis	Cases n	Mean	Std. dev.	Min.	Max.	10%	25%	Median 50%	75%	90%
1998	904	72.2	12.7	13.2	102	54.7	63.3	74.1	81.5	87.4
1999	897	72.4	13.0	26.9	102	54.3	63.5	74.5	82.0	88.0
2000	815	73.1	12.6	24.7	103	56.3	63.6	75.3	81.9	88.5
2001	875	71.9	13.2	28.3	103	54.8	62.4	74.6	81.1	88.4
2002	1476	73.2	12.9	17.7	104	55.3	63.9	75.3	82.2	88.9
2003	1477	73.1	12.3	23.5	101	56.4	64.3	74.5	82.5	88.5
2004	1415	72.2	13.4	13.8	100	54.7	64.0	74.0	82.7	87.8
2005	1391	73.7	12.8	15.1	99.9	57.0	65.4	75.5	83.3	89.7
2006	1389	72.4	13.0	21.2	98.7	54.8	64.4	74.3	82.4	86.8
2007	1517	72.8	13.0	17.8	103	54.1	65.1	74.3	82.8	87.5
2008	1489	73.5	13.2	18.9	102	55.6	65.3	74.4	83.6	88.6
2009	1437	73.4	13.1	15.9	102	55.7	65.6	75.1	83.3	88.5
2010	1341	73.6	13.3	14.9	101	55.3	66.2	75.6	83.5	88.7
2011	1352	73.0	13.8	17.1	101	53.4	63.9	74.6	84.0	88.9
2012	1305	72.3	14.4	13.7	100	53.4	64.0	74.8	83.2	88.9
2013	1193	71.9	14.4	15.7	105	50.9	63.4	74.1	82.7	88.5
2014	862	72.1	14.0	23.2	103	51.6	64.2	74.2	82.2	88.9
1998–2014	21135	72.8	13.3	13.2	105	54.7	64.3	74.7	82.8	88.4

Table 4

Age distribution by 5-year age group and gender for period 2007–2014
(incl. DCO)

Age at diagnosis Years	Cases			Males			Females		
	n	%	Cum.%	n	%	Cum.%	n	%	Cum.%
5–9	1	0.0	0.0	1	0.0	0.0			0.0
10–14	3	0.0	0.0	1	0.0	0.0	2	0.0	0.0
15–19	22	0.1	0.1	4	0.0	0.0	18	0.2	0.2
20–24	34	0.1	0.3	14	0.1	0.2	20	0.2	0.4
25–29	59	0.2	0.5	25	0.2	0.3	34	0.3	0.7
30–34	97	0.4	0.9	51	0.4	0.7	46	0.4	1.1
35–39	164	0.7	1.6	90	0.7	1.4	74	0.7	1.8
40–44	382	1.6	3.2	211	1.6	3.0	171	1.6	3.5
45–49	707	3.0	6.2	393	3.0	6.0	314	3.0	6.5
50–54	1130	4.8	11.0	654	4.9	10.9	476	4.5	11.0
55–59	1695	7.1	18.1	1042	7.9	18.8	653	6.2	17.2
60–64	2383	10.0	28.1	1538	11.6	30.4	845	8.1	25.3
65–69	3270	13.8	41.9	2098	15.9	46.3	1172	11.2	36.4
70–74	4021	16.9	58.9	2481	18.7	65.0	1540	14.7	51.1
75–79	3522	14.8	73.7	2031	15.3	80.4	1491	14.2	65.3
80–84	3138	13.2	86.9	1540	11.6	92.0	1598	15.2	80.5
85+	3102	13.1	100.0	1060	8.0	100.0	2042	19.5	100.0
All ages	23730	100.0		13234	100.0		10496	100.0	

Included in the statistics are 32.7% multiple primaries in males and 25.9% in females.

Table 5

Age-specific incidence, DCO rate and proportion of all cancers
for period 2007-2014

Age at diagnosis Years	Males n	Females n	Males Age- spec. incid. %	Females Age- spec. incid. %	Males DCO rate n=546 %	Females DCO rate n=792 %	Males Prop.all cancers n=91183 %	Females Prop.all cancers n=89596 %
0- 4			0.0	0.0				
5- 9	1		0.1	0.0			1.0	
10-14	1	2	0.1	0.2			1.0	2.2
15-19	4	18	0.4	2.0			1.9	10.9
20-24	14	20	1.3	1.8	7.1		3.8	6.4
25-29	23	34	1.9	2.8			4.1	5.1
30-34	51	45	4.1	3.6			6.6	3.9
35-39	87	73	6.7	5.8		4.1	7.5	3.7
40-44	209	171	12.8	11.2		0.6	11.4	4.6
45-49	387	312	24.5	20.6	1.0		12.0	5.7
50-54	638	472	49.3	36.9	1.1	1.3	13.1	7.0
55-59	1026	642	96.6	57.1	1.7	0.3	14.0	8.6
60-64	1508	834	153.5	78.7	1.6	1.6	14.0	9.1
65-69	2047	1154	212.8	110.6	1.8	1.9	13.1	10.1
70-74	2420	1511	266.0	144.5	2.8	2.6	14.3	12.8
75-79	1979	1466	359.4	205.5	3.8	4.8	15.9	14.6
80-84	1483	1577	424.6	281.2	7.6	8.0	17.3	17.9
85+	1039	2011	448.8	348.0	19.3	25.4	17.1	19.6
All ages	12917	10342			4.2	7.7	14.2	11.5
Incidence								
Raw			71.5	55.2				
WS			35.3	21.7				
ES			53.0	32.5				
BRD-S			68.9	42.2				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).




Figure 6. Age distribution and age-specific incidence




Figure 6a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).

Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2014, based on the November 2013 submission. <http://www.seer.cancer.gov>.




Figure 7. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.

Table 8a

Standardized incidence ratio (SIR, with 95% confidence limits),
excess absolute risk (EAR) and DCO rate of second primaries
for period 1998–2014

MALES

Diagnosis		Observed n	Expected n	SIR	LCL 95%	UCL 95%	EAR	DCO %
C00	Lip	3	1.7	1.8	0.4	5.1	0.2	
C03-C06	Oral cavity	11	10.1	1.1	0.5	1.9	0.1	
C07-C08	Salivary gland	4	3.1	1.3	0.4	3.3	0.1	
C09-C10	Oropharynx	15	12.3	1.2	0.7	2.0	0.4	
C12-C13	Hypopharynx	9	6.7	1.3	0.6	2.5	0.3	11.1
C15	Oesophagus	63	23.2	2.7	2.1	3.5	#	14.3
C16	Stomach	160	58.9	2.7	2.3	3.2	#	6.9
C17	Small intestine	70	6.7	10.5	8.2	13.3	#	1.4
C18	Colon	522	140.1	3.7	3.4	4.1	#	1.0
C19-C20	Rectum	225	73.9	3.0	2.7	3.5	#	1.3
C21	Anus/canal	9	2.7	3.3	1.5	6.3	#	0.9
C22	Liver	106	37.2	2.8	2.3	3.4	#	17.9
C23-C24	Bile	35	13.7	2.6	1.8	3.6	#	11.4
C25	Pancreas	107	50.5	2.1	1.7	2.6	#	24.3
C32	Larynx	23	13.0	1.8	1.1	2.6	#	17.4
C33-C34	Lung	319	160.0	2.0	1.8	2.2	#	14.7
C38,C45	Mesothelioma	11	9.0	1.2	0.6	2.2		0.3
C40-C41	Bone	2	1.0	2.0	0.2	7.2		0.1
C43	Malign. melanoma	100	54.4	1.8	1.5	2.2	#	2.0
C46,C49	Soft tissue	15	7.3	2.0	1.1	3.4	#	1.1
C50	Breast	9	3.6	2.5	1.2	4.8	#	11.1
C60	Penis	7	3.2	2.2	0.9	4.6		0.5
C61	Prostate	700	408.0	1.7	1.6	1.8	#	5.1
C62	Testis	6	2.5	2.4	0.9	5.2		16.7
C64	Kidney	141	46.6	3.0	2.5	3.6	#	5.0
C65	Renal pelvis	17	5.9	2.9	1.7	4.6	#	1.5
C66	Ureter	11	3.4	3.3	1.6	5.9	#	1.1
C67	Bladder	130	65.2	2.0	1.7	2.4	#	7.7
C68	Urinary org.	3	0.9	3.5	0.7	10.1		66.7
C69	Eye carcinoma	2	0.5	4.1	0.5	14.7		0.2
C70-C72	CNS cancer	36	17.3	2.1	1.5	2.9	#	25.0
C73	Thyroid	12	7.8	1.5	0.8	2.7		16.7
C76-C79	CUP	36	23.8	1.5	1.1	2.1	#	2.8
C81	Hodgkin lymphoma	5	2.7	1.9	0.6	4.4		0.3
C82-C85	NHL	114	55.7	2.0	1.7	2.5	#	4.4
C90	Mult. myeloma	31	17.9	1.7	1.2	2.5	#	25.8
C91-C96	Leukaemia	44	23.4	1.9	1.4	2.5	#	29.5
Other primaries		8	12.2	0.7	0.3	1.3	-0.6	12.5
Not observed		0	2.5	0.0	0.0	1.5	-0.3	

Table 8a

Standardized incidence ratio (SIR, with 95% confidence limits),
excess absolute risk (EAR) and DCO rate of second primaries
for period 1998-2014

MALES

Diagnosis	Observed n	Expected n	SIR	LCL 95%	UCL 95%	EAR	DCO %
All mult. primaries	3121	1388.5	2.2	2.2	2.3 #	240.4	7.3
Patients		23230					
Median age at second malignancy (years)		73.5					
Person-years		72071					
Mean observation time (years)		3.1					
Median observation time (years)		1.8					

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries"

Table 8b

Standardized incidence ratio (SIR, with 95% confidence limits),
excess absolute risk (EAR) and DCO rate of second primaries
for period 1998–2014
FEMALES


Diagnosis	Observed n	Expected n	SIR	LCL 95%	UCL 95%	EAR	DCO %
C03-C06 Oral cavity	7	4.5	1.6	0.6	3.2	0.4	
C09-C10 Oropharynx	7	2.6	2.6	1.1	5.5 #	0.7	
C15 Oesophagus	11	4.5	2.4	1.2	4.4 #	1.1	18.2
C16 Stomach	78	33.7	2.3	1.8	2.9 #	7.3	19.2
C17 Small intestine	38	3.6	10.7	7.6	14.7 #	5.7	2.6
C18 Colon	307	91.2	3.4	3.0	3.8 #	35.5	1.0
C19-C20 Rectum	108	36.7	2.9	2.4	3.5 #	11.7	1.9
C21 Anus/canal	11	4.1	2.7	1.3	4.8 #	1.1	
C22 Liver	29	10.0	2.9	1.9	4.1 #	3.1	41.4
C23-C24 Bile	26	13.3	2.0	1.3	2.9 #	2.1	15.4
C25 Pancreas	87	39.7	2.2	1.8	2.7 #	7.8	25.3
C33-C34 Lung	145	54.4	2.7	2.3	3.1 #	14.9	16.6
C43 Malign. melanoma	63	26.6	2.4	1.8	3.0 #	6.0	
C46,C49 Soft tissue	9	4.6	2.0	0.9	3.7	0.7	
C48 Peritoneal	9	2.7	3.3	1.5	6.3 #	1.0	22.2
C50 Breast	399	221.6	1.8	1.6	2.0 #	29.2	4.8
C51 Vulva	17	9.0	1.9	1.1	3.0 #	1.3	5.9
C52 Vagina	7	1.7	4.1	1.7	8.5 #	0.9	14.3
C53 Cervix uteri	18	9.4	1.9	1.1	3.0 #	1.4	11.1
C54 Corpus uteri	97	42.3	2.3	1.9	2.8 #	9.0	3.1
C55,C57 Fem. genitals un	5	2.6	1.9	0.6	4.5	0.4	20.0
C56 Ovary	103	32.6	3.2	2.6	3.8 #	11.6	28.2
C64 Kidney	67	20.3	3.3	2.6	4.2 #	7.7	11.9
C65 Renal pelvis	8	2.6	3.0	1.3	6.0 #	0.9	
C66 Ureter	4	1.3	3.0	0.8	7.8	0.4	25.0
C67 Bladder	41	17.9	2.3	1.6	3.1 #	3.8	22.0
C70-C72 CNS cancer	13	10.8	1.2	0.6	2.1	0.4	61.5
C73 Thyroid	16	10.7	1.5	0.9	2.4	0.9	6.3
C76-C79 CUP	12	16.9	0.7	0.4	1.2	-0.8	
C82-C85 NHL	60	32.6	1.8	1.4	2.4 #	4.5	16.7
C90 Mult. myeloma	21	10.5	2.0	1.2	3.0 #	1.7	28.6
C91-C96 Leukaemia	31	14.0	2.2	1.5	3.1 #	2.8	41.9
Other primaries	29	16.8	1.7	1.2	2.5 #	2.0	17.2
Not observed	0	1.5	0.0	0.0	2.5	-0.2	
All mult. primaries	1883	807.3	2.3	2.2	2.4 #	177.0	10.8

Patients 19105
 Median age at second malignancy (years) 76.0
 Person-years 60766
 Mean observation time (years) 3.2
 Median observation time (years) 1.8

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 to 3 are pooled in category "Other primaries"

Average incidence (world standard population) 2007 - 2014: Males



Average incidence (world standard population) 2007 - 2014: Females





Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 35.3/100,000 WS N=12,917, females 21.7/100,000 WS N=10,342).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 65,347 female residents (averaged) in the period from 2007 to 2014 a total of 215 women were identified with newly diagnosed colorectal cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 18.3/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 14.9 and 22.5/100,000.

Standardized incidence ratio (SIR) 2007 - 2014: Males



Standardized incidence ratio (SIR) 2007 - 2014: Females




Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=12,917, females N=10,342).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,924 female residents (averaged) in the period from 2007 to 2014 a total of 215 women were identified with newly diagnosed colorectal cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.79. Though, the value of this parameter may vary with an underlying probability of 99% between 0.66 and 0.94.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

Year of diagnosis	Incident cases n	Prop. actively followed %	Prop. DCO %	Deaths n	Prop. deaths %	Prop. deaths with death certific. %
1998	1824	97.9	5.3	1346	73.8	93.7
1999	1838	97.6	6.1	1334	72.6	94.8
2000	1686	98.3	5.6	1197	71.0	96.2
2001	1834	97.3	6.4	1222	66.6	97.3
2002	3147	97.6	11.6	2190	69.6	97.7
2003	3161	97.3	9.4	2042	64.6	98.0
2004	3038	97.1	7.9	1957	64.4	97.9
2005	2965	96.8	7.1	1894	63.9	98.0
2006	3054	94.0	5.1	1755	57.5	99.0
2007	3391	79.9	5.9	1920	56.6	98.4
2008	3341	73.3	5.7	1782	53.3	98.4
2009	3281	71.2	5.2	1636	49.9	98.4
2010	3067	69.6	5.8	1425	46.5	97.9
2011	2999	69.5	5.1	1286	42.9	97.7
2012	2917	72.2	5.4	1087	37.3	97.9
2013	2801	99.1	5.5	820	29.3	95.7
2014	1933	98.3	7.0	402	20.8	94.0
1998-2014	46277	87.2	6.6	25295	54.7	97.5

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis
(incl. DCO)
(with respect to registry area expansion from 2.51 to 3.96 m as of 2002,
and from 3.96 to 4.64 m as of 2007, respectively)

Year of diagnosis/ death	Incident cases n	Deaths n	Prop. deaths with death certific. %	Deaths in same year n	Prop. deaths in same year %
1998	1824	1041	90.3	297	16.3
1999	1838	1075	90.9	312	17.0
2000	1686	1044	93.8	282	16.7
2001	1834	1112	95.6	286	15.6
2002	3147	1582	98.0	690	21.9
2003	3161	1697	97.8	594	18.8
2004	3038	1707	98.3	552	18.2
2005	2965	1807	96.5	541	18.2
2006	3054	1872	97.5	508	16.6
2007	3391	1984	97.6	573	16.9
2008	3341	2075	98.7	604	18.1
2009	3281	2114	98.7	536	16.3
2010	3067	2181	98.6	523	17.1
2011	2999	2188	98.3	511	17.0
2012	2917	2196	98.4	520	17.8
2013	2801	2175	97.9	466	16.6
2014	1933	1961	98.5	355	18.4
1998-2014	46277	29811	97.3	8150	17.6

Table 10c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates

(incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

Year of death	Deaths n	Prop. cancer- related %	Prop. non-cancer- related %	Prop. cancer recorded on death certificate %
1998	1041	71.9	28.1	86.9
1999	1075	73.5	26.5	86.6
2000	1044	73.6	26.4	86.0
2001	1112	69.4	30.6	84.9
2002	1582	75.3	24.7	87.2
2003	1697	73.8	26.2	86.7
2004	1707	76.2	23.8	86.6
2005	1807	71.6	28.4	81.8
2006	1872	71.6	28.4	82.9
2007	1984	72.0	28.0	83.7
2008	2075	71.6	28.4	82.0
2009	2114	69.7	30.3	79.8
2010	2181	66.6	33.4	78.8
2011	2188	67.0	33.0	78.6
2012	2196	66.3	33.7	78.3
2013	2175	63.0	37.0	74.2
2014	1961	63.0	37.0	75.3
1998-2014	29811	69.8	30.2	81.6

Table 11a

Medians of age at death according to the grouping in Table 10

MALES

Year of death	Deaths n	Age at death (all causes) Years	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate) Years
1998	510	73.8	71.5	79.5	73.2
1999	531	73.4	71.4	78.7	72.8
2000	546	74.7	72.3	82.2	73.8
2001	538	74.2	71.2	80.9	72.6
2002	818	74.2	72.1	80.8	73.3
2003	874	75.0	72.7	80.8	73.9
2004	881	75.7	74.3	81.0	75.1
2005	945	75.5	73.1	81.3	73.7
2006	1029	76.5	74.4	81.0	75.4
2007	1084	76.0	73.9	80.9	74.6
2008	1164	76.7	74.6	82.1	75.5
2009	1122	76.4	73.7	81.1	74.4
2010	1186	76.8	74.4	82.1	75.4
2011	1209	76.5	73.3	82.6	75.2
2012	1208	77.4	75.4	82.3	76.2
2013	1181	78.8	76.4	83.4	77.1
2014	1077	78.1	75.6	83.0	76.6
1998-2014	15903	76.3	73.9	81.7	75.0

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 11b

Medians of age at death according to the grouping in Table 10
FEMALES

Year of death	Deaths n	Age at death (all causes) Years	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate) Years
1998	531	78.6	76.5	84.8	78.4
1999	544	79.9	78.4	85.9	79.7
2000	498	80.6	78.8	86.5	79.6
2001	574	80.9	78.0	87.1	80.0
2002	764	81.0	79.6	86.5	80.6
2003	823	81.3	78.9	85.9	80.3
2004	826	81.3	79.3	85.2	80.1
2005	862	81.9	80.0	85.2	80.7
2006	843	82.0	79.7	86.2	80.6
2007	900	82.1	79.3	86.7	80.6
2008	911	82.6	80.1	86.5	81.1
2009	992	82.8	79.3	87.5	80.5
2010	995	83.3	79.9	87.2	81.5
2011	979	83.5	79.6	88.1	81.4
2012	988	83.9	79.4	88.5	81.4
2013	994	84.0	79.1	88.5	81.1
2014	884	84.0	78.7	88.3	80.9
1998-2014	13908	82.2	79.2	87.0	80.6

By 2010, life expectancy at birth was 77.5 years for boys and 82.6 years for girls.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 12a

Mortality measures (cancer-related death) and mortality-incidence-index
by year of death

MALES

Year of death	Deaths n	Mort. raw	MI-Index raw	Mort. WS	MI-Index WS	Mort. ES	MI-Index ES	Mort. BRD-S	MI-Index BRD-S
1998	366	33.0	0.40	19.2	0.39	30.3	0.41	41.1	0.42
1999	391	34.9	0.42	20.1	0.41	31.7	0.42	44.1	0.45
2000	407	35.7	0.47	20.0	0.45	31.8	0.47	43.8	0.50
2001	385	33.2	0.41	18.8	0.39	29.4	0.41	39.4	0.43
2002	622	33.4	0.38	18.0	0.36	28.2	0.38	38.1	0.39
2003	658	35.1	0.39	18.3	0.37	28.9	0.39	40.1	0.42
2004	679	36.1	0.42	17.9	0.40	28.8	0.42	40.7	0.45
2005	693	36.6	0.45	18.1	0.42	28.4	0.44	39.4	0.47
2006	742	38.7	0.46	18.7	0.42	30.0	0.45	41.9	0.49
2007	802	36.2	0.44	16.9	0.39	27.0	0.42	37.9	0.45
2008	867	39.0	0.48	17.8	0.44	28.6	0.47	40.3	0.51
2009	791	35.4	0.44	16.3	0.41	25.7	0.43	35.3	0.45
2010	813	36.1	0.48	15.8	0.43	25.1	0.46	35.4	0.49
2011	846	37.0	0.52	16.7	0.48	26.1	0.51	35.3	0.53
2012	817	35.8	0.52	15.7	0.47	25.1	0.50	34.7	0.54
2013	775	33.9	0.50	14.5	0.44	23.4	0.48	33.0	0.51
2014	694	30.4	0.66	13.0	0.57	20.8	0.61	29.3	0.67
1998-2014	11348	35.4	0.46	16.9	0.42	26.9	0.45	37.3	0.48

Table 12b

Mortality measures (cancer-related death) and mortality-incidence-index
by year of death

FEMALES

Year of death	Deaths n	Mort. raw	MI-Index raw	Mort. WS	MI-Index WS	Mort. ES	MI-Index ES	Mort. BRD-S	MI-Index BRD-S
1998	383	32.6	0.43	12.3	0.38	19.4	0.40	27.1	0.42
1999	400	33.7	0.45	11.9	0.37	19.2	0.40	26.5	0.42
2000	361	30.1	0.45	10.8	0.39	17.2	0.41	23.1	0.42
2001	387	31.8	0.45	11.6	0.38	18.4	0.40	25.4	0.42
2002	570	29.1	0.39	9.8	0.32	15.7	0.34	21.8	0.36
2003	596	30.3	0.41	10.6	0.35	16.8	0.37	23.0	0.39
2004	621	31.4	0.44	10.5	0.36	16.9	0.39	23.7	0.42
2005	600	30.2	0.44	9.9	0.37	15.9	0.39	22.0	0.41
2006	601	29.9	0.44	9.5	0.34	15.4	0.37	22.0	0.41
2007	628	27.2	0.42	9.3	0.36	14.7	0.38	20.2	0.40
2008	621	26.8	0.42	8.6	0.35	13.8	0.37	19.1	0.39
2009	683	29.4	0.48	9.7	0.41	15.3	0.43	20.9	0.45
2010	642	27.4	0.49	8.8	0.41	13.9	0.43	18.9	0.44
2011	622	26.4	0.47	8.2	0.37	13.1	0.40	18.0	0.43
2012	638	27.0	0.50	8.6	0.39	13.7	0.43	18.9	0.45
2013	598	25.3	0.51	8.3	0.41	13.1	0.44	17.8	0.47
2014	541	22.9	0.64	7.4	0.51	11.7	0.55	16.0	0.58
1998-2014	9492	28.4	0.45	9.4	0.38	15.0	0.40	20.7	0.42

Table 13

Age distribution of age at death (cancer-related) for period 2007-2014
(incl. multiple primaries)

Age at death Years	Cases			Males			Females		
	n	%	Cum.%	n	%	Cum.%	n	%	Cum.%
15-19	1	0.0	0.0	1	0.0	0.0			0.0
20-24	3	0.0	0.0	1	0.0	0.0	2	0.0	0.0
25-29	10	0.1	0.1	4	0.1	0.1	6	0.1	0.2
30-34	13	0.1	0.2	9	0.1	0.2	4	0.1	0.2
35-39	34	0.3	0.5	17	0.3	0.5	17	0.3	0.6
40-44	100	0.8	1.4	58	0.9	1.3	42	0.8	1.4
45-49	182	1.5	2.9	95	1.4	2.7	87	1.7	3.1
50-54	329	2.8	5.7	198	2.9	5.7	131	2.5	5.6
55-59	558	4.7	10.4	349	5.2	10.9	209	4.1	9.7
60-64	890	7.5	17.8	584	8.7	19.6	306	5.9	15.6
65-69	1374	11.6	29.4	913	13.6	33.1	461	8.9	24.5
70-74	1870	15.7	45.1	1187	17.6	50.8	683	13.3	37.8
75-79	1933	16.3	61.4	1214	18.0	68.8	719	14.0	51.7
80-84	2076	17.5	78.9	1134	16.8	85.6	942	18.3	70.0
85+	2511	21.1	100.0	966	14.4	100.0	1545	30.0	100.0
All ages	11884	100.0		6730	100.0		5154	100.0	

Included in the statistics are 32.7% multiple primaries in males and 25.9% in females.

Table 14

Age-specific mortality (cancer-related) and proportion of all cancers
for period 2007-2014
(incl. multiple primaries)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	MI-index	Females Age- spec. mortal.	MI-index	Males Prop.all cancers %	Females Prop.all cancers %
0- 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19	1		0.1	0.25	0.0		2.8	
20-24	1	2	0.1	0.07	0.2	0.10	2.1	7.1
25-29	4	6	0.3	0.16	0.5	0.18	6.5	9.4
30-34	9	4	0.7	0.18	0.3	0.09	10.2	3.6
35-39	17	17	1.3	0.19	1.3	0.23	9.6	6.6
40-44	58	42	3.6	0.27	2.7	0.25	12.6	6.6
45-49	95	87	6.0	0.24	5.7	0.28	9.3	7.1
50-54	198	131	15.3	0.30	10.2	0.28	10.6	7.3
55-59	349	209	32.9	0.33	18.6	0.32	11.3	8.0
60-64	584	306	59.5	0.38	28.9	0.36	12.2	8.6
65-69	913	461	94.9	0.44	44.2	0.39	12.8	8.8
70-74	1187	683	130.5	0.48	65.3	0.44	13.0	10.4
75-79	1214	719	220.4	0.60	100.8	0.48	14.3	11.4
80-84	1134	942	324.7	0.74	168.0	0.59	15.4	14.4
85+	966	1545	417.2	0.91	267.4	0.76	16.0	17.8
All ages	6730	5154					13.5	11.8
Mortality								
Raw			37.3	0.51	27.5	0.49		
WS			16.6	0.46	8.9	0.40		
ES			26.4	0.49	14.1	0.43		
BRD-S			36.8	0.52	19.4	0.45		
PYLL-70								
per 100,000			121.0		79.0			
ES			105.4		66.7			
AYLL-70			8.7		9.9			

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15a

Multiple primaries in deaths in period 1998-2014

MALES

Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C03-C06 Oral cavity	52	1.1	39	75.0	3	5.8	10	19.2
C15 Oesophagus	77	1.7	13	16.9	15	19.5	49	63.6
C16 Stomach	242	5.2	66	27.3	58	24.0	118	48.8
C17 Small intestine	51	1.1	6	11.8	20	39.2	25	49.0
C18 Colon	352	7.6			132	37.5	220	62.5
C19-C20 Rectum	206	4.5			129	62.6	77	37.4
C22 Liver	130	2.8	5	3.8	30	23.1	95	73.1
C23-C24 Bile	44	1.0	3	6.8	6	13.6	35	79.5
C25 Pancreas	157	3.4	10	6.4	28	17.8	119	75.8
C32 Larynx	71	1.5	50	70.4	1	1.4	20	28.2
C33-C34 Lung	491	10.6	82	16.7	67	13.6	342	69.7
C43 Malign. melanoma	178	3.9	106	59.6	3	1.7	69	38.8
C44 Skin others	248	5.4	118	47.6	21	8.5	109	44.0
C61 Prostate	1015	22.0	578	56.9	83	8.2	354	34.9
C64 Kidney	174	3.8	80	46.0	40	23.0	54	31.0
C67 Bladder	366	7.9	170	46.4	28	7.7	168	45.9
C70-C72 CNS cancer	78	1.7	24	30.8	4	5.1	50	64.1
C76-C79 CUP	53	1.1	9	17.0	9	17.0	35	66.0
C82-C85 NHL	175	3.8	72	41.1	30	17.1	73	41.7
C90 Mult. myeloma	52	1.1	17	32.7	6	11.5	29	55.8
C91-C96 Leukaemia	100	2.2	28	28.0	9	9.0	63	63.0
Other primaries	302	6.5	131	43.4	20	6.6	151	50.0
All mult. primaries	4614	100.0	1607	34.8	742	16.1	2265	49.1

Multiple primaries with number of cases 1 to 33 are pooled in category "Other primaries"

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 15b

Multiple primaries in deaths in period 1998-2014
FEMALES

Diagnosis		Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C16	Stomach	168	5.1	52	31.0	34	20.2	82	48.8
C18	Colon	226	6.9			74	32.7	152	67.3
C19-C20	Rectum	122	3.7			69	56.6	53	43.4
C22	Liver	36	1.1	2	5.6	9	25.0	25	69.4
C23-C24	Bile	43	1.3	9	20.9	8	18.6	26	60.5
C25	Pancreas	140	4.2	11	7.9	19	13.6	110	78.6
C33-C34	Lung	206	6.2	33	16.0	18	8.7	155	75.2
C43	Malign. melanoma	81	2.5	53	65.4	6	7.4	22	27.2
C44	Skin others	98	3.0	54	55.1	8	8.2	36	36.7
C50	Breast	835	25.3	551	66.0	60	7.2	224	26.8
C53	Cervix uteri	116	3.5	86	74.1	10	8.6	20	17.2
C54	Corpus uteri	217	6.6	143	65.9	11	5.1	63	29.0
C56	Ovary	219	6.6	74	33.8	52	23.7	93	42.5
C64	Kidney	71	2.2	35	49.3	12	16.9	24	33.8
C67	Bladder	120	3.6	57	47.5	4	3.3	59	49.2
C70-C72	CNS cancer	60	1.8	27	45.0	8	13.3	25	41.7
C73	Thyroid	31	0.9	17	54.8	3	9.7	11	35.5
C82-C85	NHL	97	2.9	41	42.3	14	14.4	42	43.3
C90	Mult. myeloma	50	1.5	16	32.0	4	8.0	30	60.0
C91-C96	Leukaemia	70	2.1	14	20.0	9	12.9	47	67.1
Other primaries		293	8.9	111	37.9	45	15.4	137	46.8
All mult. primaries		3299	100.0	1386	42.0	477	14.5	1436	43.5

Multiple primaries with number of cases 1 to 30 are pooled in category "Other primaries"

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers
for period 2007-2014
(**First primaries only** *)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	MI-index	Females Age- spec. mortal.	MI-index	Males Prop.all cancers %	Females Prop.all cancers %
0- 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19	1		0.1	0.25	0.0		3.0	
20-24	1	2	0.1	0.07	0.2	0.10	2.3	7.7
25-29	4	6	0.3	0.19	0.5	0.18	7.3	10.2
30-34	9	4	0.7	0.18	0.3	0.10	10.5	4.2
35-39	15	13	1.2	0.19	1.0	0.19	9.1	5.7
40-44	55	39	3.4	0.28	2.5	0.25	13.0	7.0
45-49	91	76	5.8	0.26	5.0	0.27	9.9	7.4
50-54	169	116	13.1	0.29	9.1	0.27	10.6	7.8
55-59	298	175	28.1	0.32	15.6	0.31	11.4	8.2
60-64	487	245	49.6	0.37	23.1	0.35	12.4	8.6
65-69	719	384	74.7	0.43	36.8	0.41	12.7	9.4
70-74	917	528	100.8	0.50	50.5	0.44	13.2	10.4
75-79	883	547	160.3	0.60	76.7	0.47	14.2	11.3
80-84	798	726	228.5	0.78	129.5	0.56	14.9	14.4
85+	692	1205	298.9	0.96	208.5	0.75	15.7	17.8
All ages	5139	4066					13.3	11.8
Mortality								
Raw			28.4	0.50	21.7	0.48		
WS			13.0	0.45	7.2	0.39		
ES			20.4	0.48	11.3	0.42		
BRD-S			27.9	0.51	15.4	0.44		
PYLL-70								
per 100,000			104.8		67.7			
ES			91.2		57.2			
AYLL-70			9.1		10.1			

* See corresponding tables with multiple primaries.

Table 17

Age-specific mortality (cancer-related) and proportion of all cancers
for period 2007-2014
(Single primaries only *)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	MI-index	Females Age- spec. mortal.	MI-index	Males Prop.all cancers %	Females Prop.all cancers %
0- 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19	1		0.1	0.25	0.0		3.0	
20-24	1	2	0.1	0.07	0.2	0.11	2.6	8.3
25-29	4	6	0.3	0.20	0.5	0.18	7.8	10.7
30-34	9	4	0.7	0.18	0.3	0.10	10.6	4.8
35-39	15	12	1.2	0.19	1.0	0.18	9.5	5.9
40-44	53	38	3.3	0.27	2.5	0.25	13.4	7.5
45-49	88	74	5.6	0.26	4.9	0.26	10.2	8.1
50-54	159	110	12.3	0.29	8.6	0.28	11.1	8.3
55-59	264	158	24.9	0.30	14.1	0.30	11.3	8.4
60-64	437	216	44.5	0.36	20.4	0.33	12.8	8.9
65-69	610	319	63.4	0.40	30.6	0.37	12.8	9.4
70-74	718	441	78.9	0.42	42.2	0.39	12.7	10.7
75-79	680	457	123.5	0.52	64.1	0.41	14.1	11.5
80-84	574	606	164.3	0.61	108.1	0.50	14.1	14.8
85+	493	998	212.9	0.72	172.7	0.64	14.4	17.7
All ages	4106	3441					13.0	12.0
Mortality								
Raw			22.7	0.43	18.4	0.43		
WS			10.7	0.40	6.2	0.36		
ES			16.5	0.41	9.7	0.38		
BRD-S			22.2	0.44	13.1	0.39		
PYLL-70								
per 100,000			96.2		62.6			
ES			83.8		53.0			
AYLL-70			9.4		10.6			

* See corresponding tables with multiple primaries.





Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at colorectal cancer-related death (see Table 10) should be considered.

Average mortality (world standard population) 2007 - 2014: Males



Average mortality (world standard population) 2007 - 2014: Females





Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2007 to 2014. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 15.6/100,000 WS N=6,353, females 8.5/100,000 WS N=4,923).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 65,347 female residents (averaged) in the period from 2007 to 2014 a total of 107 women died from colorectal cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 7.4/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 5.4 and 10.2/100,000.

Standardized mortality ratio (SMR) 2007 - 2014: Males



Standardized mortality ratio (SMR) 2007 - 2014: Females




Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=6,353, females N=4,923).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,924 female residents (averaged) in the period from 2007 to 2014 a total of 107 women died from colorectal cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.85. Though, the value of this parameter may vary with an underlying probability of 99% between 0.65 and 1.08, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases, where applicable. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

FRG	Federal Republic of Germany
GEKID	Association of Population-based Cancer Registries in Germany (Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)
MCR	Munich Cancer Registry (Tumorregister München)
SEER	Surveillance, Epidemiology, and End Results (USA)
AYLL-70	Average years of life lost prior to age 70 given a person dies before that age
BRD-S	German standard population
DCO	Death certificate only
EAR	Excess absolute risk = excess cancer cases (O - E) per 10,000 person-years
ES	European standard population (old)
LCL	Lower confidence limit
MI-index	Ratio between mortality and incidence
PYLL-70	Potential years of life lost prior to age 70 given a person dies before that age
SIR	Standardized incidence ratio
SMR	Standardized mortality ratio
UCL	Upper confidence limit
WS	World standard population

Recommended Citation

Munich Cancer Registry. ICD-10 C18-C20: Colorectal cancer - Incidence and Mortality [Internet]. 2016 [updated 2016 Apr 13; cited 2016 Jun 1]. Available from: <http://www.tumorregister-muenchen.de/en/facts/base/bC1820E-ICD-10-C18-C20-Colorectal-cancer-incidence-and-mortality.pdf>

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

Fig./Tbl.	Page
1 Pts cohorts, DCO, mult. prim., follow-up / yr	3
1a Gender distribution by year of diagnosis	4
2 Incidence by year of diagnosis	5
3 Age distribution parameters by year of diagnosis	6
4 Age distribution by 5-year age group and gender	8
5 Age-specific incidence, DCO rate, proportion malignancies	9
6 Age distribution and age-specific incidence (chart)	10
6a Age-specific incidence internationally (chart)	11
7 Cumulative follow-up years (chart)	12
8 Standardized incidence ratio of second primaries	13
9a Map of cancer incidence (WS) by county (chart)	16
9b Standardized incidence ratio (SIR) by county (chart)	17
10a Pts incident cohorts and mortality / yr	18
10b Incidence and mortality by year of diagnosis	19
10c Cancer-related deaths, death certification available / yr	20
11 Medians of age at death / yr	21
12 Mortality by year of death	23
13 Distribution of age at death	24
14 Age-specific mortality	25
15 Multiple primaries in deaths	26
16 Age-specific mortality (first primaries)	28
17 Age-specific mortality (single primaries)	29
18 Age distribution and age-specific mortality (chart)	30
19a Map of cancer mortality (WS) by county (chart)	31
19b Standardized mortality ratio (SMR) by county (chart)	32