

Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ Deutsch

ICD-10 C26: Other digestive organ cancer

Incidence and Mortality



Year of diagnosis	1998-2014
Patients	442
Diseases	442
Creation date	04/13/2016
Export date	12/23/2015
Population	4.64 m



Munich Cancer Registry at Munich Cancer Center
Marchioninstr. 15
Munich, 81377
Germany

<http://www.tumorregister-muenchen.de/en>

http://www.tumorregister-muenchen.de/en/facts/base/bC26__E-ICD-10-C26-Other-digestive-organ-cancer-incidence-and-mortality.pdf

**Global Statements about the statistics on the Internet –
Baseline Statistics** (grey button ) , **Survival** (red button )

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.64 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases^{###} are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, April 2016

[#] Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007).

^{##} Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.

^{###} DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate.

ICD-10 codes (ICD-10 2015) used for specifying cancer site

Code	Description
C26.-	Malignant neoplasm of other and ill-defined digestive organs
C26.0	Intestinal tract, part unspecified
C26.1	Spleen
C26.8	Overlapping lesion of digestive system
C26.9	Ill-defined sites within the digestive system

INCIDENCE

Table 1

All patients with invasive cancer by year of diagnosis, proportions of DCO, multiple primaries, deaths, and active follow-up (incl. DCO)

Year of diagnosis	Cases n	DCO cases n	Prop. DCO %	Prop. mult. primaries %	Prop. deaths %	Prop. actively followed %
1998	13	12	92.3	7.7	100.0	100.0
1999	12	6	50.0	8.3	91.7	91.7
2000	11	9	81.8	9.1	100.0	100.0
2001	18	14	77.8	16.7	100.0	100.0
2002	47	34	72.3	23.4	97.9	100.0 #
2003	42	30	71.4	21.4	95.2	97.6
2004	42	25	59.5	19.0	97.6	100.0
2005	27	16	59.3	18.5	100.0	100.0
2006	25	8	32.0	24.0	100.0	100.0
2007	35	16	45.7	22.9	100.0	100.0 #
2008	27	13	48.1	22.2	92.6	100.0
2009	13	8	61.5	15.4	100.0	100.0
2010	20	17	85.0	35.0	100.0	100.0
2011	26	18	69.2	34.6	100.0	100.0
2012	33	18	54.5	18.2	97.0	97.0
2013	35	17	48.6	31.4	94.3	100.0
2014	16	13	81.3	25.0	93.8	100.0 ##
1998-2014	442	274	62.0	22.2	97.5	99.3

The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be found in the respective headings.

Table 1a

All patients with invasive cancer
by year of diagnosis and gender
(incl. DCO)

Year of diagnosis	All n	Males n	Females n	Prop. males %
1998	13	7	6	53.8
1999	12	1	11	8.3
2000	11	4	7	36.4
2001	18	8	10	44.4
2002	47	18	29	38.3
2003	42	14	28	33.3
2004	42	20	22	47.6
2005	27	7	20	25.9
2006	25	12	13	48.0
2007	35	20	15	57.1
2008	27	14	13	51.9
2009	13	4	9	30.8
2010	20	8	12	40.0
2011	26	8	18	30.8
2012	33	18	15	54.5
2013	35	17	18	48.6
2014	16	1	15	6.3
1998-2014	442	181	261	41.0

Table 2

Incidence measures by year of diagnosis including DCO cases
(with respect to registry area expansion from 2.51 to 3.96 m as of 2002,
and from 3.96 to 4.64 m as of 2007, respectively)

Year of diagnosis	Males n	Females n	Males Inc. raw	Fem. Inc. raw	Males Inc. WS	Fem. Inc. WS	Males Inc. ES	Fem. Inc. ES	Males Inc. BRD-S	Fem. Inc. BRD-S
1998	7	6	0.6	0.5	0.4	0.1	0.6	0.2	0.8	0.3
1999	1	11	0.1	0.9	0.0	0.3	0.1	0.5	0.2	0.7
2000	4	7	0.4	0.6	0.2	0.1	0.3	0.2	0.3	0.4
2001	8	10	0.7	0.8	0.4	0.2	0.6	0.3	0.9	0.5
2002	18	29	1.0	1.5	0.5	0.4	0.8	0.7	1.1	1.0
2003	14	28	0.7	1.4	0.4	0.3	0.7	0.6	0.9	1.0
2004	20	22	1.1	1.1	0.5	0.3	0.9	0.5	1.3	0.7
2005	7	20	0.4	1.0	0.2	0.3	0.3	0.5	0.4	0.6
2006	12	13	0.6	0.6	0.3	0.3	0.5	0.4	0.7	0.5
2007	20	15	0.9	0.6	0.4	0.3	0.7	0.4	1.0	0.5
2008	14	13	0.6	0.6	0.3	0.1	0.5	0.2	0.6	0.3
2009	4	9	0.2	0.4	0.1	0.1	0.1	0.2	0.2	0.3
2010	8	12	0.4	0.5	0.2	0.1	0.3	0.2	0.3	0.3
2011	8	18	0.4	0.8	0.2	0.2	0.3	0.3	0.3	0.5
2012	18	15	0.8	0.6	0.4	0.1	0.6	0.3	0.8	0.4
2013	17	18	0.7	0.8	0.3	0.2	0.5	0.3	0.8	0.5
2014	1	15	0.0	0.6	0.0	0.2	0.0	0.3	0.0	0.4
1998-2014	181	261	0.6	0.8	0.3	0.2	0.4	0.4	0.6	0.5

The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

Table 3

Age distribution parameters by year of diagnosis (All patients)
(incl. DCO)

Year of diagnosis	Cases n	Std.		Median						
		Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	13	76.8	13.5	50.3	97.7	54.2	72.9	77.6	86.3	89.7
1999	12	76.7	17.0	30.8	95.5	60.3	76.2	79.6	85.2	91.4
2000	11	78.5	11.1	63.6	94.3	64.5	68.2	79.2	89.8	90.8
2001	18	80.9	14.9	42.6	97.1	50.7	77.9	85.4	89.7	95.2
2002	47	77.6	15.7	40.8	96.4	48.0	72.2	82.1	89.2	92.9
2003	42	79.6	10.0	57.0	94.7	62.5	75.8	80.2	88.5	89.4
2004	42	80.2	11.9	49.7	96.9	64.3	71.2	82.7	89.6	95.0
2005	27	79.2	11.9	52.2	95.9	63.4	69.4	80.5	89.8	93.5
2006	25	74.9	12.4	49.9	92.9	60.7	64.9	74.5	85.4	92.5
2007	35	71.4	12.4	40.7	94.0	53.8	64.5	71.5	81.4	85.8
2008	27	78.2	11.4	45.1	94.1	61.0	72.5	78.9	88.3	89.2
2009	13	79.8	14.6	48.0	99.7	62.3	68.1	83.3	88.5	95.4
2010	20	80.0	12.9	51.8	95.6	55.3	72.7	84.3	89.2	91.5
2011	26	81.6	11.5	49.8	96.2	64.5	73.2	84.7	89.6	92.7
2012	33	80.2	10.6	56.9	96.5	63.4	74.1	83.3	86.8	91.9
2013	35	81.0	8.4	61.1	93.2	70.5	75.4	81.2	87.7	92.4
2014	16	81.2	8.0	68.0	96.1	72.0	74.4	80.1	87.0	91.8
1998-2014	442	78.6	12.3	30.8	99.7	61.8	71.9	81.1	88.3	92.5

Table 3a

Age distribution parameters by year of diagnosis (MALES)
(incl. DCO)

Year of diagnosis	Cases n	Std.		Median						
		Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	7	68.4	11.5	50.3	79.6	50.3	54.2	72.9	77.6	79.6
1999	1	83.6		83.6	83.6	83.6	83.6	83.6	83.6	83.6
2000	4	66.3	2.6	63.6	68.8	63.6	64.1	66.3	68.5	68.8
2001	8	72.5	18.7	42.6	97.1	42.6	56.9	78.4	84.9	97.1
2002	18	72.0	15.9	40.8	93.6	44.8	61.6	75.1	84.1	90.1
2003	14	75.2	13.0	57.0	94.7	58.7	63.7	76.2	88.3	93.9
2004	20	77.9	12.0	49.7	96.1	61.3	70.5	79.1	86.5	92.1
2005	7	71.8	5.3	63.4	80.4	63.4	69.2	73.1	74.2	80.4
2006	12	75.3	11.0	49.9	85.8	64.9	68.4	77.5	85.3	85.5
2007	20	71.5	13.2	40.7	90.9	53.5	64.6	75.3	81.4	84.8
2008	14	72.0	11.1	45.1	86.3	60.2	66.2	73.7	78.9	85.1
2009	4	75.6	22.2	48.0	95.4	48.0	57.7	79.5	93.5	95.4
2010	8	74.4	13.1	53.9	88.8	53.9	64.3	77.8	84.3	88.8
2011	8	80.8	11.3	64.5	94.2	64.5	70.2	83.3	90.1	94.2
2012	18	77.7	11.4	56.9	91.9	62.5	66.1	81.7	86.8	91.1
2013	17	79.6	7.4	61.8	93.0	71.6	75.4	79.9	85.3	88.5
2014	1	77.4		77.4	77.4	77.4	77.4	77.4	77.4	77.4
1998-2014	181	74.8	12.4	40.7	97.1	57.0	67.3	76.3	84.1	88.9

Table 3b

Age distribution parameters by year of diagnosis (FEMALES)
(incl. DCO)

Year of diagnosis	Cases n	Std.		Median						
		Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	6	86.5	8.1	73.0	97.7	73.0	83.7	87.5	89.7	97.7
1999	11	76.1	17.6	30.8	95.5	60.3	75.1	77.7	86.7	91.4
2000	7	85.5	6.7	75.5	94.3	75.5	79.2	85.7	90.8	94.3
2001	10	87.6	5.7	77.9	95.2	78.7	84.0	88.3	92.3	94.5
2002	29	81.1	14.8	42.2	96.4	50.4	80.1	84.1	89.9	94.6
2003	28	81.8	7.5	62.1	90.1	75.7	77.1	81.7	88.7	89.4
2004	22	82.2	11.7	60.7	96.9	66.3	74.4	84.6	94.5	95.5
2005	20	81.8	12.5	52.2	95.9	61.2	78.0	83.8	92.2	94.0
2006	13	74.5	14.1	54.6	92.9	60.7	61.9	73.8	90.6	92.6
2007	15	71.3	11.8	46.5	94.0	57.9	64.3	71.3	78.1	85.8
2008	13	84.8	7.4	70.9	94.1	72.6	84.0	88.3	89.1	90.4
2009	9	81.7	11.0	62.3	99.7	62.3	81.2	83.3	87.0	99.7
2010	12	83.7	11.9	51.8	95.6	70.5	81.9	86.6	90.7	91.8
2011	18	81.9	11.8	49.8	96.2	63.3	78.9	84.7	89.6	92.7
2012	15	83.2	9.1	60.2	96.5	74.1	79.5	83.7	89.1	94.9
2013	18	82.2	9.3	61.1	93.2	68.4	77.6	82.3	90.4	92.8
2014	15	81.4	8.2	68.0	96.1	72.0	74.2	80.9	88.0	91.8
1998-2014	261	81.3	11.5	30.8	99.7	64.3	75.9	83.7	89.4	93.1

Table 4

Age distribution by 5-year age group and gender for period 2007–2014
(incl. DCO)

Age at diagnosis Years	Cases			Males			Females		
	n	%	Cum.%	n	%	Cum.%	n	%	Cum.%
40-44	1	0.5	0.5	1	1.1	1.1			0.0
45-49	4	2.0	2.4	2	2.2	3.3	2	1.7	1.7
50-54	5	2.4	4.9	4	4.4	7.8	1	0.9	2.6
55-59	3	1.5	6.3	2	2.2	10.0	1	0.9	3.5
60-64	14	6.8	13.2	8	8.9	18.9	6	5.2	8.7
65-69	14	6.8	20.0	8	8.9	27.8	6	5.2	13.9
70-74	28	13.7	33.7	11	12.2	40.0	17	14.8	28.7
75-79	23	11.2	44.9	15	16.7	56.7	8	7.0	35.7
80-84	40	19.5	64.4	15	16.7	73.3	25	21.7	57.4
85+	73	35.6	100.0	24	26.7	100.0	49	42.6	100.0
All ages	205	100.0		90	100.0		115	100.0	

Included in the statistics are 31.1% multiple primaries in males and 31.3% in females.

Table 5

Age-specific incidence, DCO rate and proportion of all cancers
for period 2007-2014

Age at diagnosis Years	Males n	Females n	Males Age- spec. incid.	Females Age- spec. incid.	Males DCO rate n=48 %	Females DCO rate n=72 %	Males	Females
							Prop.all cancers %	Prop.all cancers %
0- 4			0.0	0.0				
5- 9			0.0	0.0				
10-14			0.0	0.0				
15-19			0.0	0.0				
20-24			0.0	0.0				
25-29			0.0	0.0				
30-34			0.0	0.0				
35-39			0.0	0.0				
40-44	1		0.1	0.0			0.1	
45-49	2	2	0.1	0.1			0.1	0.0
50-54	4	1	0.3	0.1	75.0	100.0	0.1	0.0
55-59	2	1	0.2	0.1	100.0		0.0	0.0
60-64	8	6	0.8	0.6	25.0	33.3	0.1	0.1
65-69	8	6	0.8	0.6	37.5	33.3	0.1	0.1
70-74	11	17	1.2	1.6	45.5	23.5	0.1	0.1
75-79	15	8	2.7	1.1	40.0	50.0	0.1	0.1
80-84	15	25	4.3	4.5	66.7	68.0	0.2	0.3
85+	24	49	10.4	8.5	70.8	85.7	0.4	0.5
All ages	90	115			53.3	62.6	0.1	0.1
Incidence								
Raw			0.5	0.6				
WS			0.2	0.2				
ES			0.4	0.3				
BRD-S			0.5	0.4				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).

ICD-10 C26: Malignant neoplasm of other and ill-defined digestive organs

Age distribution and age-specific incidence 2007 - 2014 (Males: 90, Females: 115)




Figure 6. Age distribution and age-specific incidence

ICD-10 C26: Malignant neoplasm of other and ill-defined digestive organs

Age-specific incidence rates: international comparison




Figure 6a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).

Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2014, based on the November 2013 submission. <http://www.seer.cancer.gov>.

ICD-10 C26: Malignant neoplasm of other and ill-defined digestive organs

Cumulative follow-up years since diagnosis for period 1998 - 2014 (excl. DCO)




Figure 7. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.

Table 8a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2014

MALES

Diagnosis	Observed n	Expected n	SIR	LCL 95%	UCL 95%	EAR	DCO %
C33-C34 Lung	2	0.2	9.2	1.1	33.1 #	150.5	
C61 Prostate	2	0.6	3.5	0.4	12.8	121.2	
Other primaries	6	0.4	13.5	5.0	29.4 #	469.4	50.0
Not observed	0	0.7	0.0	0.0	5.4	-57.4	
All mult. primaries	10	1.9	5.2	2.5	9.6 #	683.8	30.0
Patients			105				
Median age at second malignancy (years)			68.4				
Person-years			118				
Mean observation time (years)			1.1				
Median observation time (years)			0.2				

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries"

Table 8b


Standardized incidence ratio (SIR, with 95% confidence limits),
 excess absolute risk (EAR) and DCO rate of second primaries
 for period 1998-2014
 FEMALES

Diagnosis	Observed n	Expected n	SIR	LCL 95%	UCL 95%	EAR	DCO %
C76-C79 CUP	3	0.0	91.3	18.8	266.8 #	395.7	33.3
Other primaries	8	0.4	19.3	8.3	37.9 #	1011	25.0
Not observed	0	0.7	0.0	0.0	5.0	-97.5	
All mult. primaries	11	1.2	9.3	4.7	16.7 #	1310	27.3
Patients			137				
Median age at second malignancy (years)			79.4				
Person-years			75				
Mean observation time (years)			0.5				
Median observation time (years)			0.2				

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries"

Average incidence (world standard population) 2007 - 2014: Males



Average incidence (world standard population) 2007 - 2014: Females





Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.2/100,000 WS N=90, females 0.2/100,000 WS N=115).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 65,347 female residents (averaged) in the period from 2007 to 2014 a total of 0 women were identified with newly diagnosed other digestive organ cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.0/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.0/100,000.

Standardized incidence ratio (SIR) 2007 - 2014: Males



Standardized incidence ratio (SIR) 2007 - 2014: Females




Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=90, females N=115).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,924 female residents (averaged) in the period from 2007 to 2014 a total of 0 women were identified with newly diagnosed other digestive organ cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.00. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 1.84, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

Year of diagnosis	Incident cases n	Prop. actively followed %	Prop. DCO %	Deaths n	Prop. deaths %	Prop. deaths with death certific. %
1998	13	100.0	92.3	13	100.0	100.0
1999	12	91.7	50.0	11	91.7	100.0
2000	11	100.0	81.8	11	100.0	100.0
2001	18	100.0	77.8	18	100.0	100.0
2002	47	100.0	72.3	46	97.9	100.0
2003	42	97.6	71.4	40	95.2	100.0
2004	42	100.0	59.5	41	97.6	97.6
2005	27	100.0	59.3	27	100.0	100.0
2006	25	100.0	32.0	25	100.0	100.0
2007	35	100.0	45.7	35	100.0	97.1
2008	27	100.0	48.1	25	92.6	96.0
2009	13	100.0	61.5	13	100.0	84.6
2010	20	100.0	85.0	20	100.0	100.0
2011	26	100.0	69.2	26	100.0	96.2
2012	33	97.0	54.5	32	97.0	100.0
2013	35	100.0	48.6	33	94.3	100.0
2014	16	100.0	81.3	15	93.8	100.0
1998-2014	442	99.3	62.0	431	97.5	98.6

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)
(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

Year of diagnosis/ death	Incident cases n	Deaths n	Prop. deaths with death certific. %	Deaths in same year n	Prop. deaths in same year %
1998	13	15	100.0	9	69.2
1999	12	12	100.0	9	75.0
2000	11	12	100.0	9	81.8
2001	18	17	100.0	14	77.8
2002	47	30	100.0	35	74.5
2003	42	24	100.0	29	69.0
2004	42	36	97.2	34	81.0
2005	27	27	100.0	22	81.5
2006	25	24	100.0	19	76.0
2007	35	28	96.4	29	82.9
2008	27	18	100.0	21	77.8
2009	13	11	90.9	11	84.6
2010	20	12	100.0	18	90.0
2011	26	19	84.2	23	88.5
2012	33	22	100.0	29	87.9
2013	35	28	100.0	29	82.9
2014	16	15	100.0	15	93.8
1998-2014	442	350	98.3	355	80.3

Table 10c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates
(incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

Year of death	Deaths n	Prop. cancer- related %	Prop. non-cancer- related %	Prop. cancer recorded on death certificate %
1998	15	80.0	20.0	100.0
1999	12	75.0	25.0	91.7
2000	12	100.0		100.0
2001	17	58.8	41.2	100.0
2002	30	86.7	13.3	100.0
2003	24	87.5	12.5	95.8
2004	36	94.4	5.6	97.1
2005	27	96.3	3.7	100.0
2006	24	87.5	12.5	100.0
2007	28	100.0		100.0
2008	18	100.0		100.0
2009	11	90.9	9.1	100.0
2010	12	100.0		91.7
2011	19	73.7	26.3	100.0
2012	22	100.0		100.0
2013	28	89.3	10.7	92.9
2014	15	100.0		100.0
1998-2014	350	90.0	10.0	98.3

Table 11a

Medians of age at death according to the grouping in Table 10

MALES

Year of death	Deaths n	Age at death (all causes) Years	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate) Years
1998	8	75.2	73.0	77.6	75.2
1999	3	71.1	67.8	71.1	71.1
2000	3	68.2	68.2		68.2
2001	7	79.3	79.3		79.3
2002	10	81.7	81.3	90.1	81.7
2003	9	62.4	62.4		62.4
2004	11	77.3	77.3	79.4	77.3
2005	10	77.8	77.5	84.2	77.8
2006	11	76.0	78.3	74.5	76.0
2007	14	73.4	73.4		71.6
2008	11	75.0	75.0		75.0
2009	5	72.8	72.8		72.8
2010	3	73.5	73.5		81.2
2011	10	72.2	73.2	71.1	83.4
2012	9	83.6	83.6		83.6
2013	16	76.2	76.2	81.4	76.1
2014	3	66.2	66.2		66.2
1998-2014	143	75.1	75.0	76.3	75.4

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 11b

Medians of age at death according to the grouping in Table 10
FEMALES

Year of death	Deaths n	Age at death (all causes) Years	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate) Years
1998	7	88.7	88.7	92.0	88.7
1999	9	81.6	86.8	77.5	84.2
2000	9	84.3	84.3		84.3
2001	10	88.3	87.2	88.4	88.3
2002	20	88.3	89.2	81.1	88.3
2003	15	87.6	88.4	75.9	87.9
2004	25	86.1	86.1		86.1
2005	17	83.1	83.1		83.1
2006	13	78.3	78.3	79.4	78.3
2007	14	69.2	69.2		69.2
2008	7	88.4	88.4		88.4
2009	6	82.5	83.3	24.5	83.3
2010	9	81.8	81.8		81.8
2011	9	80.8	79.4	84.4	80.8
2012	13	84.5	84.5		84.5
2013	12	82.4	81.2	93.1	81.2
2014	12	82.8	82.8		82.8
1998-2014	207	84.0	84.0	84.0	84.0

By 2010, life expectancy at birth was 77.5 years for boys and 82.6 years for girls.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 12a

Mortality measures (cancer-related death) and mortality-incidence-index
by year of death

MALES

Year of death	Deaths n	Mort. raw	MI-Index raw	Mort. WS	MI-Index WS	Mort. ES	MI-Index ES	Mort. BRD-S	MI-Index BRD-S
1998	7	0.6	1.00	0.4	0.93	0.6	0.99	0.8	1.06
1999	2	0.2	2.00	0.1	2.51	0.2	2.05	0.3	1.45
2000	3	0.3	0.75	0.2	0.77	0.2	0.78	0.2	0.78
2001	7	0.6	0.88	0.3	0.84	0.6	0.88	0.8	0.91
2002	9	0.5	0.50	0.2	0.45	0.4	0.49	0.6	0.55
2003	9	0.5	0.64	0.3	0.76	0.4	0.65	0.5	0.55
2004	9	0.5	0.45	0.2	0.47	0.4	0.46	0.5	0.43
2005	9	0.5	1.29	0.2	1.11	0.4	1.32	0.5	1.52
2006	10	0.5	0.83	0.2	0.79	0.4	0.81	0.6	0.88
2007	14	0.6	0.70	0.3	0.70	0.5	0.71	0.7	0.70
2008	11	0.5	0.79	0.2	0.70	0.3	0.74	0.5	0.79
2009	5	0.2	1.25	0.1	1.11	0.2	1.09	0.2	1.09
2010	3	0.1	0.38	0.1	0.34	0.1	0.34	0.1	0.32
2011	7	0.3	0.88	0.2	1.03	0.2	0.94	0.3	0.84
2012	9	0.4	0.50	0.2	0.48	0.3	0.49	0.4	0.51
2013	14	0.6	0.82	0.2	0.90	0.4	0.84	0.6	0.78
2014	3	0.1	3.00	0.1	5.31	0.1	3.78	0.1	2.65
1998-2014	131	0.4	0.72	0.2	0.72	0.3	0.72	0.4	0.72

Table 12b

Mortality measures (cancer-related death) and mortality-incidence-index
by year of death

FEMALES

Year of death	Deaths n	Mort. raw	MI-Index raw	Mort. WS	MI-Index WS	Mort. ES	MI-Index ES	Mort. BRD-S	MI-Index BRD-S
1998	5	0.4	0.83	0.1	0.87	0.2	0.86	0.3	0.88
1999	7	0.6	0.64	0.1	0.44	0.2	0.52	0.4	0.54
2000	9	0.7	1.29	0.2	1.62	0.3	1.42	0.5	1.38
2001	3	0.2	0.30	0.1	0.31	0.1	0.31	0.2	0.31
2002	17	0.9	0.59	0.3	0.72	0.5	0.67	0.5	0.56
2003	12	0.6	0.43	0.2	0.46	0.3	0.46	0.4	0.39
2004	25	1.3	1.14	0.4	1.12	0.6	1.11	0.8	1.06
2005	17	0.9	0.85	0.2	0.72	0.3	0.76	0.6	0.88
2006	11	0.5	0.85	0.2	0.77	0.3	0.80	0.4	0.85
2007	14	0.6	0.93	0.3	0.97	0.4	0.99	0.5	0.92
2008	7	0.3	0.54	0.1	0.69	0.1	0.62	0.2	0.61
2009	5	0.2	0.56	0.0	0.49	0.1	0.51	0.1	0.51
2010	9	0.4	0.75	0.1	1.10	0.2	0.95	0.3	0.94
2011	7	0.3	0.39	0.1	0.59	0.2	0.52	0.2	0.51
2012	13	0.6	0.87	0.1	0.90	0.2	0.88	0.3	0.81
2013	11	0.5	0.61	0.1	0.68	0.2	0.65	0.3	0.65
2014	12	0.5	0.80	0.1	0.73	0.2	0.75	0.3	0.77
1998-2014	184	0.6	0.70	0.2	0.75	0.3	0.73	0.4	0.71

Table 13

Age distribution of age at death (cancer-related) for period 2007-2014
(incl. multiple primaries)

Age at death Years	Cases			Males			Females		
	n	%	Cum.%	n	%	Cum.%	n	%	Cum.%
40-44	1	0.7	0.7	1	1.5	1.5			0.0
45-49	3	2.1	2.8	1	1.5	3.0	2	2.6	2.6
50-54	5	3.5	6.3	3	4.5	7.6	2	2.6	5.1
55-59	3	2.1	8.3	1	1.5	9.1	2	2.6	7.7
60-64	11	7.6	16.0	5	7.6	16.7	6	7.7	15.4
65-69	14	9.7	25.7	9	13.6	30.3	5	6.4	21.8
70-74	24	16.7	42.4	12	18.2	48.5	12	15.4	37.2
75-79	18	12.5	54.9	12	18.2	66.7	6	7.7	44.9
80-84	24	16.7	71.5	8	12.1	78.8	16	20.5	65.4
85+	41	28.5	100.0	14	21.2	100.0	27	34.6	100.0
All ages	144	100.0		66	100.0		78	100.0	

Included in the statistics are 31.1% multiple primaries in males and 31.3% in females.

Table 14

Age-specific mortality (cancer-related) and proportion of all cancers
for period 2007-2014
(incl. multiple primaries)

Age at death Years	Males n	Females n	Males Age- spec. mortal. MI-index	Females Age- spec. mortal. MI-index	Males Prop.all cancers %	Females Prop.all cancers %		
0- 4			0.0	0.0				
5- 9			0.0	0.0				
10-14			0.0	0.0				
15-19			0.0	0.0				
20-24			0.0	0.0				
25-29			0.0	0.0				
30-34			0.0	0.0				
35-39			0.0	0.0				
40-44	1		0.1	1.00	0.0	0.2		
45-49	1	2	0.1	0.50	0.1	1.00	0.1	0.2
50-54	3	2	0.2	0.75	0.2	2.00	0.2	0.1
55-59	1	2	0.1	0.50	0.2	2.00	0.0	0.1
60-64	5	6	0.5	0.63	0.6	1.00	0.1	0.2
65-69	9	5	0.9	1.13	0.5	0.83	0.1	0.1
70-74	12	12	1.3	1.09	1.1	0.71	0.1	0.2
75-79	12	6	2.2	0.80	0.8	0.75	0.1	0.1
80-84	8	16	2.3	0.53	2.9	0.64	0.1	0.2
85+	14	27	6.0	0.58	4.7	0.55	0.2	0.3
All ages	66	78					0.1	0.2
Mortality								
Raw			0.4	0.73	0.4	0.68		
WS			0.2	0.74	0.1	0.79		
ES			0.3	0.73	0.2	0.75		
BRD-S			0.4	0.71	0.3	0.73		
PYLL-70								
per 100,000			1.1		1.0			
ES			0.9		0.9			
AYLL-70			8.8		9.6			

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15a

Multiple primaries in deaths in period 1998-2014
MALES

Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C12-C13 Hypopharynx	1	2.3			1	100.0		
C16 Stomach	1	2.3	1	100.0				
C17 Small intestine	1	2.3	1	100.0				
C18 Colon	3	6.8	2	66.7			1	33.3
C23-C24 Bile	2	4.5	1	50.0	1	50.0		
C25 Pancreas	1	2.3			1	100.0		
C30-C31 Sinuses	1	2.3	1	100.0				
C32 Larynx	1	2.3	1	100.0				
C33-C34 Lung	4	9.1	1	25.0	2	50.0	1	25.0
C43 Malign. melanoma	2	4.5					2	100.0
C44 Skin others	4	9.1	1	25.0			3	75.0
C61 Prostate	13	29.5	11	84.6	2	15.4		
C64 Kidney	2	4.5			2	100.0		
C67 Bladder	2	4.5	2	100.0				
C76-C79 CUP	5	11.4	2	40.0	3	60.0		
C82-C85 NHL	1	2.3			1	100.0		
All mult. primaries	44	100.0	24	54.5	13	29.5	7	15.9

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 15b

Multiple primaries in deaths in period 1998-2014
FEMALES

Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C03-C06 Oral cavity	1	2.0	1	100.0				
C07-C08 Salivary gland	1	2.0			1	100.0		
C09-C10 Oropharynx	1	2.0	1	100.0				
C16 Stomach	2	4.0	2	100.0				
C17 Small intestine	1	2.0					1	100.0
C18 Colon	3	6.0	3	100.0				
C22 Liver	1	2.0			1	100.0		
C40-C41 Bone	1	2.0	1	100.0				
C44 Skin others	1	2.0	1	100.0				
C48 Peritoneal	1	2.0	1	100.0				
C50 Breast	11	22.0	7	63.6	4	36.4		
C53 Cervix uteri	2	4.0	1	50.0			1	50.0
C54 Corpus uteri	3	6.0	3	100.0				
C56 Ovary	4	8.0	2	50.0	1	25.0	1	25.0
C64 Kidney	2	4.0	2	100.0				
C67 Bladder	3	6.0	2	66.7	1	33.3		
C70-C72 CNS cancer	2	4.0	1	50.0	1	50.0		
C73 Thyroid	2	4.0	2	100.0				
C76-C79 CUP	4	8.0			2	50.0	2	50.0
C82-C85 NHL	1	2.0	1	100.0				
C90 Mult. myeloma	2	4.0	1	50.0	1	50.0		
C91-C96 Leukaemia	1	2.0					1	100.0
All mult. primaries	50	100.0	32	64.0	12	24.0	6	12.0

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers
for period 2007-2014
(**First primaries only** *)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	MI-index	Females Age- spec. mortal.	MI-index	Males Prop.all cancers %	Females Prop.all cancers %
0- 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29			0.0		0.0			
30-34			0.0		0.0			
35-39			0.0		0.0			
40-44	1		0.1	1.00	0.0		0.2	
45-49	1	2	0.1	0.50	0.1	1.00	0.1	0.2
50-54	2	2	0.2	0.67	0.2	2.00	0.1	0.1
55-59	1	2	0.1	1.00	0.2	2.00	0.0	0.1
60-64	4	4	0.4	0.57	0.4	0.80	0.1	0.1
65-69	7	3	0.7	1.40	0.3	0.75	0.1	0.1
70-74	9	9	1.0	1.50	0.9	0.69	0.1	0.2
75-79	9	3	1.6	0.90	0.4	0.60	0.1	0.1
80-84	6	9	1.7	0.43	1.6	0.56	0.1	0.2
85+	8	24	3.5	0.44	4.2	0.60	0.2	0.4
All ages	48	58					0.1	0.2
Mortality								
Raw			0.3	0.72	0.3	0.67		
WS			0.1	0.74	0.1	0.77		
ES			0.2	0.71	0.2	0.74		
BRD-S			0.3	0.69	0.2	0.70		
PYLL-70								
per 100,000			0.9		0.9			
ES			0.8		0.8			
AYLL-70			9.1		11.0			

* See corresponding tables with multiple primaries.

Table 17

Age-specific mortality (cancer-related) and proportion of all cancers
for period 2007-2014
(Single primaries only *)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	MI-index	Females Age- spec. mortal.	MI-index	Males Prop.all cancers %	Females Prop.all cancers %
0- 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29			0.0		0.0			
30-34			0.0		0.0			
35-39			0.0		0.0			
40-44	1		0.1	1.00	0.0		0.3	
45-49	1	2	0.1	1.00	0.1	1.00	0.1	0.2
50-54	2	2	0.2	0.67	0.2	2.00	0.1	0.2
55-59	1	2	0.1	1.00	0.2	2.00	0.0	0.1
60-64	4	4	0.4	0.57	0.4	0.80	0.1	0.2
65-69	6	3	0.6	1.20	0.3	0.75	0.1	0.1
70-74	9	9	1.0	1.50	0.9	0.69	0.2	0.2
75-79	9	3	1.6	0.90	0.4	0.60	0.2	0.1
80-84	5	9	1.4	0.38	1.6	0.56	0.1	0.2
85+	8	24	3.5	0.44	4.2	0.60	0.2	0.4
All ages	46	58					0.1	0.2
Mortality								
Raw			0.3	0.71	0.3	0.67		
WS			0.1	0.74	0.1	0.77		
ES			0.2	0.70	0.2	0.74		
BRD-S			0.2	0.68	0.2	0.70		
PYLL-70								
per 100,000			0.9		0.9			
ES			0.7		0.8			
AYLL-70			9.5		11.0			

* See corresponding tables with multiple primaries.

ICD-10 C26: Malignant neoplasm of other and ill-defined digestive organs

Age distribution and age-specific mortality 2007 - 2014 (Males: 66, Females: 78)





Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at other digestive organ cancer-related death (see Table 10) should be considered.

Average mortality (world standard population) 2007 - 2014: Males



Average mortality (world standard population) 2007 - 2014: Females





Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2007 to 2014. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.2/100,000 WS N=66, females 0.1/100,000 WS N=78).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 65,347 female residents (averaged) in the period from 2007 to 2014 a total of 0 women died from other digestive organ cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.0/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.0/100,000.

Standardized mortality ratio (SMR) 2007 - 2014: Males



Standardized mortality ratio (SMR) 2007 - 2014: Females




Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=66, females N=78).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,924 female residents (averaged) in the period from 2007 to 2014 a total of 0 women died from other digestive organ cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.00. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 2.67, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases, where applicable. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

FRG	Federal Republic of Germany
GEKID	Association of Population-based Cancer Registries in Germany (Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)
MCR	Munich Cancer Registry (Tumorregister München)
SEER	Surveillance, Epidemiology, and End Results (USA)
AYLL-70	Average years of life lost prior to age 70 given a person dies before that age
BRD-S	German standard population
DCO	Death certificate only
EAR	Excess absolute risk = excess cancer cases (O - E) per 10,000 person-years
ES	European standard population (old)
LCL	Lower confidence limit
MI-index	Ratio between mortality and incidence
PYLL-70	Potential years of life lost prior to age 70 given a person dies before that age
SIR	Standardized incidence ratio
SMR	Standardized mortality ratio
UCL	Upper confidence limit
WS	World standard population

Recommended Citation

Munich Cancer Registry. ICD-10 C26: Other digestive organ cancer - Incidence and Mortality [Internet]. 2016 [updated 2016 Apr 13; cited 2016 Jun 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/base/bC26__E-ICD-10-C26-Other-digestive-organ-cancer-incidence-and-mortality.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

Fig./Tbl.		Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	3
1a	Gender distribution by year of diagnosis	4
2	Incidence by year of diagnosis	5
3	Age distribution parameters by year of diagnosis	6
4	Age distribution by 5-year age group and gender	8
5	Age-specific incidence, DCO rate, proportion malignancies	9
6	Age distribution and age-specific incidence (chart)	10
6a	Age-specific incidence internationally (chart)	11
7	Cumulative follow-up years (chart)	12
8	Standardized incidence ratio of second primaries	12
9a	Map of cancer incidence (WS) by county (chart)	14
9b	Standardized incidence ratio (SIR) by county (chart)	15
10a	Pts incident cohorts and mortality / yr	16
10b	Incidence and mortality by year of diagnosis	17
10c	Cancer-related deaths, death certification available / yr	18
11	Medians of age at death / yr	19
12	Mortality by year of death	21
13	Distribution of age at death	22
14	Age-specific mortality	23
15	Multiple primaries in deaths	24
16	Age-specific mortality (first primaries)	26
17	Age-specific mortality (single primaries)	27
18	Age distribution and age-specific mortality (chart)	28
19a	Map of cancer mortality (WS) by county (chart)	29
19b	Standardized mortality ratio (SMR) by county (chart)	30