Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ Deutsch

ICD-10 C40, C41: Bone cancer

Incidence and Mortality

Year of diagnosis	1998-2014
Patients	520
Diseases	522
Creation date	04/13/2016
Export date	12/23/2015
Population	4.64 m



Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

http://www.tumorregister-muenchen.de/en/facts/base/bC4041E-ICD-10-C40-C41-Bone-cancer-incidence-and-mortality.pdf

Global Statements about the statistics on the Internet – Baseline Statistics (grey button ——), Survival (red button ——)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.64 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases**** are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, April 2016

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007).
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.

ICD-10 codes (ICD-10 2015) used for specifying cancer site

Code	Description
C40 C40.0 C40.1 C40.2 C40.3 C40.8 C40.9	Malignant neoplasm of bone and articular cartilage of limbs Scapula and long bones of upper limb Short bones of upper limb Long bones of lower limb Short bones of lower limb Overlapping lesion of bone and articular cartilage of limbs Bone and articular cartilage of limb, unspecified
C41 C41.0 C41.1 C41.2 C41.3 C41.4 C41.8 C41.9	Malignant neoplasm of bone and articular cartilage of other and unspecified sites Bones of skull and face Mandible Vertebral column Ribs, sternum and clavicle Pelvic bones, sacrum and coccyx Overlapping lesion of bone and articular cartilage Bone and articular cartilage, unspecified

DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate.

INCIDENCE

Table 1

All patients with invasive cancer by year of diagnosis, proportions of DCO, multiple primaries, deaths, and active follow-up (incl. DCO)

				Prop.		Prop.
		/DCO	Prop.	mult.	Prop.	actively
Year of	Cases	cases	DCO	primaries	deaths	followed
diagnosis	n	n	%	્	용	ે
1998	21			28.6	33.3	85.7
1999	17	1	5.9	29.4	47.1	100.0
2000	16	1	6.3	18.8	31.3	93.8
2001	14	1	7.1	7.1	42.9	92.9
2002	30	2	6.7	30.0	50.0	93.3 #
2003	27	3	11.1	7.4	48.1	92.6
2004	37	8	21.6	16.2	59.5	97.3
2005	26	1	3.8	19.2	50.0	76.9
2006	24	1	4.2	20.8	33.3	75.0
2007	52	1	1.9	21.2	30.8	67.3 #
2008	36	3	8.3	11.1	55.6	66.7
2009	39	2	5.1	12.8	35.9	61.5
2010	44	2	4.5	20.5	43.2	65.9
2011	45	4	8.9	28.9	44.4	66.7
2012	48	2	4.2	14.6	25.0	70.8
2013	37			8.1	16.2	100.0
2014	9	4	44.4	44.4	55.6	100.0 ##
1998-2014	522	36	6.9	18.8	40.0	78.9

[#] The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

^{##} Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be found in the respective headings.

Table 1a

All patients with invasive cancer by year of diagnosis and gender (incl. DCO)

Year of	All/	Males	Females	Prop. males
diagnosis	n/	n	n	ଚ
1998	21	10	11	47.6
1999	/17	/ 7	10	41.2
2000	/ 16	5	11/	31.3
2001	14	8	6	57.1
2002	30	17	13	56.7
2003	27	14	13	51.9
2004	37	21	16	56.8
2005	26	16	10	61.5
2006	24	12	12	50.0
2007	52	33	19	63.5
2008	36	20	16	55.6
2009	39	26	13	66.7
2010	44	21	23	47.7
2011	45	35	10	77.8
2012	48	31	17	64.6
2013	37	17	20	45.9
2014	9	4	5	44.4
1998-2014	522	297	225	56.9

Table 2

Incidence measures by year of diagnosis including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

			Males	Fem.	Males	Fem.	Males	Fem.	Males	Fem.
Year of	Males	Females	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	10	11	0.9	0.9	0.9	0.7	0.9	0.8	1.0	0.9
1999	7	10	0.6	0.8	0.6	0.8	0.6	0.8	0.6	0.8
2000	5	11 /	0.4	0.9	0.3	0.9	0.4	0.9	0.4	0.9
2001	8	6 <	0.7	0.5	0.9	0.6	0.8	0.5	0.8	0.6
2002	17	13	0.9	0.7	0.7	0.7	0.8	0.7	1.0	0.7
2003	14	13	0.7	0.7	0.7	0.5	0.7	0.5	0.8	0.6
2004	21	16	1.1	0.8	0.9	0.7	1.0	0.7	1.2	0.9
2005	16	10	0.8	0.5	1.0	0.4	0.9	0.4	0.9	0.5
2006	12	12	0.6	0.6	0.6	0.5	0.6	0.5	0.6	0.5
2007	33	19	1.5	0.8	1.5	0.9	1.5	0.9	1.6	0.8
2008	20	16	0.9	0.7	1.0	0.6	0.9	0.6	1.0	0.6
2009	26	13	1.2	0.6	1.1	0.6	1.1	0.6	1.2	0.6
2010	21/	23	0.9	1.0	1.0	1.0	1.0	1.0	1.0	1.0
2011	35	10	1.5	0.4	1.2	0.2	1.4	0.3	1.4	0.4
2012	31	17	1.4	0.7	1.3	0.6	1.3	0.6	1.4	0.7
2013	17	20	0.7	0.8	0.6	1.0	0.7	0.9	0.7	0.9
2014	4	\5	0.2	0.2	0.1	0.1	0.1	0.1	0.2	0.2
1998-2014	297	225	0.9	0.7	0.9	0.6	0.9	0.6	1.0	0.7




Table 3

Age distribution parameters by year of diagnosis (All patients) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	21	47.2	20.1	14.1	77.1	18.6	29.6	51.8	62.4	71.8
1999	17	46.4	26.3	5.5	87.5	8.2	29.1	47.5	58.7	87.1
2000	16	43.8	20.1	7.9	87.0	14.8	31.7	45.0	54.2	75.2
2001	14	37.2	27.3	6.5	78.6	11.4	17.9	24.1	73.0	75.4
2002	30	46.6	23.9	12.6	87.2	16,3	23.3	42.7	65.6	78.2
2003	27	49.7	24.9	10.0	89.2	13.0	27.7	50.9	69.4	86.0
2004	37	52.1	24.2	12.5	92.9	17.8	28.3	56.9	73.7	82.4
2005	26	43.0	26.5	8.7	91.8	12.2	16.6	42.2	63.6	76.3
2006	24	43.9	22.4	9.5	80.5	13.2	30.0	38.9	64.0	74.7
2007	52	40.3	24.2	6.7	85.6	12.3	16.7	37.6	63.2	70.6
2008	36	44.8	28.0	8.3	88.9	12.2	17.7	41.8	71.2	84.2
2009	39	41.6	21.4	11.4	81.0	15.2	21.2	42.4	57.1	76.5
2010	44	43.8	25.4	5.6	78.4	11.6	17.7	49.5	67.3	75.2
2011	45	56.6	21.0	5.2	91.8	16.0	48.7	58.4	70.8	82.0
2012	48	49.1	24.6	7.4	87.1	14.7	22.5	52.7	70.6	80.0
2013	37	42.1	24.7	3.6	86.7	11.4	23.0	40.8	56.0	79.0
2014	9	68.4	22.0	26.8	100	26.8	57.3	75.1	84.0	100
1998-2014	522	46.3	24.4	3.6	100	13.3	22.8	47.7	67.3	78.6

Table 3a

Age distribution parameters by year of diagnosis (MALES)

(incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	10	42.7	20.9	17.0	73.9	17.8	22.8	41.4	59.1	70.8
1999	7	40.0	18.5	5.5	58.7	5.5	29.1	47.5	52.8	58.7
2000	5	45.0	12.4	31.7	56.1	31.7	31.8	50.1	55.4	56.1
2001	8	37.6	25.5	11.4	74.0	11.4	18.7	26.2	62.6	74.0
2002	17	49.7	21.0	18.9	87.2	21.4	32.5	45.9	64.2	75.3
2003	14	42.7	21.9	10.0	76.4	11.2	25.2	44.4	58.4	70.2
2004	21	52.4	24.4	12.5	91.9	15.7	33.3	60.9	67.5	79.5
2005	16	37.1	25.7	8.7	91.8	9.9	13.4	28.6	56.8	68.7
2006	12	40.1	20.6	13.2	73.3	14.0	26.7	35.2	58.0	70.7
2007	33	40.4	23.0	9.8	81.9	13.4	20.4	36.2	63.4	69.7
2008	20	39.0	24.5	11.9	88.9	14.9	18.4	28.7	60.5	73.1
2009	26	43.2	19.9	13.3	78.2	16.2	26.2	44.1	56.4	72.6
2010	21	41.1	26.3	7.5	78.4	10.7	14.7	33.1	67.2	75.2
2011	35	52.2	21.0	5.2	91.8	15.3	43.0	53.4	68.7	72.6
2012	31	47.5	27.0	7.4	87.1	13.3	16.6	52.1	73.4	80.0
2013	17	46.3	22.1	11.4	86.7	13.1	32.7	45.2	56.0	80.0
2014	4	66.9	15.2	51.3	84.0	51.3	54.3	66.2	79.6	84.0
1998-2014	297	44.8	23.1	5.2	91.9	13.3	22.8	47.1	64.0	75.1

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	11	51.3	19.5	14.1	77.1	26.8	32.3	56.3	65.1	71.8
1999	10	50.9	30.8	8.2	87.5	11.9	19.4	51.0	83.2	87.3
2000	11	43.2	23.4	7.9	87.0	14.8	27.5	44.9	52.9	75.2
2001	6	36.8	32.0	6.5	78.6	6.5	11.7	24.1	75.4	78.6
2002	13	42.6	27.6	12.6	85.8	14,4	16.5	33.8	67.3	81.1
2003	13	57.1	26.6	13.0	89.2	21.8	39.4	67.1	83.0	86.9
2004	16	51.7	24.7	17.8	92.9	19.9	26.9	55.9	73.9	84.0
2005	10	52.2	26.4	12.2	88.2	14.4	35.3	54.3	74.1	82.3
2006	12	47.7	24.3	9.5	80.5	12.6	33.4	42.0	69.4	78.6
2007	19	40.2	26.9	6.7	85.6	10.7	14.6	39.1	62.9	83.1
2008	16	51.9	31.1	8.3	88.2	9.1	15.4	66.4	77.6	84.3
2009	13	38.3	24.6	11.4	81.0	12.7	15.3	29.4	57.1	77.0
2010	23	46.3	24.9	5.6	78.2	13.0	20.4	59.2	67.5	73.7
2011	10 /	71.7	12.3	51.5	84.4	53.4	59.1	75.0	82.3	84.2
2012	17/	51.9	20.2	16.3	80.7	22.0	44.0	57.2	62.8	77.6
2013	20	38.5	26.8	3.6	83.5	9.4	14.6	33.5	61.2	78.8
2014	5	69.6	28.0	26.8	100	26.8	60.0	76.0	85.4	100
1998-2014	225	48.3	26.0	3.6	100	13.0	23.3	50.6	72.5	82.3

Table 4

Age distribution by 5-year age group and gender for period 2007-2014 (incl. DCO)

Age at								
diagnosis	Cases		Males			Females		
Years	n	% Cum.%	n	용	Cum.%	n	%	Cum.%
0 - 4	1	0.3 /0.3			0.0	1	0.8	0.8
5-9	11	3.5 3.9	6	3.2	3.2	5	4.1	4.9
10-14	31	10.0 / 13.9/	18	9.6	12.8	13	10.6	15.4
15-19	28	9.0 / 22.9	17	9.1	21.9	11	8.9	24.4
20-24	19	6.1 29.0	13	7.0	28.9	6	4.9	29.3
25-29	16	5.2 34.2	8	4.3	33.2	8	6.5	35.8
30-34	9	2.9 37.1	7	3.7	36.9	2	1.6	37.4
35-39	8	2.6 39.7	6	3.2	40.1	2	1.6	39.0
40 - 44	19	6.1 45.8	13	7.0	47.1	6	4.9	43.9
45-49	17	5.5 51.3	12	6.4	53.5	5	4.1	48.0
50-54	21	6.8 58.1	17	9.1	62.6	4	3.3	51.2
55-59	23	7.4 65.5	10	5.3	67.9	13	10.6	61.8
60-64	19	6.1 71.6	13	7.0	74.9	6	4.9	66.7
65-69	21 /	6.8 78.4	13	7.0	81.8	8	6.5	73.2
70-74	21	6.8 85.2	13	7.0	88.8	8	6.5	79.7
75-79	22	7.1 92.3	11	5.9	94.7	11	8.9	88.6
80-84	14	4.5 96.8	4	2.1	96.8	10	8.1	96.7
85+	10	3.2 100.0	6	3.2	100.0	4	3.3	100.0
All ages	310	100.0	187	100.0		123	100.0	

Included in the statistics are 23.0% multiple primaries in males and 25.2% in females.



							Males	Females
			Males	Females	Males	Females	Prop.all	Prop.all
Age at			Age-	Age-	DCO rate	DCO rate	cancers	cancers
diagnosis	Males	Females	spec.	spec.	n=7	n=11	n=91183	n=89596
Years	n	n	incid.	/incid.	용	90	ଚ	양
0- 4		1	0.0	0.1				0.7
5- 9	6	5	0.7	0.6			6.3	6.4
10-14	18	13	1.9	1.5			18.0	14.6
15-19	17	11	1.8	1.2			7.9	6.7
20-24	13	6	1.2	0.6			3.5	1.9
25-29	8	8	0.7	0.6			1.4	1.2
30-34	7	2	0.6	0.2			0.9	0.2
35-39	6	2	0.5	0.2			0.5	0.1
40 - 44	13	6	0.8	0.4			0.7	0.2
45-49	12	5	0.8	0.3			0.4	0.1
50-54	17	4	1.3	0.3	11.8		0.3	0.1
55-59	10	13 /	0.9	1.2		7.7	0.1	0.2
60-64	13	6	1.3	0.6	7.7		0.1	0.1
65-69	13	8	1.4	0.8		12.5	0.1	0.1
70 - 74	13	8	1.4	0.8		25.0	0.1	0.1
75-79	11	\ 11\	2.0	1.5	9.1	9.1	0.1	0.1
80-84	4	10	1.1	1.8	25.0	30.0	0.0	0.1
85+	6	4	2.6	0.7	33.3	75.0	0.1	0.0
All ages	187	123			3.7	8.9	0.2	0.1
Incidence								
Raw			1.0	0.7				
WS			1.0	0.6				
ES			1.0	0.6				
BRD-S			1.1	0.7				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).



ICD-10 C40, C41: Malignant neoplasm of bone and articular cartilage




Figure 6. Age distribution and age-specific incidence



ICD-10 C40, C41: Malignant neoplasm of bone and articular cartilage




Figure 6a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).



Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2014, based on the November 2013 submission. http://www.seer.cancer.gov.




Figure 7. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.

		Table 8	a				
Standardized i	ncidence r	catio (SIR,	with 95	% conf	idence	limits),	
excess absol							
	for	period 19	98-2014				
		MALES					
	Observed	Expected		LCL	UCL		DCO
Diagnosis	n	n	SIR	95%	95%	EAR	용
C33-C34 Lung	5	0.7	7.4	2.4	17.3 #	53.5	40.0
C91-C96 Leukaemia	3	0.1	32.8	6.8	95.7 #	35.9	
Other primaries	10	2.7	3.7	1.8	6.7 #	89.8	
Not observed	0	2.3	0.0	0.0	1.6	-28.8	
	1.0	F 0	2 1	1 0	4 0 "	150 5	
All mult. primaries	18	5.8	3.1	1.8	4.9 #	150.5	11.1
Patients			282				
Median age at second m	alignancy	(years)	67.9				
Person-years			809				
Mean observation time	(years)		2.9				
Median observation tim	e (years)		1.5				

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries"

Table 8b

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2014


FEMALES

	Observed Ex	pected		LCL	UCL		DCO
Diagnosis	/ n /	n	SIR	95%	95%	EAR	왕
C33-C34 Lung	2	0.3	7.8	0.9 2	28.1	23.4	
C43 Malign. melanor	2 na 3	0.2	17.4	3.6	50.8 #	38.0	
C50 Breast	3	1.2	2.5	0.5	7.3	24.1	
Other primaries	9	0.7	/12.7/	5.8 2	24.2 #	111.4	11.1
Not observed	0	1.6	0.0	0.0	2.4	-21.0	
All mult. primaries	17	3.9	4.3	2.5	7.0 #	175.9	5.9
Patients		20	6				
Median age at second malig	gnancy (years)	64.	0				
Person-years		74	4				
Mean observation time (year	ars)	3.	6				
Median observation time (years)	1.	5				

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries"

Average incidence (world standard population) 2007 - 2014: Males



Average incidence (world standard population) 2007 - 2014: Females





Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 1.0/100,000 WS N=187, females 0.6/100,000 WS N=123).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 65,347 female residents (averaged) in the period from 2007 to 2014 a total of 2 women were identified with newly diagnosed bone cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.4/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 2.8/100,000.

Standardized incidence ratio (SIR) 2007 - 2014: Males



Standardized incidence ratio (SIR) 2007 - 2014: Females




Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=187, females N=123).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,924 female residents (averaged) in the period from 2007 to 2014 a total of 2 women were identified with newly diagnosed bone cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.58. Though, the value of this parameter may vary with an underlying probability of 99% between 0.03 and 2.71, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

						Prop.
		Prop.				deaths
	Incident	actively	Prop.		Prop.	with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	용	용	n	%	용
1998	21	85.7		7	33.3	71.4
1999	17	100.0	5.9	8	47.1	75.0
2000	16	93.8	6.3	5	31.3	100.0
2001	1.4	92.9	7.1	6	42.9	100.0
2002	30	93.3	6.7	15	50.0	86.7
2003	27	92.6	11.1	13	48.1	100.0
2004	37	97.3	21.6	22	59.5	95.5
2005	26	76.9	3.8	13	50.0	100.0
2006	24	75.0	4.2	8	33.3	100.0
2007	52	67.3	1.9	16	30.8	100.0
2008	36	66.7	8.3	20	55.6	95.0
2009	39	61.5	5.1	14	35.9	100.0
2010	44	65.9	4.5	19	43.2	100.0
2011	45	66.7	8.9	20	44.4	100.0
2012	48	70.8	4.2	12	25.0	100.0
2013	37	100.0		6	16.2	83.3
2014	9	100.0	44.4	5	55.6	100.0
1998-2014	522	78.9	6.9	209	40.0	95.7

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

			D		
			Prop.		Drop
V	T		deaths	Darkha da	Prop.
Year of	Incident	/ _ /,	with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	'n	96	n	%
1998	21	11	90.9	4	19.0
1999	17	6	66.7	2	11.8
2000	16	7	85.7	1	6.3
2001	14	10	100.0	2	14.3
2002	30	13	92.3	2	6.7
2003	27	18	100.0	4	14.8
2004	37	15	86.7	10	27.0
2005	26	14	92.9	2	7.7
2006	24	14	92.9	1	4.2
2007	52	16	100.0	5	9.6
2008	36	29	96.6	11	30.6
2009	39	13	100.0	3	7.7
2010	44	22	100.0	7	15.9
2011	45	18	100.0	5	11.1
2012	48	23	100.0	7	14.6
2013	37	25	100.0	1	2.7
2014	9	20	95.0	5	55.6
1998-2014	522	274	96.0	72	13.8

Table 10c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

				Prop.
				cancer
		Prop.	Prop.	recorded
		cancer-	non-cancer-	on death
Year of	Deaths	related	related	certificate
death	n	%	/ %	%
1998	11	81.8	18.2	80.0
1999	6	83.3	16.7	100.0
2000	7	85.7	14.3	100.0
2001	10	70.0	30.0	80.0
2002	13	84.6	15.4	91.7
2003	18	77.8	22.2	88.9
2004	15	80.0	20.0	92.3
2005	14	85.7	14.3	92.3
2006	14	92.9	7.1	100.0
2007	16	81.3	18.8	81.3
2008	29	93.1	6.9	92.9
2009	\ 13	92.3	7.7	92.3
2010	22	95.5	4.5	95.5
2011	18	77.8	22.2	77.8
2012	23	100.0		100.0
2013	25	92.0	8.0	96.0
2014	20	90.0	10.0	94.7
1998-2014	274	87.6	12.4	91.6

 $$\operatorname{\textsc{Table 11a}}$$ Medians of age at death according to the grouping in Table 10 $$\operatorname{\textsc{MALES}}$$

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(non-cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	7	55.5	50.7	59.3	50.7
1999	2	44.4	44.4		44.4
2000	4	49.3	30.2	68.4	30.2
2001	7	56.8	65.4	49.1	65.4
2002	7	50.6	49.9	60.1	50.6
2003	9	35.8	35.8	59.1	35.8
2004	8	67.7	67.7		73.8
2005	9	41.2	36.6	60.7	36.6
2006	7/	58.5	58.5		63.7
2007	1,1	58.6	58.6	49.9	58.6
2008	15	53.7	49.2	86.5	53.7
2009	9	65.2	65.2		65.2
2010	10	60.8	60.8		60.8
2011	9	67.2	68.1	56.9	68.1
2012	18	68.1	68.1		68.1
2013	14	61.0	61.0		61.0
2014	5	45.6	45.6		45.6
1998-2014	151	59.1	58.5	60.1	59.6

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

 $\begin{array}{c} \text{Table 11b} \\ \text{Medians of age at death according to the grouping in Table 10} \\ \text{FEMALES} \end{array}$

					Age at
		Age at	Age at	Age at	death
		death	death	death	(according
		(all	(cancer-	(non-cancer-	to death
Year of	Deaths	causes)	related)	related)	certificate)
death	n	Years	Years	Years	Years
1998	4	61.4	57.2	65.7	57.8
1999	4	68.8	66.1	88.2	65.7
2000	3	75.2	75.2		75.2
2001	3	78.6	89.9	63.7	84.2
2002	6	75.1	70.7	82.9	75.1
2003	9	69.4	68.5	85.0	69.4
2004	7	84.0	71.9	88.7	74.7
2005	5	37.1	35.1	88.3	35.1
2006	7/	70.0	60.6	84.1	70.0
2007	5	74.7	60.8	94.3	60.8
2008	14	71.4	69.2	89.0	69.2
2009	4	65.2	50.6	96.6	50.6
2010	12	67.5	67.1	93.3	67.1
2011	9	70.6	63.4	94.3	63.4
2012	\5	57.9	57.9		57.9
2013	11	75.9	56.0	80.6	65.9
2014	15	71.4	71.4	76.7	76.4
1998-2014	123	70.6	67.0	86.9	69.4

By 2010, life expectancy at birth was 77.5 years for boys and 82.6 years for girls.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	6	0.5	0.60	0.4	0.42	0.5	0.51	0.5	0.52
1999	2	0.2	0.29	0.1	0.21	0.2	0.26	0.2	0.26
2000	3	0.3	0.60	0.2	0.75	0.2	0.62	0.3	0.73
2001	6	0.5	0.75	0.4	0.48	0.5	0.65	0.6	0.72
2002	6	0.3	0.35	0.3	0.34	0.3	0.36	0.3	0.33
2003	7	0.4	0.50	0.3	0.46	0.4	0.48	0.4	0.59
2004	8	0.4	0.38	0.4	0.43	0.4	0.42	0.5	0.42
2005	8	0.4	0.50	0.4	0.38	0.4	0.44	0.5	0.53
2006	7	0.4	0.58	0.2	0.37	0.3	0.49	0.4	0.61
2007	9	0.4	0.27	0.4	0.24	0.4	0.27	0.4	0.28
2008	14	0.6	0.70	0.5	0.55	0.6	0.60	0.6	0.64
2009	9	0.4	0.35	0.3	0.26	0.3	0.30	0.4	0.32
2010	10	0.4	0.48	0.3	0.31	0.4	0.39	0.4	0.46
2011	8	0.4	0.23	0.2	0.18	0.3	0.20	0.4	0.25
2012	18	0.8	0.58	0.6	0.43	0.7	0.50	0.8	0.58
2013	14	0.6	0.82	0.5	0.76	0.5	0.79	0.6	0.82
2014	5	0.2	1.25	0.2	1.98	0.2	1.45	0.2	1.23
1998-2014	140	0.4	0.47	0.3	0.38	0.4	0.43	0.5	0.47

Table 12b

Mortality measures (cancer-related death) and mortality-incidence-index by year of death FEMALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	3	0.3	0.27	0.1	0.17	0.2	0.21	0.2	0.24
1999	3	0.3	0.30	0.1	0.18	0.2	0.26	0.2	0.27
2000	3	0.2	0.27	0.1	0.11	0.2	0.17	0.2	0.26
2001	1	0.1	0.17	0.0	0.03	0.0	0.06	0.0	0.07
2002	5	0.3	0.38	0.2	0.22	0.2	0.29	0.3	0.36
2003	7	0.4	0.54	0.2	0.38	0.2	0.47	0.3	0.53
2004	4	0.2	0.25	0.1	0.14	0.1	0.19	0.2	0.19
2005	4	0.2	0.40	0.2	0.44	0.2	0.44	0.2	0.43
2006	6	0.3	0.50	0.2	0.36	0.2	0.45	0.3	0.48
2007	4	0.2	0.21	0.1	0.13	0.1	0.15	0.2	0.22
2008	13	0.6	0.81	0.3	0.52	0.4	0.67	0.5	0.80
2009	3	0.1	0.23	0.1	0.13	0.1	0.19	0.1	0.21
2010	11	0.5	0.48	0.3	0.32	0.4	0.37	0.4	0.41
2011	6	0.3	0.60	0.2	1.12	0.2	0.82	0.3	0.71
2012	5	0.2	0.29	0.1	0.25	0.2	0.26	0.2	0.26
2013	9	0.4	0.45	0.2	0.23	0.3	0.31	0.3	0.38
2014	13	0.6	2.60	0.5	4.46	0.5	3.34	0.5	3.02
1998-2014	100	0.3	0.44	0.2	0.30	0.2	0.36	0.3	0.41

Table 13

Age distribution of age at death (cancer-related) for period 2007-2014

(incl. multiple primaries)

Age at	~					- 1		
death	Cases	. / .	Males		\ \	Females		
Years	n	% Cum.%	n	%	Cum.%	n	૾ૢ	Cum.%
0 - 4	1	0.7 0.7			0.0	1	1.6	1.6/
5-9	2	1.3 / 2.0 /	1	1.1	/ 1.1	1	1.6	3.1
10-14	3	2.0 4.0	2	2.3	3.4	1	1.6	4.7
15-19	7	4.6 8.6	5	5.7	9.2/	2	3.1	7.8
20-24	11	7.3 15.9	7	8.0	17.2	4	6.3	14.1
25-29	6	4.0 19.9	3	3.4	20.7	3	4.7	18.8
30-34	4	2.6 22.5	2	2.3	23.0	2	3.1	21.9
35-39	4	2.6 25.2	1	1.1	24.1	3	4.7	26.6
40 - 44	6	4.0 29.1	5	5.7	29.9	1	1.6	28.1
45-49	7	4.6 33.8	5	5.7	35.6	2	3.1	31.3
50-54	7	4.6 38.4	3	3.4	39.1	4	6.3	37.5
55-59	9	6.0 44.4	6	6.9	46.0	3	4.7	42.2
60-64	13	8.6 53.0	9	10.3	56.3	4	6.3	48.4
65-69	18	11.9 64.9	13	14.9	71.3	5	7.8	56.3
70-74	16	10.6 75.5	10	11.5	82.8	6	9.4	65.6
75-79	15	9.9 85.4	7	8.0	90.8	8	12.5	78.1
80-84	12	7.9 93.4	4	4.6	95.4	8	12.5	90.6
85+	10	6.6 100.0	4	4.6	100.0	6	9.4	100.0
All ages	151	100.0	87	100.0		64	100.0	

Included in the statistics are 23.0% multiple primaries in males and 25.2% in females.



Table 14

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2014 (incl. multiple primaries)

		/.		\			_
_		Males		Females		Males	Females
Age at		Age-		Age-		_	Prop.all
death	Males Femal	/ = /		spec.		cancers	cancers
Years	n n	mortal.	MI-index	mortal.	MI-index	96	olo
0- 4		L 0.0		0.1	1.00		6.7
5- 9	1	1 / 0.1	0.17	0.1	0.20	4.8	5.6
10-14	2	0.2	0.11	0.1/	0.08	11.1	5.0
15-19	5 2	0.5	0.29	0.2	0.18	13.9	9.1
20-24	7	0.6	0.54	0.4	0.67	14.6	14.3
25-29	3	0.2	0.38	0.2	0.38	4.8	4.7
30-34	2 2	0.2	0.29	0.2	1.00	2.3	1.8
35-39	1 :	0.1	0.17	0.2	1.50	0.6	1.2
40 - 44	5	0.3	0.38	0.1	0.17	1.1	0.2
45-49	5 / 2	0.3	0.42	0.1	0.40	0.5	0.2
50-54		0.2		0.3	1.00	0.2	0.2
55-59		0.6		0.3	0.23	0.2	0.1
60-64		0.9		0.4	0.67	0.2	0.1
65-69		5 1.4		0.5	0.63	0.2	0.1
70-74		5 1.1		0.6	0.75	0.1	0.1
75-79		3 1.3		1.1	0.73	0.1	0.1
80-84		1.1		1.4	0.80	0.1	0.1
85+		5 1.7		1.0	1.50	0.1	0.1
001	•	1.7	0.07	1.0	1.30	0.1	0.1
All ages	87 64	1				0.2	0.1
TILL ages	0,					/ 0.2	· · ·
Mortality /							
Raw		0.5	0.47	0.3	0.52		
WS		0.4	0.37	0.2	0.37		
ES		0.4		0.3	0.43		
BRD-S		0.5		0.3	0.48		
DIAD 5		0.3	0.40	0.5	0.40		
PYLL-70							
per 100,000		9.3		6.3			
ES ES		9.6		6.9			
AYLL-70		24.0		27.9			
171111 / /		24.0		27.3			

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15a $\begin{tabular}{ll} Multiple primaries in deaths in period 1998-2014 \\ \hline MALES \end{tabular}$

					Syn-	Syn-		
	/ _	_/	_		chron	chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n	%↓	n	← %	n	olo →	n	← %
C09-C10 Oropharynx	/ 1 /	2.0					1	100.0
C18 Colon	2 /	3.9	2	100.0				
C19-C20 Rectum	4	7.8	3	75.0			1	25.0
C22 Liver	_ 1	2.0					1	100.0
C33-C34 Lung	9	17.6	4	44.4	1	11.1	4	44.4
C44 Skin others	1	2.0	1	100.0				
C46,C49 Soft tissue	5	9.8	3	60.0			2	40.0
C61 Prostate	9	17.6	7	77.8	1	11.1	1	11.1
C62 Testis	2	3.9	1	50.0	1	50.0		
C64 Kidney	2	3.9	1	50.0			1,	50.0
C67 Bladder	1	2.0	1	100.0				
C70-C72 CNS cancer	3	5.9	2	66.7	_ 1	33.3		
C76-C79 CUP	1	2.0					1	100.0
C81 Hodgkin lymphoma	1	2.0	1	100.0				
C82-C85 NHL	3	5.9					3	100.0
C90 Mult. myeloma	5	9.8	4	80.0	1	20.0		
C91-C96 Leukaemia	1	2.0					1	100.0
All mult. primaries	51	100.0	30	58.8	5	9.8	16	31.4




Table 15b

Multiple primaries in deaths in period 1998-2014
FEMALES

					Syn-	Syn-		
					chron	chron		
	Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis	n	~ 응 ↓	n	← %	n	← %	n	← %
C16 Stomach	/ 3	6.5					3	100.0
C18 Colon	/ 1	2.2	1	100.0				
C19-C20 Rectum	/ 3	6.5	2	66.7			1	33.3
C22 Liver	1	2.2	1	100.0				
C25 Pancreas	2	4.3					2	100.0
C26 GI cancer	1	2.2					1	100.0
C33-C34 Lung	3	6.5	1	33.3			2	66.7
C40-C41 Bone	1	2.2					1	100.0
C43 Malign. me	lanoma 3	6.5	2	66.7			1	33.3
C46,C49 Soft tissue	e 2	4.3	1	50.0			1	50.0
C48 Peritoneal	1	2.2					1	100.0
C50 Breast	10	21.7	4	40.0	_ 1	10.0	5	50.0
C53 Cervix ute	ri / 4	8.7	3	75.0			1	25.0
C54 Corpus ute:	ri / 2	4.3	1	50.0			1	50.0
C56 Ovary	3	6.5	3	100.0				
C64 Kidney	1	2.2					1	100.0
C67 Bladder	1	2.2					1	100.0
C70-C72 CNS cancer	\ 1	2.2	1	100.0				
C81 Hodgkin ly	mphoma 1	2.2	1	100.0				
C91-C96 Leukaemia	1	2.2					1	100.0
C96 Systemic	1	2.2					1	100.0
All mult. primarie	s 46	100.0	21	45.7	1	2.2	24	52.2

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2014

(First primaries only *)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males	Females	spec.		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	용	용
0 - 4		1 /	0.0		0.1	1.00		7.7
5- 9	1	1 /	0.1	0.17	0.1	0.20	5.0	5.6
10-14	2	1 <	0.2	0.11	0.1	0.08	11.1	5.3
15-19	4	2	0.4	0.25	0.2	0.18	12.1	10.0
20-24	6	4	0.5	0.50	0.4	1.00	14.0	15.4
25-29	2	3	0.2	0.25	0.2	0.38	3.6	5.1
30-34	2	2	0.2	0.29	0.2	1.00	2.3	2.1
35-39	1	2	0.1	0.17	0.2	1.00	0.6	0.9
40 - 44	5		0.3	0.42	0.0		1.2	
45-49	5	2	0.3	0.45	0.1	0.40	0.5	0.2
50-54	2	4	0.2	0.13	0.3	1.00	0.1	0.3
55-59	6	3	0.6	0.67	0.3	0.33	0.2	0.1
60-64	7	2	0.7	0.88	0.2	0.50	0.2	0.1
65-69	7	4	0.7	0.88	0.4	0.67	0.1	0.1
70-74	6	4	0.7	0.67	0.4	0.80	0.1	0.1
75-79	5	\5	0.9	0.63	0.7	0.71	0.1	0.1
80-84	2	7	0.6	1.00	1.2	0.88	0.0	0.1
85+	4	5	1.7	0.80	0.9	1.25	0.1	0.1
All ages	67	52					0.2	0.2
Mortality								
Raw			0.4	0.42	0.3	0.50		
WS			0.3	0.33	0.2	0.36		
ES			0.3	0.38	0.2	0.41		
BRD-S			0.4	0.41	0.3	0.47		
PYLL-70								
per 100,000			8.1		5.9			
ES			8.4		6.5			
AYLL-70			25.9		29.9			

^{*} See corresponding tables with multiple primaries.

Table 17

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2014

(Single primaries only *)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
death	Males	Females	spec.		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	용	%
0- 4		1 /	0.0		0.1	1.00		7.7
5- 9	1	1 /	0.1	0.17	0.1	0.20	5.0	5.6
10-14	2	1 <	0.2	0.12	0.1/	0.08	11.1	5.6
15-19	4	2	0.4	0.25	0.2	0.18	12.1	11.1
20-24	6	4	0.5	0.50	0.4	1.00	15.4	16.7
25-29	2	3	0.2	0.29	0.2	0.38	3.9	5.4
30-34	2	1	0.2	0.29	0.1	0.50	2.4	1.2
35-39	1	1	0.1	0.17	0.1	0.50	0.6	0.5
40 - 44	5		0.3	0.42	0.0		1.3	
45-49	5	2	0.3	0.45	0.1	0.40	0.6	0.2
50-54	2 /	4	0.2	0.14	0.3	1.33	0.1	0.3
55-59	4	2	0.4	0.44	0.2	0.22	0.2	0.1
60-64	5	/ 2	0.5	0.63	0.2	0.50	0.1	0.1
65-69	3	3	0.3	0.75	0.3	0.50	0.1	0.1
70-74	5	2	0.5	0.56	0.2	0.40	0.1	0.0
75-79	5	4	0.9	0.63	0.6	0.67	0.1	0.1
80-84	2	4	0.6	1.00	0.7	0.50	0.0	0.1
85+	4	5	1.7	0.80	0.9	1.25	0.1	0.1
All ages	58	42					0.2	0.1
Mortality /								
Raw			0.3	0.38	0.2	0.42		
WS			0.3	0.31	0.2	0.32		
ES			0.3	0.34	0.2	0.35		
BRD-S			0.3	0.38	0.2	0.39		
PYLL-70								
per 100,000			7.8		5.3			
ES			8.1		6.0			
AYLL-70			29.6		31.2			

^{*} See corresponding tables with multiple primaries.

ICD-10 C40, C41: Malignant neoplasm of bone and articular cartilage Age distribution and age-specific mortality 2007 - 2014 (Males: 87, Females: 64)





Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at bone cancer-related death (see Table 10) should be considered.



Average mortality (world standard population) 2007 - 2014: Males



Average mortality (world standard population) 2007 - 2014: Females





Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2007 to 2014. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.4/100,000 WS N=84, females 0.2/100,000 WS N=64).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 65,347 female residents (averaged) in the period from 2007 to 2014 a total of 1 women died from bone cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.4/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 2.8/100,000.

Standardized mortality ratio (SMR) 2007 - 2014: Males



Standardized mortality ratio (SMR) 2007 - 2014: Females




Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=84, females N=64).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,924 female residents (averaged) in the period from 2007 to 2014 a total of 1 women died from bone cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.59. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 4.37, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases, where applicable. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

FRG Federal Republic of Germany

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

MCR Munich Cancer Registry (Tumorregister München)
SEER Surveillance, Epidemiology, and End Results (USA)

AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

BRD-S German standard population

DCO Death certificate only EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

ES European standard population (old)

LCL Lower confidence limit

MI-index Ratio between mortality and incidence

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age

SIR Standardized incidence ratio
SMR Standardized mortality ratio
UCL Upper confidence limit
WS World standard population

Recommended Citation

Munich Cancer Registry. ICD-10 C40, C41: Bone cancer - Incidence and Mortality [Internet]. 2016 [updated 2016 Apr 13; cited 2016 Jun 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/base/bC4041E-ICD-10-C40-C41-Bone-cancer-incidence-and-mortality.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

Fig./Tb	l.	Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	3
1a	Gender distribution by year of diagnosis	4
2	Incidence by year of diagnosis	5
3	Age distribution parameters by year of diagnosis	6
4	Age distribution by 5-year age group and gender	8
5	Age-specific incidence, DCO rate, proportion malignancies	9
6	Age distribution and age-specific incidence (chart)	10
6a	Age-specific incidence internationally (chart)	11
7	Cumulative follow-up years (chart)	12
8	Standardized incidence ratio of second primaries	12
9a	Map of cancer incidence (WS) by county (chart)	14
9b	Standardized incidence ratio (SIR) by county (chart)	15
10a	Pts incident cohorts and mortality / yr	16
10b	Incidence and mortality by year of diagnosis	17
10c	Cancer-related deaths, death certification available / yr	18
11	Medians of age at death / yr	19
12	Mortality by year of death	21
13	Distribution of age at death	22
14	Age-specific mortality	23
15	Multiple primaries in deaths	24
16	Age-specific mortality (first primaries)	26
17	Age-specific mortality (single primaries)	27
18	Age distribution and age-specific mortality (chart)	28
19a	Map of cancer mortality (WS) by county (chart)	29
19b	Standardized mortality ratio (SMR) by county (chart)	30