Munich Cancer Registry



- Survival
- Selection Matrix
- Homepage
- Deutsch

ICD-10 C46: Kaposi sarcoma

Incidence and Mortality

Year of diagnosis	1998-2014
Patients	75
Diseases	75
Creation date	04/13/2016
Export date	12/23/2015
Population	4.64 m



Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

http://www.tumorregister-muenchen.de/en/facts/base/bC46__E-ICD-10-C46-Kaposi-sarcoma-incidence-and-mortality.pdf

Global Statements about the statistics on the Internet -

Baseline Statistics (grey button ____), Survival (red button ____)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.64 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases^{###} are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, April 2016

- [#] Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007).
- ^{##} Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- ### DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate.

Some remarks regarding this cancer type

Kaposi's sarcoma was once a rare disease. With an increased occurrence of HIV and AIDS the incidence of Kaposi's sarcoma also increased particularly in men. Incidence and prognosis of Kaposi's sarcoma as an HIV-associated disease are highly dependent on the therapeutic success of the infection itself. Progress in this field is reflected in a change of global epidemiological parameters such as incidence, mortality or survival.

ICD-10 codes (ICD-10 2015) used for specifying cancer site

Code	Description
C46 C46.0 C46.1 C46.2 C46.3 C46.7 C46.8 C46.9	Kaposi sarcoma Skin Soft tissue Palate Lymph nodes Other sites Multiple organs Kaposi sarcoma, unspecified

INCIDENCE

Table 1


All patients with invasive cancer by year of diagnosis, proportions of DCO, multiple primaries, deaths, and active follow-up (incl. DCO)


				Prop.		Prop.
		DCO	Prop.	mult.	Prop.	actively
Year of	Cases	cases	DCO	primaries	deaths	followed
diagnosis	n	n	010	90	୍ଚ	00
1998	8	4	50.0	25.0	87.5	100.0
1999	1			100.0	100.0	100.0
2000	9	2	22.2	33.3	44.4	88.9
2001	2	1	50.0		50.0	50.0
2002	4	2	50.0	75.0	100.0	100.0 #
2003	2				100.0	100.0
2004	2	1	50.0	100.0	50.0	100.0
2005	4			25.0	50.0	100.0
2006	6			50.0	16.7	83.3
2007	6	2	33.3	16.7	33.3	83.3 #
2008	7	1	14.3	28.6	57.1	100.0
2009	4			25.0		100.0
2010	1					100.0
2011	7			14.3	57.1	71.4
2012	6			16.7	33.3	83.3
2013	3					100.0
2014	3	1	33.3		33.3	100.0 ##
1998-2014	75	14	18.7	28.0	48.0	90.7

- # The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.
- ## Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be found in the respective headings.



Table la





Incidence measures by year of diagnosis including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

			Males	Fem.	Males	Fem.	Males	Fem.	Males	Fem.
Year of	Males	Females	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
2										
1998	8		0.7		0.5		0.6		0.6	
1999	1		0.1		0.1		0.1		0.1	
2000	8	1 /	0.7	0.1	0.5	0.0	0.6	0.0	0.7	0.0
2001	2		0.2		0.1		0.2		0.2	
2002	3	1	0.2	0.1	0.1	0.0	0.2	0.0	0.1	0.0
2003	2		0.1		0.1		0.1		0.1	
2004	1	1	0.1	0.1	0.0	0.0	0.1	0.0	0.1	0.0
2005	4		0.2		0.1		0.2		0.2	
2006	5	1	0.3	0.0	0.2	0.0	0.2	0.0	0.2	0.0
2007	5	1	0.2	0.0	0.1	0.0	0.2	0.0	0.2	0.0
2008	7		0.3		0.2		0.3		0.3	
2009	4		0.2		0.1		0.1		0.2	
2010	1 /		0.0		0.0		0.0		0.0	
2011	5	2	0.2	0.1	0.1	0.0	0.2	0.1	0.2	0.1
2012	5	1	0.2	0.0	0.1	0.0	0.1	0.0	0.2	0.0
2013	3		0.1		0.1		0.1		0.1	
2014	3		0.1		0.1		0.1		0.1	
1998-2014	67	8	0.2	0.0	0.1	0.0	0.2	0.0	0.2	0.0

The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

				(/				
Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	8	49.5	10.7	36.2	60.4	36.2	37.4	53.1	59.2	60.4
1999	1	70.2		70.2	70.2	70.2	70.2	70.2	70.2	70.2
2000	9	60.3	23.5	31.2	92.9	31.2	44.4	53.4	85.3	92.9
2001	2	63.3	11.6	55.1	71.5	55.1	55.1	63.3	71.5	71.5
2002	4	57.3	11.1	44.5	71.4	44.5	49.6	56.7	65.0	71.4
2003	2	57.9	35.7	32.6	83.1	32.6	32.6	57.9	83.1	83.1
2004	2	67.9	29.3	47.1	88.6	47.1	47.1	67.9	88.6	88.6
2005	4	44.6	11.1	35.1	58.9	35.1	35.8	42.2	53.4	58.9
2006	6	53.7	19.3	29.8	84.6	29.8	43.9	49.0	65.8	84.6
2007	6	46.6	12.1	36.1	70.3	36.1	41.5	43.1	45.4	70.3
2008	7	68.8	17.2	37.8	88.1	37.8	60.4	73.2	85.1	88.1
2009	4	55.3	21.5	35.8	83.7	35.8	39.0	50.8	71.6	83.7
2010	1	50.5		50.5	50.5	50.5	50.5	50.5	50.5	50.5
2011	7 /	66.1	21.8	29.6	95.3	29.6	47.6	68.6	80.7	95.3
2012	6	73.6	16.0	43.0	87.7	43.0	74.5	75.6	85.3	87.7
2013	3	51.7	23.4	28.2	75.1	28.2	28.2	51.8	75.1	75.1
2014	3	66.3	25.9	37.5	87.8	37.5	37.5	73.7	87.8	87.8
1998-2014	75	58.8	18.9	28.2	95.3	36.1	43.0	55.1	74.8	85.3

Age distribution parameters by year of diagnosis (All patients) (incl. DCO)

Table 3a

Age distribution parameters by year of diagnosis (MALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	8	49.5	10.7	36.2	60.4	36.2	37.4	53.1	59.2	60.4
1999	1	70.2		70.2	70.2	70.2	70.2	70.2	70.2	70.2
2000	8	56.4	21.8	31.2	92.9	31.2	40.9	53.0	69.7	92.9
2001	2	63.3	11.6	55.1	71.5	55.1	55.1	63.3	71.5	71.5
2002	3	52.6	7.3	44.5	58.6	44.5	44.5	54.8	58.6	58.6
2003	2	57.9	35.7	32.6	83.1	32.6	32.6	57.9	83.1	83.1
2004	1	47.1		47.1	47.1	47.1	47.1	47.1	47.1	47.1
2005	4	44.6	11.1	35.1	58.9	35.1	35.8	42.2	53.4	58.9
2006	5	47.5	13.4	29.8	65.8	29.8	43.9	44.1	53.9	65.8
2007	5	47.5	13.3	36.1	70.3	36.1	41.5	44.1	45.4	70.3
2008	7	68.8	17.2	37.8	88.1	37.8	60.4	73.2	85.1	88.1
2009	4	55.3	21.5	35.8	83.7	35.8	39.0	50.8	71.6	83.7
2010	1	50.5		50.5	50.5	50.5	50.5	50.5	50.5	50.5
2011	5	63.5	25.5	29.6	95.3	29.6	47.6	68.6	76.2	95.3
2012	5	70.8	16.2	43.0	85.3	43.0	74.5	74.8	76.5	85.3
2013	3	51.7	23.4	28.2	75.1	28.2	28.2	51.8	75.1	75.1
2014	3	66.3	25.9	37.5	87.8	37.5	37.5	73.7	87.8	87.8
1998-2014	67	56.7	18.1	28.2	95.3	35.8	42.1	54.1	73.2	85.1

1190	arberr	Ducion	param		1. DCO		Lagnob	10 (1111)		
Year of diagnosis	Cases n	Mean	Std. dev.	Min.	Max.	10%	25%	Median 50%	75%	90%
2000 2002 2004 2006 2007 2011 2012	1 1 1 1 2 1	91.6 71.4 88.6 84.6 42.1 72.7 87.7	11.3	91.6 71.4 88.6 84.6 42.1 64.7 87.7	84.6	42.1 64.7	91.6 71.4 88.6 84.6 42.1 64.7 87.7	88.6	42.1	71.4 88.6 84.6 42.1
1998-2014	8	76.4	16.6	42.1	91.6	42.1	68.0	82.7	88.2	91.6

Table 3b

Age distribution parameters by year of diagnosis (FEMALES)

Age at	a								
diagnosis	Cases			Males			Females		
Years	n	00	Cum.%	n	9	Cum.%	n	00	Cum.%
25-29	2	5.4	5.4	2	6.1	6.1			0.0
30-34	0	0.0	5.4			6.1			0.0
35-39	4	10.8	16.2	4	12.1	18.2			0.0
40 - 44	5	13.5	29.7	4	12.1	30.3	1	25.0	25.0
45-49	2	5.4	35.1	2	6.1	36.4			25.0
50-54	2	5.4	40.5	2	6.1	42.4			25.0
55-59	1	2.7	43.2	1	3.0	45.5			25.0
60-64	3	8.1	51.4	2	6.1	51.5	1	25.0	50.0
65-69	1	2.7	54.1	1	3.0	54.5			50.0
70-74	6	16.2	70.3	6	18.2	72.7			50.0
75-79	3	8.1	78.4	3	9.1	81.8			50.0
80-84	2	5.4	83.8	1	3.0	84.8	1	25.0	75.0
85+	6	16.2	100.0	5	15.2	100.0	1	25.0	100.0
All ages	37	100.0		33	100.0		4	100.0	

Age distribution by 5-year age group and gender for period 2007-2014 (incl. DCO)

Included in the statistics are 24.2% multiple primaries in males and 0.0% in females.

Age-specific incidence, DCO rate and proportion of all cancers for period 2007-2014

							Males	Females
			Males	Females	Males	Females	Prop.all	Prop.all
Age at			Age-	Age-	DCO rate	DCO rate	cancers	cancers
diagnosis	Males	Females	spec.	spec.	n=3	n=1	n=91183	n=89596
Years	n	n	incid.	incid.	00	010	olo	00
0- 4			0.0	0.0				
5- 9			0.0	0.0				
10-14			0.0	0.0				
15-19			0.0	0.0				
20-24			0.0	0.0				
25-29	2		0.2	0.0			0.4	
30-34			0.0	0.0				
35-39	4		0.3	0.0	25.0		0.3	
40 - 44	4	1	0.2	0.1		100.0	0.2	0.0
45-49	2		0.1	0.0	50.0		0.1	
50-54	2		0.2	0.0			0.0	
55-59	1		0.1	0.0			0.0	
60-64	2	1	0.2	0.1			0.0	0.0
65-69	1		0.1	0.0			0.0	
70-74	6		0.7	0.0			0.0	
75-79	3		0.5	0.0			0.0	
80-84	1	1	0.3	0.2			0.0	0.0
85+	5	1	2.2	0.2	20.0		0.1	0.0
All ages	33	4			9.1	25.0	0.0	0.0
Incidence								
Raw			0.2	0.0				
WS			0.1	0.0				
ES			0.1	0.0				
BRD-S			0.2	0.0				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).






Figure 6. Age distribution and age-specific incidence

MCR





Figure 6a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).



Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2014, based on the November 2013 submission. http://www.seer.cancer.gov.



ICD-10 C46: Kaposi sarcoma

Figure 7. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.

Table 8a Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2014 MALES Observed Expected LCL UCL DCO Diagnosis n SIR 95% 95% EAR 8 n Other primaries 11 1.4 7.6 3.8 13.7 # 422.3 Not observed 0 0.7 0.0 -31.5 0.0 5.2 2.2 9.1 # 390.8 All mult. primaries 11 5.1 2.6

Patients50Median age at second malignancy (years)65.5Person-years226Mean observation time (years)4.5Median observation time (years)2.6


The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries"

		Table 8b					
Standardized in excess absolut	te risk (E		rate of				
Diagnosis	Observed n	Expected n	SIR		UCL 95%	EAR	DCO %
Not observed	0	0.3	0.0	0.0	13.8	-187	
All mult. primaries	0	0.3	0.0	0.0	13.8	-187	
Patients		6					
Person-years		14					
Mean observation time (years)	2.4					
Median observation time		1.7					

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries"



Average incidence (world standard population) 2007 - 2014: Males







Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.1/100,000 WS N=33, females 0.0/100,000 WS N=4).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 65,347 female residents (averaged) in the period from 2007 to 2014 a total of 0 women were identified with newly diagnosed kaposi sarcoma. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.0/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.0/100,000.



Standardized incidence ratio (SIR) 2007 - 2014: Males






Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=33, females N=4).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,924 female residents (averaged) in the period from 2007 to 2014 a total of 0 women were identified with newly diagnosed kaposi sarcoma. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.00. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 50.15, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

						Prop.
		Prop.				deaths
	Incident	actively	Prop.		Prop.	with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	olo	00	n	00	010
1998	8	100.0	50.0	7	87.5	100.0
1999	1	100.0		1	100.0	
2000	9	88.9	22.2	4	44.4	100.0
2001	2 4	50.0	50.0	1 🔨	50.0	100.0
2002	4	100.0	50.0	4	100.0	100.0
2003	2	100.0		4 2	100.0	100.0
2004	2	100.0	50.0	1	50.0	100.0
2005	2 4	100.0		2	50.0	100.0
2006	6	83.3		1	16.7	100.0
2007	6	83.3	33.3	2	33.3	100.0
2008	7	100.0	14.3	4	57.1	100.0
2009	4	100.0				
2010	1	100.0				
2011	7	71.4		4	57.1	75.0
2012	6	83.3		2	33.3	100.0
2013	3	100.0				
2014	3	100.0	33.3	1	33.3	100.0
1998-2014	75	90.7	18.7	36	48.0	94.4



Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)


			Duon		
			Prop. deaths		Prop.
Year of	Incident			Doothe in	
	/	Deethe	with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	'n	90	n	010
1998	8	5	100.0	4	50.0
1999	1	2	50.0	1	100.0
2000	9	3	100.0	2	22.2
2001	2	1	100.0		
2002	4	4	100.0	3	75.0
2003	2	1	100.0		
2004	2	3	100.0	1	50.0
2005	4				
2006	6				
2007	6	4	100.0	2	33.3
2008	7	2	100.0	1	14.3
2009	4	2	100.0	-	1110
2010	1	2	100.0		
2011	7	2	100.0		
2012		4	75.0		
2012	6 3	4	100.0		
	3	4		1	22.2
2014	3	3	100.0	/1 /	33.3
1998-2014	75	42	95.2	15	20.0



Table 10c

Annual cohorts of deaths, proportion of cancer-related and non-cancerrelated deaths, and cancer recorded on death certificates (incl. DCO) (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

				Prop.	
				cancer	
		Prop.	Prop.	recorded	
		cancer-	non-cancer-	on death	
Year of	Deaths				
	/ -	related	related	certificate	
death	n	010	90	00	
1998	5	40.0	60.0	100.0	
1999	2	100.0		100.0	
2000	3	66.7	33.3	100.0	
2001	1	100.0		100.0	
2002	4	50.0	50.0	100.0	
2003	1		100.0		
2004	3	100.0		66.7	
2005					
2006					
2007	4	25.0	75.0	75.0	
2008	2	20.0	100.0	50.0	
2009	2	100.0	100.0	100.0	
	2	100.0	100 0	100.0	
2010		100 0	100.0	100.0	
2011	2	100.0		100.0	
2012	4		100.0	33.3	
2013	4	100.0		100.0	
2014	3	33.3	66.7	66.7	
1998-2014	42	52.4	47.6	77.5	



Year of death	Deaths	Age at death (all causes) Years	Age at death (cancer- related) Years	Age at death (non-cancer- related) Years	Age at death (according to death certificate) Years
ueath	11	IEals	IEals	IEals	lears
1998	5	53.2	52.7	54.2	53.2
1999	2	53.3	53.3		36.1
2000	2	57.9	61.8	54.1	57.9
2001	1	93.6	93.6		93.6
2002	4	56.8	63.3	51.6	56.8
2003	1	88.3		88.3	
2004	2	68.8	68.8		47.1
2005					
2006					
2007	3	45.4	49.7	41.0	47.5
2008	2	54.2		54.2	37.8
2009	3/ 2 1 2 4	71.8	71.8		71.8
2010	1	89.7		89.7	
2011	2	56.0	56.0		56.0
2012		63.9		63.9	91.4
2013	2	86.4	86.4		86.4
2014	3	88.8	65.1	89.7	77.0
1998-2014	36	60.2	65.0	56.3	56.6

Table 11a

Medians of age at death according to the grouping in Table 10 $$\rm MALES$$

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Year of death	Deaths n	Age at death (all causes) Years	Age at death (cancer- related) Years	Age at death (non-cancer- related) Years	Age at death (according to death certificate) Years
1998					
1999					
2000 2001	1	91.6	91.6		91.6
2002 2003					
2004 2005	1	73.0	73.0		73.0
2006					
2007 2008	1	42.1		42.1	42.1
2009 2010 2011	1	89.5		89.5	
2012 2013 2014	2	85.5	85.5		85.5
1998-2014	6	85.5	85.5	65.8	82.4

Table 11b

Medians of age at death according to the grouping in Table 10 $$\operatorname{FEMALES}$

By 2010, life expectancy at birth was 77.5 years for boys and 82.6 years for girls.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Year of

1998

1999

2000

2001

2002

2003 2004

2005 2006

2007

2008 2009

2010 2011

2012

2013

2014

1998-2014

2

1

2

2

2

1

18

0.1

0.0

0.1

0.1

0.1

0.0

0.1

2.00

0.20

0.40

0.67

0.33

0.27

0.50

death

2.19

0.25

0.40

0.37

0.64

0.33

0.27

0.1

0.0

0.1

0.1

0.1

0.0

0.1

MALES												
Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index				
n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S				
2	0.2	0.25	0.1	0.28	0.2	0.30	0.2	0.29				
2	0.2	2.00	0.1	2.03	0.1	1.80	0.2	1.62				
1	0.1	0.13	0.1	0.11	0.1	0.10	0.1	0.10				
1	0.1	0.50	0.0	0.43	0.1	0.57	0.1	0.65				
2	0.1	0.67	0.1	0.63	0.1	0.67	0.1	0.78				

1.63

0.23

0.38

0.44

0.31

0.38

0.25

0.1

0.0

0.1

0.1

0.1

0.0

0.0

2.08

0.22

0.40

0.42

0.52

0.26

0.35

0.1

0.0

0.0

0.1

0.0

0.0

0.0

Table 12a

Mortality measures (cancer-related death) and mortality-incidence-index by year of death

Table	12b
TUDIC	

Mortality measures (cancer-related death) and mortality-incidence-index by year of death FEMALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998									
1999									
2000	1	0.1	1.00	0.0	1.00	0.0	1.00	0.0	1.00
2001									
2002									
2003						7			
2004	1	0.1	1.00	0.0	2.47	0.0	1.85	0.0	1.85
2005									
2006									
2007									
2008									
2009									
2010									
2011									
2012	0	0 1		0 0		0 0		0 0	
2013	2	0.1		0.0		0.0		0.0	
2014									
1998-2014	4	0.0	0.50	0.0	0.34	0.0	0.41	0.0	0.46
1)90-2014	4	0.0	0.00	0.0	0.54	0.0	0.41	0.0	0.40

Age at death	Cases			Males			Females		
Years	n	00	Cum.%	n	00	Cum.%	n	00	Cum.%
45-49 50-54	2 0	20.0	20.0	2	25.0	25.0 25.0			0.0
55-59	0	0.0	20.0			25.0			0.0
60-64	1	10.0	30.0	1	12.5	37.5			0.0
65-69	2	20.0	50.0	2	25.0	62.5			0.0
70-74	1	10.0	60.0	1	12.5	75.0			0.0
75-79	1	10.0	70.0	1	12.5	87.5			0.0
80-84	1	10.0	80.0			87.5	1	50.0	50.0
85+	2	20.0	100.0	1	12.5	100.0	1	50.0	100.0
All ages	10	100.0		8	100.0		2	100.0	

Age distribution of age at death (cancer-related) for period 2007-2014 (incl. multiple primaries)

Table 13

Included in the statistics are 24.2% multiple primaries in males and 0.0% in females.

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2014 (incl. multiple primaries)

Age at death Years	Males Females n n	/ + /	Females Age- spec. .ndex mortal.	MI-index	Males Prop.all cancers %	Females Prop.all cancers %
0- 4 5- 9 10-14 15-19 20-24 25-29 30-34 35-39		0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0	0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0			
40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85+	2 1 2 1 1 1 1 1	0.0 0.0 0.1 0. 0.2 2. 0.1 0. 0.2 0. 0.0	0.0 00 0.0 0.0 50 0.0 50 0.0 0.0 17 0.0 33 0.0 0.2 20 0.2	1.00 1.00	0.2 0.0 0.0 0.0 0.0 0.0	0.0 0.0
All ages	8 2				0.0	0.0
Mortality Raw WS ES BRD-S		0.0 0. 0.0 0.	24 0.0 22 0.0 23 0.0 24 0.0	0.50 0.19 0.27 0.40		
PYLL-70 per 100,000 ES AYLL-70		0.4 0.3 11.5				

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Multiple primaries in deaths in period 1998-2014 MALES

Diagnosis C16 Stomach C19-C20 Rectum C25 Pancreas C33-C34 Lung C44 Skin others C82-C85 NHL C90 Mult. myeloma C91-C96 Leukaemia	Total n 1 1 1 1 1 4 1 2	Total %1 8.3 8.3 8.3 8.3 8.3 33.3 8.3 16.7	Pre n 1 1	Pre ←% 100.0 25.0 50.0	Syn- chron ±30d n 1 1		Post n 1 1 1 2 1	Post ←% 100.0 100.0 100.0 50.0 50.0
All mult. primaries	12	100.0	3	25.0	3	25.0	6	50.0

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

THAT CIPTO	p mar		'EMALE		1100 13	50 2011		
Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	chron	Post n	Post ←%
C82-C85 NHL	3	100.0	1	33.3	1	33.3	1	33.3
All mult. primaries	3	100.0	1	33.3	1	33.3	1	33.3

Table 15b

Multiple primaries in deaths in period 1998-2014 FEMALES

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2014 (First primaries only *)

		Males		Females		Males	Females
Age at		Age-		Age-		Prop.all	Prop.all
death	Males Females	spec.		spec.		cancers	cancers
Years	n n	mortal.	MI-index	mortal.	MI-index	00	90
0- 4		0.0		0.0			
5- 9		0.0		0.0			
10-14		0.0		0.0			
15-19		0.0		0.0			
20-24		0.0		0.0			
25-29		0.0		0.0			
30-34		0.0		0.0			
35-39		0.0		0.0			
40-44		0.0		0.0			
45-49	2	0.1	1.00	0.0		0.2	
50-54		0.0		0.0			
55-59		0.0		0.0			
60-64		0.0		0.0			
65-69	2	0.2	2.00	0.0		0.0	
70-74	1	0.1	0.17	0.0		0.0	
75-79	1	0.2	0.33	0.0		0.0	
80-84	1	0.0		0.2	1.00		0.0
85+	1 1	0.4	0.20	0.2	1.00	0.0	0.0
All ages	7 2					0.0	0.0
Mortality							
Raw		0.0	0.22	0.0	0.50		
WS		0.0	0.19	0.0	0.19		
ES		0.0	0.20	0.0	0.27		
BRD-S		0.0	0.22	0.0	0.40		
DIE		0.0	0.22	0.0	0.10		
PYLL-70							
per 100,000		0.3					
ES ES		0.2					
AYLL-70		12.5					
		+2.5					

* See corresponding tables with multiple primaries.

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2013 (Single primaries only *)

Age at death Years	Males Females n n	/ = /	Females Age- spec. mortal. MI-index	Males Females Prop.all Prop.all cancers cancers
$\begin{array}{c} 0-4\\ 5-9\\ 10-14\\ 15-19\\ 20-24\\ 25-29\\ 30-34\\ 35-39\\ 40-44\\ 45-49 \end{array}$	1	0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0	0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0	0.1
43-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85+		$\begin{array}{c} 0.1 & 0.37 \\ 0.0 \\ 0.0 \\ 0.0 \\ 0.0 \\ 0.1 & 0.23 \\ 0.2 & 0.38 \\ 0.0 \\ 0.0 \end{array}$	$\begin{array}{c} 0.0\\ 0.0\\ 0.0\\ 0.0\\ 0.0\\ 0.0\\ 0.0\\ 0.2\\ 1.14\\ 0.2\\ 1.15 \end{array}$	0.0 0.0 0.0 0.0
All ages Mortality	3 2			0.0 0.0
Raw WS ES BRD-S		$\begin{array}{ccc} 0.0 & 0.13 \\ 0.0 & 0.10 \\ 0.0 & 0.11 \\ 0.0 & 0.13 \end{array}$	0.0 0.57 0.0 0.21 0.0 0.31 0.0 0.46	
PYLL-70 per 100,000 ES AYLL-70		0.2 0.1 22.5		

* See corresponding tables with multiple primaries.





Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at kaposi sarcoma-related death (see Table 10) should be considered.






Average mortality (world standard population) 2007 - 2014: Males

Average mortality (world standard population) 2007 - 2014: Females



Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2007 to 2014. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.0/100,000 WS N=8, females 0.0/100,000 WS N=2).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 65,347 female residents (averaged) in the period from 2007 to 2014 a total of 0 women died from kaposi sarcoma. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.0/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.0/100,000.



Standardized mortality ratio (SMR) 2007 - 2014: Males

Standardized mortality ratio (SMR) 2007 - 2014: Females



Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=8, females N=2).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,924 female residents (averaged) in the period from 2007 to 2014 a total of 0 women died from kaposi sarcoma. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.00. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 110.43, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases, where applicable. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

FRG GEKID	Federal Republic of Germany Association of Population-based Cancer Registries in Germany (Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)
MCR	Munich Cancer Registry (Tumorregister München)
SEER	Surveillance, Epidemiology, and End Results (USA)
AYLL-70 BRD-S	Average years of life lost prior to age 70 given a person dies before that age German standard population
DCO	Death certificate only
EAR	Excess absolute risk
50	= excess cancer cases (O - E) per 10,000 person-years
ES	European standard population (old)
LCL	Lower confidence limit
MI-index	Ratio between mortality and incidence
PYLL-70	Potential years of life lost prior to age 70 given a person dies before that age
SIR	Standardized incidence ratio
SMR	Standardized mortality ratio
UCL	Upper confidence limit
WS	World standard population

Recommended Citation

Munich Cancer Registry. ICD-10 C46: Kaposi sarcoma - Incidence and Mortality [Internet]. 2016 [updated 2016 Apr 13; cited 2016 Jun 1]. Available from: http://www.tumorregister-

muenchen.de/en/facts/base/bC46_E-ICD-10-C46-Kaposi-sarcoma-incidence-and-mortality.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

Fig./Tb	l.	Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	4
1a	Gender distribution by year of diagnosis	5
2	Incidence by year of diagnosis	6
3	Age distribution parameters by year of diagnosis	7
4	Age distribution by 5-year age group and gender	9
5	Age-specific incidence, DCO rate, proportion malignancies	10
6	Age distribution and age-specific incidence (chart)	11
6a	Age-specific incidence internationally (chart)	12
7	Cumulative follow-up years (chart)	13
8	Standardized incidence ratio of second primaries	13
9a	Map of cancer incidence (WS) by county (chart)	15
9b	Standardized incidence ratio (SIR) by county (chart)	16
10a	Pts incident cohorts and mortality / yr	17
10b	Incidence and mortality by year of diagnosis	18
10c	Cancer-related deaths, death certification available / yr	19
11	Medians of age at death / yr	20
12	Mortality by year of death	22
13	Distribution of age at death	23
14	Age-specific mortality	24
15	Multiple primaries in deaths	25
16	Age-specific mortality (first primaries)	27
17	Age-specific mortality (single primaries)	28
18	Age distribution and age-specific mortality (chart)	29
19a	Map of cancer mortality (WS) by county (chart)	30
19b	Standardized mortality ratio (SMR) by county (chart)	31

