Munich Cancer Registry



- Survival
- Selection Matrix
- Homepage
- ▶ Deutsch

ICD-10 C50: Breast cancer (men)

Incidence and Mortality

| Year of diagnosis | 1998-2014 |
|--------------------|------------|
| Patients | 423 |
| Diseases | 427 |
| Creation date | 04/13/2016 |
| Export date | 12/23/2015 |
| Population (males) | 2.28 m |



Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

http://www.tumorregister-muenchen.de/en/facts/base/bC50m_E-ICD-10-C50-Breast-cancer-men-incidence-and-mortality.pdf

Global Statements about the statistics on the Internet -

Baseline Statistics (grey button ____), Survival (red button ____)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.64 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases^{###} are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, April 2016

- [#] Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007).
- ^{##} Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- ### DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate.

ICD-10 codes (ICD-10 2015) used for specifying cancer site

| Code | Description |
|-------|--------------------------------|
| C50 | Malignant neoplasm of breast |
| C50.0 | Nipple and areola |
| C50.1 | Central portion of breast |
| C50.2 | Upper-inner quadrant of breast |
| C50.3 | Lower-inner quadrant of breast |
| C50.4 | Upper-outer quadrant of breast |
| C50.5 | Lower-outer quadrant of breast |
| C50.6 | Axillary tail of breast |
| C50.8 | Overlapping lesion of breast |
| C50.9 | Breast, unspecified |
| | |

Gender: Male

INCIDENCE

Table 1

All patients with invasive cancer by year of diagnosis, proportions of DCO, multiple primaries, deaths, and active follow-up (incl. DCO)

| | | | | Prop. | | Prop. | |
|-----------|-------|-------|-------|-----------|--------|----------|--|
| | | DCO | Prop. | mult. | Prop. | actively | |
| Year of | Cases | cases | DCO | primaries | deaths | followed | |
| diagnosis | n | n | 00 | 00 | 00 | 90 | |
| | | | | | | | |
| 1998 | 7 | 1 | 14.3 | 42.9 | 57.1 | 100.0 | |
| 1999 | 12 | 2 | 16.7 | 58.3 | 83.3 | 100.0 | |
| 2000 | 13 | 3 | 23.1 | 38.5 | 69.2 | 92.3 | |
| 2001 | 13 | | | 30.8 | 46.2 | 84.6 | |
| 2002 | 20 | 3 | 15.0 | 35.0 | 75.0 | 95.0 # | |
| 2003 | 35 | 1 | 2.9 | 40.0 | 54.3 | 100.0 | |
| 2004 | 27 | 3 | 11.1 | 33.3 | 55.6 | 96.3 | |
| 2005 | 23 | | | 26.1 | 34.8 | 95.7 | |
| 2006 | 24 | 2 | 8.3 | 33.3 | 45.8 | 95.8 | |
| 2007 | 41 | 2 | 4.9 | 41.5 | 53.7 | 80.5 # | |
| 2008 | 26 | 5 | 19.2 | 50.0 | 46.2 | 84.6 | |
| 2009 | 26 | 1 | 3.8 | 11.5 | 23.1 | 53.8 | |
| 2010 | 28 | | | 25.0 | 28.6 | 53.6 | |
| 2011 | 32 | 2 | 6.3 | 21.9 | 34.4 | 65.6 | |
| 2012 | 34 | 2 | 5.9 | 29.4 | 29.4 | 82.4 | |
| 2013 | 40 | 2 | 5.0 | 45.0 | 25.0 | 100.0 | |
| 2014 | 26 | | | 26.9 | | 100.0 ## | |
| | | | | | | | |
| 1998-2014 | 427 | 29 | 6.8 | 34.0 | 41.2 | 85.7 | |
| | | | | | | | |

- # The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.
- ## Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be found in the respective headings.



Incidence measures by year of diagnosis including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

| Year of | Cases | Incidence | Incidence | Incidence | Incidence |
|-----------|-------|-----------|-----------|-----------|-----------|
| diagnosis | n | raw | WS | ES | BRD-S |
| | _ | | | | |
| 1998 | 7 | 0.6 | 0.4 | 0.5 | 0.7 |
| 1999 | 12 | 1.1 | 0.6 | 1.0 | 1.2 |
| 2000 | 13 | 1.1 | 0.7 | 1.0 | 1.4 |
| 2001 | 13 | 1.1 | 0.7 | 1.0 | 1.2 |
| 2002 | 20 | 1.1 | 0.6 | 0.9 | 1.2 |
| 2003 | 35 | 1.9 | 1.0 | 1.5 | 2.0 |
| 2004 | 27 | 1.4 | 0.8 | 1.2 | 1.5 |
| 2005 | 23 | 1.2 | 0.6 | 1.0 | 1.3 |
| 2006 | 24 | 1.3 | 0.7 | 1.0 | 1.3 |
| 2007 | 41 | 1.9 | 0.9 | 1.4 | 1.9 |
| 2008 | 26 | 1.2 | 0.6 | 0.9 | 1.2 |
| 2009 | 26 | 1.2 | 0.6 | 0.9 | 1.1 |
| 2010 | 28 | 1.2 | 0.6 | 0.9 | 1.2 |
| 2011 | 32 | 1.4 | 0.7 | 1.0 | 1.4 |
| 2012 | 34 | 1.5 | 0.8 | 1.1 | 1.5 |
| 2013 | 40 | 1.8 | 0.8 | 1.2 | 1.6 |
| 2014 | 26 | 1.1 | 0.5 | 0.8 | 1.1 |
| | | | | | |
| 1998-2014 | 427 | 1.3 | 0.7 | 1.0 | 1.3 |
| | | | | | |

The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

| Year of | Cases | | Std. | | | | | Median | | |
|-----------|-------|------|------|------|------|------|------|--------|------|------|
| diagnosis | n | Mean | dev. | Min. | Max. | 10% | 25% | 50% | 75% | 90% |
| 1998 | 7 | 59.8 | 9.9 | 47.9 | 77.4 | 47.9 | 49.1 | 59.8 | 63.6 | 77.4 |
| 1999 | 12 | 68.8 | 10.3 | 52.8 | 85.2 | 55.5 | 62.0 | 66.1 | 78.6 | 79.6 |
| 2000 | 13 | 66.7 | 15.5 | 41.2 | 87.8 | 44.4 | 56.6 | 68.8 | 79.7 | 84.7 |
| 2001 | 13 | 63.4 | 9.9 | 48.9 | 84.7 | 50.4 | 58.4 | 62.6 | 67.0 | 77.8 |
| 2002 | 20 | 66.1 | 12.1 | 34.5 | 82.5 | 49.4 | 60.7 | 67.5 | 75.7 | 79.8 |
| 2003 | 35 | 66.5 | 13.5 | 30.3 | 89.6 | 46.2 | 59.6 | 68.3 | 76.0 | 82.5 |
| 2004 | 27 | 68.2 | 10.7 | 45.7 | 89.6 | 55.1 | 62.7 | 70.3 | 74.3 | 85.4 |
| 2005 | 23 | 71.2 | 8.6 | 52.6 | 90.9 | 56.5 | 68.1 | 71.8 | 76.5 | 81.1 |
| 2006 | 24 | 65.5 | 12.5 | 45.7 | 86.1 | 47.4 | 57.2 | 66.7 | 73.5 | 84.7 |
| 2007 | 41 | 69.3 | 11.6 | 41.3 | 96.1 | 58.2 | 60.7 | 71.7 | 76.4 | 80.6 |
| 2008 | 26 | 68.8 | 9.2 | 44.9 | 83.8 | 60.3 | 62.6 | 66.1 | 77.1 | 80.6 |
| 2009 | 26 | 69.1 | 11.5 | 46.0 | 89.5 | 54.4 | 61.4 | 70.4 | 77.3 | 85.9 |
| 2010 | 28 | 69.4 | 12.0 | 44.0 | 91.0 | 47.7 | 62.9 | 69.9 | 76.1 | 87.6 |
| 2011 | 32 | 70.8 | 11.5 | 48.6 | 90.6 | 55.3 | 60.1 | 71.6 | 81.3 | 84.0 |
| 2012 | 34 | 66.9 | 10.5 | 46.8 | 83.7 | 49.9 | 59.8 | 67.6 | 77.1 | 78.6 |
| 2013 | 40 | 70.7 | 12.9 | 45.1 | 89.3 | 49.9 | 65.6 | 72.4 | 78.8 | 88.4 |
| 2014 | 26 | 69.2 | 11.1 | 41.7 | 83.6 | 54.2 | 58.6 | 72.0 | 77.3 | 81.9 |
| 1998-2014 | 427 | 68.3 | 11.6 | 30.3 | 96.1 | 52.2 | 60.6 | 69.8 | 76.9 | 82.5 |

Age distribution parameters by year of diagnosis (incl. DCO)

Age distribution by 5-year age group for period 2007-2014 (incl. DCO)

| Age at | | | | |
|-----------|-------|-------|-------|--|
| diagnosis | Cases | | | |
| Years | n | 00 | Cum.% | |
| 40-44 | 4 | 1.6 | 1.6 | |
| 45-49 | 16 | 6.3 | 7.9 | |
| 50-54 | 10 | 4.0 | 11.9 | |
| 55-59 | 20 | 7.9 | 19.8 | |
| 60-64 | 36 | 14.2 | 34.0 | |
| 65-69 | 32 | 12.6 | 46.6 | |
| 70-74 | 47 | 18.6 | 65.2 | |
| 75-79 | 45 | 17.8 | 83.0 | |
| 80-84 | 27 | 10.7 | 93.7 | |
| 85+ | 16 | 6.3 | 100.0 | |
| | | | | |
| All ages | 253 | 100.0 | | |
| | | | | |

Included in the statistics are 38.6% multiple primaries.


Age-specific incidence, DCO rate and proportion of all cancers for period 2007-2014

| | | | | Prop. all | |
|-----------|-------|-----------|----------|-----------|--|
| Age at | | | DCO rate | cancers | |
| diagnosis | Cases | Age-spec. | n=14 | n=91183 | |
| Years | n | incidence | 8 | % | |
| IEals | 11 | THETGENCE | ō | 0 | |
| 0 1 | | 0.0 | | | |
| 0-4 | | 0.0 | | | |
| 5-9 | | 0.0 | | | |
| 10-14 | | 0.0 | | | |
| 15-19 | | 0.0 | | | |
| 20-24 | | 0.0 | | | |
| 25-29 | | 0.0 | | | |
| 30-34 | | 0.0 | | | |
| 35-39 | | 0.0 | | | |
| 40-44 | 4 | 0.2 | 25.0 | 0.2 | |
| 45-49 | 16 | 1.0 | | 0.5 | |
| 50-54 | 10 | 0.8 | | 0.2 | |
| 55-59 | 19 | 1.8 | | 0.3 | |
| 60-64 | 36 | 3.7 | 5.6 | 0.3 | |
| 65-69 | 32 | 3.3 | 3.1 | 0.2 | |
| 70-74 | 46 | 5.1 | | 0.3 | |
| 75-79 | 45 | 8.2 | 6.7 | 0.4 | |
| 80-84 | 27 | 7.7 | 14.8 | 0.3 | |
| 85+ | 15 | 6.5 | 20.0 | 0.2 | |
| | | | | | |
| All ages | 250 | | 5.6 | 0.3 | |
| 2 | | | | | |
| Incidence | | | | | |
| Raw | | 1.4 | | | |
| WS | | 0.7 | | | |
| ES | | 1.0 | | | |
| BRD-S | | 1.3 | | | |
| | | | | | |
| | | | | | |

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).



Table 5



ICD-10 C50: Malignant neoplasm of breast (men)

Figure 6. Age distribution and age-specific incidence





Figure 6a. Age-specific incidence in MCR registry areas compared to Germany (FRG, GEKID extrapolation) and SEER (Surveillance, Epidemiology, and End Results, USA).



Reference:

Extrapolated age-specific patient population of Germany, data status middle of 2010. Association of Population-based Cancer Registries in Germany (GEKID e.V.). Berlin, 2014. http://www.gekid.de. Last access: 02/11/2015

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2014, based on the November 2013 submission. http://www.seer.cancer.gov.



ICD-10 C50: Malignant neoplasm of breast (men)

Figure 7. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.




Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2014

| | Observed | Expected | | LCL | UCL | | DCO |
|-----------------------|-------------|----------|------|------|---------|-------|------|
| Diagnosis | n | n | SIR | 95% | 95% | EAR | 010 |
| | | | | | | | |
| C09-C10 Oropharynx | 2 | 0.2 | 8.1 | 1.0 | 29.2 | 12.4 | |
| C15 Oesophagus | 2 | 0.5 | 4.4 | 0.5 | 16.0 | 10.9 | |
| C16 Stomach | 5 5 5 | 1.0 | 4.8 | 1.6 | 11.3 # | 28.0 | 20.0 |
| C18 Colon | 5 | 2.5 | 2.0 | 0.6 | 4.7 | 17.6 | |
| C25 Pancreas | 5 | 0.9 | 5.3 | 1.7 | 12.4 # | 28.6 | |
| C33-C34 Lung | 7 | 3.0 | 2.4 | 0.9 | 4.8 | 28.4 | 14.3 |
| C50 Breast | 4 | 0.1 | 57.8 | 15.8 | 148.1 # | 27.8 | |
| C61 Prostate | 23 | 7.4 | 3.1 | 2.0 | 4.7 # | 110.0 | 13.0 |
| C64 Kidney | 2 | 0.9 | 2.3 | 0.3 | 8.2 | 7.9 | |
| C67 Bladder | 3 | 1.2 | 2.6 | 0.5 | 7.5 | 13.0 | |
| C82-C85 NHL | 2 | 1.0 | 2.0 | 0.2 | 7.1 | 6.9 | |
| | | | | | | | |
| Other primaries | 6 | 3.2 | 1.9 | 0.7 | 4.1 | 19.6 | 16.7 |
| Not observed | 0 | 3.7 | 0.0 | 0.0 | 1.0 | -25.8 | |
| | | | | | | | |
| All mult. primaries | 66 | 25.6 | 2.6 | 2.0 | 3.3 # | 285.3 | 9.1 |
| | | | | | | | |
| | | | | | | | |
| Patients | | | 404 | | | | |
| Median age at second | malignancy | (years) | 71.4 | | | | |
| Person-years | | | 1416 | | | | |
| Mean observation time | (years) | | 3.5 | | | | |
| Median observation ti | me (years) | | 2.2 | | | | |
| | | | | | | | |

The occurrence of second malignancy is statistically significant.


Observed second primaries with count 1 are pooled in category "Other primaries"



Average incidence (world standard population) 2007 - 2014

Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (0.7/100,000 WS N=250).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,886 male residents (averaged) in the period from 2007 to 2014 a total of 8 men were identified with newly diagnosed breast cancer (men). Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.6/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.2 and 1.6/100,000.



Standardized incidence ratio (SIR) 2007 - 2014

Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (N=250).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,358 male residents (averaged) in the period from 2007 to 2014 a total of 8 men were identified with newly diagnosed breast cancer (men). Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 1.12. Though, the value of this parameter may vary with an underlying probability of 99% between 0.36 and 2.59, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

| Year of diagnosis | Incident cases n | Prop. actively followed % | Prop. DCO % | Deaths n | Prop. deaths % | Prop. deaths with death certific. % |
|----------------------|------------------------|------------------------------------|-------------------|-------------|----------------------|---|
| 1998 | 7 | 100.0 | 14.3 | 4 | 57.1 | 75.0 |
| 1999 | 12 | 100.0 | 16.7 | 10 | 83.3 | 100.0 |
| 2000 | 13 | 92.3 | 23.1 | 9 | 69.2 | 100.0 |
| 2001 | 13 | 84.6 | | 6 | 46.2 | 83.3 |
| 2002 | 20 | 95.0 | 15.0 | 15 | 75.0 | 100.0 |
| 2003 | 35 | 100.0 | 2.9 | 19 | 54.3 | 94.7 |
| 2004 | 27 | 96.3 | 11.1 | 15 | 55.6 | 100.0 |
| 2005 | 23 | 95.7 | | 8 | 34.8 | 100.0 |
| 2006 | 24 | 95.8 | 8.3 | 11 | 45.8 | 100.0 |
| 2007 | 41 | 80.5 | 4.9 | 22 | 53.7 | 95.5 |
| 2008 | 26 | 84.6 | 19.2 | 12 | 46.2 | 100.0 |
| 2009 | 26 | 53.8 | 3.8 | 6 | 23.1 | 100.0 |
| 2010 | 28 | 53.6 | | 8 | 28.6 | 100.0 |
| 2011 | 32 | 65.6 | 6.3 | 11 | 34.4 | 100.0 |
| 2012 | 34 | 82.4 | 5.9 | 10 | 29.4 | 100.0 |
| 2013 | 40 | 100.0 | 5.0 | 10 | 25.0 | 100.0 |
| 2014 | 26 | 100.0 | | | | |
| 1998-2014 | 427 | 85.7 | 6.8 | 176 | 41.2 | 97.7 |



Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

| | | | Prop. | | |
|------------|----------|--------|------------|-------------|-----------|
| | | | deaths | | Prop. |
| Year of | Incident | | with death | Deaths in | deaths in |
| diagnosis/ | cases | Deaths | certific. | same year | same year |
| death | n | n | 90 | n | 00 |
| 1.0.0.0 | | | | | |
| 1998 | 7 | 4 | 100.0 | | |
| 1999 | 12 | 3 | 100.0 | 1 | 8.3 |
| 2000 | 13 | 7 | 100.0 | 4 | 30.8 |
| 2001 | 13 | 12 | 83.3 | | |
| 2002 | 20 | 9 | 88.9 | 4 | 20.0 |
| 2003 | 35 | 10 | 100.0 | 5 | 14.3 |
| 2004 | 27 | 13 | 100.0 | 2 | 7.4 |
| 2005 | 23 | 10 | 100.0 | _ 1 | 4.3 |
| 2006 | 24 | 12 | 100.0 | 2 | 8.3 |
| 2007 | 41 | 14 | 100.0 | 5 | 12.2 |
| 2008 | 26 | 13 | 100.0 | 5 | 19.2 |
| 2009 | 26 | 16 | 100.0 | 2 | 7.7 |
| 2010 | 28 | 13 | 100.0 | | |
| 2011 | 32 | 21 | 100.0 | 2 | 6.3 |
| 2012 | 34 | 21 | 95.2 | 2 2 7 | 5.9 |
| 2013 | 40 | 24 | 100.0 | 7 | 17.5 |
| 2014 | 26 | 15 | 100.0 | | |
| 1998-2014 | 427 | 217 | 98.2 | 42 | 9.8 |



Table 10c

Annual cohorts of deaths, proportion of cancer-related and non-cancerrelated deaths, and cancer recorded on death certificates (incl. DCO) (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

| | | | | Prop. | |
|-----------|--------|---------|-------------|-------------|--|
| | | | | cancer | |
| | | Prop. | Prop. | recorded | |
| | | cancer- | non-cancer- | on death | |
| Year of | Deaths | related | related | certificate | |
| death | n | 010 | 00 | 00 | |
| | | | | | |
| 1998 | 4 | 75.0 | 25.0 | 75.0 | |
| 1999 | 3 | 66.7 | 33.3 | 66.7 | |
| 2000 | 7 | 85.7 | 14.3 | 85.7 | |
| 2001 | 12 | 83.3 | 16.7 | 90.0 | |
| 2002 | 9 | 66.7 | 33.3 | 75.0 | |
| 2003 | 10 | 70.0 | 30.0 | 90.0 | |
| 2004 | 13 | 69.2 | 30.8 | 100.0 | |
| 2005 | 10 | 70.0 | 30.0 | 90.0 | |
| 2006 | 12 | 83.3 | 16.7 | 91.7 | |
| 2007 | 14 | 50.0 | 50.0 | 71.4 | |
| 2008 | 13 | 38.5 | 61.5 | 61.5 | |
| 2009 | 16 | 81.3 | 18.8 | 93.8 | |
| 2010 | 13 | 61.5 | 38.5 | 61.5 | |
| 2011 | 21 | 90.5 | 9.5 | 85.7 | |
| 2012 | 21 | 76.2 | 23.8 | 70.0 | |
| 2013 | 24 | 37.5 | 62.5 | 62.5 | |
| 2014 | 15 | 86.7 | 13.3 | 93.3 | |
| | | | | | |
| 1998-2014 | 217 | 69.1 | 30.9 | 79.8 | |
| | | | | | |

Medians of age at death according to the grouping in Table 10

| Year of death | Deaths n | Age at death (all causes) Years | Age at death (cancer- related) Years | Age at death (non-cancer- related) Years | Age at death (according to death certificate) Years |
|------------------|-------------|---|--|--|--|
| 1998 | 4 | 81.1 | 82.8 | 79.4 | 82.8 |
| 1999 | 3 | 68.0 | 67.3 | 79.7 | 67.3 |
| 2000 | 7 | 79.1 | 78.0 | 87.8 | 78.2 |
| 2001 | 12 | 71.9 | 69.1 | 84.5 | 70.2 |
| 2002 | 9 | 75.9 | 76.1 | 69.3 | 76.1 |
| 2003 | 10 | 71.9 | 70.2 | 88.4 | 71.4 |
| 2004 | 13 | 79.9 | 78.9 | 86.4 | 79.9 |
| 2005 | 10 | 80.8 | 79.5 | 84.5 | 82.1 |
| 2006 | 12 | 70.6 | 70.6 | 72.9 | 68.0 |
| 2007 | 14 | 78.4 | 69.1 | 87.6 | 76.4 |
| 2008 | 13 | 82.0 | 70.8 | 86.1 | 78.3 |
| 2009 | 16 | 70.7 | 72.1 | 66.9 | 69.3 |
| 2010 | 13 | 71.4 | 73.0 | 71.4 | 73.0 |
| 2011 | 21 | 74.1 | 74.1 | 84.6 | 74.6 |
| 2012 | 21 | 79.2 | 75.7 | 80.8 | 77.7 |
| 2013 | 24 | 80.3 | 76.9 | 80.9 | 82.8 |
| 2014 | 15 | 78.2 | 78.2 | 77.8 | 75.8 |
| 1998-2014 | 217 | 76.8 | 74.6 | 81.4 | 76.0 |



By 2010, life expectancy at birth was 77.5 years for boys and 82.6 years for girls.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Mortality measures (cancer-related death) and mortality-incidence-index by year of death $% \left({\left({{{\mathbf{x}}_{i}} \right)} \right)$

| Year of | Deaths | Mort. | MI-Index | Mort. | MI-Index | Mort. | MI-Index | Mort. | MI-Index |
|-----------|--------|-------|----------|-------|----------|-------|----------|-------|----------|
| death | n | raw | raw | WS | WS | ES | ES | BRD-S | BRD-S |
| | | | | | | | | | |
| 1998 | 3 | 0.3 | 0.43 | 0.2 | 0.38 | 0.3 | 0.48 | 0.4 | 0.62 |
| 1999 | 2 | 0.2 | 0.17 | 0.1 | 0.20 | 0.2 | 0.18 | 0.2 | 0.15 |
| 2000 | 6 | 0.5 | 0.46 | 0.3 | 0.42 | 0.5 | 0.46 | 0.8 | 0.57 |
| 2001 | 10 | 0.9 | 0.77 | 0.5 | 0.74 | 0.8 | 0.79 | 1.0 | 0.89 |
| 2002 | 6 | 0.3 | 0.30 | 0.2 | 0.27 | 0.3 | 0.32 | 0.4 | 0.35 |
| 2003 | 7 | 0.4 | 0.20 | 0.2 | 0.21 | 0.3 | 0.21 | 0.4 | 0.22 |
| 2004 | 9 | 0.5 | 0.33 | 0.2 | 0.28 | 0.4 | 0.31 | 0.5 | 0.36 |
| 2005 | 7 | 0.4 | 0.30 | 0.2 | 0.28 | 0.3 | 0.32 | 0.5 | 0.35 |
| 2006 | 10 | 0.5 | 0.42 | 0.3 | 0.39 | 0.4 | 0.40 | 0.5 | 0.42 |
| 2007 | 7 | 0.3 | 0.17 | 0.2 | 0.17 | 0.2 | 0.18 | 0.3 | 0.16 |
| 2008 | 5 | 0.2 | 0.19 | 0.1 | 0.16 | 0.2 | 0.18 | 0.2 | 0.20 |
| 2009 | 13 | 0.6 | 0.50 | 0.3 | 0.48 | 0.5 | 0.51 | 0.6 | 0.53 |
| 2010 | 8 | 0.4 | 0.29 | 0.2 | 0.25 | 0.2 | 0.27 | 0.4 | 0.30 |
| 2011 | 19 | 0.8 | 0.61 | 0.4 | 0.60 | 0.6 | 0.61 | 0.8 | 0.60 |
| 2012 | 16 | 0.7 | 0.47 | 0.3 | 0.35 | 0.4 | 0.39 | 0.7 | 0.47 |
| 2013 | 9 | 0.4 | 0.24 | 0.1 | 0.19 | 0.3 | 0.23 | 0.4 | 0.26 |
| 2014 | 13 | 0.6 | 0.50 | 0.2 | 0.44 | 0.4 | 0.47 | 0.5 | 0.47 |
| | | | | | | | | | |
| 1998-2014 | 150 | 0.5 | 0.35 | 0.2 | 0.32 | 0.4 | 0.34 | 0.5 | 0.37 |

Age distribution of age at death (cancer-related) for period 2007-2014 (incl. multiple primaries)

| Age at | | | | |
|----------|-------|-------|-------|--|
| death | Cases | | | |
| Years | n | 90 | Cum.% | |
| 40-44 | 2 | 2.2 | 2.2 | |
| 45-49 | 2 | 2.2 | 4.3 | |
| 50-54 | 2 | 2.2 | 6.5 | |
| 55-59 | 4 | 4.3 | 10.8 | |
| 60-64 | 6 | 6.5 | 17.2 | |
| 65-69 | 13 | 14.0 | 31.2 | |
| 70-74 | 15 | 16.1 | 47.3 | |
| 75-79 | 20 | 21.5 | 68.8 | |
| 80-84 | 12 | 12.9 | 81.7 | |
| 85+ | 17 | 18.3 | 100.0 | |
| | | | | |
| All ages | 93 | 100.0 | | |
| | | | | |

Included in the statistics are 38.6% multiple primaries.

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2014 (incl. multiple primaries)

| Age at | | | | Prop. all | |
|-------------|-------|-----------|----------|-----------|--|
| death | Cases | Age-spec. | | cancers | |
| Years | n | mortality | MI-index | 00 | |
| | | | | | |
| 0- 4 | | 0.0 | | | |
| 5- 9 | | 0.0 | | | |
| 10-14 | | 0.0 | | | |
| 15-19 | | 0.0 | | | |
| 20-24 | | 0.0 | | | |
| 25-29 | | 0.0 | | | |
| 30-34 | | 0.0 | | | |
| 35-39 | | 0.0 | | | |
| 40 - 44 | 2 | 0.1 | 0.50 | 0.4 | |
| 45-49 | 2 | 0.1 | 0.13 | 0.2 | |
| 50-54 | 2 | 0.2 | 0.20 | 0.1 | |
| 55-59 | 4 | 0.4 | 0.20 | 0.1 | |
| 60-64 | 6 | 0.6 | 0.17 | 0.1 | |
| 65-69 | 13 | 1.4 | 0.41 | 0.2 | |
| 70-74 | 15 | 1.4 | 0.32 | 0.2 | |
| 75-79 | | | | | |
| | 20 | 3.6 | 0.44 | 0.2 | |
| 80-84 | 12 | 3.4 | 0.44 | 0.2 | |
| 85+ | 17 | 7.3 | 1.06 | 0.3 | |
| | | | | | |
| All ages | 93 | | | 0.2 | |
| | | | | | |
| Mortality | | | | | |
| Raw | | 0.5 | 0.37 | | |
| WS | | 0.2 | 0.33 | | |
| ES | | 0.4 | 0.35 | | |
| BRD-S | | 0.5 | 0.37 | | |
| | | | | | |
| PYLL-70 | | | | | |
| per 100,000 | | 1.6 | | | |
| ËS | | 1.4 | | | |
| AYLL-70 | | 9.1 | | | |
| | | | | | |

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Multiple primaries in deaths in period 1998-2014

| | | | | | Syn- | Syn- | | |
|----------------------|------------|-------|-----|-------|-------|-------|------|-------|
| | / . | | - | | chron | chron | - · | |
| | Total | Total | Pre | Pre | ±30d | ±30d | Post | Post |
| Diagnosis | n | %↓ | n | ÷→ | n | ~⁰ | n | ⇔⊖ |
| | | | | | | | | |
| C09-C10 Oropharynx | 3 | 3.5 | | | | | 3 | 100.0 |
| C12-C13 Hypopharynx | / 1 / | 1.2 | | | | | 1 | 100.0 |
| C15 Oesophagus | 3 | 3.5 | 3 | 100.0 | | | | |
| C16 Stomach | 5 | 5.9 | 3 | 60.0 | | | 2 | 40.0 |
| C18 Colon | 5 | 5.9 | 2 | 40.0 | | | 3 | 60.0 |
| C19-C20 Rectum | 4 | 4.7 | 4 | 100.0 | | | | |
| C22 Liver | 2 | 2.4 | 1 | 50.0 | | | 1 | 50.0 |
| C25 Pancreas | 5 | 5.9 | | | | | 5 | 100.0 |
| C30-C31 Sinuses | 1 | 1.2 | 1 | 100.0 | | | | |
| C32 Larynx | 2 | 2.4 | 2 | 100.0 | | | | |
| C33-C34 Lung | 13 | 15.3 | 1 | 7.7 | 3 | 23.1 | 9 | 69.2 |
| C43 Malign. melanoma | 1 | 1.2 | 1 | 100.0 | | | | |
| C44 Skin others | 6 | 7.1 | 2 | 33.3 | _ 1 | 16.7 | 3 | 50.0 |
| C46,C49 Soft tissue | 1 | 1.2 | | | | | 1 | 100.0 |
| C50 Breast | 6 | 7.1 | | | 4 | 66.7 | 2 | 33.3 |
| C61 Prostate | 14 | 16.5 | 7 | 50.0 | 1 | 7.1 | 6 | 42.9 |
| C64 Kidney | 2 | 2.4 | | | | | 2 | 100.0 |
| C67 Bladder | 4 | 4.7 | 3 | 75.0 | | | 1 | 25.0 |
| C76-C79 CUP | 1 | 1.2 | 1 | 100.0 | | | | |
| C81 Hodgkin lymphoma | 2 | 2.4 | 1 | 50.0 | | | 1 | 50.0 |
| C82-C85 NHL | 3 | 3.5 | 3 | 100.0 | | | - | 00.0 |
| C91-C96 Leukaemia | 1 | 1.2 | 5 | 100.0 | | | 1 | 100.0 |
| ost oso leakaemia | _ <u> </u> | 1.2 | | | | | - | -00.0 |
| All mult. primaries | 85 | 100.0 | 35 | 41.2 | 9 | 10.6 | 41 | 48.2 |

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2014 (First primaries only *)

| Age at | | | | Prop. all | |
|----------------|---------|-----------|----------|-----------|--|
| death | Cases | Age-spec. | | cancers | |
| Years | n | mortality | MI-index | equicer 5 | |
| ICALS | 11 | morearrey | MI INGEX | 0 | |
| 0- 4 | | 0.0 | | | |
| 5-9 | | 0.0 | | | |
| 10-14 | | 0.0 | | | |
| 15-19 | | 0.0 | | | |
| 20-24 | | 0.0 | | | |
| 25-29 | | 0.0 | | | |
| 30-34 | | 0.0 | | | |
| 35-39 | | 0.0 | | | |
| 40-44 | 1 | 0.0 | 0.33 | 0.2 | |
| 45-49 | 1 | 0.1 | 0.07 | 0.1 | |
| 50-54 | 2 | 0.1 | 0.20 | 0.1 | |
| 55-59 | 3 | 0.2 | 0.20 | 0.1 | |
| 60-64 | 5 | 0.5 | 0.17 | 0.1 | |
| 65-69 | 10 | 1.0 | 0.43 | 0.2 | |
| 70-74 | 8 | 0.9 | 0.24 | 0.2 | |
| 75-79 | ° 14 | 2.5 | 0.42 | 0.1 | |
| 80-84 | 14 | 2.5 | 0.42 | 0.2 | |
| 85+ | 10 | 5.2 | 1.20 | 0.2 | |
| 0.5- | 12 | J.Z | 1.20 | 0.5 | |
| All area | 66 | | | 0.2 | |
| All ages | 00 | | | 0.2 | |
| Montolity | | | | | |
| Mortality | | 0.4 | 0.35 | | |
| Raw | | | | | |
| WS | | 0.2 | 0.30 | | |
| ES | | 0.3 | 0.33 | | |
| BRD-S | | 0.4 | 0.36 | | |
| DVII 70 | | | | | |
| PYLL-70 | | 1 0 | | | |
| per 100,000 | | 1.2 | | | |
| ES DVIII 70 | | 1.0 | | | |
| AYLL-70 | | 8.4 | | | |
| | | | | | |

* See corresponding tables with multiple primaries.

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2014 (Single primaries only *)

| Age at | | | | Prop. all | |
|---------------|-------|-----------|----------|-----------|--|
| death | Cases | Age-spec. | | cancers | |
| Years | n | mortality | MI-index | 00 | |
| | | | | | |
| 0- 4 | | 0.0 | | | |
| 5-9 | | 0.0 | | | |
| 10-14 | | 0.0 | | | |
| 15-19 | | 0.0 | | | |
| 20-24 | | 0.0 | | | |
| 25-29 | | 0.0 | | | |
| 30-34 | | 0.0 | | | |
| 35-39 | | 0.0 | | | |
| 40-44 | 1 | 0.1 | 0.33 | 0.3 | |
| 45-49 | 1 | 0.1 | 0.07 | 0.1 | |
| 50-54 | 2 | 0.2 | 0.20 | 0.1 | |
| 55-59 | 3 | 0.3 | 0.21 | 0.1 | |
| 60-64 | 3 | 0.3 | 0.12 | 0.1 | |
| 65-69 | 8 | 0.8 | 0.42 | 0.2 | |
| 70-74 | 7 | 0.8 | 0.24 | 0.1 | |
| 75-79 | 9 | 1.6 | 0.29 | 0.2 | |
| 80-84 | 4 | 1.1 | 0.22 | 0.1 | |
| 85+ | 9 | 3.9 | 1.29 | 0.3 | |
| 0.5 1 | | 5.5 | 1.29 | 0.5 | |
| All ages | 47 | | | 0.1 | |
| AII ayes | · | | | 0.1 | |
| Mortality | | | | | |
| Raw | | 0.3 | 0.27 | | |
| WS | | 0.1 | 0.27 | | |
| ES | | 0.1 | 0.23 | | |
| BRD-S | | 0.2 | 0.27 | | |
| BRD-3 | | 0.5 | 0.27 | | |
| PYLL-70 | | | | | |
| | | 1 0 | | | |
| per 100,000 | | 1.0 | | | |
| ES NVLL 70 | | 0.9 | | | |
| AYLL-70 | | 9.2 | | | |
| | | | | | |

* See corresponding tables with multiple primaries.





Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at breast cancer (men)-related death (see Table 10) should be considered.






Average mortality (world standard population) 2007 - 2014

Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2007 to 2014. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (0.2/100,000 WS N=88).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,886 male residents (averaged) in the period from 2007 to 2014 a total of 1 men died from breast cancer (men). Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.1/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 0.6/100,000.



Standardized mortality ratio (SMR) 2007 - 2014

Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (N=88).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 63,358 male residents (averaged) in the period from 2007 to 2014 a total of 1 men died from breast cancer (men). Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.40. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 3.00, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases, where applicable. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

| FRG GEKID | Federal Republic of Germany Association of Population-based Cancer Registries in Germany (Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.) |
|--------------|---|
| MCR | Munich Cancer Registry (Tumorregister München) |
| SEER | Surveillance, Epidemiology, and End Results (USA) |
| | |
| AYLL-70 | Average years of life lost prior to age 70 given a person dies before that age |
| BRD-S | German standard population |
| DCO | Death certificate only |
| EAR | Excess absolute risk |
| | = excess cancer cases (O - E) per 10,000 person-years |
| ES | European standard population (old) |
| LCL | Lower confidence limit |
| MI-index | Ratio between mortality and incidence |
| PYLL-70 | Potential years of life lost prior to age 70 given a person dies before that age |
| SIR | Standardized incidence ratio |
| SMR | Standardized mortality ratio |
| UCL | Upper confidence limit |
| WS | World standard population |
| | |

Recommended Citation

Munich Cancer Registry. ICD-10 C50: Breast cancer (men) - Incidence and Mortality [Internet]. 2016 [updated 2016 Apr 13; cited 2016 Jun 1]. Available from: http://www.tumorregistermuenchen.de/en/facts/base/bC50m_E-ICD-10-C50-Breast-cancer-men-incidence-and-mortality.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

| Fig./Tb | l. | Page |
|---------|---|------|
| 1 | Pts cohorts, DCO, mult. prim., follow-up / yr | 3 |
| 2 | Incidence by year of diagnosis | 4 |
| 3 | Age distribution parameters by year of diagnosis | 5 |
| 4 | Age distribution by 5-year age group | 6 |
| 5 | Age-specific incidence, DCO rate, proportion malignancies | 7 |
| 6 | Age distribution and age-specific incidence (chart) | 8 |
| 6a | Age-specific incidence internationally (chart) | 9 |
| 7 | Cumulative follow-up years (chart) | 10 |
| 8 | Standardized incidence ratio of second primaries | 11 |
| 9a | Map of cancer incidence (WS) by county (chart) | 12 |
| 9b | Standardized incidence ratio (SIR) by county (chart) | 13 |
| 10a | Pts incident cohorts and mortality / yr | 14 |
| 10b | Incidence and mortality by year of diagnosis | 15 |
| 10c | Cancer-related deaths, death certification available / yr | 16 |
| 11 | Medians of age at death / yr | 17 |
| 12 | Mortality by year of death | 18 |
| 13 | Distribution of age at death | 19 |
| 14 | Age-specific mortality | 20 |
| 15 | Multiple primaries in deaths | 21 |
| 16 | Age-specific mortality (first primaries) | 22 |
| 17 | Age-specific mortality (single primaries) | 23 |
| 18 | Age distribution and age-specific mortality (chart) | 24 |
| 19a | Map of cancer mortality (WS) by county (chart) | 25 |
| 19b | Standardized mortality ratio (SMR) by county (chart) | 26 |