

Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ Deutsch

ICD-10 C64-C68: Urinary tract cancer

Incidence and Mortality

Year of diagnosis	1998-2014
Patients	20,431
Diseases	21,231
Creation date	04/13/2016
Export date	12/23/2015
Population	4.64 m



Munich Cancer Registry at Munich Cancer Center
Marchioninstr. 15
Munich, 81377
Germany

<http://www.tumorregister-muenchen.de/en>

<http://www.tumorregister-muenchen.de/en/facts/base/bC6468E-ICD-10-C64-C68-Urinary-tract-cancer-incidence-and-mortality.pdf>

Global Statements about the statistics on the Internet –
Baseline Statistics (grey button) , **Survival** (red button)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.64 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases^{###} are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, April 2016

[#] Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007).

^{##} Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.

^{###} DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate.

ICD-10 codes (ICD-10 2015) used for specifying cancer site

Code	Description
C64	Malignant neoplasm of kidney, except renal pelvis
C65	Malignant neoplasm of renal pelvis
C66	Malignant neoplasm of ureter
C67.-	Malignant neoplasm of bladder
C68.-	Malignant neoplasm of other and unspecified urinary organs

INCIDENCE

Table 1

All patients with invasive cancer by year of diagnosis, proportions of DCO, multiple primaries, deaths, and active follow-up (incl. DCO)

Year of diagnosis	Cases n	DCO cases n	Prop. DCO %	Prop. mult. primaries %	Prop. deaths %	Prop. actively followed %
1998	756	59	7.8	33.3	69.4	97.9
1999	724	53	7.3	34.4	69.3	97.8
2000	698	69	9.9	35.2	68.6	97.6
2001	713	66	9.3	35.5	67.7	97.9
2002	1327	181	13.6	37.8	73.2	98.2 #
2003	1301	150	11.5	35.4	67.2	96.9
2004	1292	147	11.4	38.5	63.2	97.4
2005	1316	96	7.3	38.7	57.0	95.7
2006	1350	98	7.3	37.7	59.9	92.5
2007	1473	123	8.4	37.1	55.5	80.7 #
2008	1559	132	8.5	38.9	53.2	73.0
2009	1568	127	8.1	41.0	52.4	72.1
2010	1561	124	7.9	40.4	48.4	70.1
2011	1513	105	6.9	39.1	47.1	71.7
2012	1522	112	7.4	39.8	42.6	71.4
2013	1438	119	8.3	40.9	36.0	99.0
2014	1120	109	9.7	33.8	21.5	97.9 ##
1998–2014	21231	1870	8.8	38.0	54.4	86.7

The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be found in the respective headings.

Table 1a

All patients with invasive cancer
by year of diagnosis and gender
(incl. DCO)

Year of diagnosis	All n	Males n	Females n	Prop. males %
1998	756	491	265	64.9
1999	724	487	237	67.3
2000	698	462	236	66.2
2001	713	441	272	61.9
2002	1327	866	461	65.3
2003	1301	877	424	67.4
2004	1292	857	435	66.3
2005	1316	871	445	66.2
2006	1350	896	454	66.4
2007	1473	988	485	67.1
2008	1559	1050	509	67.4
2009	1568	1052	516	67.1
2010	1561	1043	518	66.8
2011	1513	1020	493	67.4
2012	1522	1055	467	69.3
2013	1438	984	454	68.4
2014	1120	747	373	66.7
1998–2014	21231	14187	7044	66.8

Table 2

Incidence measures by year of diagnosis including DCO cases
(with respect to registry area expansion from 2.51 to 3.96 m as of 2002,
and from 3.96 to 4.64 m as of 2007, respectively)

Year of diagnosis	Males n	Females n	Males Inc. raw	Fem. Inc. raw	Males Inc. WS	Fem. Inc. WS	Males Inc. ES	Fem. Inc. ES	Males Inc. BRD-S	Fem. Inc. BRD-S
1998	491	265	44.3	22.5	27.3	10.1	40.5	14.9	52.5	19.0
1999	487	237	43.5	20.0	26.2	9.5	39.0	13.7	50.4	17.2
2000	462	236	40.6	19.6	24.3	8.3	36.2	12.5	47.5	16.3
2001	441	272	38.1	22.4	22.5	9.5	33.5	14.4	42.8	18.6
2002	866	461	46.5	23.5	25.5	9.9	39.2	14.7	52.3	19.1
2003	877	424	46.8	21.5	25.9	8.8	38.7	13.1	50.9	17.1
2004	857	435	45.6	22.0	24.9	8.9	37.1	13.5	48.6	17.7
2005	871	445	46.0	22.4	24.6	9.4	36.8	13.7	47.3	17.9
2006	896	454	46.8	22.6	24.7	10.0	36.8	14.3	48.2	18.1
2007	988	485	44.6	21.0	23.3	8.6	34.7	12.6	44.7	16.6
2008	1050	509	47.2	21.9	24.1	9.4	36.1	13.6	46.7	17.5
2009	1052	516	47.1	22.2	23.8	9.1	35.4	13.2	46.4	17.4
2010	1043	518	46.3	22.1	22.8	8.0	34.0	12.4	44.0	16.4
2011	1020	493	44.6	20.9	22.0	9.2	32.7	12.9	42.1	16.1
2012	1055	467	46.2	19.8	22.0	7.5	33.3	11.4	44.1	15.4
2013	984	454	43.1	19.2	20.5	7.8	31.0	11.4	40.8	14.7
2014	747	373	32.7	15.8	15.7	6.4	23.8	9.5	31.2	12.4
1998-2014	14187	7044	44.3	21.1	23.1	8.7	34.6	12.8	45.0	16.7

The computation of the incidence measures includes all primaries, irrespective of first or subsequent malignancy.

Table 3

Age distribution parameters by year of diagnosis (All patients)
(incl. DCO)

Year of diagnosis	Cases n	Mean	Std. dev.	Min.	Max.	10%	25%	Median 50%	75%	90%
1998	756	68.6	13.6	2.8	99.7	51.6	61.0	69.9	77.9	85.3
1999	724	67.9	12.7	1.1	94.3	52.6	59.5	68.3	77.3	84.1
2000	698	69.2	12.6	0.3	99.7	53.6	61.4	70.0	78.2	85.2
2001	713	69.2	12.3	1.9	96.4	53.5	61.5	69.4	78.3	85.1
2002	1327	70.9	12.4	0.1	99.5	55.2	63.5	72.2	79.6	86.0
2003	1301	70.4	13.0	0.4	103	54.5	63.2	71.3	79.3	85.4
2004	1292	70.1	13.1	0.0	99.0	54.2	62.7	71.2	79.2	85.1
2005	1316	69.9	12.8	0.7	101	54.6	62.6	70.7	79.1	84.6
2006	1350	69.9	13.5	0.2	101	53.5	63.1	71.2	78.8	85.1
2007	1473	70.1	13.4	1.2	101	53.4	63.9	71.3	79.3	85.3
2008	1559	70.3	13.2	0.2	100	53.0	63.4	71.5	79.6	85.7
2009	1568	70.5	13.3	0.5	103	53.7	63.5	71.8	79.8	85.1
2010	1561	71.1	12.9	5.4	100	53.7	63.4	72.3	80.6	86.5
2011	1513	70.6	13.9	0.5	97.6	53.3	63.3	72.1	79.9	86.5
2012	1522	71.4	12.4	1.4	103	54.9	64.5	72.9	80.1	85.0
2013	1438	71.3	13.0	0.3	101	54.4	64.7	72.5	80.5	86.0
2014	1120	71.0	12.5	1.2	107	54.7	63.3	72.7	79.6	85.8
1998-2014	21231	70.3	13.0	0.0	107	53.9	63.0	71.6	79.4	85.5

Table 3a

Age distribution parameters by year of diagnosis (MALES)
(incl. DCO)

Year of diagnosis	Cases n	Mean	Std. dev.	Min.	Max.	10%	25%	Median 50%	75%	90%
1998	491	67.5	13.5	5.0	95.6	50.4	59.9	68.9	76.6	84.3
1999	487	67.1	12.3	2.3	94.1	52.6	59.4	66.9	76.0	82.7
2000	462	67.6	12.7	0.3	99.7	52.3	60.7	68.1	76.6	81.6
2001	441	67.5	11.3	1.9	95.1	53.3	60.7	67.1	75.6	81.4
2002	866	69.9	12.0	0.1	97.6	54.7	62.8	71.0	78.2	83.8
2003	877	69.0	12.8	0.4	101	53.3	62.1	69.7	77.6	83.6
2004	857	68.7	12.9	0.0	98.8	53.5	61.5	69.7	77.6	82.8
2005	871	68.7	11.7	0.7	101	54.6	61.7	68.9	76.9	82.9
2006	896	69.2	12.6	0.8	101	53.9	62.6	70.0	77.9	83.6
2007	988	69.0	12.7	1.3	101	53.2	62.7	70.1	77.3	83.7
2008	1050	69.7	12.8	0.2	100	52.5	62.8	70.7	78.5	85.0
2009	1052	69.6	12.8	0.5	97.4	53.3	62.6	70.8	78.4	84.1
2010	1043	69.3	12.6	5.4	99.1	52.4	61.2	70.8	78.3	84.1
2011	1020	70.2	12.8	1.5	96.9	52.7	62.7	71.4	79.0	85.9
2012	1055	70.4	12.5	1.4	103	53.8	62.9	72.3	79.1	84.4
2013	984	70.7	12.4	0.9	98.6	54.4	64.1	71.9	79.5	84.9
2014	747	70.4	12.5	1.2	97.0	54.6	62.4	72.0	79.3	84.9
1998-2014	14187	69.3	12.6	0.0	103	53.3	62.1	70.4	78.0	84.1

Table 3b

Age distribution parameters by year of diagnosis (FEMALES)
(incl. DCO)

Year of diagnosis	Cases n	Mean	Std. dev.	Min.	Max.	10%	25%	Median 50%	75%	90%
1998	265	70.6	13.5	2.8	99.7	55.0	62.6	72.4	79.2	86.2
1999	237	69.4	13.5	1.1	94.3	52.6	60.8	71.6	78.7	85.6
2000	236	72.2	11.9	37.2	94.5	58.7	63.4	73.7	81.0	87.4
2001	272	71.9	13.4	30.6	96.4	54.2	64.1	73.7	81.2	88.2
2002	461	72.8	12.8	2.4	99.5	57.9	65.3	74.1	81.9	87.7
2003	424	73.3	13.0	2.5	103	57.3	66.0	75.3	82.6	87.9
2004	435	72.9	13.1	18.5	99.0	56.5	64.9	74.8	82.4	88.3
2005	445	72.2	14.6	4.2	98.8	54.0	64.4	74.8	81.9	88.5
2006	454	71.4	15.1	0.2	96.7	52.5	64.6	74.0	81.8	87.5
2007	485	72.4	14.7	1.2	99.1	55.6	67.1	74.7	82.3	87.1
2008	509	71.8	13.9	0.6	97.0	55.7	64.5	73.7	81.9	86.9
2009	516	72.3	14.0	1.7	103	55.5	66.1	74.3	82.1	87.0
2010	518	74.6	12.8	5.4	100	56.4	68.4	75.8	84.3	89.6
2011	493	71.2	16.0	0.5	97.6	53.6	64.7	73.8	81.5	88.0
2012	467	73.6	11.8	9.7	96.4	58.2	67.4	75.5	82.1	86.7
2013	454	72.4	14.3	0.3	101	54.8	66.0	74.3	81.6	88.1
2014	373	72.3	12.5	25.7	107	55.6	64.7	74.3	81.1	87.6
1998–2014	7044	72.3	13.7	0.2	107	55.5	65.2	74.2	81.8	87.7

Table 4

Age distribution by 5-year age group and gender for period 2007–2014
(incl. DCO)

Age at diagnosis Years	Cases			Males			Females		
	n	%	Cum.%	n	%	Cum.%	n	%	Cum.%
0–4	43	0.4	0.4	21	0.3	0.3	22	0.6	0.6
5–9	16	0.1	0.5	8	0.1	0.4	8	0.2	0.8
10–14	4	0.0	0.5	2	0.0	0.4	2	0.1	0.8
15–19	2	0.0	0.6	1	0.0	0.4	1	0.0	0.9
20–24	7	0.1	0.6	3	0.0	0.4	4	0.1	1.0
25–29	13	0.1	0.7	8	0.1	0.5	5	0.1	1.1
30–34	36	0.3	1.0	21	0.3	0.8	15	0.4	1.5
35–39	97	0.8	1.9	68	0.9	1.7	29	0.8	2.3
40–44	182	1.5	3.4	127	1.6	3.3	55	1.4	3.7
45–49	346	2.9	6.3	278	3.5	6.8	68	1.8	5.5
50–54	572	4.9	11.2	430	5.4	12.2	142	3.7	9.2
55–59	825	7.0	18.2	600	7.6	19.7	225	5.9	15.1
60–64	1086	9.2	27.5	795	10.0	29.8	291	7.6	22.7
65–69	1738	14.8	42.3	1229	15.5	45.2	509	13.3	36.1
70–74	2040	17.4	59.6	1445	18.2	63.4	595	15.6	51.7
75–79	1859	15.8	75.4	1211	15.3	78.7	648	17.0	68.7
80–84	1537	13.1	88.5	963	12.1	90.8	574	15.0	83.7
85+	1351	11.5	100.0	729	9.2	100.0	622	16.3	100.0
All ages	11754	100.0		7939	100.0		3815	100.0	

Included in the statistics are 54.2% multiple primaries in males and 38.1% in females.

Table 5

Age-specific incidence, DCO rate and proportion of all cancers
for period 2007-2014

Age at diagnosis Years	Males n	Females n	Males Age- spec. incid. %	Females Age- spec. incid. %	Males DCO rate n=505 %	Females DCO rate n=423 %	Males Prop.all cancers n=91183 %	Females Prop.all cancers n=89596 %
0- 4	19	22	2.2	2.7			10.7	15.9
5- 9	8	8	0.9	1.0			8.3	10.3
10-14	2	2	0.2	0.2			2.0	2.2
15-19	1	1	0.1	0.1			0.5	0.6
20-24	3	4	0.3	0.4			0.8	1.3
25-29	8	5	0.7	0.4			1.4	0.8
30-34	21	15	1.7	1.2			2.7	1.3
35-39	68	29	5.2	2.3			5.9	1.5
40-44	124	54	7.6	3.5	0.8		6.8	1.4
45-49	273	68	17.3	4.5	0.4	1.5	8.5	1.2
50-54	418	142	32.3	11.1	1.7	0.7	8.6	2.1
55-59	582	222	54.8	19.8	1.9	1.8	7.9	3.0
60-64	779	288	79.3	27.2	1.9	2.4	7.2	3.1
65-69	1198	495	124.5	47.4	2.8	2.4	7.7	4.3
70-74	1409	582	154.9	55.7	3.7	4.5	8.3	4.9
75-79	1188	634	215.7	88.9	7.5	6.2	9.5	6.3
80-84	938	559	268.5	99.7	11.7	16.5	10.9	6.4
85+	722	617	311.8	106.8	25.6	39.1	11.8	6.0
All ages	7761	3747			6.5	11.3	8.5	4.2
Incidence								
Raw			43.0	20.0				
WS			21.2	8.1				
ES			31.8	11.9				
BRD-S			41.4	15.5				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).




Figure 6. Age distribution and age-specific incidence




Figure 6a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).

Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2014, based on the November 2013 submission. <http://www.seer.cancer.gov>.




Figure 7. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.

Table 8a

Standardized incidence ratio (SIR, with 95% confidence limits),
excess absolute risk (EAR) and DCO rate of second primaries
for period 1998–2014

MALES

Diagnosis	Observed n	Expected n	SIR	LCL 95%	UCL 95%	EAR	DCO %
C03-C06 Oral cavity	11	5.2	2.1	1.1	3.8 #	1.5	9.1
C09-C10 Oropharynx	17	6.3	2.7	1.6	4.3 #	2.8	
C12-C13 Hypopharynx	9	3.5	2.6	1.2	4.9 #	1.5	
C15 Oesophagus	28	11.9	2.4	1.6	3.4 #	4.3	14.3
C16 Stomach	54	29.7	1.8	1.4	2.4 #	6.4	7.4
C17 Small intestine	13	3.4	3.8	2.0	6.5 #	2.5	
C18 Colon	174	70.7	2.5	2.1	2.9 #	27.4	6.9
C19-C20 Rectum	65	37.4	1.7	1.3	2.2 #	7.3	3.1
C21 Anus/canal	4	1.4	2.8	0.8	7.3	0.7	25.0
C22 Liver	43	18.9	2.3	1.6	3.1 #	6.4	14.0
C23-C24 Bile	9	6.9	1.3	0.6	2.5	0.5	22.2
C25 Pancreas	58	25.7	2.3	1.7	2.9 #	8.6	25.9
C32 Larynx	14	6.6	2.1	1.2	3.5 #	2.0	
C33-C34 Lung	306	81.4	3.8	3.4	4.2 #	59.6	13.1
C38,C45 Mesothelioma	4	4.6	0.9	0.2	2.2	-0.2	50.0
C43 Malign. melanoma	69	28.0	2.5	1.9	3.1 #	10.9	1.4
C46,C49 Soft tissue	13	3.7	3.5	1.9	6.0 #	2.5	
C48 Peritoneal	5	0.5	10.1	3.3	23.7 #	1.2	40.0
C60 Penis	5	1.6	3.1	1.0	7.3 #	0.9	
C61 Prostate	1269	205.6	6.2	5.8	6.5 #	282.0	4.4
C62 Testis	7	1.4	5.0	2.0	10.4 #	1.5	
C64 Kidney	208	23.8	8.8	7.6	10.0 #	48.9	11.1
C65 Renal pelvis	79	3.0	26.2	20.7	32.6 #	20.2	
C66 Ureter	65	1.7	38.0	29.3	48.4 #	16.8	
C67 Bladder	147	33.0	4.5	3.8	5.2 #	30.2	8.8
C68 Urethra	49	0.5	93.5	69.2	123.6 #	12.9	
C68 Urinary org.	6	0.4	13.7	5.0	29.9 #	1.5	83.3
C70-C72 CNS cancer	17	8.9	1.9	1.1	3.1 #	2.2	11.8
C73 Thyroid	13	4.0	3.2	1.7	5.5 #	2.4	7.7
C74-C80 Cancer others	3	2.1	1.5	0.3	4.3	0.3	
C76-C79 CUP	27	12.1	2.2	1.5	3.3 #	4.0	11.1
C81 Hodgkin lymphoma	3	1.4	2.2	0.4	6.3	0.4	
C82-C85 NHL	76	28.3	2.7	2.1	3.4 #	12.7	7.9
C90 Mult. myeloma	19	9.1	2.1	1.3	3.3 #	2.6	10.5
C91-C96 Leukaemia	25	11.9	2.1	1.4	3.1 #	3.5	28.0
Other primaries	13	7.8	1.7	0.9	2.9	1.4	15.4
Not observed	0	1.6	0.0	0.0	2.3	-0.4	
All mult. primaries	2927	704.0	4.2	4.0	4.3 #	589.6	7.2

Patients 12676

Median age at second malignancy (years) 72.1

Person-years 37707

Mean observation time (years) 3.0

Median observation time (years) 1.5

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 to 2 are pooled in category "Other primaries"

Table 8b

Standardized incidence ratio (SIR, with 95% confidence limits),
excess absolute risk (EAR) and DCO rate of second primaries
for period 1998-2014

FEMALES

Diagnosis	Observed n	Expected n	SIR	LCL 95%	UCL 95%	EAR	DCO %
C03-C06 Oral cavity	2	1.3	1.5	0.2	5.6	0.4	
C15 Oesophagus	6	1.3	4.5	1.7	9.9 #	2.6	16.7
C16 Stomach	19	9.2	2.1	1.2	3.2 #	5.4	5.3
C17 Small intestine	5	1.1	4.8	1.5	11.1 #	2.2	
C18 Colon	52	25.3	2.1	1.5	2.7 #	14.8	1.9
C19-C20 Rectum	24	10.4	2.3	1.5	3.4 #	7.5	8.3
C21 Anus/canal	2	1.2	1.7	0.2	6.1	0.5	
C22 Liver	6	2.9	2.1	0.8	4.6	1.7	16.7
C23-C24 Bile	12	3.7	3.2	1.7	5.7 #	4.6	25.0
C25 Pancreas	34	11.2	3.0	2.1	4.2 #	12.6	32.4
C26 GI cancer	2	0.5	4.0	0.5	14.4	0.8	
C32 Larynx	2	0.4	5.2	0.6	18.9	0.9	
C33-C34 Lung	71	16.2	4.4	3.4	5.5 #	30.4	15.5
C43 Malign. melanoma	13	7.8	1.7	0.9	2.8	2.9	15.4
C46,C49 Soft tissue	2	1.3	1.5	0.2	5.5	0.4	
C48 Peritoneal	2	0.8	2.4	0.3	8.5	0.6	
C50 Breast	142	65.0	2.2	1.8	2.6 #	42.6	9.2
C51 Vulva	5	2.5	2.0	0.6	4.6	1.4	20.0
C52 Vagina	2	0.5	4.2	0.5	15.3	0.8	
C53 Cervix uteri	17	2.7	6.3	3.7	10.1 #	7.9	5.9
C54 Corpus uteri	23	12.6	1.8	1.2	2.7 #	5.7	8.7
C55,C57 Fem. genitals un	4	0.6	6.2	1.7	15.8 #	1.9	25.0
C56 Ovary	16	9.5	1.7	1.0	2.7	3.6	25.0
C64 Kidney	85	6.0	14.1	11.3	17.4 #	43.7	15.3
C65 Renal pelvis	36	0.8	46.2	32.3	63.9 #	19.5	
C66 Ureter	30	0.4	74.3	50.1	106.0 #	16.4	
C67 Bladder	78	4.9	15.9	12.6	19.9 #	40.5	11.5
C68 Urethra	3	0.1	40.2	8.3	117.4 #	1.6	
C68 Urinary org.	4	0.1	39.3	10.7	100.7 #	2.2	75.0
C70-C72 CNS cancer	5	3.2	1.6	0.5	3.7	1.0	40.0
C73 Thyroid	20	3.2	6.2	3.8	9.6 #	9.3	5.0
C76-C79 CUP	12	4.6	2.6	1.3	4.6 #	4.1	8.3
C82-C85 NHL	28	9.4	3.0	2.0	4.3 #	10.3	14.3
C90 Mult. myeloma	5	3.1	1.6	0.5	3.8	1.1	
C91-C96 Leukaemia	15	4.0	3.8	2.1	6.2 #	6.1	20.0
Other primaries	5	2.3	2.2	0.7	5.1	1.5	40.0
Not observed	0	2.4	0.0	0.0	1.5	-1.4	
All mult. primaries	789	232.4	3.4	3.2	3.6 #	308.1	11.8

Patients 6063

Median age at second malignancy (years) 74.8

Person-years 18066


Mean observation time (years) 3.0

Median observation time (years) 1.3

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries"

Average incidence (world standard population) 2007 - 2014: Males



Average incidence (world standard population) 2007 - 2014: Females





Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 21.2/100,000 WS N=7,761, females 8.1/100,000 WS N=3,747).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 65,347 female residents (averaged) in the period from 2007 to 2014 a total of 111 women were identified with newly diagnosed urinary tract cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 8.5/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 6.3 and 11.5/100,000.

Standardized incidence ratio (SIR) 2007 - 2014: Males



Standardized incidence ratio (SIR) 2007 - 2014: Females




Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=7,761, females N=3,747).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,924 female residents (averaged) in the period from 2007 to 2014 a total of 111 women were identified with newly diagnosed urinary tract cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 1.12. Though, the value of this parameter may vary with an underlying probability of 99% between 0.87 and 1.43, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

Year of diagnosis	Incident cases n	Prop. actively followed %	Prop. DCO %	Deaths n	Prop. deaths %	Prop. deaths with death certific. %
1998	756	97.9	7.8	525	69.4	94.1
1999	724	97.8	7.3	502	69.3	95.2
2000	698	97.6	9.9	479	68.6	96.7
2001	713	97.9	9.3	483	67.7	96.7
2002	1327	98.2	13.6	972	73.2	97.1
2003	1301	96.9	11.5	874	67.2	98.2
2004	1292	97.4	11.4	816	63.2	98.0
2005	1316	95.7	7.3	750	57.0	97.7
2006	1350	92.5	7.3	809	59.9	99.0
2007	1473	80.7	8.4	817	55.5	98.4
2008	1559	73.0	8.5	829	53.2	99.3
2009	1568	72.1	8.1	822	52.4	98.9
2010	1561	70.1	7.9	755	48.4	98.9
2011	1513	71.7	6.9	713	47.1	97.8
2012	1522	71.4	7.4	649	42.6	97.8
2013	1438	99.0	8.3	517	36.0	97.5
2014	1120	97.9	9.7	241	21.5	92.1
1998-2014	21231	86.7	8.8	11553	54.4	97.7

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis
(incl. DCO)
(with respect to registry area expansion from 2.51 to 3.96 m as of 2002,
and from 3.96 to 4.64 m as of 2007, respectively)

Year of diagnosis/ death	Incident cases n	Deaths n	Prop. deaths with death certific. %	Deaths in same year n	Prop. deaths in same year %
1998	756	478	92.5	132	17.5
1999	724	456	95.2	141	19.5
2000	698	464	95.0	130	18.6
2001	713	472	96.0	123	17.3
2002	1327	694	96.7	301	22.7
2003	1301	804	96.8	290	22.3
2004	1292	786	96.9	265	20.5
2005	1316	772	96.5	218	16.6
2006	1350	821	97.6	237	17.6
2007	1473	937	97.8	272	18.5
2008	1559	933	98.8	277	17.8
2009	1568	1016	99.2	324	20.7
2010	1561	1056	98.7	295	18.9
2011	1513	1011	98.4	283	18.7
2012	1522	1092	98.4	299	19.6
2013	1438	1078	99.0	284	19.7
2014	1120	995	98.8	205	18.3
1998-2014	21231	13865	97.6	4076	19.2

Table 10c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates

(incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

Year of death	Deaths n	Prop. cancer- related %	Prop. non-cancer- related %	Prop. cancer recorded on death certificate %
1998	478	64.4	35.6	81.7
1999	456	70.2	29.8	83.2
2000	464	69.2	30.8	83.7
2001	472	69.7	30.3	85.2
2002	694	72.0	28.0	85.8
2003	804	70.9	29.1	85.1
2004	786	70.7	29.3	84.3
2005	772	71.8	28.2	83.5
2006	821	71.7	28.3	81.1
2007	937	73.1	26.9	82.9
2008	933	71.5	28.5	82.5
2009	1016	71.8	28.2	82.8
2010	1056	68.4	31.6	80.5
2011	1011	68.2	31.8	82.7
2012	1092	65.4	34.6	77.3
2013	1078	66.4	33.6	78.4
2014	995	65.9	34.1	77.1
1998-2014	13865	69.4	30.6	81.8

Table 11a

Medians of age at death according to the grouping in Table 10

MALES

Year of death	Deaths n	Age at death (all causes) Years	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate) Years
1998	316	75.9	74.3	77.9	75.4
1999	299	76.5	74.3	81.3	75.4
2000	305	76.6	73.5	80.1	76.0
2001	308	75.7	73.5	80.2	74.6
2002	437	76.3	75.1	78.4	75.8
2003	544	76.0	74.9	80.3	75.7
2004	502	76.5	75.6	79.9	76.3
2005	505	76.7	75.4	80.0	75.9
2006	530	75.6	74.1	79.2	75.0
2007	616	76.9	75.4	80.0	75.8
2008	628	77.0	75.5	80.3	76.1
2009	686	77.3	74.9	81.7	75.9
2010	686	77.7	76.1	81.5	77.1
2011	679	76.9	74.9	81.9	76.2
2012	716	78.6	77.1	81.8	77.7
2013	727	78.3	76.5	81.2	77.4
2014	676	78.8	76.1	83.7	77.2
1998-2014	9160	77.1	75.4	80.7	76.3

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 11b

Medians of age at death according to the grouping in Table 10
FEMALES

Year of death	Deaths n	Age at death (all causes) Years	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate) Years
1998	162	81.7	79.4	82.7	81.7
1999	157	78.9	78.5	80.0	79.2
2000	159	78.7	78.7	80.4	79.1
2001	164	80.7	79.0	87.3	80.3
2002	257	79.8	78.5	83.3	79.4
2003	260	80.0	79.1	80.7	79.9
2004	284	81.1	80.0	83.6	80.8
2005	267	81.3	79.1	83.5	80.6
2006	291	81.1	79.7	83.9	80.0
2007	321	80.8	79.7	84.0	80.6
2008	305	80.8	79.0	85.4	80.0
2009	330	80.8	78.6	84.3	79.5
2010	370	82.2	80.2	85.8	81.2
2011	332	82.2	79.5	87.8	80.5
2012	376	81.4	78.4	85.5	79.6
2013	351	80.6	78.6	84.8	79.4
2014	319	82.3	80.0	87.2	81.0
1998-2014	4705	81.0	79.1	84.7	80.1

By 2010, life expectancy at birth was 77.5 years for boys and 82.6 years for girls.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Table 12a

Mortality measures (cancer-related death) and mortality-incidence-index
by year of death

MALES

Year of death	Deaths n	Mort. raw	MI-Index raw	Mort. WS	MI-Index WS	Mort. ES	MI-Index ES	Mort. BRD-S	MI-Index BRD-S
1998	204	18.4	0.42	10.3	0.38	17.0	0.42	24.1	0.46
1999	204	18.2	0.42	10.2	0.39	16.7	0.43	24.2	0.48
2000	204	17.9	0.45	9.9	0.42	16.1	0.45	23.0	0.49
2001	217	18.7	0.50	10.3	0.46	16.5	0.50	23.2	0.55
2002	314	16.9	0.37	8.6	0.34	14.2	0.37	20.4	0.40
2003	383	20.4	0.44	10.5	0.41	16.8	0.44	23.9	0.48
2004	355	18.9	0.42	9.2	0.38	15.2	0.42	21.7	0.46
2005	355	18.7	0.42	8.9	0.37	14.5	0.40	21.0	0.46
2006	385	20.1	0.44	9.7	0.40	15.4	0.43	21.6	0.46
2007	458	20.7	0.48	9.6	0.42	15.7	0.46	22.2	0.51
2008	445	20.0	0.43	8.8	0.37	14.5	0.41	21.0	0.46
2009	491	22.0	0.48	9.7	0.42	15.6	0.45	22.0	0.49
2010	473	21.0	0.47	9.0	0.41	14.6	0.44	20.8	0.49
2011	476	20.8	0.47	9.0	0.42	14.4	0.45	19.8	0.48
2012	472	20.7	0.46	8.5	0.40	14.0	0.43	19.8	0.46
2013	492	21.5	0.51	8.9	0.44	14.5	0.48	20.6	0.52
2014	441	19.3	0.60	8.0	0.52	13.2	0.56	18.6	0.61
1998-2014	6369	19.9	0.46	9.2	0.41	15.0	0.44	21.2	0.48

Table 12b

Mortality measures (cancer-related death) and mortality-incidence-index
by year of death

FEMALES

Year of death	Deaths n	Mort. raw	MI-Index raw	Mort. WS	MI-Index WS	Mort. ES	MI-Index ES	Mort. BRD-S	MI-Index BRD-S
1998	105	8.9	0.40	2.9	0.29	4.9	0.33	7.1	0.37
1999	116	9.8	0.49	3.5	0.37	5.6	0.41	7.9	0.46
2000	117	9.7	0.50	3.3	0.40	5.4	0.44	7.7	0.48
2001	112	9.2	0.42	3.1	0.33	5.0	0.35	7.3	0.40
2002	186	9.5	0.41	3.3	0.34	5.3	0.36	7.3	0.39
2003	187	9.5	0.45	3.2	0.37	5.1	0.40	7.2	0.42
2004	201	10.2	0.47	3.2	0.37	5.2	0.39	7.4	0.43
2005	200	10.1	0.46	3.3	0.36	5.2	0.39	7.3	0.42
2006	204	10.2	0.46	3.3	0.34	5.3	0.38	7.4	0.42
2007	228	9.9	0.47	3.0	0.35	5.0	0.40	7.4	0.45
2008	222	9.6	0.45	3.1	0.34	5.0	0.37	7.0	0.41
2009	239	10.3	0.47	3.4	0.38	5.4	0.41	7.5	0.44
2010	250	10.7	0.49	3.2	0.41	5.2	0.43	7.6	0.47
2011	214	9.1	0.45	2.9	0.33	4.7	0.37	6.5	0.42
2012	242	10.3	0.53	3.2	0.44	5.2	0.47	7.5	0.50
2013	224	9.5	0.50	3.0	0.39	4.8	0.42	6.7	0.47
2014	215	9.1	0.59	2.7	0.44	4.5	0.48	6.4	0.53
1998-2014	3262	9.8	0.47	3.1	0.37	5.1	0.40	7.2	0.44

Table 13

Age distribution of age at death (cancer-related) for period 2007-2014
(incl. multiple primaries)

Age at death Years	Cases			Males			Females		
	n	%	Cum.%	n	%	Cum.%	n	%	Cum.%
5-9	2	0.0	0.0	2	0.0	0.0			0.0
10-14	1	0.0	0.1			0.0	1	0.1	0.1
15-19	0	0.0	0.1			0.0			0.1
20-24	2	0.0	0.1	1	0.0	0.1	1	0.1	0.1
25-29	2	0.0	0.1	1	0.0	0.1	1	0.1	0.2
30-34	2	0.0	0.2	2	0.0	0.1			0.2
35-39	8	0.1	0.3	2	0.0	0.2	6	0.3	0.5
40-44	23	0.4	0.7	12	0.3	0.5	11	0.6	1.0
45-49	72	1.2	1.9	50	1.2	1.7	22	1.1	2.2
50-54	158	2.6	4.5	114	2.8	4.6	44	2.3	4.4
55-59	263	4.4	8.9	198	4.9	9.5	65	3.3	7.8
60-64	394	6.6	15.5	296	7.3	16.8	98	5.0	12.8
65-69	632	10.6	26.1	445	11.0	27.9	187	9.6	22.5
70-74	1038	17.4	43.5	768	19.1	46.9	270	13.9	36.4
75-79	1063	17.8	61.3	721	17.9	64.8	342	17.6	54.0
80-84	1144	19.2	80.5	731	18.1	83.0	413	21.3	75.3
85+	1165	19.5	100.0	685	17.0	100.0	480	24.7	100.0
All ages	5969	100.0		4028	100.0		1941	100.0	

Included in the statistics are 54.2% multiple primaries in males and 38.1% in females.

Table 14

Age-specific mortality (cancer-related) and proportion of all cancers
for period 2007-2014
(incl. multiple primaries)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	MI-index	Females Age- spec. mortal.	MI-index	Males Prop.all cancers %	Females Prop.all cancers %
0- 4			0.0		0.0			
5- 9	2		0.2	0.25	0.0		9.5	
10-14		1	0.0		0.1	0.50		5.0
15-19			0.0		0.0			
20-24	1	1	0.1	0.33	0.1	0.25	2.1	3.6
25-29	1	1	0.1	0.13	0.1	0.20	1.6	1.6
30-34	2		0.2	0.10	0.0		2.3	
35-39	2	6	0.2	0.03	0.5	0.21	1.1	2.3
40-44	12	11	0.7	0.09	0.7	0.20	2.6	1.7
45-49	50	22	3.2	0.18	1.5	0.32	4.9	1.8
50-54	114	44	8.8	0.27	3.4	0.31	6.1	2.5
55-59	198	65	18.6	0.33	5.8	0.29	6.4	2.5
60-64	296	98	30.1	0.37	9.2	0.34	6.2	2.8
65-69	445	187	46.3	0.36	17.9	0.37	6.2	3.6
70-74	768	270	84.4	0.53	25.8	0.45	8.4	4.1
75-79	721	342	130.9	0.60	47.9	0.53	8.5	5.4
80-84	731	413	209.3	0.76	73.7	0.72	9.9	6.3
85+	685	480	295.9	0.94	83.1	0.77	11.3	5.5
All ages	4028	1941					8.1	4.4
Mortality								
Raw			22.3	0.51	10.4	0.51		
WS			9.6	0.44	3.3	0.40		
ES			15.6	0.48	5.3	0.44		
BRD-S			22.1	0.52	7.5	0.48		
PYLL-70								
per 100,000			60.0		24.8			
ES			53.1		21.0			
AYLL-70			8.6		9.0			

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15a

Multiple primaries in deaths in period 1998-2014
MALES

Diagnosis		Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C15	Oesophagus	48	1.2	11	22.9	2	4.2	35	72.9
C16	Stomach	114	2.7	37	32.5	8	7.0	69	60.5
C18	Colon	286	6.9	133	46.5	39	13.6	114	39.9
C19-C20	Rectum	139	3.3	63	45.3	18	12.9	58	41.7
C22	Liver	64	1.5	12	18.8	12	18.8	40	62.5
C25	Pancreas	84	2.0	5	6.0	9	10.7	70	83.3
C33-C34	Lung	485	11.7	73	15.1	44	9.1	368	75.9
C43	Malign. melanoma	110	2.6	65	59.1	6	5.5	39	35.5
C44	Skin others	160	3.8	75	46.9	8	5.0	77	48.1
C61	Prostate	1162	27.9	376	32.4	282	24.3	504	43.4
C64	Kidney	168	4.0			38	22.6	130	77.4
C65	Renal pelvis	147	3.5			19	12.9	128	87.1
C66	Ureter	116	2.8			25	21.6	91	78.4
C67	Bladder	359	8.6			46	12.8	313	87.2
C68	Urethra	63	1.5			14	22.2	49	77.8
C70-C72	CNS cancer	59	1.4	17	28.8	7	11.9	35	59.3
C76-C79	CUP	54	1.3	18	33.3	6	11.1	30	55.6
C82-C85	NHL	105	2.5	37	35.2	14	13.3	54	51.4
C90	Mult. myeloma	46	1.1	14	30.4	7	15.2	25	54.3
C91-C96	Leukaemia	66	1.6	12	18.2	4	6.1	50	75.8
Other primaries		327	7.9	147	45.0	21	6.4	159	48.6
All mult. primaries		4162	100.0	1095	26.3	629	15.1	2438	58.6

Multiple primaries with number of cases 1 to 40 are pooled in category "Other primaries"

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 15b

Multiple primaries in deaths in period 1998-2014
FEMALES

Diagnosis		Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C16	Stomach	42	2.7	13	31.0	9	21.4	20	47.6
C18	Colon	117	7.5	50	42.7	14	12.0	53	45.3
C19-C20	Rectum	52	3.3	24	46.2	6	11.5	22	42.3
C22	Liver	12	0.8	2	16.7	4	33.3	6	50.0
C23-C24	Bile	17	1.1			4	23.5	13	76.5
C25	Pancreas	58	3.7	2	3.4	6	10.3	50	86.2
C33-C34	Lung	123	7.9	14	11.4	19	15.4	90	73.2
C43	Malign. melanoma	37	2.4	22	59.5	2	5.4	13	35.1
C44	Skin others	41	2.6	20	48.8	2	4.9	19	46.3
C50	Breast	288	18.4	173	60.1	20	6.9	95	33.0
C51	Vulva	13	0.8	9	69.2	1	7.7	3	23.1
C53	Cervix uteri	80	5.1	61	76.3	8	10.0	11	13.8
C54	Corpus uteri	71	4.5	49	69.0	10	14.1	12	16.9
C56	Ovary	56	3.6	23	41.1	8	14.3	25	44.6
C64	Kidney	65	4.2			15	23.1	50	76.9
C65	Renal pelvis	49	3.1			12	24.5	37	75.5
C66	Ureter	42	2.7			18	42.9	24	57.1
C67	Bladder	146	9.4			17	11.6	129	88.4
C68	Urinary org.	11	0.7			1	9.1	10	90.9
C70-C72	CNS cancer	31	2.0	8	25.8	5	16.1	18	58.1
C73	Thyroid	27	1.7	13	48.1	1	3.7	13	48.1
C76-C79	CUP	28	1.8	4	14.3	2	7.1	22	78.6
C82-C85	NHL	41	2.6	15	36.6	9	22.0	17	41.5
C91-C96	Leukaemia	30	1.9	2	6.7	5	16.7	23	76.7
Other primaries		84	5.4	29	34.5	15	17.9	40	47.6
All mult. primaries		1561	100.0	533	34.1	213	13.6	815	52.2

Multiple primaries with number of cases 1 to 10 are pooled in category "Other primaries"

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers
for period 2007-2014
(First primaries only *)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	MI-index	Females Age- spec. mortal.	MI-index	Males Prop.all cancers %	Females Prop.all cancers %
0- 4			0.0		0.0			
5- 9	2		0.2	0.25	0.0		10.0	
10-14		1	0.0		0.1	0.50		5.3
15-19			0.0		0.0			
20-24	1	1	0.1	0.33	0.1	0.25	2.3	3.8
25-29	1	1	0.1	0.13	0.1	0.20	1.8	1.7
30-34	2		0.2	0.11	0.0		2.3	
35-39	2	6	0.2	0.03	0.5	0.21	1.2	2.6
40-44	11	9	0.7	0.10	0.6	0.19	2.6	1.6
45-49	47	19	3.0	0.18	1.3	0.32	5.1	1.9
50-54	80	35	6.2	0.23	2.7	0.30	5.0	2.4
55-59	148	46	13.9	0.32	4.1	0.26	5.7	2.1
60-64	229	61	23.3	0.37	5.8	0.28	5.8	2.1
65-69	313	147	32.5	0.38	14.1	0.38	5.5	3.6
70-74	496	179	54.5	0.53	17.1	0.43	7.1	3.5
75-79	458	254	83.2	0.63	35.6	0.54	7.3	5.2
80-84	437	284	125.1	0.82	50.6	0.73	8.2	5.6
85+	424	368	183.1	0.96	63.7	0.77	9.6	5.4
All ages	2651	1411					6.9	4.1
Mortality								
Raw			14.7	0.49	7.5	0.50		
WS			6.5	0.42	2.4	0.38		
ES			10.4	0.46	3.8	0.42		
BRD-S			14.5	0.51	5.4	0.46		
PYLL-70								
per 100,000			46.6		19.1			
ES			41.2		16.3			
AYLL-70			8.9		9.3			

* See corresponding tables with multiple primaries.

Table 17

Age-specific mortality (cancer-related) and proportion of all cancers
for period 2007-2014
(Single primaries only *)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	MI-index	Females Age- spec. mortal.	MI-index	Males Prop.all cancers %	Females Prop.all cancers %
0- 4			0.0		0.0			
5- 9	2		0.2	0.25	0.0		10.0	
10-14		1	0.0		0.1	0.50		5.6
15-19			0.0		0.0			
20-24	1	1	0.1	0.33	0.1	0.25	2.6	4.2
25-29	1	1	0.1	0.13	0.1	0.20	2.0	1.8
30-34	2		0.2	0.11	0.0		2.4	
35-39	2	4	0.2	0.03	0.3	0.14	1.3	2.0
40-44	11	7	0.7	0.10	0.5	0.15	2.8	1.4
45-49	41	16	2.6	0.17	1.1	0.29	4.8	1.8
50-54	66	32	5.1	0.22	2.5	0.29	4.6	2.4
55-59	111	38	10.5	0.28	3.4	0.23	4.8	2.0
60-64	177	47	18.0	0.34	4.4	0.24	5.2	1.9
65-69	206	124	21.4	0.31	11.9	0.35	4.3	3.7
70-74	322	131	35.4	0.43	12.5	0.35	5.7	3.2
75-79	286	194	51.9	0.48	27.2	0.45	5.9	4.9
80-84	282	225	80.7	0.62	40.1	0.63	6.9	5.5
85+	299	298	129.1	0.73	51.6	0.65	8.8	5.3
All ages	1809	1119					5.7	3.9
Mortality								
Raw			10.0	0.40	6.0	0.43		
WS			4.6	0.35	1.9	0.32		
ES			7.2	0.38	3.1	0.36		
BRD-S			9.9	0.41	4.3	0.40		
PYLL-70								
per 100,000			37.2		16.0			
ES			33.0		13.7			
AYLL-70			9.6		9.3			

* See corresponding tables with multiple primaries.





Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at urinary tract cancer-related death (see Table 10) should be considered.

Average mortality (world standard population) 2007 - 2014: Males



Average mortality (world standard population) 2007 - 2014: Females





Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2007 to 2014. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 8.8/100,000 WS N=3,692, females 3.0/100,000 WS N=1,810).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 65,347 female residents (averaged) in the period from 2007 to 2014 a total of 60 women died from urinary tract cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 3.6/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 2.3 and 5.9/100,000.

Standardized mortality ratio (SMR) 2007 - 2014: Males



Standardized mortality ratio (SMR) 2007 - 2014: Females




Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=3,692, females N=1,810).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,924 female residents (averaged) in the period from 2007 to 2014 a total of 60 women died from urinary tract cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 1.29. Though, the value of this parameter may vary with an underlying probability of 99% between 0.90 and 1.78, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases, where applicable. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

FRG	Federal Republic of Germany
GEKID	Association of Population-based Cancer Registries in Germany (Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)
MCR	Munich Cancer Registry (Tumorregister München)
SEER	Surveillance, Epidemiology, and End Results (USA)
AYLL-70	Average years of life lost prior to age 70 given a person dies before that age
BRD-S	German standard population
DCO	Death certificate only
EAR	Excess absolute risk = excess cancer cases (O - E) per 10,000 person-years
ES	European standard population (old)
LCL	Lower confidence limit
MI-index	Ratio between mortality and incidence
PYLL-70	Potential years of life lost prior to age 70 given a person dies before that age
SIR	Standardized incidence ratio
SMR	Standardized mortality ratio
UCL	Upper confidence limit
WS	World standard population

Recommended Citation

Munich Cancer Registry. ICD-10 C64-C68: Urinary tract cancer - Incidence and Mortality [Internet]. 2016 [updated 2016 Apr 13; cited 2016 Jun 1]. Available from: <http://www.tumorregister-muenchen.de/en/facts/base/bC6468E-ICD-10-C64-C68-Urinary-tract-cancer-incidence-and-mortality.pdf>

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

Fig./Tbl.		Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	3
1a	Gender distribution by year of diagnosis	4
2	Incidence by year of diagnosis	5
3	Age distribution parameters by year of diagnosis	6
4	Age distribution by 5-year age group and gender	8
5	Age-specific incidence, DCO rate, proportion malignancies	9
6	Age distribution and age-specific incidence (chart)	10
6a	Age-specific incidence internationally (chart)	11
7	Cumulative follow-up years (chart)	12
8	Standardized incidence ratio of second primaries	13
9a	Map of cancer incidence (WS) by county (chart)	15
9b	Standardized incidence ratio (SIR) by county (chart)	16
10a	Pts incident cohorts and mortality / yr	17
10b	Incidence and mortality by year of diagnosis	18
10c	Cancer-related deaths, death certification available / yr	19
11	Medians of age at death / yr	20
12	Mortality by year of death	22
13	Distribution of age at death	23
14	Age-specific mortality	24
15	Multiple primaries in deaths	25
16	Age-specific mortality (first primaries)	27
17	Age-specific mortality (single primaries)	28
18	Age distribution and age-specific mortality (chart)	29
19a	Map of cancer mortality (WS) by county (chart)	30
19b	Standardized mortality ratio (SMR) by county (chart)	31