Munich Cancer Registry



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ Deutsch

ICD-10 C66: Ureteral cancer

Incidence and Mortality

Year of diagnosis	1998-2014
Patients	534
Diseases	541
Creation date	04/13/2016
Export date	12/23/2015
Population	4.64 m



Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

http://www.tumorregister-muenchen.de/en/facts/base/bC66__E-ICD-10-C66-Ureteral-cancer-incidence-and-mortality.pdf

Global Statements about the statistics on the Internet – Baseline Statistics (grey button ——), Survival (red button ——)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut[#], with a total of 4.64 million inhabitants, account for the frequency of cancer diseases^{##} and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases**** are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, April 2016

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007).
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate.

ICD-10 codes (ICD-10 2015) used for specifying cancer site

Code	Description
C66	Malignant neoplasm of ureter

INCIDENCE

Table 1

All patients with invasive cancer by year of diagnosis, proportions of DCO, multiple primaries, deaths, and active follow-up (incl. DCO)

				Prop.		Prop.
		DCO	Prop.	mult.	Prop.	actively
Year of	Cases	cases	DCO	primaries	deaths	followed
diagnosis	n	n	%	_ / %	왕	%
1998	19			84.2	89.5	100.0
1999	11			81.8	81.8	100.0
2000	11			63.6	63.6	100.0
2001	24			62.5	95.8	100.0
2002	30			60.0	83.3	100.0 #
2003	19			57.9	57.9	100.0
2004	24			70.8	75.0	91.7
2005	38			68.4	71.1	94.7
2006	34			76.5	73.5	94.1
2007	34			70.6	55.9	79.4 #
2008	39			69.2	53.8	71.8
2009	43			74.4	62.8	79.1
2010	48	1	2.1	81.3	62.5	81.3
2011	44			79.5	50.0	79.5
2012	42	1	2.4	73.8	45.2	81.0
2013	47			72.3	46.8	97.9
2014	34			64.7	11.8	94.1 ##
1998-2014	541	2	0.4	71.9	60.3	88.5

[#] The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

^{##} Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be found in the respective headings.

Table 1a

All patients with invasive cancer by year of diagnosis and gender (incl. DCO)

Year of	All/	Males	Females	Prop. males	
diagnosis	n/	n	n	용	
1998	19	12	7	63.2	
1999	/11	/ 7	4	63.6	
2000	/ 11	6	5	54.5	
2001	24	13	11	54.2	
2002	30	17	13	56.7	
2003	19	10	9 /	52.6	
2004	24	15	9	62.5	
2005	38	27	11	71.1	
2006	34	22	12	64.7	
2007	34	26	8	76.5	
2008	39	19	20	48.7	
2009	43	29	14	67.4	
2010	48	34	14	70.8	
2011	44	26	18	59.1	
2012	42	25	17 <	59.5	
2013	47	24	23	51.1	
2014	34	24	10	70.6	
1998-2014	541	336	205	62.1	

Table 2

Incidence measures by year of diagnosis including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

			Males	Fem.	Males	Fem.	Males	Fem.	Males	Fem.
Year of	Males	Females	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	12	7	/ 1.1/	0.6	0.7	0.3	1.0	0.4	1.4	0.5
1999	7	4	0.6	0.3	0.4	0.1	0.6	0.2	0.8	0.2
2000	6	5	0.5	0.4	0.3	0.2	0.5	0.2	0.7	0.4
2001	13	11 <	1.1	0.9	0.6	0.4	1.0	0.6	1.4	0.8
2002	17	13	0.9	0.7	0.5	0.3	0.8	0.5	1.0	0.6
2003	10	9	0.5	0.5	0.2	0.2	0.4	0.3	0.6	0.4
2004	15	9	0.8	0.5	0.4	0.2	0.6	0.3	0.9	0.4
2005	27	11	1.4	0.6	0.7	0.2	1.1	0.4	1.5	0.4
2006	22	12	1.1	0.6	0.5	0.2	0.8	0.3	1.2	0.5
2007	26	8	1.2	0.3	0.6	0.1	0.9	0.2	1.1	0.3
2008	19	20	0.9	0.9	0.5	0.3	0.7	0.5	0.9	0.7
2009	29	14	1.3	0.6	0.6	0.2	0.9	0.3	1.3	0.5
2010	34	1.4	1.5	0.6	0.6	0.3	1.0	0.4	1.5	0.5
2011	26	18	1.1	0.8	0.6	0.2	0.9	0.4	1.1	0.5
2012	25	17	1.1	0.7	0.5	0.2	0.7	0.4	1.1	0.6
2013	24	23	1.1	1.0	0.4	0.3	0.7	0.5	1.0	0.7
2014	24	10	1.1	0.4	0.4	0.2	0.7	0.3	1.1	0.3
1998-2014	336	205	1.0	0.6	0.5	0.2	0.8	0.3	1.1	0.5




Table 3

Age distribution parameters by year of diagnosis (All patients) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	19	70.5	9.2	49.0	83.3	55.6	66.5	70.1	79.0	81.7
1999	11	74.4	12.9	49.4	89.5	55.0	69.7	76.5	87.7	87.8
2000	11	71.6	10.7	45.2	81.7	62.8	67.3	73.2	80.3	80.6
2001	24	73.6	7.4	57.2	87.5	61.8	70.4	74.6	77.9	80.5
2002	30	70.0	9.9	51.7	91.1	56,6	62.3	71.5	75.3	82.7
2003	19	74.5	8.6	56.6	87.3	60.9	69.8	77.4	80.7	85.0
2004	24	72.8	11.4	46.3	94.6	54.9	69.6	74.2	79.8	84.2
2005	38	70.0	12.4	38.2	93.3	53.1	65.8	69.5	78.2	89.2
2006	34	73.3	9.2	50.5	88.7	62.3	67.7	73.8	80.4	84.6
2007	34	72.5	8.2	53.2	87.8	62.9	67.6	73.9	76.3	83.7
2008	39	72.1	9.4	49.3	87.2	61.2	64.5	73.5	79.7	84.9
2009	43	73.1	10.7	29.1	89.0	60.2	68.6	72.8	81.3	83.9
2010	48	74.0	8.8	52.4	92.7	60.0	70.0	74.0	80.4	84.6
2011	44	72.0	10.1	50.3	89.7	58.5	64.5	72.1	79.9	84.8
2012	42	75.5	9.9	49.0	92.8	63.2	68.5	77.5	82.7	83.9
2013	47	75.1	9.7	40.0	90.2	64.6	71.7	76.8	81.7	84.0
2014	34	75.3	9.9	54.0	90.8	59.6	68.5	76.6	82.0	88.0
1998-2014	541	73.1	9.9	29.1	94.6	59.5	67.6	73.9	80.2	84.6

Table 3a

Age distribution parameters by year of diagnosis (MALES)

(incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	12	69.4	9.5	49.0	81.7	55.6	67.2	69.8	76.3	80.4
1999	7	70.5	14.1	49.4	89.5	49.4	55.0	72.3	80.9	89.5
2000	6	69.3	12.5	45.2	80.3	45.2	67.7	73.1	76.6	80.3
2001	13	73.7	6.6	61.6	86.8	64.0	70.5	74.8	76.9	78.0
2002	17	69.2	9.9	52.0	91.1	55.1	62.3	70.9	74.4	80.3
2003 —	10	74.4	10.8	56.6	87.3	58.8	63.9	79.1	81.2	86.1
2004	15	73.4	9.8	54.9	86.9	55.9	69.8	74.9	80.1	84.2
2005	27	70.3	12.2	47.6	93.3	53.1	65.8	68.6	78.5	89.2
2006	22	71.7	8.1	50.5	84.3	64.5	67.0	71.2	79.7	80.5
2007	26	71.9	8.6	53.2	87.8	62.9	66.9	71.2	76.3	86.4
2008	19	69.2	10.7	49.3	87.2	51.3	61.7	67.8	78.3	85.9
2009	29	70.5	11.4	29.1	89.0	59.5	65.0	70.2	77.6	82.9
2010	34	76.3	7.6	60.0	92.7	67.4	70.4	76.8	82.1	85.1
2011	26	68.7	9.5	50.3	84.9	58.0	62.0	67.4	76.4	82.6
2012	25	73.2	9.8	49.0	89.1	61.7	67.5	75.0	82.4	83.4
2013	24	73.6	10.9	40.0	86.3	64.6	67.4	75.9	82.1	83.4
2014	24	76.4	9.9	54.0	90.8	63.6	68.6	78.7	83.2	88.4
1998-2014	336	72.1	10.0	29.1	93.3	58.7	66.5	72.8	79.6	84.0

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	7	72.2	9.1	58.5	83.3	58.5	64.7	72.4	80.2	83.3
1999	4	81.2	7.8	72.3	87.8	72.3	74.6	82.3	87.7	87.8
2000	5	74.3	8.7	62.8	81.7	62.8	67.3	79.2	80.6	81.7
2001	11	73.5	8.6	57.2	87.5	61.8	70.4	73.7	79.9	80.5
2002	13	71.1	10.1	51.7	87.0	58.6	65.3	72.7	76.0	85.0
2003	9	74.6	6.0	64.1	83.1	64.1	71.0	75.0	79.2	83.1
2004	9	71.8	14.3	46.3	94.6	46.3	69.4	73.5	79.4	94.6
2005	11	69.3	13.2	38.2	91.0	59.5	65.4	71.6	78.2	78.2
2006	12	76.3	10.8	52.1	88.7	62.3	73.1	75.8	85.4	87.4
2007	8	74.7	6.8	60.3	82.8	60.3	72.6	75.7	78.8	82.8
2008	20	74.7	7.3	63.4	85.7	64.1	68.6	74.5	81.0	84.5
2009	14	78.4	6.6	66.0	87.2	67.9	74.3	79.7	83.9	84.0
2010	14	68.5	9.4	52.4	79.8	54.6	59.4	72.4	75.6	78.5
2011	18 /	76.7	9.2	56.8	89.7	59.1	71.6	76.6	84.6	88.0
2012	17/	78.9	9.3	56.7	92.8	67.2	77.7	80.0	83.3	91.5
2013	23	76.7	8.3	52.6	90.2	65.6	74.3	79.0	81.7	84.8
2014	10	72.6	10.1	59.5	88.0	59.5	62.0	74.6	76.8	86.7
1998-2014	205	74.6	9.5	38.2	94.6	61.5	69.9	75.3	80.6	85.4

Table 4

Age distribution by 5-year age group and gender for period 2007-2014 (incl. DCO)

Age at									
diagnosis	Cases			Males			Females		
Years	n	%	Cum.%	n	엉	Cum.%	n	양	Cum.%
25-29	1	0.3	0.3	/ 1	0.5	0.5			0.0
30-34	0	0.0	0.3			0.5			0.0
35-39	1	0.3	0.6	1	0.5	1.0			0.0
40 - 44	0	0.0	0.6			1.0/			0.0
45-49	2	0.6	1.2	2	1.0	1.9			0.0
50-54	10	3.0	4.2	7	3.4	5.3	3	2.4	2.4
55-59	15	4.5	8.8	8	3.9	9.2	7	5.6	8.1
60-64	32	9.7	18.4	25	12.1	21.3	7	5.6	13.7
65-69	46	13.9	32.3	35	16.9	38.2	11	8.9	22.6
70-74	61	18.4	50.8	37	17.9	56.0	24	19.4	41.9
75-79	70	21.1	71.9	37	17.9	73.9	33	26.6	68.5
80-84	63	19.0	90.9	36	17.4	91.3	27	21.8	90.3
85+	30	9.1	100.0	18	8.7	100.0	12	9.7	100.0
All ages	331	100.0		207	100.0		124	100.0	

Included in the statistics are 143.8% multiple primaries in males and 122.1% in females.

Table 5

Age-specific incidence, DCO rate and proportion of all cancers for period_2007-2014

							Males	Females
			Males	Females	Males	Females	Prop.all	Prop.all
Age at			Age-	Age-	DCO rate	DCO rate	cancers	cancers
diagnosis	Males	Females		spec.	n=0	n=2	n=91183	n=89596
Years	n	n	incid.	incid.	%	90	્ર	૾ૢ
0- 4			0.0	0.0				
5- 9			0.0	0.0				
10-14			0.0	0.0				
15-19			0.0	0.0				
20-24			0.0	0.0				
25-29	1		0.1	0.0			0.2	
30-34			0.0	0.0				
35-39	1		0.1	0.0			0.1	
40 - 44			0.0	0.0				
45-49	2		0.1	0.0			0.1	
50-54	7	3	0.5	0.2			0.1	0.0
55-59	7	7 /	0.7	0.6			0.1	0.1
60-64	25	7	2.5	0.7			0.2	0.1
65-69	35	11	3.6	1.1			0.2	0.1
70-74	35	24	3.8	2.3			0.2	0.2
75-79	37	32	6.7	4.5		3.1	0.3	0.3
80-84	36	27	10.3	4.8		3.7	0.4	0.3
85+	18	12	7.8	2.1			0.3	0.1
All ages	204	123			0.0	1.6	0.2	0.1
Incidence								
Raw			1.1	0.7				
WS			0.5	0.2				
ES			0.8	0.4				
BRD-S			1.1	0.5				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).






Figure 6. Age distribution and age-specific incidence



ICD-10 C66: Malignant neoplasm of ureter Age-specific incidence rates: international comparison Average 11 Region MCR Period population 2007-2014 4.6 m 10 SEER 2007-2011 64.6 m 1-year averaged incidence (per 100,000) 9 8 6 5 4 3 2 0 94 35-39

Figure 6a. Age-specific incidence in MCR registry areas compared to SEER (Surveillance, Epidemiology, and End Results, USA).

Age at diagnosis (years)

MALES



Reference:

Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence - SEER 18 Regs Research Data, released April 2014, based on the November 2013 submission. http://www.seer.cancer.gov.




Figure 7. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.




Table 8a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2014

MALES

		Observed	Expected		LCL	UCL		DCO
Diagnos	is	n /	n	SIR	95%	95%	EAR	응
C15	Oesophagus	/3	0.3	10.3	2.1	30.2 #	31.2	33.3
C18	Colon	/ 2	1.9	1.1/	0.1	3.9	1.6	
C33-C34	Lung	6	2.1	2.9	1,1	6.3 #	45.3	16.7
C61	Prostate	25	5.1	4.9	3.2	7.2 #	228.6	4.0
C64	Kidney	11	0.6	18.8	9.4	33.7 #	119.8	54.5
C65	Renal pelvis	31	0.1	380.6	258.6	540.3 #	355.7	
C66	Ureter	5	0.0	105.0	34.1	245.1 #	57.0	
C67	Bladder	53	0.9	58.7	44.0	76.8 #	599.4	
C68	Urethra	7	0.0	460.6	185.2	948.9 #	80.4	
C68	Urinary org.	2	0.0	148.3	18.0	535.7 #	22.9	50.0
Other p	rimaries	6	2.6	2.3	0.8	4.9	38.6	33.3
Not obs	erved	0	4.3	0.0	0.0	0.9 #	-49.6	
All mul	t. primaries	151	17.9	8.4	7.1	9.9 #	1531	7.9
Patients				328				
Median a	ge at second m	alignancy	(years)	73.8				
Person-y	rears			869				
Mean obs	ervation time	(years)		2.6				
Median o	bservation tim	e (years)		1.8				

The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries"

Table 8b

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2014

FEMALES

		Observed	Expected		LCL	UCL		DCO
Diagnos	sis	n /	/ n	SIR	95%	95%	EAR	%
C16	Stomach	2 2	0.3	6.3	0.8	22.7	31.0	
C18	Colon		0.9	2.3/	0.3	8.2	20.7	
C50	Breast	3	2.0	1.5	0.3	4.4	18.2	
C64	Kidney	47	0.2	34.2	13.8	70.5 #	125.2	57.1
C65	Renal pelvis	14	0.0	509.4	278.5	854.8 #	257.5	
C66	Ureter	2	0.0	128.0	15.5	462.2 #	36.6	
C67	Bladder	23	0.2	133.6	84.7	200.4 #	420.7	4.3
Other p	primaries	4	0.8	4.9	1.3	12.5 #	58.6	
Not obs	served	0	3.2	0.0	0.0	1.2	-58.8	
All mul	lt. primaries	57	7.6	7.5	5.7	9.7 #	909.6	8.8
Patients	-			200				
Median a	age at second m	alignancy	(years)	76.5				
Person-y	years			543				
Mean obs	servation time	(years)		2.7				


1.3

The occurrence of second malignancy is statistically significant.

Median observation time (years)

Observed second primaries with count 1 are pooled in category "Other primaries"

Average incidence (world standard population) 2007 - 2014: Males



Average incidence (world standard population) 2007 - 2014: Females





Figure 9a. Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.5/100,000 WS N=204, females 0.2/100,000 WS N=123).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 65,347 female residents (averaged) in the period from 2007 to 2014 a total of 4 women were identified with newly diagnosed ureteral cancer. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.3/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.2/100,000.

Standardized incidence ratio (SIR) 2007 - 2014: Males



Standardized incidence ratio (SIR) 2007 - 2014: Females




Figure 9b. Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=204, females N=123).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,924 female residents (averaged) in the period from 2007 to 2014 a total of 4 women were identified with newly diagnosed ureteral cancer. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 1.24. Though, the value of this parameter may vary with an underlying probability of 99% between 0.21 and 3.91, and is therefore not statistically striking.

MORTALITY

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

		Prop.				Prop. deaths
	Incident	actively	Prop.		Prop.	with death
Year of	cases	followed	DCO	Deaths	deaths	certific.
diagnosis	n	90	00	n	90	%
1998	19	100.0		17	89.5	100.0
1999	11	100.0		9	81.8	100.0
2000	11/	100.0		7	63.6	100.0
2001	24	100.0		23/	95.8	100.0
2002	30	100.0		25	83.3	96.0
2003	19	100.0		11	57.9	100.0
2004	24	91.7		18	75.0	94.4
2005	38	94.7		27	71.1	96.3
2006	34	94.1		25	73.5	100.0
2007	34	79.4		19	55.9	100.0
2008	39	71.8		21	53.8	95.2
2009	43	79.1		27	62.8	100.0
2010	48	81.3	2.1	30	62.5	100.0
2011	44	79.5		22	50.0	100.0
2012	42	81.0	2.4	19	45.2	100.0
2013	47	97.9		22	46.8	100.0
2014	34	94.1		4	11.8	25.0
1998-2014	541	88.5	0.4	326	60.3	97.9

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

			Prop.		
			deaths		Prop.
Year of	Incident		with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n /	n	%	n	%
1998	19	13	84.6	3	15.8
1999	11	8	100.0	/ 1	9.1
2000	11	9	100.0	2	18.2
2001	24	12	100.0	3	12.5
2002	30	10	100.0	1	3.3
2003	19	25	96.0	4	21.1
2004	24	13	100.0	1	4.2
2005	38	15	100.0	4	10.5
2006	34	23	95.7	2	5.9
2007	34	21	95.2	1	2.9
2008	39	33	100.0	2	5.1
2009	43	33	100.0	7	16.3
2010	48	27	100.0	3	6.3
2011	44	43	97.7	3 5 3 9	11.4
2012	42	24	100.0	3	7.1
2013	47	31	100.0	9	19.1
2014	34	44	100.0	/1 /	2.9
1998-2014	541	384	98.4	52	9.6

Table 10c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

				Prop.
				cancer
		D	D == 0 ==	
		Prop.	Prop.	recorded
	/	cancer-	non-cancer-	on death
Year of	Deaths	related	related	certificate
death	ń	90	8	%
1998	13	61.5	38.5	81.8
1999	8	75.0	25.0	87.5
2000	9	77.8	22.2	88.9
2001	12	75.0	25.0	83.3
2002	10	80.0	20.0	100.0
2003	25	84.0	16.0	87.5
2004	13	69.2	30.8	69.2
2005	/15	86.7	13.3	86.7
2006	23	73.9	26.1	81.8
2007	21	85.7	14.3	95.0
2008	33	69.7	30.3	72.7
2009	33	84.8	15.2	90.9
2010	27	81.5	18.5	88.9
2011	43	81.4	18.6	88.1
2012	24	75.0	25.0	83.3
2013	31	71.0	29.0	83.9
2014	44	86.4	13.6	93.2
2011	11	00.4	13.0	33.2
1998-2014	384	78.6	21.4	86.2
1990-2014	304	10.0	∠⊥.4	00.2

 $\begin{array}{c} \text{Table 11a} \\ \text{Medians of age at death according to the grouping in Table 10} \\ \text{MALES} \end{array}$

Year of death	Deaths n	Age at death (all causes)	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate)
1998	4	65.2	66.6	61.3	66.6
1999	5	75.8	76.4	69.1	75.8
2000	6	71.8	67.0	71.8	70.6
2001	8	75.2	74.9	75.6	75.2
2002	6	74.6	73.4	78.8	74.6
2002	17	76.3	76.1	85.0	75.9
2003	7	75.6	74.6	94.7	74.6
2004	5		72.1	91.9	74.6
		72.5			
2006	15	78.8	77.6	82.0	78.2
2007	18	75.2	73.4	80.9	75.2
2008	19	77.6	76.1	79.7	76.8
2009	24	74.1	73.9	76.5	73.7
2010	16	75.8	75.2	78.1	75.8
2011	27	80.6	79.2	85.9	79.9
2012	17	75.8	75.8	78.1	78.6
2013	20	80.4	80.4	83.4	80.4
2014	25	78.8	78.8	85.3	78.8
1998-2014	239	76.5	76.1	79.8	76.4

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Year of death	Deaths n	Age at death (all causes)	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate)
	_	/ /			
1998	9	78.0	67.4	84.8	72.7
1999	3	77.2	79.5	77.2	79.5
2000	3	73.2	73.2		73.2
2001	4	75.0	69.6	83.7	69.6
2002	4	73.6	73.6		73.6
2003	8	82.4	80.7	89.2	82.4
2004	6	78.3	80.3	76.2	80.3
2005	10	76.2	75.2	77.2	74.7
2006	8	81.6	82.4	79.0	77.5
2007	3	83.3	83.3		83.3
2008	14	79.5	78.6	84.9	79.5
2009	9	76.3	78.5	65.0	76.3
2010	11	77.6	77.3	78.8	77.3
2011	16	81.5	80.8	84.4	81.5
2012	\7	83.3	76.2	83.9	79.8
2013	11	80.9	80.9	82.1	80.4
2014	19	81.9	80.2	86.1	80.6
1998-2014	145	79.9	79.1	83.1	79.4

By 2010, life expectancy at birth was 77.5 years for boys and 82.6 years for girls.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort. N	II-Index	Mort. N	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	3	0.3	0.25	0.2	0.26	0.2	0.24	0.3	0.20
1999	4	0.4	0.57	0.2	0.43	0.3	0.52	0.6	0.66
2000	4	0.4	0.67	0.2	0.55	0.3	0.63	0.4	0.61
2001	7	0.6	0.54	0.3	0.54	0.5	0.53	0.8	0.57
2002	4	0.2	0.24	0.1	0.23	0.2	0.23	0.2	0.23
2003	14	0.7	1.40	0.4	1.43	0.6	1.46	0.9	1.46
2004	6	0.3	0.40	0.2	0.40	0.3	0.42	0.4	0.41
2005	4	0.2	0.15	0.1	0.15	0.2	0.15	0.2	0.14
2006	11	0.6	0.52	0.2	0.47	0.5	0.57	0.7	0.64
2007	15	0.7	0.60	0.3	0.60	0.5	0.64	0.7	0.68
2008	11	0.5	0.58	0.2	0.44	0.4	0.51	0.6	0.64
2009	20	0.9	0.69	0.4	0.62	0.6	0.67	0.8	0.67
2010	14	0.6	0.41	0.3	0.46	0.4	0.45	0.6	0.43
2011	22	1.0	0.88	0.3	0.61	0.6	0.73	0.9	0.82
2012	15	0.7	0.63	0.3	0.59	0.4	0.61	0.6	0.58
2013	15	0.7	0.63	0.3	0.62	0.5	0.65	0.7	0.63
2014	23	1.0	0.96	0.4	0.87	0.7	0.91	1.0	0.97
1998-2014	192	0.6	0.58	0.3	0.54	0.4	0.58	0.7	0.60

Table 12b

Mortality measures (cancer-related death) and mortality-incidence-index by year of death FEMALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	${\tt MI-Index}$
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	5	0.4	0.71	0.2	0.82	0.3	0.76	0.4	0.72
1999	2	0.2	0.50	0.1	0.66	0.1	0.62	0.2	0.72
2000	3	0.2	0.60	0.1	0.70	0.2	0.67	0.2	0.52
2001	2	0.2	0.18	0.1	0.18	0.1	0.17	0.1	0.18
2002	4	0.2	0.31	0.1	0.23	0.1	0.25	0.2	0.32
2003	7	0.4	0.78	0.1	0.67	0.2	0.67	0.2	0.64
2004	3	0.2	0.33	0.0	0.19	0.1	0.22	0.1	0.28
2005	9	0.5	0.82	0.2	0.62	0.3	0.68	0.3	0.73
2006	6	0.3	0.50	0.1	0.45	0.2	0.46	0.2	0.45
2007	3	0.1	0.38	0.0	0.37	0.1	0.33	0.1	0.33
2008	12	0.5	0.60	0.2	0.51	0.3	0.54	0.4	0.56
2009	8	0.3	0.57	0.1	0.66	0.2	0.64	0.2	0.50
2010	8	0.3	0.57	0.1	0.45	0.2	0.47	0.3	0.50
2011	13	0.6	0.72	0.1	0.54	0.2	0.60	0.4	0.72
2012	3	0.1	0.19	0.0	0.21	0.1	0.20	0.1	0.17
2013	7	0.3	0.30	0.1	0.24	0.1	0.25	0.2	0.28
2014	15	0.6	1.50	0.2	1.12	0.3	1.19	0.5	1.39
1998-2014	110	0.3	0.54	0.1	0.47	0.2	0.49	0.2	0.50

Table 13

Age distribution of age at death (cancer-related) for period 2007-2014

(incl. multiple primaries)

Age at death	Cases			Males			Females		
Years	n	용	Cum.%	n	왕	Cum.%	n	용	Cum.%
50-54	6	2.9	2.9	5	3.6	3.6	1	1.4	1.4
55-59	8	3.9	6.8	5	3.6	7.2	3	4.3	5.8
60-64	12	5.8	12.6	10	7.2	14.5	2	2.9	8.7
65-69	21	10.1	22.7	15	10.9	25.4	6	8.7	17.4
70 - 74	35	16.9	39.6	24	17.4	42.8	11	15.9	33.3
75-79	37	17.9	57.5	25	18.1	60.9	12	17.4	50.7
80-84	47	22.7	80.2	27	19.6	80.4	20	29.0	79.7
85+	41	19.8	100.0	27	19.6	100.0	14	20.3	100.0
All ages	207	100.0		138	100.0		69	100.0	

Included in the statistics are 143.8% multiple primaries in males and 122.1% in females.



Table 14

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2014 (incl. multiple primaries)

			Males		Females		Males	Females
Age at			Age-		Age-		Prop.all	Prop.all
	Males	Females	spec.		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	용	9
0- 4			0.0					
0- 4 5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29			0.0		0.0			
30-34			0.0		0.0			
35-39			0.0		0.0			
40 - 44			0.0		0.0			
45-49			0.0		0.0			
50-54	5 /	1	0.4	0.71	0.1	0.33	0.3	0.1
55-59	5	3	0.5	0.63	0.3	0.43	0.2	0.1
60-64	10	2	1.0	0.40	0.2	0.29	0.2	0.1
65-69	15	6	1.6	0.43	0.6	0.55	0.2	0.1
70-74	24	11	2.6	0.65	1.1	0.46	0.3	0.2
75-79	25	12	4.5	0.68	1.7		0.3	0.2
80-84	27	20	7.7		3.6	0.74	0.4	0.3
85+	27	14	11.7	1.50	2.4	1.17	0.4	0.2
All ages	138	69					0.3	0.2
AII ages	130	09					0.3	0.2
Mortality								
Raw			0.8	0.67	0.4	0.56		
WS			0.3	0.61	0.1	0.48		
ES			0.5	0.65	0.2	0.50		
BRD-S			0.8	0.68	0.3	0.51		
PYLL-70								
per 100,000			1.6		0.5			
ES			1.5		0.5			
AYLL-70			7.5		7.1			

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15a $\begin{tabular}{ll} Multiple primaries in deaths in period 1998-2014 \\ \hline MALES \\ \end{tabular}$

						Syn-	Syn-		
						chron	chron		
		Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis		n	%↓	n	← %	n	← %	n	← %
C15 Oeso	phagus	4	1.4	1	25.0			3	75.0
C18 Colo	n	/ 12 /	4.2	9	75.0	1	8.3	2	16.7
C19-C20 Rect	um	5	1.7	3	60.0	1	20.0	1	20.0
C32 Lary	nx	3	1.0	1	33.3	/ 1	33.3	1	33.3
C33-C34 Lung		8	2.8	1	12.5	1	12.5	6	75.0
C43 Mali	gn. melanoma	6	2.1	4	66.7			2	33.3
C44 Skin	others	8	2.8	7	87.5			1	12.5
C61 Pros	tate	27	9.3	15	55.6	5	18.5	7	25.9
C64 Kidn	ey	19	6.6	7	36.8	4	21.1	8	42.1
C65 Rena	l pelvis	33	11.4	4	12.1	20	60.6	9	27.3
C66 Uret	er	10	3.5			3	30.0	7	70.0
C67 Blad	lder /	116	40.1	52	44.8	_ 15	12.9	49	42.2
C68 Uret	hra /	8	2.8	2	25.0	1	12.5	5	62.5
C68 Urin	ary org.	8	2.8	1	12.5	2	25.0	5	62.5
C82-C85 NHL		5	1.7	3	60.0	2	40.0		
C91-C96 Leuk	aemia	3	1.0	1	33.3			2	66.7
Other primar	ies \	14	4.8	8	57.1			6	42.9
All mult. pr	imaries	289	100.0	119	41.2	56	19.4	114	39.4

Multiple primaries with number of cases 1 to 2 are pooled in category "Other primaries"

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 15b

Multiple primaries in deaths in period 1998-2014
FEMALES

						Syn-	Syn-		
						chron	chron		
		Total	Total	Pre	Pre	±30d	±30d	Post	Post
Diagnosis		n /	%↓	n	← %	n	← %	n	← %
C16 St	tomach	/2	1.6					2	100.0
C18 C	olon	10	7.8	6	60.0	2	20.0	2	20.0
C19-C20 Re	ectum	/ 3	2.3	2	66.7			1	33.3
C33-C34 L1	ung	3	2.3			/ 1	33.3	2	66.7
C44 S1	kin others	3	2.3	2	66.7			1	33.3
C48 P6	eritoneal	2	1.6	1	50.0	/ 1	50.0		
C50 B:	reast	11	8.5	7	63.6			4	36.4
C53 C6	ervix uteri	5	3.9	5	100.0				
C54 C	orpus uteri	3	2.3	3	100.0				
C56 0	vary	2	1.6	1	50.0			1	50.0
C64 K:	idney	7	5.4	1	14.3	2	28.6	4	57.1
C65 R6	enal pelvis	22	17.1	5	22.7	12	54.5	5	22.7
C66 U:	reter	3	2.3			1	33.3	2	66.7
C67 B3	ladder	44	34.1	12	27.3	9	20.5	23	52.3
C68 U:	rinary org.	3	2.3	1	33.3	1	33.3	1	33.3
Other prin	maries \	6	4.7	3	50.0			3	50.0
All mult.	primaries	129	100.0	49	38.0	29	22.5	51	39.5

Multiple primaries with number of cases 1 are pooled in category "Other primaries"

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2014

(First primaries only *)

Age at death Years	Males Female	/ = /	MI-index	Females Age- spec. mortal.	MI-index	cancers	Females Prop.all cancers
0- 4 5- 9 10-14 15-19 20-24 25-29 30-34 35-39		0.0 0.0 0.0 0.0 0.0 0.0		0.0 0.0 0.0 0.0 0.0 0.0			
33-39 40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85+	3 1 3 1 4 8 8 4 9 4 10 8 8 9	0.0 0.0 0.2 0.3 0.4 0.8 1.0 1.8 2.3 4.3	0.60 1.50 0.29 0.50 1.00 0.63 0.53	0.0 0.0 0.1 0.1 0.0 0.4 0.4 1.1 1.6	1.00 0.50 0.80 0.50 0.50 0.69 1.33	0.2 0.1 0.1 0.1 0.1 0.2 0.1	0.1 0.0 0.1 0.1 0.2 0.2
All ages	55 35					0.1	0.1
Mortality Raw WS ES BRD-S		0.3 0.1 0.2 0.3	0.59	0.2 0.1 0.1 0.1	0.65 0.58 0.60 0.60		
PYLL-70 per 100,000 ES AYLL-70		0.9 0.8 7.8		0.3 0.2 6.7			

^{*} See corresponding tables with multiple primaries.

Table 17

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2014 (Single primaries only *)

Age at death Years	Males n	Females n	Males Age- spec. mortal.	MI-index	Females Age- spec. mortal.	MI-index	cancers	Females Prop.all cancers
0 - 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29			0.0		0.0			
30-34			0.0		0.0			
35-39			0.0		0.0			
40 - 44			0.0		0.0			
45-49			0.0		0.0			
50-54	2	/ 1/	0.2		0.1	1.00	0.1	0.1
55-59	1/		0.1		0.0		0.0	
60-64	2		0.2		0.0		0.1	
65-69	3	4	0.3		0.4		0.1	0.1
70-74	3	\ 1	0.3		0.1	0.20	0.1	0.0
75-79	3 \	5	0.5		0.7		0.1	0.1
80-84	5	6	1.4		1.1		0.1	0.1
85+	5	3	2.2	1.67	0.5	0.50	0.1	0.1
All ages	24	20					0.1	0.1
Mortality						/		
Raw			0.1		0.1			
WS			0.1	0.42	0.0	0.54		
ES			0.1	0.46	0.1			
BRD-S			0.1	0.47	0.1	0.56		
PYLL-70			0.4		2 2			
per 100,000			0.4		0.2			
ES 70			0.4		0.1			
AYLL-70			8.8		5.5			

^{*} See corresponding tables with multiple primaries.





Figure 18. Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).

The difference between age at diagnosis (Table 3) and age at ureteral cancer-related death (see Table 10) should be considered.



Average mortality (world standard population) 2007 - 2014: Males



Average mortality (world standard population) 2007 - 2014: Females





Figure 19a. Map of cancer mortality (world standard population) by county averaged for period 2007 to 2014. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.3/100,000 WS N=130, females 0.1/100,000 WS N=65).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 65,347 female residents (averaged) in the period from 2007 to 2014 a total of 2 women died from ureteral cancer. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.1/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 0.5/100,000.

Standardized mortality ratio (SMR) 2007 - 2014: Males



Standardized mortality ratio (SMR) 2007 - 2014: Females




Figure 19b. Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=130, females N=65).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,924 female residents (averaged) in the period from 2007 to 2014 a total of 2 women died from ureteral cancer. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 1.20. Though, the value of this parameter may vary with an underlying probability of 99% between 0.06 and 5.58, and is therefore not statistically striking.

Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases, where applicable. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

3. Single primary (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

Shortcuts

FRG Federal Republic of Germany

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

MCR Munich Cancer Registry (Tumorregister München)
SEER Surveillance, Epidemiology, and End Results (USA)

AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

BRD-S German standard population

DCO Death certificate only EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

ES European standard population (old)

LCL Lower confidence limit

MI-index Ratio between mortality and incidence

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age

SIR Standardized incidence ratio
SMR Standardized mortality ratio
UCL Upper confidence limit
WS World standard population

Recommended Citation

Munich Cancer Registry. ICD-10 C66: Ureteral cancer - Incidence and Mortality [Internet]. 2016 [updated 2016 Apr 13; cited 2016 Jun 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/base/bC66 E-ICD-10-C66-Ureteral-cancer-incidence-and-mortality.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

Fig./Tb	l.	Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	3
1a	Gender distribution by year of diagnosis	4
2	Incidence by year of diagnosis	5
3	Age distribution parameters by year of diagnosis	6
4	Age distribution by 5-year age group and gender	8
5	Age-specific incidence, DCO rate, proportion malignancies	9
6	Age distribution and age-specific incidence (chart)	10
6a	Age-specific incidence internationally (chart)	11
7	Cumulative follow-up years (chart)	12
8	Standardized incidence ratio of second primaries	13
9a	Map of cancer incidence (WS) by county (chart)	15
9b	Standardized incidence ratio (SIR) by county (chart)	16
10a	Pts incident cohorts and mortality / yr	17
10b	Incidence and mortality by year of diagnosis	18
10c	Cancer-related deaths, death certification available / yr	19
11	Medians of age at death / yr	20
12	Mortality by year of death	22
13	Distribution of age at death	23
14	Age-specific mortality	24
15	Multiple primaries in deaths	25
16	Age-specific mortality (first primaries)	27
17	Age-specific mortality (single primaries)	28
18	Age distribution and age-specific mortality (chart)	29
19a	Map of cancer mortality (WS) by county (chart)	30
19b	Standardized mortality ratio (SMR) by county (chart)	31