# **Munich Cancer Registry**



- ▶ Survival
- ▶ Selection Matrix
- ▶ Homepage
- ▶ Deutsch

## **ICD-10 C96: Systemic Disease NOS**

## **Incidence and Mortality**

Year of diagnosis	1998-2014
Patients	195
Diseases	195
Creation date	04/13/2016
Export date	12/23/2015
Population	4.64 m



Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

http://www.tumorregister-muenchen.de/en/facts/base/bC96\_\_E-ICD-10-C96-Systemic-Disease-NOS-incidence-and-mortality.pdf

# Global Statements about the statistics on the Internet – Baseline Statistics (grey button ——), Survival (red button ——)

In these analyses, the clinics and physicians of Upper Bavaria and the city and county of Landshut<sup>#</sup>, with a total of 4.64 million inhabitants, account for the frequency of cancer diseases<sup>##</sup> and the achieved long term results. Additionally, the long term survival evaluated by the Munich Cancer Registry (MCR) is compared with the results of the population-based registry in the USA (SEER), which is useful for checking the consistency of the data on an international level.

In comparing several tables, inconsistent figures may be detected. This is based on the fact that different patient cohorts are included in the base calculation, for example when proportions of multiple tumors or DCO-cases\*\*\*\* are concerned. In other cases the individual tumor diagnosis is the basis for calculation, for example with incidence.

The foot notes describe the currentness of the data. The baseline statistics and survival data are updated annually. This yearly analysis comprises the Annual Report of the MCR.

Clinics and physicians have access to essentially more detailed data, with which they can check, compare and in the best case optimize their own data and results.

We would be pleased to receive corrections, critique and useful suggestions. Just send an e-mail to tumor@ibe.med.uni-muenchen.de.

Munich Cancer Registry, April 2016

- Base data has been collected since 1998. An increase in new diseases is apparent, which is an effect of two extensions in the MCR catchment area (from a base population of 2.51 million to 3.96 in 2002, and to 4.52 million in 2007).
- Due to the high frequency and good prognosis of non-malignant skin cancer (C44), no systematic ascertainment is performed for this diagnosis. C44 is not designated as a primary, but rather as a secondary tumor.
- DCO (death certificate only) identifies a cancer case that first becomes available to the MCR through the death certificate.

#### ICD-10 codes (ICD-10 2015) used for specifying cancer site

Code	Description
C96	Other and unspecified malignant neoplasms of lymphoid, haematopoietic and related tissue
C96.0	Multifocal and multisystemic (disseminated) Langerhans-cell histiocytosis [Letterer-Siwe disease]
C96.2	Malignant mast cell tumour
C96.4	Sarcoma of dendritic cells (accessory cells)
C96.5	Multifocal and unisystemic Langerhans-cell histiocytosis
C96.6	Unifocal Langerhans-cell histiocytosis
C96.7	Other specified malignant neoplasms of lymphoid, haematopoietic and related tissue
C96.8	Histiocytic sarcoma
C96.9	Malignant neoplasm of lymphoid, haematopoietic and related tissue, unspecified

#### **INCIDENCE**

Table 1

All patients by year of diagnosis, proportions of DCO, multiple primaries, deaths, and active follow-up (incl. DCO)

				Prop.		Prop.
		DCO	Prop.	mult.	Prop.	actively
Year of	Cases	cases	DCO	primaries	deaths	followed
diagnosis	n	n	olo	용	용	ું ર
1998	6	2	33.3		33.3	66.7
1999	8	5	62.5		62.5	100.0
2000	5	2	40.0		60.0	100.0
2001	6	3	50.0	50.0	83.3	100.0
2002	7			28.6	28.6	85.7 #
2003	14			50.0	35.7	85.7
2004	/11			9.1	36.4	90.9
2005	13 /	1	7.7	15.4	15.4	69.2
2006	8			12.5	50.0	75.0
2007	16	2	12.5	6.3	18.8	56.3 #
2008	\15 \			20.0	26.7	40.0
2009	7	2	28.6	28.6	71.4	85.7
2010	10			30.0	20.0	40.0
2011	13			15.4	30.8	61.5
2012	16			37.5	37.5	81.3
2013	13	3	23.1	15.4	30.8	100.0
2014	27	22	81.5	37.0	85.2	100.0 ##
1998-2014	195	42	21.5	23.1	42.6	77.9

<sup>#</sup> The increases of incident cases in 2002 and 2007 reflect the expansion to additional registry areas.

<sup>##</sup> Please be aware that data of recent annual patient cohorts may not yet be fully processed. The years under evaluation can be found in the respective headings.

Table 1a

All patients
by year of diagnosis and gender
(incl. DCO)

Year of	All/	Males	Female	s Prop. mai	Les
diagnosi	s n	/ n	n	\	
1998	/ 6	/ 2	4	33.3	
1999	/ 8	/ 7	1 /	87.5	
2000	/ 5		3 4 3	40.0	
2001	/ 6	2	4	33.3	
2002	7	4		57.1	
2003	14	9	5	64.3	
2004	11	5	6	45.5	
2005	13	8	5	61.5	
2006	8	4	4	50.0	
2007	16	15	1	93.8	
2008	15	6	9	40.0	
2009	7	4	3	57.1	
2010	10	7	3	70.0	
2011	13	7	6	53.8	
2012	16	12	4	75.0	
2013	13	6	7	46.2	
2014	27	16	11	59.3	
1998-201	4 195	116	79	59.5	

Table 2

Incidence measures by year of diagnosis including DCO cases (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

			Males	Fem.	Males	Fem.	Males	Fem.	Males	Fem.
Year of	Males	Females	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.	Inc.
diagnosis	n	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	2	4	0.2	0.3	0.3	0.5	0.2	0.4	0.2	0.4
1999	7	1 /	0.6	0.1	0.8	0.0	0.7	0.0	0.7	0.1
2000	2	3 /	0.2	0.2	0.3	0.1	0.2	0.1	0.2	0.2
2001	2	4 <	0.2	0.3	0.1	0.1	0.2	0.2	0.2	0.2
2002	4	3	0.2	0.2	0.2	0.1	0.2	0.1	0.2	0.1
2003	9	5	0.5	0.3	0.5	0.2	0.5	0.3	0.5	0.3
2004	5	6	0.3	0.3	0.2	0.5	0.2	0.4	0.2	0.3
2005	8	5	0.4	0.3	0.6	0.3	0.5	0.3	0.4	0.3
2006	4	4	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2007	15	1	0.7	0.0	0.7	0.0	0.7	0.0	0.7	0.0
2008	6	9	0.3	0.4	0.2	0.3	0.3	0.4	0.3	0.4
2009	4	3	0.2	0.1	0.1	0.0	0.1	0.1	0.2	0.1
2010	7 /	3	0.3	0.1	0.3	0.2	0.3	0.2	0.3	0.1
2011	7	6	0.3	0.3	0.2	0.2	0.3	0.2	0.3	0.2
2012	12	4	0.5	0.2	0.4	0.1	0.4	0.1	0.6	0.1
2013	6	7	0.3	0.3	0.2	0.1	0.2	0.2	0.3	0.2
2014	16	11	0.7	0.5	0.3	0.1	0.5	0.2	0.7	0.3
1998-2014	116	79	0.4	0.2	0.3	0.2	0.3	0.2	0.4	0.2




Table 3

Age distribution parameters by year of diagnosis (All patients) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	6	34.8	26.4	7.8	74.3	7.8	11.9	32.4	50.1	74.3
1999	8	47.0	34.4	2.4	84.4	2.4	9.2	60.3	75.2	84.4
2000	5	51.3	37.9	3.8	92.7	3.8	20.8	59.9	79.3	92.7
2001	6	70.0	22.6	35.0	87.8	35.0	48.6	80.7	87.4	87.8
2002	7	39.8	24.7	9.4	80.2	9,4	11.2	39.9	53.1	80.2
2003	14	44.9	26.9	2.2	90.2	14.6	22.9	42.8	71.6	78.4
2004	11	37.0	31.8	0.8	85.4	1.3	3.8	38.0	75.8	77.8
2005	13	30.6	23.8	5.1	68.9	7.9	9.7	27.0	50.4	68.9
2006	8	47.5	27.4	6.6	76.8	6.6	22.8	52.3	73.1	76.8
2007	16	35.5	23.6	0.1	81.7	11.6	18.3	32.5	43.8	79.8
2008	15	45.4	20.0	8.4	88.9	22.5	30.9	45.9	56.1	73.2
2009	7	65.3	21.1	20.7	84.0	20.7	63.2	70.6	81.5	84.0
2010	10	38.4	27.1	0.3	73.3	0.4	15.4	41.1	62.7	70.9
2011	13 /	53.6	15.4	24.1	78.5	37.7	43.7	56.0	61.1	77.5
2012	16/	58.5	27.5	1.5	85.1	18.4	34.1	71.5	80.2	83.2
2013	13	60.7	24.9	22.5	93.5	34.5	42.7	48.9	84.6	91.6
2014	27	76.9	15.9	39.5	91.6	43.5	75.2	82.8	88.4	91.2
1998-2014	195	51.0	27.6	0.1	93.5	9.5	28.1	50.1	76.8	85.2

Table 3a

Age distribution parameters by year of diagnosis (MALES)

(incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	2	27.2	27.5	7.8	46.7	7.8	7.8	27.2	46.7	46.7
1999	7	41.7	33.4	2.4	79.1	2.4	8.9	54.8	71.3	79.1
2000	2	12.3	12.0	3.8	20.8	3.8	3.8	12.3	20.8	20.8
2001	2	68.0	27.4	48.6	87.4	48.6	48.6	68.0	87.4	87.4
2002	4	37.3	18.8	11.2	53.1	11.2	23.5	42.4	51.0	53.1
2003	9	42.2	27.5	2.2	90.2	2.2	28.5	36.7	54.8	90.2
2004	5	39.9	24.3	8.4	75.8	8.4	32.9	38.0	44.5	75.8
2005	8	28.4	24.5	5.1	68.9	5.1	9.0	21.4	46.1	68.9
2006	4	52.5	31.7	6.6	76.8	6.6	31.7	63.2	73.2	76.8
2007	15	35.5	24.5	0.1	81.7	11.6	17.2	30.3	45.5	79.8
2008	6	41.7	21.1	8.4	73.2	8.4	34.4	42.7	48.9	73.2
2009	4	56.7	24.3	20.7	72.2	20.7	41.9	66.9	71.4	72.2
2010	7	38.0	26.9	0.3	73.3	0.3	15.4	35.4	68.5	73.3
2011	7	53.7	17.8	24.1	78.5	24.1	40.3	59.8	63.8	78.5
2012	12	54.6	29.2	1.5	83.1	18.4	25.3	70.4	78.8	81.4
2013	6	47.0	20.4	22.5	84.6	22.5	39.8	44.2	46.9	84.6
2014	16	72.5	18.6	39.5	91.2	40.5	60.3	77.3	86.8	89.0
1998-2014	116	46.7	26.8	0.1	91.2	8.4	25.2	46.3	72.1	81.7

Table 3b

Age distribution parameters by year of diagnosis (FEMALES) (incl. DCO)

Year of	Cases		Std.					Median		
diagnosis	n	Mean	dev.	Min.	Max.	10%	25%	50%	75%	90%
1998	4	38.6	29.1	11.9	74.3	11.9	15.0	34.1	62.2	74.3
1999	1	84.4		84.4	84.4	84.4	84.4	84.4	84.4	84.4
2000	3	77.3	16.5	59.9	92.7	59.9	59.9	79.3	92.7	92.7
2001	4	71.0	24.4	35.0	87.8	35.0	56.0	80.7	86.0	87.8
2002	3	43.2	35.5	9.4	80.2	9,4	9.4	39.9	80.2	80.2
2003	5	49.8	28.2	16.9	78.4	16.9	22.9	59.1	71.6	78.4
2004	6	34.6	39.2	0.8	85.4	0.8	1.3	21.2	77.8	85.4
2005	5	34.1	25.1	7.9	68.9	7.9	16.4	27.0	50.4	68.9
2006	4	42.5	26.1	16.8	76.6	16.8	22.8	38.3	62.2	76.6
2007	1	36.8		36.8	36.8	36.8	36.8	36.8	36.8	36.8
2008	9	47.9	20.2	22.5	88.9	22.5	30.9	47.1	56.1	88.9
2009	3	76.8	10.4	64.9	84.0	64.9	64.9	81.5	84.0	84.0
2010	3	39.3	33.8	0.6	62.7	0.6	0.6	54.6	62.7	62.7
2011	6 /	53.5	13.8	37.7	77.5	37.7	43.7	53.0	56.2	77.5
2012	4	70.2	20.2	41.3	85.1	41.3	56.3	77.3	84.2	85.1
2013	7	72.3	23.3	34.5	93.5	34.5	48.9	84.1	91.6	93.5
2014	11	83.2	8.0	65.7	91.6	75.2	76.6	85.7	89.5	91.2
1998-2014	79	57.2	27.6	0.6	93.5	16.4	36.8	59.9	82.8	88.9

Table 4

Age distribution by 5-year age group and gender for period 2007-2014 (incl. DCO)

Age at									
diagnosis	Cases			Males			Females		
Years	n	% Cu	m.%	n	양	Cum.%	n	%	Cum.%
0 - 4	4	3.4	3.4	3	4.1	4.1	1	2.3	2.3
5-9	1	0.9	4.3	1	1.4	5.5			2.3
10-14	2	1.7	6.0	2	2.7	8.2			2.3
15-19	4	3.4	9.4	4	5.5	13.7			2.3
20-24	5	4.3 1	3.7	4	5.5	19.2	1	2.3	4.5
25-29	5	4.3 1	7.9	4	5.5	24.7	1	2.3	6.8
30-34	5	4.3 2	2.2	3	4.1	28.8	2	4.5	11.4
35-39	7	6.0 2	8.2	5	6.8	35.6	2	4.5	15.9
40 - 44	8	6.8 3	5.0	5	6.8	42.5	3	6.8	22.7
45-49	10	8.5 4	3.6	8	11.0	53.4	2	4.5	27.3
50-54	4	3.4 4	7.0	1	1.4	54.8	3	6.8	34.1
55-59	5	4.3 5	1.3	1	1.4	56.2	4	9.1	43.2
60-64	6	5.1 5	6.4	3	4.1	60.3	3	6.8	50.0
65-69	4	3.4 5	9.8	3	4.1	64.4	1	2.3	52.3
70 - 74	8	6.8 6	6.7	7	9.6	74.0	1	2.3	54.5
75-79	11	9.4 7	6.1	8	11.0	84.9	3	6.8	61.4
80 - 84	12	10.3 8	6.3	6	8.2	93.2	6	13.6	75.0
85+	16	13.7 10	0.0	5	6.8	100.0	\11	25.0	100.0
All ages	117	100.0		73	100.0		4 4	100.0	

Included in the statistics are 34.2% multiple primaries in males and 27.3% in females.



							Males	Females
			Males	Females	Males	Females	Prop.all	Prop.all
Age at			Age-	Age-	DCO rate	DCO rate	_	cancers
diagnosis	Males	Females	/ =	spec.	n=14	n=15	n=91183	n=89596
Years	n	n	incid.	incid.	%	90	%	%
0- 4	3	1	0.3	0.1			1.7	0.7
5- 9	1		0.1	0.0			1.0	
10-14	2		0.2	0.0			2.0	
15-19	4		0.4	0.0			1.9	
20-24	4	1	0.4	0.1			1.1	0.3
25-29	4	1	0.3	0.1			0.7	0.2
30-34	3	2	0.2	0.2			0.4	0.2
35-39	5	2	0.4	0.2			0.4	0.1
40 - 44	5	3	0.3	0.2			0.3	0.1
45-49	8	2	0.5	0.1			0.2	0.0
50-54	1	3	0.1	0.2			0.0	0.0
55-59	1	4	0.1	0.4			0.0	0.1
60-64	3	3	0.3	0.3			0.0	0.0
65-69	3	1	0.3	0.1			0.0	0.0
70-74	7	1	0.8	0.1	14.3		0.0	0.0
75-79	8	3	1.5	0.4	62.5	66.7	0.1	0.0
80-84	6	6	1.7	1.1	50.0	66.7	0.1	0.1
85+	5	11	2.2	1.9	100.0	81.8	0.1	0.1
All ages	73	44			19.2	34.1	0.1	0.0
Incidence								
Raw			0.4	0.2				
WS			0.3	0.1				
ES			0.4	0.2				
BRD-S			0.4	0.2				

The age-specific incidence characterizes the disease risk in a particular age group. The age distribution depends on the patient population frequency in each age group and reflects the tangible clinical picture of everyday patients care (see following chart).



CD-10 C96: Other and unspecified malignant neoplasms of lymphoid, haematopoietic and related tissue Age distribution and age-specific incidence 2007 - 2014 (Males: 73, Females: 44)




Figure 6. Age distribution and age-specific incidence






Figure 6a. Age-specific incidence in MCR registry areas.




Figure 7. Cumulative follow-up years depending on time since diagnosis

The increase of the lost to follow-up rate can be interpreted as a consequence of a declining number of survivors over time.

Table 8a

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2014

M	Δ	Т	F	C

	Observed	Expected		LCL	UCL		DCO
Diagnosis	n /	n	SIR	95%	95%	EAR	%
C61 Prostate	2 /	0.4	5.0	0.6	17.9	49.5	50.0
C91-C96 Leukaemia	2	0.0	63.9	7.7	230.8	# 61.0	50.0
Other primaries	2	0.1	29.7	3.6	107.4	# 59.9	
Not observed	0	1.1	0.0	0.0	3.4	-33.8	
All mult. primaries	6	1.6	3.8	1.4	8.2	# 136.5	33.3
Patients			98				
Median age at second	malignancy	(years)	76.9				
Person-years			323				
Mean observation time	(years)		3.3				
Median observation time	me/(years)		1.4				

### # The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries"

Table 8b

Standardized incidence ratio (SIR, with 95% confidence limits), excess absolute risk (EAR) and DCO rate of second primaries for period 1998-2014

FEMALES


	Observed	Expected		LCL	UCL		DCO
Diagnosis	n	n	SIR	95%	95%	EAR	읭
C91-C96 Leukaemia	2	0.0	88.8	10.8	320.8 #	98.3	100.0
Other primaries	2	0.1	24.5	3.0	88.6 #	95.4	50.0
Not observed	0	1.1	0.0	0.0	3.4	-54.4	
All mult. primaries	4	1.2	3.3	0.9	8.5	139.2	75.0
Patients			59				
Median age at second	malignancy	(vears)	86 2				

Patients			59
Median age at second	malignancy	(years)	86.2
Person-years			201
Mean observation time	e (years)		3.4
Median observation to	ime (years)		2.1


# The occurrence of second malignancy is statistically significant.

Observed second primaries with count 1 are pooled in category "Other primaries"

### Average incidence (world standard population) 2007 - 2014: Males




#### Average incidence (world standard population) 2007 - 2014: Females




**Figure 9a.** Map of cancer incidence (world standard population, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual incidence rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.3/100,000 WS N=73, females 0.1/100,000 WS N=44).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 65,347 female residents (averaged) in the period from 2007 to 2014 a total of 0 women were identified with newly diagnosed systemic Disease NOS. Therefore, the mean incidence rate for this cancer type in this area can be calculated at 0.0/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.0/100,000.

#### Standardized incidence ratio (SIR) 2007 - 2014: Males



#### Standardized incidence ratio (SIR) 2007 - 2014: Females



**Figure 9b.** Map of standardized incidence ratio (SIR, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual SIR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=73, females N=44).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,924 female residents (averaged) in the period from 2007 to 2014 a total of 0 women were identified with newly diagnosed systemic Disease NOS. Therefore, the mean standardized incidence ratio (SIR) for this cancer type in this area can be calculated at 0.00. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 4.61, and is therefore not statistically striking.

#### **MORTALITY**

Table 10a

Patient cohorts of incident cancers by year of diagnosis, follow-up status, proportion of DCO, deaths among the annual cohorts and proportion of available death certificates (with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

Year of diagnosis	Incident cases n	Prop. actively followed %	Prop. DCO	Deaths n	Prop. deaths	Prop. deaths with death certific. %
1998	6	66.7	33.3	2	33.3	100.0
1999	8	100.0	62.5	5	62.5	100.0
2000		100.0	40.0	3	60.0	100.0
2000	5 6	100.0	50.0	5 /	83.3	100.0
2002	7	85.7	30.0		28.6	100.0
2003	14	85.7		2 5	35.7	60.0
2003	11	90.9		4	36.4	75.0
2005	13	69.2	7.7	2	15.4	100.0
2006	8	75.0	/ • /	4	50.0	100.0
2007	16	56.3	12.5	3	18.8	100.0
2008	15	40.0	12.0	4	26.7	100.0
2009	7	85.7	28.6	5	71.4	80.0
2010	10	40.0	20.0	2	20.0	100.0
2011	13	61.5		4	30.8	75.0
2012	16	81.3		6	37.5	100.0
2013	13	100.0	23.1	4	30.8	100.0
2014	27	100.0	81.5	23	85.2	95.7
1998-2014	195	77.9	21.5	83	42.6	92.8

Table 10b

Annual cohorts of incident cancers and deaths, proportion of death certificates and cases deceased the same year of cancer diagnosis (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

			Duran		
			Prop.		Dron
			deaths		Prop.
Year of	Incident	_ /,	with death	Deaths in	deaths in
diagnosis/	cases	Deaths	certific.	same year	same year
death	n	n	90	n	90
1998	6	4	100.0	2	33.3
1999	8	6	100.0	5	62.5
2000	5	2	100.0	2	40.0
2001	6	5	80.0	4	66.7
2002	7	3	100.0	1	14.3
2003	1.4	2	100.0		
2004	/11	5	100.0		
2005	13	3	66.7	_ 1	7.7
2006	8	4	100.0	1	12.5
2007	16	6	100.0	3	18.8
2008	15	6	100.0	2	13.3
2009	7	5	100.0	3	42.9
2010	10	3	66.7		
2011	13	7	85.7	3	23.1
2012	16	3 3	100.0	3 1 2	6.3
2013	13	3	100.0	2	15.4
2014	27	20	100.0	22	81.5
1998-2014	195	87	95.4	52	26.7

Table 10c

Annual cohorts of deaths, proportion of cancer-related and non-cancer-related deaths, and cancer recorded on death certificates (incl. DCO)

(with respect to registry area expansion from 2.51 to 3.96 m as of 2002, and from 3.96 to 4.64 m as of 2007, respectively)

				Prop.
				cancer
		Prop.	Prop.	recorded
		cancer-	non-cancer-	on death
Year of	Deaths	related	related	certificate
death	n	90	%	%
1998	4	25.0	75.0	100.0
1999	6	33.3	66.7	83.3
2000	2	50.0	50.0	100.0
2001	5	80.0	20.0	100.0
2002	3	66.7	33.3	100.0
2003	2	100.0		100.0
2004	5	80.0	20.0	60.0
2005	3	33.3	66.7	50.0
2006	4	75.0	25.0	75.0
2007	6	83.3	16.7	100.0
2008	6	83.3	16.7	83.3
2009	5	40.0	60.0	60.0
2010	3 7	100.0		100.0
2011		57.1	42.9	66.7
2012	3	66.7	33.3	66.7
2013	3	100.0		100.0
2014	20	75.0	25.0	95.0
1998-2014	87	67.8	32.2	85.5

 $$\operatorname{\textsc{Table}}$$  11a  $$\operatorname{\textsc{Medians}}$$  of age at death according to the grouping in Table 10  $$\operatorname{\textsc{MALES}}$$ 

Year of death	Deaths n	Age at death (all causes)	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate)
1998	2	70.1	59.7	80.5	70.1
1999	5	65.8	75.2	54.8	68.5
2000					
2001	1	87.4		87.4	87.4
2002	2	17.3	32.7	1.9	17.3
2003	2	30.8	30.8		30.8
2004	4	63.9	46.1	81.8	68.3
2005	1	85.1		85.1	85.1
2006	2	61.2	61.2		61.2
2007	6 2 3 2 6	56.8	55.5	79.8	56.8
2008	2	67.6	67.6		67.6
2009	3	63.5	66.7	63.3	66.7
2010	2	66.7	66.7		78.3
2011	6	58.5	54.2	61.7	54.2
2012	2	72.2	74.5	70.0	72.2
2013	3	75.4	75.4		75.4
2014	10	84.4	85.3	82.3	84.4
1998-2014	53	71.3	69.8	77.6	73.5

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

 $\begin{tabular}{ll} Table 11b \\ \hline \begin{tabular}{ll} Medians of age at death according to the grouping in Table 10 \\ \hline \begin{tabular}{ll} FEMALES \end{tabular}$ 

Year of death	Deaths n	Age at death (all causes)	Age at death (cancer-related) Years	Age at death (non-cancer-related) Years	Age at death (according to death certificate)
	_	//			
1998	2	62.2		62.2	62.2
1999	1	84.4		84.4	84.4
2000	2	86.0	79.3	92.7	86.0
2001	4	80.7	80.7		84.3
2002	1	80.6	80.6		80.6
2003					
2004	1	64.1	64.1		64.1
2005	2	84.3	90.2	78.4	
2006	2	37.0	2.1	71.9	2.1
2007					
2008	4	75.9	72.5	79.2	72.5
2009	2	79.4		79.4	81.1
2010	4 2 1 1	61.9	61.9		61.9
2011	1	30.8		30.8	
2012	1	48.7	48.7		
2013					
2014	10	85.7	84.4	87.9	87.0
1998-2014	34	79.3	80.0	78.4	81.1

By 2010, life expectancy at birth was 77.5 years for boys and 82.6 years for girls.

Deaths of patients are considered to be cancer-related, in case that fact was recorded on the death certificate, or patients had suffered from metastasis or recurrence.

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998	1	0.1	0.50	0.0	0.17	0.1	0.31	0.1	0.34
1999	2	0.2	0.29	0.1	0.12	0.2	0.23	0.3	0.37
2000									
2001									
2002	1	0.1	0.25	0.0	0.18	0.0	0.20	0.0	0.21
2003	2	0.1	0.22	0.1	0.26	0.1	0.25	0.1	0.21
2004	3	0.2	0.60	0.1	0.49	0.2	0.65	0.2	0.73
2005									
2006	2	0.1	0.50	0.1	0.37	0.1	0.47	0.1	0.45
2007	5	0.2	0.33	0.1	0.19	0.2	0.29	0.2	0.33
2008	2	0.1	0.33	0.1	0.21	0.1	0.28	0.1	0.33
2009	2	0.1	0.50	0.1	0.44	0.1	0.49	0.1	0.44
2010	2	0.1	0.29	0.0	0.13	0.1	0.24	0.1	0.32
2011	4	0.2	0.57	0.1	0.55	0.1	0.53	0.2	0.55
2012	1	0.0	0.08	0.0	0.04	0.0	0.05	0.0	0.06
2013	3	0.1	0.50	0.0	0.23	0.1	0.36	0.1	0.50
2014	6	0.3	0.38	0.1	0.31	0.2	0.38	0.3	0.39
1998-2014	36	0.1	0.31	0.1	0.20	0.1	0.27	0.1	0.31

Table 12b

Mortality measures (cancer-related death) and mortality-incidence-index by year of death FEMALES

Year of	Deaths	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index	Mort.	MI-Index
death	n	raw	raw	WS	WS	ES	ES	BRD-S	BRD-S
1998									
1999									
2000	1	0.1	0.33	0.0	0.25	0.0	0.29	0.1	0.41
2001	4	0.3	1.00	0.1	0.68	0.1	0.87	0.3	1.06
2002	111	0.1	0.33	0.0	0.05	0.0	0.12	0.0	0.29
2003									
2004	1	0.1	0.17	0.0	0.07	0.0	0.11	0.0	0.15
2005	1	0.1	0.20	0.0	0.03	0.0	0.06	0.0	0.09
2006	1	0.0	0.25	0.1	0.65	0.1	0.45	0.1	0.23
2007									
2008	3	0.1	0.33	0.1	0.21	0.1	0.24	0.1	0.25
2009									
2010	1	0.0	0.33	0.0	0.17	0.0	0.24	0.0	0.31
2011									
2012	1	0.0	0.25	0.0	0.51	0.0	0.41	0.0	0.37
2013									
2014	9	0.4	0.82	0.1	0.69	0.1	0.76	0.2	0.87
1998-2014	23	0.1	0.29	0.0	0.16	0.0	0.20	0.1	0.25

Table 13

Age distribution of age at death (cancer-related) for period 2007-2014

(incl. multiple primaries)

Age at death	Cases		Males			Females		
Years	n	% Cum.	% n	ଚ	Cum.%	n	90	Cum.%
25-29	1	2.6 /2.	6 / 1	4.0	4.0			0.0
30-34	1	2.6 / 5.	1 /		4.0	1	7.1	7.1
35-39	0	0.0 / 5.	1/		4.0			7.1
40 - 44	1	2.6 / 7.	7 1	4.0	8.0/			7.1
45-49	3	7.7 15.	4 2	8.0	16.0	1	7.1	14.3
50-54	0	0.0 15.	4		16.0			14.3
55-59	3	7.7 23.	1 3	12.0	28.0			14.3
60-64	3	7.7 30.	8 2	8.0	36.0	1	7.1	21.4
65-69	2	5.1 35.	9 2	8.0	44.0			21.4
70-74	4	10.3 46.	2 3	12.0	56.0	1	7.1	28.6
75-79	9	23.1 69.	2 7	28.0	84.0	2	14.3	42.9
80-84	3	7.7 76.	9		84.0	3	21.4	64.3
85+	9	23.1 100.	0 4	16.0	100.0	5	35.7	100.0
All ages	39	100.0	25	100.0		14	100.0	

Included in the statistics are 34.2% multiple primaries in males and 27.3% in females.




Table 14

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2014 (incl. multiple primaries)

			Males		Females		Males	Females
Age at			Age-		Age-			Prop.all
death	Males	Females	spec.		spec.		cancers	cancers
Years	n	n	mortal.	MI-index	mortal.	MI-index	90	90
0 - 4			0.0		0.0			
5- 9			0.0		0.0			
10-14			0.0		0.0			
15-19			0.0		0.0			
20-24			0.0		0.0			
25-29	1		0.1	0.25	0.0		1.6	
30-34		1	0.0		0.1	0.50		0.9
35-39			0.0		0.0			
40 - 44	1		0.1	0.20	0.0		0.2	
45-49	2	1	0.1	0.25	0.1	0.50	0.2	0.1
50-54			0.0		0.0			
55-59	3 /		0.3	3.00	0.0		0.1	
60-64	2	/1	0.2	0.67	0.1	0.33	0.0	0.0
65-69	2		0.2	0.67	0.0		0.0	
70-74	3	\ 1	0.3	0.43	0.1	1.00	0.0	0.0
75-79	7	2	1.3	0.88	0.3	0.67	0.1	0.0
80-84		3	0.0		0.5	0.50		0.0
85+	4	5	1.7	0.80	0.9	0.45	0.1	0.1
All ages	25	14					0.1	0.0
Mortality /								
Raw			0.1	0.34	0.1	0.32		
WS			0.1	0.22	0.0	0.19		
ES			0.1	0.30	0.0	0.23		
BRD-S			0.1	0.33	0.1	0.28		
DIAD 5			0.1	0.33	0.1	0.20		
PYLL-70								
per 100,000	)		1.1		0.4			
ES			0.9		0.4			
AYLL-70			15.7		22.5			

The rates underestimate the prognosis if other synchronous cancers are prognostic unfavorable.

Table 15a

Multiple primaries in deaths in period 1998-2014

MALES

Multiple pr	ımarıes	ın dea MAI		n perio	d 1998-	-2014		
Diagnosis	Total	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C12 C12 Hereanhammer	1	5.6	1	100.0				
C12-C13 Hypopharynx C16 Stomach C18 Colon	1 1	5.6 5.6	1	100.0			1	100.0
C44 Skin others C61 Prostate	2 2	11.1	1	50.0	1	50.0	2	100.0
C64 Kidney C70-C72 CNS cancer C81 Hodgkin lymphoma	2 1 1	11.1 5.6 5.6	2	100.0			1	100.0
C90 Mult. myeloma C91-C96 Leukaemia	1 5	5.6 27.8			1 2	100.0	3	60.0
C96 Systemic	1	5.6					1	100.0
All mult. primaries	18	100.0	6	33.3	4	22.2	8	44.4

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 15b

Multiple primaries in deaths in period 1998-2014
FEMALES

Multiple p	rimaries		aths ALES	in peri	od 1998	-2014		
Diagnosis	Total n	Total %↓	Pre n	Pre ←%	Syn- chron ±30d n	Syn- chron ±30d ←%	Post n	Post ←%
C17 Small intestine C19-C20 Rectum	1 1	12.5 12.5	1 1	100.0				
C50 Breast C54 Corpus uteri	1 1	12.5 12.5 12.5	1	100.0			1	100.0
C56 Ovary C91-C96 Leukaemia	1 2	12.5 25.0					1 2	100.0
C96 Systemic	1	12.5					1	100.0

ICD-10 C44 (Other malignant neoplasms of skin) is not systematically recorded by MCR and therefore not considered for evaluation as a particular primary but at least as a multiple malignancy.

Table 16

Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2014

(First primaries only \*)

Age at death Years	Males Fema	/ = /	MI-index	Females Age- spec. mortal.	MI-index	cancers	Females Prop.all cancers
0- 4 5- 9 10-14 15-19 20-24 25-29	1	0.0 0.0 0.0 0.0 0.0 0.0	0.25	0.0 0.0 0.0 0.0 0.0		1.8	
30-34 35-39 40-44 45-49 50-54 55-59	1	1 0.0 0.0 0.1 1 0.1 0.0 0.2	0.20 0.25	0.1 0.0 0.0 0.1 0.0	0.50	0.2 0.2 0.1	0.1
60-64 65-69 70-74 75-79 80-84 85+	2 1 2 5	1 0.2 0.1 1 0.2 0.9 3 0.0 5 0.9	1.00 1.00 0.40 0.83	0.1 0.0 0.1 0.0 0.5	0.50 1.00 0.60 0.63	0.1 0.0 0.0 0.1	0.0 0.0 0.1 0.1
All ages	18 1:		1.00	0.9	0.03	0.0	0.0
Mortality Raw WS ES BRD-S		0.1 0.1 0.1 0.1	0.19	0.1 0.0 0.0 0.0	0.34 0.19 0.23 0.28		
PYLL-70 per 100,000 ES AYLL-70		1.0 0.9 17.5		0.4 0.4 22.5			

<sup>\*</sup> See corresponding tables with multiple primaries.

Table 17


Age-specific mortality (cancer-related) and proportion of all cancers for period 2007-2014

(Single primaries only \*)

	Males		Females		Males	Females
Age at	Age-		Age-			Prop.all
death Males Females	/ = /		spec.		cancers	cancers
Years n n		MI-index		MI-index	୬	%
0- 4	0.0		0.0			
5- 9	0.0		0.0			
10-14	0.0		0.0			
15-19	0.0		0.0			
20-24	0.0		0.0			
25-29 1	0.1	0.25	0.0		2.0	
30-34	0.0		0.1	0.50		1.2
35-39	0.0		0.0			
40-44 1	0.1	0.20	0.0		0.3	
45-49 2 1	0.1	0.25	0.1	0.50	0.2	0.1
50-54	0.0		0.0			
55-59 1	0.1	1.00	0.0		0.0	
60-64 2	0.2	1.00	0.0		0.1	
65-69	0.0		0.0			
70-74	0.1	0.25	0.0		0.0	
75-79 3	0.5	0.60	0.0		0.1	
80-84	0.0		0.5	0.60		0.1
85+ 1 4	0.4	0.50	0.7	0.57	0.0	0.1
All ages 12 9					0.0	0.0
-						
Mortality	0 1	0.00	0 0	0.07		
Raw	0.1	0.22	0.0	0.27		
WS	0.0	0.14	0.0	0.14		
ES	0.1	0.19	0.0	0.17		
BRD-S	0.1	0.21	0.0	0.22		
PYLL-70						
per 100,000	0.9		0.4			
ES	0.8		0.3			
AYLL-70	20.4		30.0			

<sup>\*</sup> See corresponding tables with multiple primaries.

CD-10 C96: Other and unspecified malignant neoplasms of lymphoid, haematopoietic and related tissue Age distribution and age-specific mortality 2007 - 2014 (Males: 25, Females: 14)




**Figure 18.** Distribution of age at death (bars) and age-specific mortality (all patients: solid line, patients with single primaries: dotted line). The age-specific incidence is additionally plotted for comparison (dashed line).


The difference between age at diagnosis (Table 3) and age at systemic Disease NOS-related death (see Table 10) should be considered.



### Average mortality (world standard population) 2007 - 2014: Males




#### Average mortality (world standard population) 2007 - 2014: Females




**Figure 19a.** Map of cancer mortality (world standard population) by county averaged for period 2007 to 2014. According to their individual mortality rates, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population mean (males 0.1/100,000 WS N=23, females 0.0/100,000 WS N=14).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 65,347 female residents (averaged) in the period from 2007 to 2014 a total of 0 women died from systemic Disease NOS. Therefore, the mean mortality rate for this cancer type in this area can be calculated at 0.0/100,000 (world standard population). Though, the value of this parameter may vary with an underlying probability of 99% between 0.0 and 1.0/100,000.

### Standardized mortality ratio (SMR) 2007 - 2014: Males



#### Standardized mortality ratio (SMR) 2007 - 2014: Females



**Figure 19b.** Map of standardized mortality ratio (SMR, incl. DCO cases) by county averaged for period 2007 to 2014. According to their individual SMR values, the counties are displayed in different red and blue color temperatures where the fine white color indicates the population overall of 1.0 (males N=23, females N=14).

The results should be interpreted with caution! E.g., in county Ebersberg with a population of 64,924 female residents (averaged) in the period from 2007 to 2014 a total of 0 women died from systemic Disease NOS. Therefore, the mean standardized mortality ratio (SMR) for this cancer type in this area can be calculated at 0.00. Though, the value of this parameter may vary with an underlying probability of 99% between 0.00 and 15.06, and is therefore not statistically striking.

#### Statistical Notes

In all tables and figures the respective reference values should be carefully considered. The incidence rates include diagnoses (with multiple primary), and death certificate only (DCO) cases, where applicable. For mortality statistics patients, diagnoses and progressive course of disease are presented. In the calculations, all courses of disease are considered whereby progressions occurred and/or death certificate identified progressive cancers were ascertained. Additionally there are three groups of disease course to consider:

#### 1. All multiple primaries included

The mortality statistic describes the tumor-specific death, independent of any malignancy. The patient perspective, induced secondary malignancies, and the problem of multiple malignancies from the same primary tumor all have reasons for their inclusion.

#### 2. First singular primary (no information about other prior or synchronous malignancy)

The mortality statistic describes the cancer-related death for patients who have no therapeutic restrictions due to a previous or synchronous cancer. These statistics are comparable to studies that have exclusion criteria based on a second malignancy.

### **3. Single primary** (no information about other prior, syn- or metachronous malignancy)

The mortality statistic describes the tumor-specific death that occurs without any impact through secondary primaries, earlier, synchronous, later or induced. Precisely the difference between disease group 1 and 2 highlight the magnitude of the problem of secondary malignancies.

For this reason differences appear concerning official mono-causal mortality statistics. To judge the maximum deviation, 2 further tables are presented. In the first table the distribution of secondary malignancies before, at or after the described cancer are shown, that could be an alternative cause of death. In the second table, the age-specific mortality rates for all courses of disease, without designation of secondary malignancies are shown.

A previously minimally acknowledged statistic is the **age at death**, which allows for a good assessment of the quality of classification of the apparent tumor-specific death. For assumed tumor-independent deaths, the age of death should be estimated from the age of diagnosis and the normal life expectancy, whereas tumor-dependent deaths can be estimated from the age of diagnosis plus the average tumor-specific life expectancy. The comparison of different tumors demonstrates this association, if the causes of cancer and the competing cause of death are independent of each other (e.g. breast and colon versus head/neck and lung).

The index from mortality and incidence (Mortality-Incidence ratio, **MI-index**) is a statistic that allows for the evaluation of the quality of data. For diseases with poor prognoses, comparable values are obtained from all age groups, because to a large extent, the numerator and denominator contain the same cases. For tumors with a good prognosis, increasing and decreasing incidence and age-specific differences in prognosis can more strongly alter the MI- index. Additionally, attention should be paid to the confidence intervals where fewer cases are reported.

The complexity of problems identified here emphasizes the importance of relative survival data for the appropriate analysis of long term results.

As a measurement of the burden of disease, the number of potential life years loss due to premature deaths in a cohort can be calculated (**PYLL**, potential years of life lost, standardized per 100,000 persons or per European standard) as well as the average loss of life years per individual (**AYLL**, average years of life lost). Depending upon the analytic aim (health economy, prevention, health care research) different methods exist for the generation of these measurements. In the results presented here, the age for a premature death is considered to be before 70 years, according to the guidelines of the OECD and the WHO (as seen in the abbreviation PYLL-70 or AYLL-70).

#### **Shortcuts**

FRG Federal Republic of Germany

GEKID Association of Population-based Cancer Registries in Germany

(Gesellschaft der epidemiologischen Krebsregister in Deutschland e.V.)

MCR Munich Cancer Registry (Tumorregister München)
SEER Surveillance, Epidemiology, and End Results (USA)

AYLL-70 Average years of life lost prior to age 70 given a person dies before that age

BRD-S German standard population

DCO Death certificate only EAR Excess absolute risk

= excess cancer cases (O - E) per 10,000 person-years

ES European standard population (old)

LCL Lower confidence limit

MI-index Ratio between mortality and incidence

PYLL-70 Potential years of life lost prior to age 70 given a person dies before that age

SIR Standardized incidence ratio SMR Standardized mortality ratio UCL Upper confidence limit WS World standard population

#### **Recommended Citation**

Munich Cancer Registry. ICD-10 C96: Systemic Disease NOS - Incidence and Mortality [Internet]. 2016 [updated 2016 Apr 13; cited 2016 Jun 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/base/bC96\_\_E-ICD-10-C96-Systemic-Disease-NOS-incidence-and-mortality.pdf

#### Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

#### **Disclaimer**

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

## Index of figures and tables

Fig./Tb	l.	Page
1	Pts cohorts, DCO, mult. prim., follow-up / yr	3
1a	Gender distribution by year of diagnosis	4
2	Incidence by year of diagnosis	5
3	Age distribution parameters by year of diagnosis	6
4	Age distribution by 5-year age group and gender	8
5	Age-specific incidence, DCO rate, proportion malignancies	9
6	Age distribution and age-specific incidence (chart)	10
6a	Age-specific incidence (chart)	11
7	Cumulative follow-up years (chart)	11
8	Standardized incidence ratio of second primaries	12
9a	Map of cancer incidence (WS) by county (chart)	13
9b	Standardized incidence ratio (SIR) by county (chart)	14
10a	Pts incident cohorts and mortality / yr	15
10b	Incidence and mortality by year of diagnosis	16
10c	Cancer-related deaths, death certification available / yr	17
11	Medians of age at death / yr	18
12	Mortality by year of death	20
13	Distribution of age at death	21
14	Age-specific mortality	22
15	Multiple primaries in deaths	23
16	Age-specific mortality (first primaries)	25
17	Age-specific mortality (single primaries)	26
18	Age distribution and age-specific mortality (chart)	27
19a	Map of cancer mortality (WS) by county (chart)	28
19b	Standardized mortality ratio (SMR) by county (chart)	29