

Munich Cancer Registry



- Incidence and Mortality
- Selection Matrix
- Homepage
- *Deutsch*

ICD-10 C09-C14: Pharynx cancer

Survival

Year of diagnosis	1988-1997	1998-2020
Patients	1,381	5,878
Diseases	1,401	6,008
Cases evaluated	1,239	4,579
Creation date	04/15/2022	
Database export	12/20/2021	
Population	4.92 m	



Munich Cancer Registry
Cancer Registry Bavaria - Upper Bavaria Regional Center
at Klinikum Grosshadern/IBE
Marchioninstr. 15
Munich, 81377
Germany

<https://www.tumorregister-muenchen.de/en>

<https://www.tumorregister-muenchen.de/en/facts/surv/sC0914E-ICD-10-C09-C14-Pharynx-cancer-survival.pdf>

Index of figures and tables

Fig./Tbl.	Page
1a Relative survival by period of diagnosis (chart)	3
1b Survival by period of diagnosis (table)	3
2a Survival by sex (chart)	4
2b Survival by sex (table)	4
3a Observed survival by age category (chart)	5
3b Relative survival by age category (chart)	5
3c Survival by age category (table)	6
4a Relative survival by extent of disease (chart)	7
4b Survival by extent of disease (table)	7
4c Conditional survival by extent of disease (chart)	8
4d Conditional survival by extent of disease (table)	8
5a Time to first progression (chart)	9
5b Time to first progression (table)	9
5c Observed post-progression survival (chart)	11
5d Observed post-progression survival (table)	11
5e Observed post-progression survival by period of progression (chart)	12
5f Observed post-progression survival by period of progression (table)	12




Figure 1a. Relative survival of patients with pharynx cancer by period of diagnosis. Included in the evaluation are 5,808 cases diagnosed between 1988 and 2020.

The survival results of the SEER program (Surveillance, Epidemiology, and End Results) of the American National Cancer Institute (NCI) are summarized as the period of diagnosis from 2000 to 2018, and are represented by colored diamonds in order to facilitate comparisons between MCR and SEER.

The presented survival curves are derived from clinical records with valid follow-up informations, which means that death certificate cases (DCO) cases are omitted from the analysis. With this one restriction, the MCR has provided population-based statistics since 1998, collecting data on all cancer cases in the region of southern Bavaria. Historical data of previous time periods can be heavily selected, therefore, univariate survival comparisons of the presented time periods must be carefully considered. Nonetheless, all calculable survival curves are depicted to facilitate the comparison of long time follow-up analyses of relative survival between particular cancers.

Years	Period of diagnosis							
	1988-1997 n=1,235		1998-2006 n=1,781		2007-2015 n=2,220		2016+ n=572	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	71.8	72.8	72.6	73.6	76.2	77.4	79.4	80.8
2	53.1	54.6	56.3	57.8	62.4	64.3	69.2	71.6
3	44.0	45.9	46.3	48.1	54.9	57.4	61.4	64.7
4	38.2	40.5	41.5	43.8	49.8	52.8	54.0	57.9
5	34.8	37.4	38.2	40.8	46.4	50.1	51.4	55.8
6	31.1	33.9	35.4	38.4	42.8	47.0		
7	28.4	31.5	33.0	36.3	39.6	44.1		
8	26.1	29.4	31.3	35.0	37.6	42.7		
9	24.5	28.0	28.9	32.8	34.9	40.4		
10	22.8	26.4	27.4	31.6	32.8	38.8		
11	20.9	24.6	25.6	30.0	29.9	36.0		
12	19.4	23.2	23.6	28.2	29.1	35.9		
13	17.8	21.7	21.9	26.7	26.6	33.5		
14	16.9	20.9	20.2	25.2				
15	14.9	18.9	18.9	24.1				
Median	2.3		2.6		3.9			

Table 1b. Observed (obs.) and relative (rel.) survival of patients with pharynx cancer by period of diagnosis for period 1988-2020 (N=5,808).




Figure 2a. Survival of patients with pharynx cancer by sex. Included in the evaluation are 4,573 cases diagnosed between 1998 and 2020.

Years	Sex			
	Male n=3,638		Female n=935	
	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0
1	73.9	75.0	80.5	81.5
2	58.3	60.1	70.6	72.2
3	49.8	52.1	61.9	64.2
4	44.5	47.4	57.0	59.8
5	41.0	44.4	54.3	57.7
6	37.9	41.7	50.4	54.2
7	35.1	39.3	47.0	51.2
8	33.2	37.8	45.4	50.0
9	30.7	35.7	42.0	47.0
10	29.0	34.3	39.7	44.9
11	26.8	32.4	36.6	42.1
12	25.1	31.0	34.6	40.3
13	23.1	29.2	32.3	38.1
14	21.4	27.7	30.3	36.4
15	19.9	26.3	29.1	35.3
Median	3.0		6.2	

Table 2b. Observed (obs.) and relative (rel.) survival of patients with pharynx cancer by sex for period 1998-2020 (N=4,573).




Figure 3a. Observed survival of patients with pharynx cancer by age category. Included in the evaluation are 4,573 cases diagnosed between 1998 and 2020.




Figure 3b. Relative survival of patients with pharynx cancer by age category. Included in the evaluation are 4,573 cases diagnosed between 1998 and 2020.

Years	Age category							
	0-49 yrs n=667		50-59 yrs n=1,592		60-69 yrs n=1,522		70+ yrs n=792	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	84.0	84.2	78.1	78.5	74.3	75.3	64.0	67.0
2	72.8	73.1	60.9	61.7	60.9	62.6	50.5	55.4
3	63.6	64.1	52.3	53.4	52.9	55.3	41.3	47.4
4	58.9	59.5	47.7	48.9	47.0	49.9	36.0	43.5
5	56.2	57.0	44.3	45.9	43.8	47.4	31.8	40.6
6	53.3	54.1	41.1	43.0	40.3	44.4	28.2	37.9
7	51.4	52.5	38.3	40.4	37.3	41.9	24.0	34.4
8	49.8	51.0	36.9	39.3	35.2	40.5	21.1	32.1
9	48.3	49.7	34.0	36.6	32.4	38.3	18.1	29.4
10	46.9	48.4	32.8	35.8	29.5	35.6	16.5	28.9
11	44.5	46.1	30.3	33.5	26.8	33.3	15.0	28.3
12	41.9	43.9	29.0	32.5	24.7	31.6	13.7	28.0
13	40.1	42.2	27.3	30.9	21.6	28.5	12.5	27.9
14	38.0	40.4	25.1	29.0	20.1	27.6		
15	36.5	38.9	23.8	27.9	18.4	26.2		
Median	7.8		3.4		3.4		2.1	

Table 3c. Observed (obs.) and relative (rel.) survival of patients with pharynx cancer by age category for period 1998-2020 (N=4,573).




Figure 4a. Relative survival of patients with pharynx cancer by extent of disease. For 4,329 of 4,573 cases diagnosed between 1998 and 2020 valid data could be obtained for this item. For a total of 4,321 cases an evaluable classification was established. The grey line represents the subgroup of 252 patients with missing values regarding extent of disease (5.5 % of 4,573 patients, the percent values of all other categories are related to n=4,321).

Years	Extent of disease							
	Local n=1,035		Regional n=3,009		Distant n=277		NA/NOS n=252	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	86.2	87.5	76.7	77.7	40.0	40.2	51.6	52.7
2	74.8	77.1	60.9	62.7	26.8	27.4	39.2	40.9
3	66.9	69.8	51.8	54.0	19.8	20.7	33.4	35.4
4	62.1	65.7	46.5	49.2	15.7	16.4	27.1	29.0
5	59.1	63.6	42.9	46.2	13.5	14.6	23.8	26.4
6	54.4	59.3	39.8	43.5	11.7	12.8	22.0	24.4
7	50.2	55.7	37.0	41.1	11.2	12.3	20.0	22.8
8	48.1	54.4	35.1	39.6			18.4	21.0
9	44.8	51.5	32.3	37.1			17.8	19.7
10	41.1	48.2	31.1	36.3			16.6	19.6
Median	7.0		3.2		0.7		1.1	

Table 4b. Observed (obs.) and relative (rel.) survival of patients with pharynx cancer by extent of disease for period 1998-2020 (N=4,321).




Figure 4c. Conditional relative 2-year survival of patients with pharynx cancer by extent of disease. For 4,329 of 4,573 cases diagnosed between 1998 and 2020 valid data could be obtained for this item. For a total of 4,321 cases an evaluable classification was established. The grey line represents the subgroup of 252 patients with missing values regarding extent of disease (5.5 % of 4,573 patients, the percent values of all other categories are related to n=4,321).

Extent of disease								
Years	Local		Regional		Distant		NA/NOS	
	n	Cond. surv. % 2 yrs	n	Cond. surv. % 2 yrs	n	Cond. surv. % 2 yrs	n	Cond. surv. % 2 yrs
0	1,035	77.1	3,009	62.7	277	27.4	252	40.9
1	868	79.8	2,271	69.5	108	51.0	125	66.9
2	737	85.4	1,768	78.6	70	59.6	95	70.9
3	641	91.2	1,459	85.5	49	70.5	75	74.5
4	568	90.3	1,265	88.6	36	76.9	60	83.3
5	524	87.8	1,099	89.3	31	84.4	51	87.1

Table 4d. Conditional relative 2-year survival of patients with pharynx cancer by extent of disease for period 1998-2020 (N=4,321).

Conditional relative survival rates refer to the relative survival probability, in this case for 2 years after cancer diagnosis, compared to the age- and sex-matched population (=100 %) under the condition of being alive for a certain time period (x-axis in Figure 4a). The results illustrate to what extent the cancer induced mortality of particular subgroups declines in the subsequent years after detection of the malignancy. For instance, according to the presented survival statistics, patients in the subgroup extent of disease="Local", who are alive at least 3 years after cancer diagnosis, the conditional relative 2-year survival rate is 91.2% (n=641).




Figure 5a. Time to first progression of 4,563 patients with pharynx cancer diagnosed between 1998 and 2020 (in solid cancers M0 only) estimated by cumulative incidence function (CI, solid line) accounting for death as competing risk and by inverse Kaplan-Meier estimate (1-KM, dashed line). The frequency of events may be underestimated due to underreporting.

		Type of progression						Distant metastasis (CI)
		Any type (CI)	Any type (1-KM)	Local relapse (CI)	Local relapse (1-KM)	Lymph node (CI)	Lymph node (1-KM)	
N	Events	4,302	4,302	4,563	4,563	4,563	4,563	4,305
compet.	Years	2,379	2,379	651	651	259	259	1,011
		728		2,529		2,871		1,910
		%	%	%	%	%	%	%
0		0.0	0.0	0.0	0.0	0.0	0.0	0.0
1		33.3	34.1	5.6	6.8	2.8	3.3	10.1
2		44.3	45.9	10.0	13.0	4.3	5.5	17.3
3		48.8	50.8	11.4	15.3	5.0	6.7	20.5
4		51.3	53.7	12.5	17.3	5.2	7.1	21.8
5		52.6	55.2	13.0	18.2	5.4	7.5	22.6
6		53.8	56.6	13.5	19.3	5.5	7.7	23.1
7		54.6	57.6	13.9	20.1	5.7	8.2	23.4
8		55.2	58.5	14.1	20.6	5.7	8.3	23.7
9		55.9	59.5	14.4	21.4	5.8	8.6	24.0
10		56.5	60.2	14.8	22.3	5.9	8.8	24.0
11		57.0	61.0	15.0	22.9	5.9	8.9	24.2
12		57.2	61.4	15.2	23.4	6.0	9.1	24.5
13		57.7	62.2	15.3	24.0	6.0	9.1	24.7
14		58.2	63.1	15.5	24.6	6.0	9.3	24.8
15		58.6	63.8	15.7	25.1	6.0	9.3	25.0

<i>cont'd</i>	Type of progression		
	Distant metast. (1- KM)	NOS (CI)	NOS (1-KM)
N	4,305	4,560	4,560
Events	1,011	1,183	1,183
compet.		2,028	
Years	%	%	%
0	0.0	0.0	0.0
1	11.3	18.6	19.8
2	20.4	22.1	24.2
3	24.8	23.3	26.0
4	26.8	24.2	27.4
5	28.1	24.7	28.3
6	29.0	25.2	29.1
7	29.4	25.5	29.8
8	30.0	25.8	30.5
9	30.6	26.1	31.2
10	30.7	26.3	31.6
11	31.2	26.6	32.2
12	31.9	26.6	32.4
13	32.3	26.9	33.1
14	32.7	27.2	34.1
15	33.1	27.4	34.7

Table 5b. Time to first progression of patients with pharynx cancer for period 1998-2020 (N=4,563), also showing the total of progression events (Events) and of deaths as competing risk (compet.).




Figure 5c. Observed post-progression survival of 1,771 patients with pharynx cancer diagnosed between 1998 and 2020. These 1,771 patients with documented progression events during their course of disease represent 38.8 % of the totally 4,563 evaluated cases (incl. M1, n=261, 5.7 %). Patients with cancer relapse documented via death certificates only were excluded (n=885, 19.4 %). Multiple progression types on different sites are included in the evaluation even when not occurring synchronously. The NOS (not otherwise specified) class is included under the condition, that it is the one and only progression type during the course of disease.

Medical record documentation often lacks the linguistic severity to distinguish between local relapse, regional lymph node metastasis and distant spread in solid cancers. Frequently, the statement “not specified” is the only information in registries regarding relapse of the disease. The category “Any type” denotes all cases who suffered from at least one relapse during the course of disease (incl. primary M1-status). Although, the real number of relapsed patients is likely to be much higher. The accumulated percentage of patients with local relapse or distant metastasis exceeds the 100 % value because patients are potentially considered in more than one subgroup.

Years	Type of progression				
	Any type n=1,771 %	Local relapse n=657 %	Lymph node n=257 %	Distant metastasis n=865 %	NOS n=570 %
0	100.0	100.0	100.0	100.0	100.0
1	48.1	45.7	43.9	34.9	59.2
2	31.3	27.3	27.5	20.4	41.9
3	25.1	20.1	23.6	15.0	37.1
4	21.1	17.5	17.7	11.8	32.7
5	18.5	15.9	14.0	10.2	28.2
6	16.4	13.3	14.0	8.6	26.0
7	14.8	11.7	12.8	7.7	23.1
8	13.8	9.8		7.3	22.0
9	12.6	8.9		7.3	18.9
10	11.3	8.0		6.4	17.6

Table 5d. Observed post-progression survival of patients with pharynx cancer for period 1998-2020 (N=1,771).




Figure 5e. Observed post-progression (distant metastasis) survival of 1,027 patients with pharynx cancer diagnosed between 1988 and 2020 by period of progression.

Years	Period of progression			
	1988-1997 n=111 %	1998-2006 n=328 %	2007-2015 n=443 %	2016+ n=145 %
0	100.0	100.0	100.0	100.0
1	24.6	27.6	36.5	46.6
2		14.1	21.8	29.6
3		8.6	16.3	
4		6.4	13.2	
5			11.1	
6			9.0	
7			7.7	
8			7.1	
9			7.1	

Table 5f. Observed post-progression (distant metastasis) survival of patients with pharynx cancer for period 1988-2020 by period of progression (N=1,027).

Shortcuts

MCR Munich Cancer Registry, Germany

NCI National Cancer Institute, USA

SEER Surveillance, Epidemiology, and End Results, USA

UICC Union for International Cancer Control, Geneva

DCO Death certificate only Death certificate provides the only notification to the registry.

NA Not available

NOS Not otherwise specified

OS Overall/Observed survival Overall/Observed survival (Kaplan-Meier estimate)

Date of entry: diagnosis

Event: death from any cause

RS Relative survival

Survival compared to “general population”, ratio of observed to expected survival (Ederer II method), reflecting cancer specific survival

AS Assembled survival

Assembled chart of observed, expected, relative survival

CS Conditional survival

Survival probability under the condition of surviving a given period of time

TTP Time to progression

Time to first progression / relapse

Date of entry: diagnosis

Event: (progression / relapse): first local-, lymph node recurrence, distant metastasis or unspecified progression

1-KM

1 minus Kaplan-Meier estimator (“inverse” Kaplan-Meier estimator)

CI

Cumulative incidence

Death as competing risk (according to Kalbfleisch und Prentice)

PPS Post-progression survival

Survival since first progression / relapse (Kaplan-Meier estimate)

Date of entry (progression / relapse): first local-, lymph node recurrence, distant metastasis or unspecified progression

Event: death from any cause

Recommended Citation

Munich Cancer Registry. Survival ICD-10 C09-C14: Pharynx cancer [Internet]. 2022 [updated 2022 Apr 15; cited 2022 Jun 1]. Available from: <https://www.tumorregister-muenchen.de/en/facts/surv/sC0914E-ICD-10-C09-C14-Pharynx-cancer-survival.pdf>

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.