

# Munich Cancer Registry



- ▶ Incidence and Mortality
- ▶ Selection Matrix
- ▶ Homepage
- ▶ *Deutsch*

## ICD-10 C25: Pancreas cancer

### Survival

Year of diagnosis	1988-1997	1998-2019
Patients	868	16,391
Diseases	868	16,394
Cases evaluated	773	9,712
Creation date	01/27/2021	
Database export	01/07/2021	
Population	4.92 m	




Munich Cancer Registry  
Cancer Registry Bavaria - Upper Bavaria Regional Center  
at Klinikum Grosshadern/IBE  
Marchioninstr. 15  
Munich, 81377  
Germany

<https://www.tumorregister-muenchen.de/en>

[https://www.tumorregister-muenchen.de/en/facts/surv/sC25\\_\\_E-ICD-10-C25-Pancreas-cancer-survival.pdf](https://www.tumorregister-muenchen.de/en/facts/surv/sC25__E-ICD-10-C25-Pancreas-cancer-survival.pdf)

## Index of figures and tables

Fig./Tbl.	Page
1a Relative survival by period of diagnosis (chart)	3
1b Survival by period of diagnosis (table)	3
2a Survival by sex (chart)	4
2b Survival by sex (table)	4
3a Observed survival by age category (chart)	5
3b Relative survival by age category (chart)	5
3c Survival by age category (table)	6
4a Relative survival by TNM staging (chart)	7
4b Survival by TNM staging (table)	7
4c Conditional survival by extent of disease (chart)	8
4d Conditional survival by extent of disease (table)	8
5a Time to first progression (chart)	9
5b Time to first progression (table)	9
5c Observed post-progression survival (chart)	11
5d Observed post-progression survival (table)	11
5e Observed post-progression survival by period of progression (chart)	12
5f Observed post-progression survival by period of progression (table)	12




**Figure 1a.** Relative survival of patients with pancreas cancer by period of diagnosis. Included in the evaluation are 10,132 cases diagnosed between 1988 and 2019.

The survival results of the SEER program (Surveillance, Epidemiology, and End Results) of the American National Cancer Institute (NCI) are summarized as the period of diagnosis from 2000 to 2017, and are represented by colored diamonds in order to facilitate comparisons between MCR and SEER.

The presented survival curves are derived from clinical records with valid follow-up informations, which means that death certificate cases (DCO) cases are omitted from the analysis. With this one restriction, the MCR has provided population-based statistics since 1998, collecting data on all cancer cases in the region of southern Bavaria. Historical data of previous time periods can be heavily selected, therefore, univariate survival comparisons of the presented time periods must be carefully considered. Nonetheless, all calculable survival curves are depicted to facilitate the comparison of long time follow-up analyses of relative survival between particular cancers.

Years	Period of diagnosis					
	1988-1997 n=769		1998-2006 n=2,658		2007+ n=6,705	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0
1	31.4	32.2	33.2	34.1	40.8	42.0
2	16.4	17.1	16.2	17.1	23.4	24.5
3	12.0	12.8	11.4	12.2	16.5	17.7
4	9.8	10.7	9.0	9.9	13.5	14.8
5	8.2	9.1	7.7	8.6	11.7	13.1
6	7.7	8.8	6.8	7.7	10.6	12.2
7	7.2	8.3	6.0	7.0	9.8	11.5
8	6.7	7.9	5.4	6.5	9.2	11.0
9	6.0	7.4	5.0	6.2	8.9	10.9
10	5.7	7.1	4.7	5.9	8.5	10.7
Median	0.6		0.6		0.7	


**Table 1b.** Observed (obs.) and relative (rel.) survival of patients with pancreas cancer by period of diagnosis for period 1988-2019 (N=10,132).




**Figure 2a.** Survival of patients with pancreas cancer by sex. Included in the evaluation are 9,363 cases diagnosed between 1998 and 2019.

Years	Sex			
	Male n=4,853		Female n=4,510	
	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0
1	39.1	40.1	38.1	39.1
2	21.3	22.5	21.2	22.1
3	14.8	16.1	15.1	16.1
4	12.0	13.4	12.2	13.3
5	10.5	11.9	10.4	11.5
6	9.4	11.0	9.4	10.6
7	8.5	10.2	8.6	9.9
8	7.7	9.5	8.0	9.4
9	7.4	9.3	7.6	9.1
10	7.0	9.1	7.0	8.6
11	6.6	8.9	6.6	8.3
12	6.4	8.8	6.6	8.5
13	6.0	8.6	6.6	8.6
14	5.7	8.3	6.4	8.8
15			6.0	8.4
Median	0.7		0.7	

**Table 2b.** Observed (obs.) and relative (rel.) survival of patients with pancreas cancer by sex for period 1998-2019 (N=9,363).




**Figure 3a.** Observed survival of patients with pancreas cancer by age category. Included in the evaluation are 9,363 cases diagnosed between 1998 and 2019.



**Figure 3b.** Relative survival of patients with pancreas cancer by age category. Included in the evaluation are 9,363 cases diagnosed between 1998 and 2019.

Years	Age category										
	0-49 yrs n=563		50-59 yrs n=1,336		60-69 yrs n=2,687		70-79 yrs n=3,326		80+ yrs n=1,451		
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	58.1	58.2	50.1	50.3	43.4	43.9	35.7	36.6	18.2	20.0	
2	38.7	38.8	26.6	26.9	24.6	25.2	18.7	19.8	9.0	10.7	
3	33.0	33.0	18.0	18.3	17.4	18.1	13.0	14.2	5.1	6.6	
4	29.4	29.4	15.5	15.8	13.7	14.5	10.3	11.7	3.6	5.2	
5	28.2	28.3	12.9	13.3	12.0	12.9	8.4	9.9	3.0	4.9	
6	26.9	27.0	12.0	12.4	10.7	11.7	7.3	9.1			
7	25.9	26.0	11.0	11.5	9.9	11.0	6.6	8.6			
8	24.2	24.6	10.2	10.8	9.0	10.3	6.0	8.1			
9	24.2	24.2	9.7	10.2	8.8	10.2	5.4	7.8			
10	23.5	24.0	9.5	10.2	8.0	9.6	5.0	7.7			
Median	1.3		1.0		0.8		0.6		0.3		

**Table 3c.** Observed (obs.) and relative (rel.) survival of patients with pancreas cancer by age category for period 1998-2019 (N=9,363).




**Figure 4a.** Relative survival of patients with pancreas cancer by TNM staging. For 8,116 of 9,363 cases diagnosed between 1998 and 2019 valid data could be obtained for this item. For a total of 8,097 cases an evaluable classification was established. The accumulated percentage exceeds the 100 % value because patients are potentially considered in more than one subgroup. The grey line represents the subgroup of 1,266 patients with missing values regarding TNM staging (13.5 % of 9,363 patients, the percent values of all other categories are related to n=8,097).

Years	TNM staging													
	T1 n=241		T2 n=679		T3 n=2,838		T4 n=1,015		N0 n=1,329		N+ n=3,073		M0 n=2,164	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	81.1	82.2	61.7	62.8	58.4	59.6	32.8	33.6	71.6	73.0	50.8	51.9	67.9	69.3
2	71.9	74.1	42.1	43.8	33.8	35.2	12.3	12.8	51.6	53.8	26.3	27.4	41.8	43.6
3	65.0	68.4	35.0	37.2	22.8	24.4	7.0	7.4	41.3	43.8	16.9	18.0	30.1	32.1
4	61.2	65.2	30.2	32.3	17.6	19.3	4.9	5.3	34.5	37.4	12.8	14.0	24.2	26.4
5	57.7	62.7	27.7	30.3	14.5	16.2	3.9	4.3	30.7	34.0	10.3	11.5	20.5	22.8
Median	7.7		1.5		1.2		0.7		2.1		1.0		1.6	

Years	TNM staging			
	M1 n=4,615		NA/NOS n=1,266	
	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0
1	21.9	22.6	28.9	30.2
2	8.4	8.8	16.2	17.5
3	5.5	5.9	11.5	12.7
4	4.5	4.9	10.1	11.5
5	3.8	4.2	9.3	10.7
Median	0.4		0.5	

**Table 4b.** Observed (obs.) and relative (rel.) survival of patients with pancreas cancer by TNM staging for period 1998-2019 (N=8,097).




**Figure 4c.** Conditional relative 2-year survival of patients with pancreas cancer by extent of disease. For 8,116 of 9,363 cases diagnosed between 1998 and 2019 valid data could be obtained for this item. For a total of 8,053 cases an evaluable classification was established. The grey line represents the subgroup of 1,310 patients with missing values regarding extent of disease (14.0 % of 9,363 patients, the percent values of all other categories are related to n=8,053).

Years	Extent of disease							
	Local		Regional		Distant		NA/NOS	
	n	Cond. surv. % 2 yrs	n	Cond. surv. % 2 yrs	n	Cond. surv. % 2 yrs	n	Cond. surv. % 2 yrs
0	1,357	52.0	2,081	35.2	4,615	8.8	1,310	17.9
1	877	58.4	1,173	37.0	924	26.1	369	43.3
2	572	69.4	591	49.9	317	55.4	192	67.5
3	420	77.9	346	63.0	186	70.8	129	83.9
4	320	83.7	226	74.0	134	78.8	106	89.1
5	256	87.6	164	80.8	96	85.2	91	91.3

**Table 4d.** Conditional relative 2-year survival of patients with pancreas cancer by extent of disease for period 1998-2019 (N=8,053).

Conditional relative survival rates refer to the relative survival probability, in this case for 2 years after cancer diagnosis, compared to the age- and sex-matched population (=100 %) under the condition of being alive for a certain time period (x-axis in Figure 4a). The results illustrate to what extent the cancer induced mortality of particular subgroups declines in the subsequent years after detection of the malignancy. For instance, according to the presented survival statistics, patients in the subgroup extent of disease="Local", who are alive at least 3 years after cancer diagnosis, the conditional relative 2-year survival rate is 77.9% (n=420).






**Figure 5a.** Time to first progression of 9,329 patients with pancreas cancer diagnosed between 1998 and 2019 (in solid cancers M0 only) estimated by cumulative incidence function (CI, solid line) accounting for death as competing risk and by inverse Kaplan-Meier estimate (1-KM, dashed line). The frequency of events may be underestimated due to underreporting.

	Type of progression							
	Any type (CI)	Any type (1-KM)	Distant metastasis (CI)	Distant metast. (1-KM)	Locoreg. relapse (CI)	Locoreg. rel. (1-KM)	NOS (CI)	
N	4,806	4,806	4,808	4,808	9,329	9,329	9,327	
Events	3,663	3,663	2,244	2,244	539	539	1,625	
compet.	383		1,695		7,505		6,418	
Years	%	%	%	%	%	%	%	%
0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1	54.0	55.9	27.8	33.8	2.8	5.3	13.9	
2	71.4	74.9	41.5	54.5	5.0	12.5	16.9	
3	76.5	80.7	46.6	63.3	5.7	16.0	17.7	
4	79.0	83.6	49.0	67.9	6.0	18.4	18.0	
5	80.2	85.0	49.8	69.6	6.2	19.7	18.3	
6	81.0	86.1	50.4	71.1	6.3	20.8	18.4	
7	81.5	86.7	50.7	71.8	6.4	21.4	18.5	
8	81.9	87.2	51.0	72.5	6.4	22.1	18.6	
9	82.4	87.9	51.3	73.3	6.6	23.3	18.6	
10	82.6	88.2	51.6	73.9	6.6	24.1	18.6	
11	82.7	88.3	51.7	74.1	6.6	24.1	18.6	
12	82.7	88.3	51.7	74.1	6.6	24.1	18.6	
13	82.8	88.5	51.7	74.1	6.6	24.1	18.7	
14	82.8	88.5	51.7	74.1	6.6	24.1	18.7	
15	83.0	88.8	51.8	74.7	6.8	27.1	18.7	

Type of progression	
<i>cont'd</i>	NOS (1-KM)
N	9,327
Events	1,625
compet.	
Years	%
0	0.0
1	20.2
2	28.7
3	32.0
4	33.8
5	35.2
6	36.2
7	37.3
8	37.7
9	38.2
10	38.2
11	38.2
12	38.2
13	38.9
14	38.9
15	38.9

**Table 5b.** Time to first progression of patients with pancreas cancer for period 1998-2019 (N=9,329), also showing the total of progression events (Events) and of deaths as competing risk (compet.).




**Figure 5c.** Observed post-progression survival of 6,183 patients with pancreas cancer diagnosed between 1998 and 2019. These 6,183 patients with documented progression events during their course of disease represent 66.3 % of the totally 9,329 evaluated cases (incl. M1, n=4,523, 48.5 %). Patients with cancer relapse documented via death certificates only were excluded (n=2,005, 21.5 %). Multiple progression types on different sites are included in the evaluation even when not occurring synchronously. The NOS (not otherwise specified) class is included under the condition, that it is the one and only progression type during the course of disease.

Medical record documentation often lacks the linguistic severity to distinguish between local relapse, regional lymph node metastasis and distant spread in solid cancers. Frequently, the statement “not specified” is the only information in registries regarding relapse of the disease. The category “Any type” denotes all cases who suffered from at least one relapse during the course of disease (incl. primary M1-status). Although, the real number of relapsed patients is likely to be much higher. The accumulated percentage of patients with local relapse or distant metastasis exceeds the 100 % value because patients are potentially considered in more than one subgroup.

Years	Type of progression			
	Any type n=6,183 %	Distant metastasis n=5,621 %	Locoregional relapse n=538 %	NOS n=575 %
0	100.0	100.0	100.0	100.0
1	26.3	23.7	35.7	40.9
2	11.9	10.1	19.4	22.6
3	7.6	6.5	13.3	13.6
4	5.8	5.0	9.0	10.4
5	4.8	4.0	6.6	8.7
6	4.3	3.6		8.2
7	4.0	3.3		7.9
8	3.7	3.1		
9	3.4	2.9		
10	3.3	2.9		

**Table 5d.** Observed post-progression survival of patients with pancreas cancer for period 1998-2019 (N=6,183).



**Figure 5e.** Observed post-progression (distant metastasis) survival of 5,974 patients with pancreas cancer diagnosed between 1988 and 2019 by period of progression.

Years	Period of progression		
	1988-1997 n=323 %	1998-2006 n=1,470 %	2007+ n=4,181 %
0	100.0	100.0	100.0
1	15.3	18.0	25.9
2		6.5	11.5
3		4.2	7.5
4		3.0	5.8
5		2.5	4.7
6		2.1	4.4
7		1.8	4.1
8		1.6	3.9
9		1.5	3.8

**Table 5f.** Observed post-progression (distant metastasis) survival of patients with pancreas cancer for period 1988-2019 by period of progression (N=5,974).

## Shortcuts

MCR Munich Cancer Registry, Germany

NCI National Cancer Institute, USA

SEER Surveillance, Epidemiology, and End Results, USA

UICC Union for International Cancer Control, Geneva

DCO Death certificate only Death certificate provides the only notification to the registry.

NA Not available

NOS Not otherwise specified

OS Overall/Observed survival Overall/Observed survival (Kaplan-Meier estimate)  
Date of entry: diagnosis  
Event: death from any cause

RS Relative survival Survival compared to “general population”,  
ratio of observed to expected survival (Ederer II method),  
reflecting cancer specific survival

AS Assembled survival Assembled chart of  
observed, expected, relative survival

CS Conditional survival Survival probability under the condition of surviving  
a given period of time

TTP Time to progression Time to first progression / relapse  
Date of entry: diagnosis  
Event: (progression / relapse): first local-, lymph node recurrence,  
distant metastasis or unspecified progression

1-KM 1 minus Kaplan-Meier estimator  
 (“inverse” Kaplan-Meier estimator)

CI Cumulative incidence  
Death as competing risk (according to Kalbfleisch und Prentice)

PPS Post-progression survival Survival since first progression / relapse (Kaplan-Meier estimate)  
Date of entry (progression / relapse): first local-, lymph node  
recurrence, distant metastasis or unspecified progression  
Event: death from any cause

## Recommended Citation

Munich Cancer Registry. Survival ICD-10 C25: Pancreas cancer [Internet]. 2021 [updated 2021 Jan 27; cited 2021 Mar 1]. Available from: [https://www.tumorregister-muenchen.de/en/facts/surv/sC25\\_\\_E-ICD-10-C25-Pancreas-cancer-survival.pdf](https://www.tumorregister-muenchen.de/en/facts/surv/sC25__E-ICD-10-C25-Pancreas-cancer-survival.pdf)

## Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

## Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.