

# Munich Cancer Registry



- ▶ Incidence and Mortality
- ▶ Selection Matrix
- ▶ Homepage
- ▶ *Deutsch*

## ICD-10 C40, C41: Bone cancer

### Survival

|                   |            |           |
|-------------------|------------|-----------|
| Year of diagnosis | 1988-1997  | 1998-2019 |
| Patients          | 131        | 664       |
| Diseases          | 131        | 667       |
| Cases evaluated   | 126        | 522       |
| Creation date     | 01/27/2021 |           |
| Database export   | 01/07/2021 |           |
| Population        | 4.92 m     |           |




Munich Cancer Registry  
Cancer Registry Bavaria - Upper Bavaria Regional Center  
at Klinikum Grosshadern/IBE  
Marchioninstr. 15  
Munich, 81377  
Germany

<https://www.tumorregister-muenchen.de/en>

<https://www.tumorregister-muenchen.de/en/facts/surv/sC4041E-ICD-10-C40-C41-Bone-cancer-survival.pdf>

## Index of figures and tables

| Fig./Tbl. |   | Page |
|-----------|---|------|
| 1a        | Relative survival by period of diagnosis (chart)                    | 3    |
| 1b        | Survival by period of diagnosis (table)                             | 3    |
| 2a        | Survival by sex (chart)   | 4    |
| 2b        | Survival by sex (table)   | 4    |
| 3a        | Relative survival by age category (chart)                           | 5    |
| 3b        | Survival by age category (table)                                    | 5    |
| 4a        | Relative survival by TNM staging (chart)                            | 6    |
| 4b        | Survival by TNM staging (table)                                     | 6    |
| 5a        | Relative Überleben nach topography (Grafik)                         | 7    |
| 5b        | Survival by topography (table)                                      | 7    |
| 6a        | Time to first progression (chart)                                   | 8    |
| 6b        | Time to first progression (table)                                   | 8    |
| 6c        | Observed post-progression survival (chart)                          | 10   |
| 6d        | Observed post-progression survival (table)                          | 10   |
| 6e        | Observed post-progression survival by period of progression (chart) | 11   |
| 6f        | Observed post-progression survival by period of progression (table) | 11   |




**Figure 1a.** Relative survival of patients with bone cancer by period of diagnosis. Included in the evaluation are 641 cases diagnosed between 1988 and 2019.

The survival results of the SEER program (Surveillance, Epidemiology, and End Results) of the American National Cancer Institute (NCI) are summarized as the period of diagnosis from 2000 to 2017, and are represented by colored diamonds in order to facilitate comparisons between MCR and SEER.

The presented survival curves are derived from clinical records with valid follow-up informations, which means that death certificate cases (DCO) cases are omitted from the analysis. With this one restriction, the MCR has provided population-based statistics since 1998, collecting data on all cancer cases in the region of southern Bavaria. Historical data of previous time periods can be heavily selected, therefore, univariate survival comparisons of the presented time periods must be carefully considered. Nonetheless, all calculable survival curves are depicted to facilitate the comparison of long time follow-up analyses of relative survival between particular cancers.

| Years  | Period of diagnosis |        |                    |        |                |        |
|--------|---------------------|--------|--------------------|--------|----------------|--------|
|        | 1988-1997<br>n=125  |        | 1998-2006<br>n=183 |        | 2007+<br>n=333 |        |
|        | obs. %              | rel. % | obs. %             | rel. % | obs. %         | rel. % |
| 0      | 100.0               | 100.0  | 100.0              | 100.0  | 100.0          | 100.0  |
| 1      | 86.3                | 86.9   | 90.8               | 91.5   | 85.7           | 86.4   |
| 2      | 76.5                | 77.8   | 77.9               | 78.9   | 76.2           | 77.6   |
| 3      | 70.8                | 72.3   | 73.1               | 74.9   | 71.4           | 73.4   |
| 4      | 66.6                | 68.6   | 70.7               | 73.1   | 68.5           | 71.0   |
| 5      | 64.9                | 66.9   | 68.9               | 71.7   | 66.3           | 69.1   |
| 6      | 58.8                | 61.7   | 65.8               | 69.3   | 63.8           | 67.1   |
| 7      | 58.8                | 61.3   | 65.2               | 69.0   | 61.8           | 65.8   |
| 8      | 54.4                | 57.8   | 63.3               | 67.6   | 60.4           | 64.9   |
| 9      | 53.5                | 56.6   | 62.7               | 67.1   | 58.7           | 63.3   |
| 10     | 52.6                | 55.8   | 62.1               | 67.0   |                |        |
| Median | 10.2                |        |                    |        |                |        |


**Table 1b.** Observed (obs.) and relative (rel.) survival of patients with bone cancer by period of diagnosis for period 1988-2019 (N=641).



**Figure 2a.** Survival of patients with bone cancer by sex. Included in the evaluation are 516 cases diagnosed between 1998 and 2019.

| Years  | Sex           |        |                 |        |
|--------|---------------|--------|-----------------|--------|
|        | Male<br>n=285 |        | Female<br>n=231 |        |
|        | obs. %        | rel. % | obs. %          | rel. % |
| 0      | 100.0         | 100.0  | 100.0           | 100.0  |
| 1      | 87.9          | 88.6   | 87.1            | 87.8   |
| 2      | 75.7          | 77.1   | 78.1            | 79.4   |
| 3      | 69.5          | 71.4   | 75.1            | 77.2   |
| 4      | 66.8          | 69.0   | 72.5            | 75.1   |
| 5      | 64.3          | 67.0   | 71.0            | 73.9   |
| 6      | 62.6          | 65.7   | 66.9            | 70.4   |
| 7      | 60.6          | 64.5   | 66.3            | 70.1   |
| 8      | 58.3          | 62.7   | 65.6            | 69.9   |
| 9      | 57.0          | 61.6   | 64.8            | 69.1   |
| 10     | 56.2          | 61.0   | 64.0            | 69.0   |
| 11     | 53.4          | 58.4   | 64.0            | 69.0   |
| 12     | 51.3          | 57.0   | 64.0            | 68.9   |
| 13     | 51.3          | 56.6   | 61.5            | 67.5   |
| 14     | 51.3          | 56.2   | 61.5            | 67.1   |
| 15     | 51.3          | 55.7   | 61.5            | 66.8   |
| Median | 19.9          |        |                 |        |


**Table 2b.** Observed (obs.) and relative (rel.) survival of patients with bone cancer by sex for period 1998-2019 (N=516).



**Figure 3a.** Relative survival of patients with bone cancer by age category. Included in the evaluation are 516 cases diagnosed between 1998 and 2019.

| Years  | Age category      |        |                    |        |                    |        |                  |        |
|--------|-------------------|--------|--------------------|--------|--------------------|--------|------------------|--------|
|        | 0-19 yrs<br>n=119 |        | 20-39 yrs<br>n=123 |        | 40-59 yrs<br>n=132 |        | 60+ yrs<br>n=142 |        |
|        | obs. %            | rel. % | obs. %             | rel. % | obs. %             | rel. % | obs. %           | rel. % |
| 0      | 100.0             | 100.0  | 100.0              | 100.0  | 100.0              | 100.0  | 100.0            | 100.0  |
| 1      | 95.4              | 94.8   | 97.3               | 96.7   | 89.5               | 89.4   | 71.6             | 73.7   |
| 2      | 88.8              | 87.9   | 80.1               | 79.6   | 81.3               | 81.8   | 60.6             | 64.7   |
| 3      | 81.1              | 80.6   | 77.2               | 77.1   | 77.1               | 77.9   | 56.2             | 62.0   |
| 4      | 78.2              | 77.4   | 75.3               | 74.6   | 75.3               | 76.4   | 52.3             | 59.6   |
| 5      | 77.2              | 76.5   | 73.3               | 72.8   | 73.4               | 74.3   | 49.1             | 57.8   |
| 6      | 76.1              | 76.0   | 70.0               | 70.2   | 72.4               | 73.5   | 44.0             | 54.1   |
| 7      | 76.1              | 75.6   | 70.0               | 69.9   | 67.8               | 70.2   | 42.9             | 54.3   |
| 8      | 76.1              | 75.2   | 70.0               | 69.5   | 67.8               | 70.1   | 37.2             | 49.3   |
| 9      | 76.1              | 74.9   | 70.0               | 69.2   | 67.8               | 70.0   | 33.4             | 45.9   |
| 10     | 74.7              | 74.2   | 68.5               | 68.3   | 67.8               | 69.9   | 33.4             | 46.3   |
| Median |                   |        |                    |        |                    |        | 4.2              |        |


**Table 3b.** Observed (obs.) and relative (rel.) survival of patients with bone cancer by age category for period 1998-2019 (N=516).



**Figure 4a.** Relative survival of patients with bone cancer by TNM staging. For 273 of 516 cases diagnosed between 1998 and 2019 valid data could be obtained for this item. For a total of 269 cases an evaluable classification was established. The accumulated percentage exceeds the 100 % value because patients are potentially considered in more than one subgroup. The grey line represents the subgroup of 247 patients with missing values regarding TNM staging (47.9 % of 516 patients, the percent values of all other categories are related to n=269).

| Years  | TNM staging |        |             |        |            |        |            |        |            |        |            |        |                 |        |
|--------|-------------|--------|-------------|--------|------------|--------|------------|--------|------------|--------|------------|--------|-----------------|--------|
|        | T1<br>n=72  |        | T2<br>n=141 |        | N0<br>n=82 |        | M0<br>n=68 |        | M1<br>n=67 |        | MX<br>n=57 |        | NA/NOS<br>n=247 |        |
|        | obs. %      | rel. % | obs. %      | rel. % | obs. %     | rel. % | obs. %     | rel. % | obs. %     | rel. % | obs. %     | rel. % | obs. %          | rel. % |
| 0      | 100.0       | 100.0  | 100.0       | 100.0  | 100.0      | 100.0  | 100.0      | 100.0  | 100.0      | 100.0  | 100.0      | 100.0  | 100.0           | 100.0  |
| 1      | 100.0       | 99.7   | 82.4        | 82.7   | 92.5       | 92.5   | 96.9       | 96.5   | 67.8       | 67.3   | 91.0       | 90.9   | 88.6            | 89.3   |
| 2      | 97.1        | 97.2   | 66.6        | 67.6   | 79.5       | 80.5   | 89.2       | 89.2   | 37.8       | 37.8   | 85.5       | 86.4   | 80.9            | 82.5   |
| 3      | 92.7        | 93.9   | 58.9        | 60.2   | 70.0       | 71.0   | 81.0       | 82.2   |            |        | 81.9       | 81.9   | 77.2            | 79.6   |
| 4      | 91.1        | 93.3   | 56.5        | 57.6   | 65.7       | 66.6   | 79.3       | 81.2   |            |        | 79.9       | 81.4   | 73.9            | 76.8   |
| 5      | 89.4        | 93.0   | 54.0        | 55.5   | 64.3       | 66.2   | 79.3       | 81.0   |            |        | 79.9       | 81.0   | 71.5            | 74.8   |
| 6      | 89.4        | 92.8   | 52.2        | 53.8   | 64.3       | 65.9   | 79.3       | 80.8   |            |        | 76.0       | 78.8   | 67.0            | 70.9   |
| 7      | 89.4        | 92.6   | 50.2        | 52.6   | 60.7       | 63.5   | 74.5       | 77.3   |            |        | 76.0       | 78.7   | 65.4            | 70.0   |
| 8      | 89.4        | 92.4   | 50.2        | 52.3   | 58.7       | 61.7   |            |        |            |        | 76.0       | 78.6   | 62.9            | 68.0   |
| 9      | 85.8        | 92.1   | 50.2        | 51.9   | 58.7       | 60.4   |            |        |            |        | 74.0       | 78.3   | 61.5            | 66.8   |
| 10     | 85.8        | 91.3   | 48.8        | 51.0   |            |        |            |        |            |        | 74.0       | 77.4   | 60.8            | 66.4   |
| Median |             |        | 9.0         |        |            |        |            |        | 1.5        |        |            |        |                 |        |


**Table 4b.** Observed (obs.) and relative (rel.) survival of patients with bone cancer by TNM staging for period 1998-2019 (N=269).



**Figure 5a.** Relative survival of patients with bone cancer by topography. For 497 of 516 cases diagnosed between 1998 and 2019 valid data could be obtained for this item.

| Years  | Topography           |        |                          |        |
|--------|----------------------|--------|--------------------------|--------|
|        | Extremities<br>n=247 |        | Non-Extremities<br>n=250 |        |
|        | obs. %               | rel. % | obs. %                   | rel. % |
| 0      | 100.0                | 100.0  | 100.0                    | 100.0  |
| 1      | 87.1                 | 87.7   | 88.3                     | 89.0   |
| 2      | 77.0                 | 78.1   | 76.4                     | 78.0   |
| 3      | 71.6                 | 73.1   | 71.9                     | 74.2   |
| 4      | 69.2                 | 71.0   | 69.2                     | 72.0   |
| 5      | 67.3                 | 69.5   | 66.8                     | 70.1   |
| 6      | 64.8                 | 67.4   | 63.6                     | 67.4   |
| 7      | 63.6                 | 66.9   | 61.8                     | 66.5   |
| 8      | 63.6                 | 66.9   | 58.8                     | 63.9   |
| 9      | 62.3                 | 65.6   | 57.9                     | 63.5   |
| 10     | 60.8                 | 64.7   | 57.9                     | 63.5   |
| Median |                      |        |                          |        |

**Table 5b.** Observed (obs.) and relative (rel.) survival of patients with bone cancer by topography for period 1998-2019 (N=497).




**Figure 6a.** Time to first progression of 508 patients with bone cancer diagnosed between 1998 and 2019 (in solid cancers M0 only) estimated by cumulative incidence function (CI, solid line) accounting for death as competing risk and by inverse Kaplan-Meier estimate (1-KM, dashed line). The frequency of events may be underestimated due to underreporting.

|         | Type of progression |                 |                    |                      |                 |                   |                         |
|---------|---------------------|-----------------|--------------------|----------------------|-----------------|-------------------|-------------------------|
|         | Any type (CI)       | Any type (1-KM) | Local relapse (CI) | Local relapse (1-KM) | Lymph node (CI) | Lymph node (1-KM) | Distant metastasis (CI) |
| N       | 442                 | 442             | 508                | 508                  | 508             | 508               | 442                     |
| Events  | 191                 | 191             | 78                 | 78                   | 7               | 7                 | 114                     |
| compet. | 21                  |                 | 143                |                      | 177             |                   | 52                      |
| Years   | %                   | %               | %                  | %                    | %               | %                 | %                       |
| 0       | 0.0                 | 0.0             | 0.0                | 0.0                  | 0.0             | 0.0               | 0.0                     |
| 1       | 22.0                | 22.3            | 5.8                | 6.3                  | 0.4             | 0.5               | 12.0                    |
| 2       | 33.2                | 33.8            | 9.8                | 11.0                 | 0.4             | 0.5               | 19.7                    |
| 3       | 38.7                | 39.3            | 12.8               | 14.9                 | 0.8             | 1.0               | 21.9                    |
| 4       | 40.7                | 41.4            | 13.9               | 16.4                 | 1.1             | 1.4               | 24.5                    |
| 5       | 43.8                | 44.6            | 15.1               | 18.1                 | 1.5             | 2.1               | 25.8                    |
| 6       | 45.4                | 46.3            | 16.1               | 19.4                 | 1.5             | 2.1               | 26.9                    |
| 7       | 45.7                | 46.6            | 16.6               | 20.3                 | 1.5             | 2.1               | 27.2                    |
| 8       | 45.7                | 46.6            | 16.6               | 20.3                 | 1.5             | 2.1               | 27.2                    |
| 9       | 46.1                | 47.0            | 16.6               | 20.3                 | 1.5             | 2.1               | 27.2                    |
| 10      | 46.5                | 47.5            | 17.0               | 20.9                 | 1.5             | 2.1               | 28.0                    |



| <i>cont'd</i> | Type of progression           |          |            |
|---------------|-------------------------------|----------|------------|
|               | Distant<br>metast. (1-<br>KM) | NOS (CI) | NOS (1-KM) |
| N             | 442                           | 508      | 508        |
| Events        | 114                           | 66       | 66         |
| compet.       |                               | 146      |            |
| Years         | %                             | %        | %          |
| 0             | 0.0                           | 0.0      | 0.0        |
| 1             | 12.4                          | 6.5      | 6.9        |
| 2             | 20.5                          | 9.7      | 10.6       |
| 3             | 22.9                          | 11.2     | 12.6       |
| 4             | 25.6                          | 11.9     | 13.5       |
| 5             | 27.1                          | 13.0     | 15.1       |
| 6             | 28.3                          | 13.8     | 16.1       |
| 7             | 28.7                          | 13.8     | 16.1       |
| 8             | 28.7                          | 13.8     | 16.1       |
| 9             | 28.7                          | 14.1     | 16.6       |
| 10            | 29.7                          | 14.1     | 16.6       |

**Table 6b.** Time to first progression of patients with bone cancer for period 1998-2019 (N=508), also showing the total of progression events (Events) and of deaths as competing risk (compet.).




**Figure 6c.** Observed post-progression survival of 220 patients with bone cancer diagnosed between 1998 and 2019. These 220 patients with documented progression events during their course of disease represent 43.3 % of the totally 508 evaluated cases (incl. M1, n=66, 13.0 %). Patients with cancer relapse documented via death certificates only were excluded (n=40, 7.9 %). Multiple progression types on different sites are included in the evaluation even when not occurring synchronously. The NOS (not otherwise specified) class is included under the condition, that it is the one and only progression type during the course of disease. Subgroups with sample size <20 are omitted from the chart.

Medical record documentation often lacks the linguistic severity to distinguish between local relapse, regional lymph node metastasis and distant spread in solid cancers. Frequently, the statement “not specified” is the only information in registries regarding relapse of the disease. The category “Any type” denotes all cases who suffered from at least one relapse during the course of disease (incl. primary M1-status). Although, the real number of relapsed patients is likely to be much higher. The accumulated percentage of patients with local relapse or distant metastasis exceeds the 100 % value because patients are potentially considered in more than one subgroup.

| Years | Type of progression    |                            |                                  |                  |
|-------|------------------------|----------------------------|----------------------------------|------------------|
|       | Any type<br>n=220<br>% | Local relapse<br>n=82<br>% | Distant metastasis<br>n=144<br>% | NOS<br>n=54<br>% |
| 0     | 100.0                  | 100.0                      | 100.0                            | 100.0            |
| 1     | 74.1                   | 75.9                       | 68.2                             | 79.2             |
| 2     | 55.8                   | 60.8                       | 43.1                             | 69.7             |
| 3     | 47.8                   | 56.7                       | 34.1                             | 60.1             |
| 4     | 43.3                   | 52.4                       | 31.5                             | 55.1             |
| 5     | 41.0                   | 49.5                       | 27.2                             | 55.1             |
| 6     | 39.6                   | 49.5                       | 27.2                             |                  |
| 7     | 39.6                   | 49.5                       |                                  |                  |
| 8     | 37.7                   | 45.1                       |                                  |                  |
| 9     | 36.6                   |                            |                                  |                  |
| 10    | 35.4                   |                            |                                  |                  |

**Table 6d.** Observed post-progression survival of patients with bone cancer for period 1998-2019 (N=220).



**Figure 6e.** Observed post-progression (distant metastasis) survival of 185 patients with bone cancer diagnosed between 1988 and 2019 by period of progression.

| Years | Period of progression  |                        |                     |
|-------|------------------------|------------------------|---------------------|
|       | 1988-1997<br>n=27<br>% | 1998-2006<br>n=48<br>% | 2007+<br>n=110<br>% |
| 0     | 100.0                  | 100.0                  | 100.0               |
| 1     |                        | 67.8                   | 69.6                |
| 2     |                        |                        | 46.7                |
| 3     |                        |                        | 38.0                |
| 4     |                        |                        | 35.6                |

**Table 6f.** Observed post-progression (distant metastasis) survival of patients with bone cancer for period 1988-2019 by period of progression (N=185).

## Shortcuts

|      |  |   |
|------|--|---|
| MCR  | Munich Cancer Registry, Germany                  |   |
| NCI  | National Cancer Institute, USA                   |   |
| SEER | Surveillance, Epidemiology, and End Results, USA |   |
| UICC | Union for International Cancer Control, Geneva   |   |
| DCO  | Death certificate only                           | Death certificate provides the only notification to the registry.   |
| NA   | Not available                                    |   |
| NOS  | Not otherwise specified                          |   |
| OS   | Overall/Observed survival                        | Overall/Observed survival (Kaplan-Meier estimate)<br>Date of entry: diagnosis<br>Event: death from any cause  |
| RS   | Relative survival                                | Survival compared to “general population”,<br>ratio of observed to expected survival (Ederer II method),<br>reflecting cancer specific survival   |
| AS   | Assembled survival                               | Assembled chart of<br>observed, expected, relative survival   |
| CS   | Conditional survival                             | Survival probability under the condition of surviving<br>a given period of time   |
| TTP  | Time to progression                              | Time to first progression / relapse<br>Date of entry: diagnosis<br>Event: (progression / relapse): first local-, lymph node recurrence,<br>distant metastasis or unspecified progression  |
| 1-KM |  | 1 minus Kaplan-Meier estimator<br>("inverse" Kaplan-Meier estimator)  |
| CI   |  | Cumulative incidence<br>Death as competing risk (according to Kalbfleisch und Prentice)   |
| PPS  | Post-progression survival                        | Survival since first progression / relapse (Kaplan-Meier estimate)<br>Date of entry (progression / relapse): first local-, lymph node<br>recurrence, distant metastasis or unspecified progression<br>Event: death from any cause |

## Recommended Citation

Munich Cancer Registry. Survival ICD-10 C40, C41: Bone cancer [Internet]. 2021 [updated 2021 Jan 27; cited 2021 Mar 1]. Available from: <https://www.tumorregister-muenchen.de/en/facts/surv/sC4041E-ICD-10-C40-C41-Bone-cancer-survival.pdf>

## Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

## Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.