

Munich Cancer Registry



- ▶ Incidence and Mortality
- ▶ Selection Matrix
- ▶ Homepage
- ▶ *Deutsch*

ICD-10 C46: Kaposi sarcoma

Survival

Year of diagnosis	1988-1997	1998-2019
Patients	56	99
Diseases	56	99
Cases evaluated	54	74
Creation date	01/27/2021	
Database export	01/07/2021	
Population	4.92 m	



Munich Cancer Registry
Cancer Registry Bavaria - Upper Bavaria Regional Center
at Klinikum Grosshadern/IBE
Marchioninstr. 15
Munich, 81377
Germany

<https://www.tumorregister-muenchen.de/en>

https://www.tumorregister-muenchen.de/en/facts/surv/sC46__E-ICD-10-C46-Kaposi-sarcoma-survival.pdf

Index of figures and tables

Fig./Tbl.		Page
1a	Relative survival by period of diagnosis (chart)	3
1b	Survival by period of diagnosis (table)	3
2a	Survival by sex (chart)	4
2b	Survival by sex (table)	4




Figure 1a. Relative survival of patients with Kaposi sarcoma by period of diagnosis. Included in the evaluation are 121 cases diagnosed between 1988 and 2017.

The survival results of the SEER program (Surveillance, Epidemiology, and End Results) of the American National Cancer Institute (NCI) are summarized as the period of diagnosis from 2000 to 2017, and are represented by colored diamonds in order to facilitate comparisons between MCR and SEER.

The presented survival curves are derived from clinical records with valid follow-up informations, which means that death certificate cases (DCO) cases are omitted from the analysis. With this one restriction, the MCR has provided population-based statistics since 1998, collecting data on all cancer cases in the region of southern Bavaria. Historical data of previous time periods can be heavily selected, therefore, univariate survival comparisons of the presented time periods must be carefully considered. Nonetheless, all calculable survival curves are depicted to facilitate the comparison of long time follow-up analyses of relative survival between particular cancers.

Years	Period of diagnosis			
	1988-1997 n=54		1998+ n=67	
	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0
1	64.2	64.0	91.5	92.8
2	49.8	49.5	84.7	88.5
3	43.5	44.7	81.2	88.6
4			79.4	87.0
5			75.4	86.0
6			73.2	84.4
7			68.0	81.0
8			68.0	80.9
9			68.0	80.9
10			68.0	80.9
Median	1.3			

Table 1b. Observed (obs.) and relative (rel.) survival of patients with Kaposi sarcoma by period of diagnosis for period 1988-2017 (N=121).




Figure 2a. Survival of patients with Kaposi sarcoma by sex. Included in the evaluation are 67 cases diagnosed between 1998 and 2017. Subgroups with sample size <20 are omitted from the chart.

Years	Sex	
	obs. %	rel. %
0	100.0	100.0
1	92.5	93.5
2	86.8	90.5
3	82.8	90.4
4	80.8	88.5
5	78.7	88.4
6	76.3	87.9
7	70.7	84.2
8	70.7	84.2
9	70.7	84.2
Median		

Table 2b. Observed (obs.) and relative (rel.) survival of patients with Kaposi sarcoma by sex for period 1998-2017 (N=67).

Shortcuts

MCR Munich Cancer Registry, Germany

NCI National Cancer Institute, USA

SEER Surveillance, Epidemiology, and End Results, USA

UICC Union for International Cancer Control, Geneva

DCO Death certificate only Death certificate provides the only notification to the registry.

NA Not available

NOS Not otherwise specified

OS Overall/Observed survival Overall/Observed survival (Kaplan-Meier estimate)
Date of entry: diagnosis
Event: death from any cause

RS Relative survival Survival compared to “general population”,
ratio of observed to expected survival (Ederer II method),
reflecting cancer specific survival

AS Assembled survival Assembled chart of
observed, expected, relative survival

CS Conditional survival Survival probability under the condition of surviving
a given period of time

TTP Time to progression Time to first progression / relapse
Date of entry: diagnosis
Event: (progression / relapse): first local-, lymph node recurrence,
distant metastasis or unspecified progression

1-KM 1 minus Kaplan-Meier estimator
 (“inverse” Kaplan-Meier estimator)

CI Cumulative incidence
Death as competing risk (according to Kalbfleisch und Prentice)

PPS Post-progression survival Survival since first progression / relapse (Kaplan-Meier estimate)
Date of entry (progression / relapse): first local-, lymph node
recurrence, distant metastasis or unspecified progression
Event: death from any cause

Recommended Citation

Munich Cancer Registry. Survival ICD-10 C46: Kaposi sarcoma [Internet]. 2021 [updated 2021 Jan 27; cited 2021 Mar 1]. Available from: https://www.tumorregister-muenchen.de/en/facts/surv/sC46__E-ICD-10-C46-Kaposi-sarcoma-survival.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.