

Munich Cancer Registry



- Incidence and Mortality
- Selection Matrix
- Homepage
- *Deutsch*

ICD-10 C73: Follicular thyroid ca.

Survival

Year of diagnosis	1988-1997	1998-2020
Patients	216	1,046
Diseases	217	1,046
Cases evaluated	202	897
Creation date	04/15/2022	
Database export	12/20/2021	
Population	4.92 m	



Munich Cancer Registry
Cancer Registry Bavaria - Upper Bavaria Regional Center
at Klinikum Grosshadern/IBE
Marchioninstr. 15
Munich, 81377
Germany

<https://www.tumorregister-muenchen.de/en>

https://www.tumorregister-muenchen.de/en/facts/surv/sC73F_E-ICD-10-C73-Follicular-thyroid-ca.-survival.pdf

Index of figures and tables

Fig./Tbl.	Page
1a Relative survival by period of diagnosis (chart)	3
1b Survival by period of diagnosis (table)	3
2a Survival by sex (chart)	4
2b Survival by sex (table)	4
3a Relative survival by age category (chart)	5
3b Survival by age category (table)	5
4c Relative survival by TNM staging 1998+ (chart)	6
4d Survival by TNM staging 1998+ (table)	6
4e Relative survival by TNM staging 2007+ (chart)	7
4f Survival by TNM staging 2007+ (table)	7
4g Conditional survival by extent of disease (chart)	8
4h Conditional survival by extent of disease (table)	8
5a Time to first progression (chart)	9
5b Time to first progression (table)	9
5c Observed post-progression survival (chart)	11
5d Observed post-progression survival (table)	11
5e Observed post-progression survival by period of progression (chart)	12
5f Observed post-progression survival by period of progression (table)	12




Figure 1a. Relative survival of patients with follicular thyroid ca. by period of diagnosis. Included in the evaluation are 1,095 cases diagnosed between 1988 and 2020.

The survival results of the SEER program (Surveillance, Epidemiology, and End Results) of the American National Cancer Institute (NCI) are summarized as the period of diagnosis from 2000 to 2018, and are represented by colored diamonds in order to facilitate comparisons between MCR and SEER.

The presented survival curves are derived from clinical records with valid follow-up informations, which means that death certificate cases (DCO) cases are omitted from the analysis. With this one restriction, the MCR has provided population-based statistics since 1998, collecting data on all cancer cases in the region of southern Bavaria. Historical data of previous time periods can be heavily selected, therefore, univariate survival comparisons of the presented time periods must be carefully considered. Nonetheless, all calculable survival curves are depicted to facilitate the comparison of long time follow-up analyses of relative survival between particular cancers.

Years	Period of diagnosis							
	1988-1997 n=201		1998-2006 n=355		2007-2015 n=398		2016+ n=141	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	92.8	94.4	94.3	95.5	94.8	95.9	97.8	98.6
2	88.2	91.2	91.6	94.3	92.9	95.3	96.3	98.0
3	82.9	87.5	89.3	93.2	90.3	93.8	96.3	98.1
4	79.2	85.1	85.5	90.7	88.1	92.8		
5	77.7	84.9	84.6	90.9	86.0	91.6		
6	72.9	80.8	82.2	89.9	83.3	90.0		
7	68.1	77.0	81.0	89.9	81.5	89.2		
8	66.5	76.3	78.2	88.3	79.8	88.5		
9	63.8	74.5	75.5	86.5	79.0	89.1		
10	59.9	71.1	72.6	85.0	76.8	87.9		
11	58.3	70.4	70.7	83.9	74.1	86.5		
12	57.8	70.8	69.1	83.5	72.2	85.4		
13	56.1	70.2	67.8	83.4	68.6	82.7		
14	54.4	69.0	66.2	83.0				
15	53.8	69.4	63.5	81.1				
Median	16.0							

Table 1b. Observed (obs.) and relative (rel.) survival of patients with follicular thyroid ca. by period of diagnosis for period 1988-2020 (N=1,095).




Figure 2a. Survival of patients with follicular thyroid ca. by sex. Included in the evaluation are 894 cases diagnosed between 1998 and 2020.

Years	Sex			
	Male n=296		Female n=598	
	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0
1	92.4	93.9	96.4	97.3
2	90.3	93.5	94.3	96.3
3	87.0	91.6	92.6	95.6
4	83.2	89.3	90.3	94.3
5	82.0	89.5	88.7	93.7
6	79.9	88.5	85.9	91.9
7	77.2	87.4	84.9	92.1
8	74.7	86.3	82.8	90.9
9	73.1	86.0	80.7	89.8
10	71.4	86.3	77.6	87.8
11	69.5	85.7	75.2	86.2
12	68.9	86.8	72.9	84.8
13	68.1	88.2	70.5	83.1
14	64.8	85.8	69.2	82.9
15	61.3	83.4	66.7	81.2
Median				

Table 2b. Observed (obs.) and relative (rel.) survival of patients with follicular thyroid ca. by sex for period 1998-2020 (N=894).




Figure 3a. Relative survival of patients with follicular thyroid ca. by age category. Included in the evaluation are 894 cases diagnosed between 1998 and 2020.

Years	Age category							
	0-49 yrs n=310		50-59 yrs n=169		60-69 yrs n=205		70+ yrs n=210	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	99.7	99.7	98.8	98.7	95.6	96.6	85.0	88.2
2	99.7	99.6	98.2	98.6	92.0	94.1	80.1	86.8
3	99.3	99.5	96.9	98.2	89.9	93.1	74.5	84.3
4	98.1	98.6	96.2	97.8	85.6	89.6	69.2	82.3
5	98.1	98.6	95.5	97.9	83.9	89.0	65.3	81.3
6	98.1	98.5	92.6	95.8	80.4	87.1	60.7	79.7
7	97.7	98.3	91.1	95.1	78.5	86.4	58.3	80.6
8	97.1	98.3	89.4	93.8	76.6	85.7	53.1	77.7
9	97.1	98.3	88.6	93.7	75.2	85.8	47.3	74.0
10	97.1	98.3	86.8	93.5	73.0	85.4	40.3	68.4
11	96.4	98.3	84.8	91.6	72.3	86.3	34.3	63.0
12	96.4	98.1	83.8	91.4	67.3	82.7	33.3	66.1
13	95.6	98.0	81.5	91.0	64.7	81.6	31.3	67.8
14	95.6	98.0	78.9	88.5	61.5	80.4	29.1	70.1
15	95.6	98.0	76.0	85.9	58.9	79.3		
Median					17.5		8.5	

Table 3b. Observed (obs.) and relative (rel.) survival of patients with follicular thyroid ca. by age category for period 1998-2020 (N=894).




Figure 4c. Relative survival of patients with follicular thyroid ca. by TNM staging. For 348 of 355 cases diagnosed between 1998 and 2006 valid data could be obtained for this item. The accumulated percentage exceeds the 100 % value because patients are potentially considered in more than one subgroup. The grey line represents the subgroup of 7 patients with missing values regarding TNM staging (2.0 % of 355 patients, the percent values of all other categories are related to n=348). Subgroups with sample size <20 are omitted from the chart.

Due to substantial changes in stage classification schemes long-term survival statistics over decades could not be created.

TNM staging														
Years	T1 n=82		T2 n=125		T3+4 n=131		N0 n=208		N+ n=42		NX n=76		M1 n=45	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	98.7	98.8	98.4	99.3	88.5	90.0	97.5	98.6	85.6	86.0	91.9	92.4	71.1	71.2
2	97.4	98.3	97.5	99.7	83.1	86.6	96.6	98.7	80.8	81.9	86.5	89.9	66.7	68.3
3	97.4	98.2	96.7	99.5	78.5	83.9	95.6	99.0	70.7	72.8	85.2	88.7	55.6	58.2
4	94.8	96.9	94.2	98.5	73.1	79.7	93.1	97.8	68.1	70.4	75.7	81.4		
5	94.8	97.0	94.2	99.1	70.7	78.7	92.6	98.6	65.6	68.6	75.7	81.5		
6	93.3	96.9	91.7	98.2	68.4	78.3	90.0	97.7	65.6	68.2	73.0	80.7		
7	91.9	96.7	91.7	98.9	66.0	77.4	87.9	96.9	65.6	67.8	73.0	81.2		
8	91.9	97.2	89.1	97.7	61.1	73.3	85.3	95.3	63.1	68.0	68.8	78.7		
9	91.9	97.7	87.4	96.6	57.1	70.2	83.2	94.4	63.1	68.5	67.4	77.7		
10	89.0	96.6	84.7	95.3	53.8	68.4	80.0	92.7	63.1	69.0	63.0	75.5		
Median					10.7						17.2		3.2	

Table 4d. Observed (obs.) and relative (rel.) survival of patients with follicular thyroid ca. by TNM staging for period 1998-2006 (N=348).




Figure 4e. Relative survival of patients with follicular thyroid ca. by TNM staging. For 531 of 539 cases diagnosed between 2007 and 2020 valid data could be obtained for this item. The accumulated percentage exceeds the 100 % value because patients are potentially considered in more than one subgroup. The grey line represents the subgroup of 8 patients with missing values regarding TNM staging (1.5 % of 539 patients, the percent values of all other categories are related to n=531). Subgroups with sample size <20 are omitted from the chart.

Due to substantial changes in stage classification schemes long-term survival statistics over decades could not be created.

TNM staging														
Years	T1 n=157		T2 n=186		T3+4 n=174		N0 n=240		N+ n=26		NX n=181		M1 n=56	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	99.3	99.7	98.4	98.9	93.0	94.2	98.7	99.4	88.5	86.0	95.4	96.8	76.0	76.9
2	98.6	99.6	97.8	99.3	89.5	92.4	98.3	99.6			94.3	96.7	66.0	69.5
3	96.4	98.4	97.2	99.7	86.3	90.5	96.0	98.4			92.3	96.1	60.0	64.0
4	94.8	97.8	95.9	99.3	82.8	88.5	94.5	97.7			89.3	94.6		
5	93.9	98.0	95.9	99.9	79.7	86.0	93.4	97.8			89.3	95.3		
6	93.9	98.1	93.6	99.3	74.6	81.9	92.3	97.4			84.5	92.4		
7	92.8	98.0	92.8	99.1	70.9	79.6	91.1	97.6			81.1	89.3		
8	91.5	97.9	89.8	97.4	70.0	79.7	89.1	96.8			78.5	88.4		
9	91.5	98.0	89.8	97.9	67.4	79.4	89.1	97.9			76.7	88.0		
10	88.4	96.4	88.5	98.4	64.6	77.2	89.1	98.9			72.8	84.8		
Median					13.5									

Table 4f. Observed (obs.) and relative (rel.) survival of patients with follicular thyroid ca. by TNM staging for period 2007-2020 (N=531).




Figure 4g. Conditional relative 5-year survival of patients with follicular thyroid ca. by extent of disease. For 879 of 894 cases diagnosed between 1998 and 2020 valid data could be obtained for this item. The grey line represents the subgroup of 15 patients with missing values regarding extent of disease (1.7 % of 894 patients, the percent values of all other categories are related to n=879). Subgroups with sample size <20 are omitted from the chart.

Years	Extent of disease							
	Local		Regional		Distant		NA/NOS	
	n	Cond. surv. % 5 yrs	n	Cond. surv. % 5 yrs	n	Cond. surv. % 5 yrs	n	Cond. surv. % 5 yrs
0	730	99.1	48	83.0	101	44.6	15	
1	689	98.9	45	85.7	70	54.4		
2	667	98.8	43	88.8	63	57.2		
3	628	98.3	39	92.4				
4	586	99.0	36	95.6				
5	553	97.6	34	94.2				
6	501	97.5	32	96.7				
7	467	97.4	32	96.4				
8	414	96.8	29	96.0				
9	381	95.2	27	98.9				
10	353	95.0						

Table 4h. Conditional relative 5-year survival of patients with follicular thyroid ca. by extent of disease for period 1998-2020 (N=879).

Conditional relative survival rates refer to the relative survival probability, in this case for 5 years after cancer diagnosis, compared to the age- and sex-matched population (=100 %) under the condition of being alive for a certain time period (x-axis in Figure 4e). The results illustrate to what extent the cancer induced mortality of particular subgroups declines in the subsequent years after detection of the malignancy. For instance, according to the presented survival statistics, patients in the subgroup extent of disease="Local", who are alive at least 3 years after cancer diagnosis, the conditional relative 5-year survival rate is 98.3% (n=628).




Figure 5a. Time to first progression of 887 patients with follicular thyroid ca. diagnosed between 1998 and 2020 (in solid cancers M0 only) estimated by cumulative incidence function (CI, solid line) accounting for death as competing risk and by inverse Kaplan-Meier estimate (1-KM, dashed line). The frequency of events may be underestimated due to underreporting.

Type of progression							
	Any type (CI)	Any type (1-KM)	Local relapse (CI)	Local relapse (1-KM)	Lymph node (CI)	Lymph node (1-KM)	Distant metastasis (CI)
N	786	786	887	887	887	887	786
Events	115	115	33	33	39	39	72
compet.	57		160		164		68
Years	%	%	%	%	%	%	%
0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1	5.2	5.2	1.2	1.2	1.6	1.7	2.5
2	7.7	7.8	1.6	1.7	2.4	2.5	4.1
3	9.5	9.6	2.1	2.2	3.2	3.3	5.0
4	10.8	11.0	3.0	3.2	3.7	3.9	6.2
5	12.2	12.4	3.1	3.4	4.1	4.4	7.1
6	13.4	13.7	3.6	3.8	4.4	4.7	7.7
7	14.8	15.2	3.7	4.0	4.5	4.9	9.1
8	15.5	16.0	3.9	4.2	4.7	5.1	9.4
9	15.7	16.2	3.9	4.2	4.7	5.1	9.6
10	16.6	17.1	4.2	4.7	4.9	5.3	10.7

<i>cont'd</i>	Type of progression		
	Distant metast. (1- KM)	NOS (CI)	NOS (1-KM)
N	786	887	887
Events	72	26	26
compet.		170	
Years	%	%	%
0	0.0	0.0	0.0
1	2.5	1.7	1.8
2	4.1	2.3	2.4
3	5.1	2.6	2.7
4	6.3	2.7	2.8
5	7.3	2.7	2.8
6	7.9	3.0	3.1
7	9.4	3.1	3.3
8	9.8	3.1	3.3
9	10.0	3.1	3.3
10	11.2	3.1	3.3

Table 5b. Time to first progression of patients with follicular thyroid ca. for period 1998-2020 (N=887), also showing the total of progression events (Events) and of deaths as competing risk (compet.).




Figure 5c. Observed post-progression survival of 209 patients with follicular thyroid ca. diagnosed between 1998 and 2020. These 209 patients with documented progression events during their course of disease represent 23.6 % of the totally 887 evaluated cases (incl. M1, n=101, 11.4 %). Patients with cancer relapse documented via death certificates only were excluded (n=14, 1.6 %). Multiple progression types on different sites are included in the evaluation even when not occurring synchronously. The NOS (not otherwise specified) class is included under the condition, that it is the one and only progression type during the course of disease.

Medical record documentation often lacks the linguistic severity to distinguish between local relapse, regional lymph node metastasis and distant spread in solid cancers. Frequently, the statement “not specified” is the only information in registries regarding relapse of the disease. The category “Any type” denotes all cases who suffered from at least one relapse during the course of disease (incl. primary M1-status). Although, the real number of relapsed patients is likely to be much higher. The accumulated percentage of patients with local relapse or distant metastasis exceeds the 100 % value because patients are potentially considered in more than one subgroup.

Years	Type of progression				
	Any type n=209 %	Local relapse n=36 %	Lymph node n=41 %	Distant metastasis n=167 %	NOS n=24 %
0	100.0	100.0	100.0	100.0	100.0
1	83.0	77.1	90.2	80.0	91.3
2	74.1	65.4	80.5	69.3	91.3
3	63.1		66.8	58.1	
4	55.7		63.7	48.9	
5	50.8			44.4	
6	47.4			41.0	
7	46.6			40.0	
8	39.7			32.6	
9	35.1			27.0	
10	33.2			24.6	

Table 5d. Observed post-progression survival of patients with follicular thyroid ca. for period 1998-2020 (N=209).




Figure 5e. Observed post-progression (distant metastasis) survival of 231 patients with follicular thyroid ca. diagnosed between 1988 and 2020 by period of progression.

Years	Period of progression			
	1988-1997 n=47 %	1998-2006 n=71 %	2007-2015 n=75 %	2016+ n=38 %
0	100.0	100.0	100.0	100.0
1	82.9	71.8	83.7	89.2
2	72.0	62.0	68.6	79.8
3	56.7	51.9	57.6	
4	54.5	41.8	50.8	
5	52.3	40.4	43.9	
6		36.0	41.1	
7		36.0	39.2	
8		31.7		

Table 5f. Observed post-progression (distant metastasis) survival of patients with follicular thyroid ca. for period 1988-2020 by period of progression (N=231).

Shortcuts

MCR Munich Cancer Registry, Germany

NCI National Cancer Institute, USA

SEER Surveillance, Epidemiology, and End Results, USA

UICC Union for International Cancer Control, Geneva

DCO Death certificate only Death certificate provides the only notification to the registry.

NA Not available

NOS Not otherwise specified

OS Overall/Observed survival Overall/Observed survival (Kaplan-Meier estimate)

Date of entry: diagnosis
Event: death from any cause

RS Relative survival Survival compared to “general population”, ratio of observed to expected survival (Ederer II method), reflecting cancer specific survival

AS Assembled survival Assembled chart of observed, expected, relative survival

CS Conditional survival Survival probability under the condition of surviving a given period of time

TTP Time to progression Time to first progression / relapse
Date of entry: diagnosis
Event: (progression / relapse): first local-, lymph node recurrence, distant metastasis or unspecified progression

1-KM 1 minus Kaplan-Meier estimator (“inverse” Kaplan-Meier estimator)

CI Cumulative incidence
Death as competing risk (according to Kalbfleisch und Prentice)

PPS Post-progression survival Survival since first progression / relapse (Kaplan-Meier estimate)
Date of entry (progression / relapse): first local-, lymph node recurrence, distant metastasis or unspecified progression
Event: death from any cause

Recommended Citation

Munich Cancer Registry. Survival ICD-10 C73: Follicular thyroid ca. [Internet]. 2022 [updated 2022 Apr 15; cited 2022 Jun 1]. Available from: https://www.tumorregister-muenchen.de/en/facts/surv/sC73F_E-ICD-10-C73-Follicular-thyroid-ca.-survival.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.