

# Munich Cancer Registry



- ▶ Incidence and Mortality
- ▶ Selection Matrix
- ▶ Homepage
- ▶ *Deutsch*

## ICD-10 C73: Thyroid cancer

### Survival

Year of diagnosis	1988-1997	1998-2016
Patients	954	7,554
Diseases	958	7,610
Cases evaluated	879	6,642
Creation date	08/22/2018	
Export date	08/09/2018	
Population	4.81 m	




Munich Cancer Registry  
Cancer Registry Bavaria - Upper Bavaria Regional Center  
at Klinikum Grosshadern/IBE  
Marchioninstr. 15  
Munich, 81377  
Germany

<https://www.tumorregister-muenchen.de/en>

[https://www.tumorregister-muenchen.de/en/facts/surv/sC73\\_\\_E-ICD-10-C73-Thyroid-cancer-survival.pdf](https://www.tumorregister-muenchen.de/en/facts/surv/sC73__E-ICD-10-C73-Thyroid-cancer-survival.pdf)

### Index of figures and tables

Fig./Tbl.		Page
1a	Relative survival by period of diagnosis (chart)	3
1b	Survival by period of diagnosis (table)	3
2a	Survival by sex (chart)	4
2b	Survival by sex (table)	4
3a	Relative survival by age category (chart)	5
3b	Survival by age category (table)	5
4a	Relative survival by histology (chart)	6
4b	Survival by histology (table)	6
4c	Relative survival by TNM staging 1998+ (chart)	7
4d	Survival by TNM staging 1998+ (table)	7
4e	Relative survival by TNM staging 2007+ (chart)	9
4f	Survival by TNM staging 2007+ (table)	9
4g	Conditional survival by extent of disease (chart)	10
4h	Conditional survival by extent of disease (table)	10
5a	Time to first progression (chart)	11
5b	Time to first progression (table)	11
5c	Observed post-progression survival (chart)	13
5d	Observed post-progression survival (table)	13
5e	Observed post-progression survival by period of progression (chart)	14
5f	Observed post-progression survival by period of progression (table)	14




**Figure 1a.** Relative survival of patients with thyroid cancer by period of diagnosis. Included in the evaluation are 7,182 cases diagnosed between 1988 and 2016.

The survival results of the SEER program (Surveillance, Epidemiology, and End Results) of the American National Cancer Institute (NCI) are summarized as the period of diagnosis from 2000 to 2014, and are represented by colored diamonds in order to facilitate comparisons between MCR and SEER.

The presented survival curves are derived from clinical records with valid follow-up informations, which means that death certificate cases (DCO) cases are omitted from the analysis. With this one restriction, the MCR has provided population-based statistics since 1998, collecting data on all cancer cases in the region of southern Bavaria. Historical data of previous time periods can be heavily selected, therefore, univariate survival comparisons of the presented time periods must be carefully considered. Nonetheless, all calculable survival curves are depicted to facilitate the comparison of long time follow-up analyses of relative survival between particular cancers.

Years	Period of diagnosis					
	1988-1997		1998-2006		2007+	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0
1	91.7	93.0	95.0	96.0	96.1	96.8
2	88.7	91.1	93.9	95.6	95.2	96.6
3	86.5	90.0	92.7	95.2	94.4	96.5
4	84.3	88.9	91.3	94.6	93.6	96.4
5	83.5	89.0	90.6	94.7	92.9	96.4
6	81.8	88.3	88.9	93.8	92.0	96.3
7	79.9	87.3	88.2	93.9	91.4	96.5
8	77.6	86.0	87.3	93.8	89.6	95.5
9	76.2	85.3	85.5	92.9		
10	73.8	83.8	84.4	92.8		
11	72.4	83.2	83.1	92.3		
12	70.7	82.2	82.2	92.3		
13	68.8	81.0	81.1	92.2		
14	66.8	79.8	80.3	92.4		
15	65.6	79.3	78.9	92.1		


**Table 1b.** Observed (obs.) and relative (rel.) survival of patients with thyroid cancer by period of diagnosis for period 1988-2016 (N=7,182).



**Figure 2a.** Survival of patients with thyroid cancer by sex. Included in the evaluation are 6,304 cases diagnosed between 1998 and 2016.

Years	Sex			
	Male n=1,721		Female n=4,583	
	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0
1	93.6	94.8	96.5	97.1
2	92.3	94.5	95.6	96.9
3	90.4	93.8	95.0	96.8
4	88.7	93.2	94.2	96.6
5	87.8	93.3	93.6	96.7
6	86.0	92.6	92.4	96.2
7	85.4	93.2	91.7	96.2
8	83.4	92.3	90.9	96.1
9	81.5	91.6	89.2	95.2
10	80.4	91.9	88.1	94.8
11	78.5	91.2	87.0	94.5
12	77.6	91.5	86.1	94.4
13	75.4	90.9	85.3	94.4
14	74.8	91.6	84.4	94.4
15	73.8	92.5	82.8	93.7


**Table 2b.** Observed (obs.) and relative (rel.) survival of patients with thyroid cancer by sex for period 1998-2016 (N=6,304).



**Figure 3a.** Relative survival of patients with thyroid cancer by age category. Included in the evaluation are 7,182 cases diagnosed between 1988 and 2016.

Years	Age category							
	0-49 yrs n=3,241		50-59 yrs n=1,685		60-69 yrs n=1,340		70+ yrs n=916	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	99.5	99.6	98.2	98.6	95.0	95.8	75.4	78.9
2	99.4	99.6	97.4	98.3	93.0	95.1	70.8	77.0
3	99.3	99.6	96.6	98.0	91.9	95.1	65.6	74.7
4	99.1	99.6	95.7	97.7	89.8	94.2	62.0	73.8
5	99.0	99.7	95.5	98.0	88.5	94.2	59.1	73.7
6	98.8	99.6	93.8	97.1	87.3	94.3	54.6	71.7
7	98.3	99.4	93.1	96.9	86.6	95.1	51.5	71.5
8	97.8	99.0	91.8	96.3	84.6	94.8	47.9	70.7
9	97.6	99.0	90.3	95.6	81.7	93.4	43.3	68.0
10	97.3	98.9	89.5	95.6	79.7	93.1	37.7	63.9
11	97.1	99.0	88.3	95.2	76.6	91.8	34.6	63.3
12	96.8	99.0	87.1	94.8	75.3	92.7	30.1	60.0
13	95.9	98.5	85.3	94.0	73.5	92.8	27.1	59.8
14	95.3	98.2	83.6	93.4	70.7	92.3	25.4	61.5
15	95.1	98.3	81.7	92.1	67.7	91.9	23.2	63.3


**Table 3b.** Observed (obs.) and relative (rel.) survival of patients with thyroid cancer by age category for period 1988-2016 (N=7,182).



**Figure 4a.** Relative survival of patients with thyroid cancer by histology. For 7,117 of 7,182 cases diagnosed between 1988 and 2016 valid data could be obtained for this item. The grey line represents the subgroup of 65 patients with missing values regarding histology (0.9 % of 7,182 patients, the percent values of all other categories are related to n=7,117).

Years	Histology											
	NOS n=328		Follicular n=816		Papillary n=5,449		Medullary n=317		Anaplastic n=207		NA n=65	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	79.5	80.7	93.5	95.0	99.4	100.0	96.6	97.2	20.8	21.4	85.6	87.2
2	75.9	77.9	90.4	93.4	98.9	100.2	93.6	94.9	13.3	13.9	85.6	87.7
3	73.5	76.7	86.9	91.2	98.4	100.4	90.8	92.8			82.2	85.9
4	72.4	76.3	82.9	88.6	97.6	100.3	89.5	92.1			78.7	83.7
5	71.7	76.4	81.1	88.1	97.1	100.5	88.6	92.1			78.7	84.1
6	69.4	74.7	78.2	86.4	96.2	100.4	84.8	89.0			75.1	82.3
7	67.4	73.5	75.8	85.3	95.7	100.7	82.4	87.1			73.2	80.3
8	67.0	73.6	72.9	83.5	94.5	100.3	81.2	86.7			69.4	78.8
9	66.1	73.1	69.8	81.5	92.9	99.6	80.4	86.3			69.4	78.5
10	64.9	73.0	65.8	78.5	91.9	99.5	79.7	85.8			65.2	76.2

**Table 4b.** Observed (obs.) and relative (rel.) survival of patients with thyroid cancer by histology for period 1988-2016 (N=7,117).



**Figure 4c.** Relative survival of patients with thyroid cancer by TNM staging. For 2,235 of 2,301 cases diagnosed between 1998 and 2006 valid data could be obtained for this item. For a total of 2,232 cases an evaluable classification was established. The accumulated percentage exceeds the 100 % value because patients are potentially considered in more than one subgroup. The grey line represents the subgroup of 69 patients with missing values regarding TNM staging (3.0 % of 2,301 patients, the percent values of all other categories are related to n=2,232).


Due to substantial changes in stage classification schemes long-term survival statistics over decades could not be created.

Years	TNM staging													
	T1 n=1,163		T2 n=422		T3+4 n=615		N0 n=1,188		N+ n=394		NX n=460		M1 n=114	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	99.7	100.2	98.8	99.5	87.2	88.7	98.1	98.9	91.8	92.4	94.6	95.4	52.6	53.5
2	99.2	100.3	98.5	100.1	84.7	87.1	97.2	98.7	90.4	91.8	93.2	94.9	50.0	51.7
3	98.9	100.6	98.0	100.3	81.4	84.9	96.8	99.2	87.1	89.1	92.0	94.6	42.8	45.6
4	98.3	100.7	96.7	100.0	78.8	83.1	95.3	98.5	86.3	88.7	90.1	93.7	34.8	37.5
5	97.8	100.9	96.1	100.4	77.5	82.7	94.8	98.8	84.9	87.9	90.1	94.5	32.1	35.2
6	97.0	100.8	92.7	97.8	75.9	82.0	93.3	98.2	83.7	87.3	88.4	93.8	28.6	32.2
7	96.4	101.0	92.4	98.4	74.9	81.8	92.7	98.4	82.3	86.3	88.2	94.5	27.7	31.6
8	96.0	101.4	91.6	98.5	72.7	80.3	91.5	98.1	81.4	85.8	87.2	94.6	25.0	28.8
9	94.8	101.1	90.5	98.2	70.0	78.5	90.4	98.0	79.3	84.5	85.5	93.8	21.4	24.7
10	94.0	101.2	88.5	97.1	68.9	78.4	89.5	98.1	78.3	83.7	83.9	93.4		

TNM staging		
<i>cont'd</i>	NA/NOS	
	n=69	
Years	obs. %	rel. %
0	100.0	100.0
1	86.2	87.1
2	83.1	84.2
3	81.4	83.1
4	78.1	81.6
5	78.1	81.7
6	74.8	80.1
7	71.4	77.8
8	71.4	78.0
9	68.0	75.8
10	66.3	75.3

**Table 4d.** Observed (obs.) and relative (rel.) survival of patients with thyroid cancer by TNM staging for period 1998-2006 (N=2,232).






**Figure 4e.** Relative survival of patients with thyroid cancer by TNM staging. For 3,932 of 4,003 cases diagnosed between 2007 and 2016 valid data could be obtained for this item. For a total of 3,929 cases an evaluable classification was established. The accumulated percentage exceeds the 100 % value because patients are potentially considered in more than one subgroup. The grey line represents the subgroup of 74 patients with missing values regarding TNM staging (1.8 % of 4,003 patients, the percent values of all other categories are related to n=3,929).

Due to substantial changes in stage classification schemes long-term survival statistics over decades could not be created.

Years	TNM staging													
	T1 n=2,513		T2 n=473		T3+4 n=890		N0 n=1,685		N+ n=702		NX n=1,168		M1 n=144	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	99.8	100.3	99.1	99.5	88.2	89.3	98.6	99.1	93.2	93.9	97.2	98.0	49.9	51.0
2	99.5	100.6	98.9	99.8	86.1	88.0	98.3	99.3	91.5	92.6	96.4	97.9	44.4	46.3
3	99.1	100.8	98.3	100.0	84.0	86.9	97.9	99.5	89.7	91.5	95.6	98.0	38.4	40.8
4	98.7	101.1	97.6	100.0	82.0	85.7	97.3	99.6	88.5	90.7	95.0	98.4	34.9	38.3
5	98.4	101.5	97.6	100.3	79.4	83.8	96.9	99.9	86.2	88.9	94.8	98.9		
6	97.9	101.8	96.3	100.0	77.7	82.9	96.3	100.0	84.5	87.5	93.8	99.0		
7	97.4	102.1	95.7	100.0	77.3	83.2	95.6	100.1	83.9	87.1	93.5	99.5		
8	96.1	101.5									92.7	99.0		

TNM staging cont'd		
NA/NOS n=74		
Years	obs. %	rel. %
0	100.0	100.0
1	80.6	81.7
2	75.7	77.7
3	73.8	77.2
4	73.8	76.8
5	71.6	75.7
6	68.5	73.6

**Table 4f.** Observed (obs.) and relative (rel.) survival of patients with thyroid cancer by TNM staging for period 2007-2016 (N=3,929).




**Figure 4g.** Conditional relative 5-year survival of patients with thyroid cancer by extent of disease. For 6,988 of 7,182 cases diagnosed between 1988 and 2016 valid data could be obtained for this item. For a total of 6,979 cases an evaluable classification was established. The grey line represents the subgroup of 203 patients with missing values regarding extent of disease (2.8 % of 7,182 patients, the percent values of all other categories are related to n=6,979).

Years	Extent of disease							
	Local		Regional		Distant		NA/NOS	
	n	Cond. surv. % 5 yrs	n	Cond. surv. % 5 yrs	n	Cond. surv. % 5 yrs	n	Cond. surv. % 5 yrs
0	5,478	99.2	1,164	93.6	337	36.4	203	83.1
1	4,863	100.2	988	95.9	165	59.9	159	94.0
2	4,547	100.6	921	96.1	143	61.9	146	94.0
3	4,194	100.0	828	96.9	117	65.0	134	95.9
4	3,871	99.8	747	96.3	98	67.1	128	92.9
5	3,557	99.3	661	95.6	84	67.8	122	90.8
6	3,164	99.0	581	96.4	69	72.0	113	93.2
7	2,727	98.5	500	97.2	60	72.3	103	95.7
8	2,245	98.7	426	96.3	51	76.5	92	94.4
9	1,823	99.0	358	96.8	44	79.5	83	95.5
10	1,561	99.1	302	96.7			79	97.7

**Table 4h.** Conditional relative 5-year survival of patients with thyroid cancer by extent of disease for period 1988-2016 (N=6,979).

Conditional relative survival rates refer to the relative survival probability, in this case for 5 years after cancer diagnosis, compared to the age- and sex-matched population (=100 %) under the condition of being alive for a certain time period (x-axis in Figure 4e). The results illustrate to what extent the cancer induced mortality of particular subgroups declines in the subsequent years after detection of the malignancy. For instance, according to the presented survival statistics, patients in the subgroup extent of disease="Local", who are alive at least 3 years after cancer diagnosis, the conditional relative 5-year survival rate is 100.0% (n=4,194).




**Figure 5a.** Time to first progression of 5,968 patients with thyroid cancer diagnosed between 1998 and 2016 (in solid cancers M0 only) estimated by cumulative incidence function (CI, solid line) accounting for death as competing risk and by inverse Kaplan-Meier estimate (1-KM, dashed line). The frequency of events may be underestimated due to underreporting.

Years	Type of progression						
	Any type (CI)	Any type (1-KM)	Local relapse (CI)	Local relapse (1-KM)	Lymph node (CI)	Lymph node (1-KM)	Distant metastasis (CI)
	n=5,968 %	n=5,968 %	n=5,968 %	n=5,968 %	n=5,968 %	n=5,968 %	n=5,968 %
0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1	4.6	4.6	0.6	0.6	1.2	1.2	1.7
2	5.8	5.8	0.8	0.8	1.7	1.7	2.3
3	6.4	6.5	1.0	1.0	2.0	2.0	2.7
4	6.9	6.9	1.2	1.2	2.1	2.2	3.0
5	7.4	7.4	1.4	1.4	2.3	2.3	3.3
6	7.9	8.0	1.6	1.6	2.3	2.4	3.6
7	8.3	8.4	1.7	1.8	2.5	2.6	4.0
8	8.7	8.8	1.8	1.8	2.7	2.8	4.2
9	9.0	9.1	1.8	1.9	2.8	2.9	4.5
10	9.2	9.3	1.9	2.0	2.9	3.0	4.9

<i>cont'd</i>	Type of progression		
	Distant metast. (1-KM) n=5,968 %	NOS (CI) n=5,968 %	NOS (1-KM) n=5,968 %
Years			
0	0.0	0.0	0.0
1	1.8	1.7	1.8
2	2.3	2.0	2.0
3	2.7	2.1	2.1
4	3.0	2.2	2.2
5	3.4	2.2	2.3
6	3.7	2.4	2.5
7	4.1	2.4	2.5
8	4.3	2.6	2.6
9	4.6	2.6	2.7
10	5.0	2.6	2.7

**Table 5b.** Time to first progression of patients with thyroid cancer for period 1998-2016 (N=5,968).




**Figure 5c.** Observed post-progression survival of 610 patients with thyroid cancer diagnosed between 1998 and 2016. These 610 patients with documented progression events during their course of disease represent 9.8 % of the totally 6,220 evaluated cases (incl. M1, n=252, 4.1 %). Patients with cancer relapse documented via death certificates only were excluded (n=102, 1.6 %). Multiple progression types on different sites are included in the evaluation even when not occurring synchronously. The NOS (not otherwise specified) class is included under the condition, that it is the one and only progression type during the course of disease.

Medical record documentation often lacks the linguistic severity to distinguish between local relapse, regional lymph node metastasis and distant spread in solid cancers. Frequently, the statement “not specified” is the only information in registries regarding relapse of the disease. The category “Any type” denotes all cases who suffered from at least one relapse during the course of disease (incl. primary M1-status). Although, the real number of relapsed patients is likely to be much higher. The accumulated percentage of patients with local relapse or distant metastasis exceeds the 100 % value because patients are potentially considered in more than one subgroup.

Years	Type of progression				
	Any type n=610 %	Local relapse n=102 %	Lymph node n=153 %	Distant metastasis n=419 %	NOS n=80 %
0	100.0	100.0	100.0	100.0	100.0
1	69.4	64.4	89.7	60.6	83.0
2	62.9	51.5	83.7	52.8	81.2
3	58.0	46.4	80.5	45.5	79.2
4	52.9	42.3	74.1	39.8	73.1
5	49.1	39.6	70.7	36.1	71.0
6	46.0	38.0	63.9	33.6	66.3
7	45.0	38.0	63.9	32.2	66.3
8	42.0		59.9	28.8	66.3
9	39.5		59.9	25.6	
10	37.3		59.9	23.1	

**Table 5d.** Observed post-progression survival of patients with thyroid cancer for period 1998-2016 (N=610).



**Figure 5e.** Observed post-progression (distant metastasis) survival of 554 patients with thyroid cancer diagnosed between 1988 and 2016 by period of progression.

Years	Period of progression		
	1988-1997 n=97 %	1998-2006 n=178 %	2007+ n=279 %
0	100.0	100.0	100.0
1	64.1	58.6	62.9
2	55.4	52.2	53.0
3	47.7	45.8	46.5
4	44.3	39.3	42.1
5	42.1	37.0	36.9
6	35.5	33.4	36.1
7	31.0	32.8	33.4
8	26.6	29.9	
9	22.2	25.2	
10		22.7	

**Table 5f.** Observed post-progression (distant metastasis) survival of patients with thyroid cancer for period 1988-2016 by period of progression (N=554).

## Shortcuts

MCR Munich Cancer Registry, Germany

NCI National Cancer Institute, USA

SEER Surveillance, Epidemiology, and End Results, USA

UICC Union for International Cancer Control, Geneva

DCO Death certificate only Death certificate provides the only notification to the registry.

NA Not available

NOS Not otherwise specified

OS Overall/Observed survival Overall/Observed survival (Kaplan-Meier estimate)  
Date of entry: diagnosis  
Event: death from any cause

RS Relative survival Survival compared to “general population”,  
ratio of observed to expected survival (Ederer II method),  
reflecting cancer specific survival

AS Assembled survival Assembled chart of  
observed, expected, relative survival

CS Conditional survival Survival probability under the condition of surviving  
a given period of time

TTP Time to progression Time to first progression / relapse  
Date of entry: diagnosis  
Event: (progression / relapse): first local-, lymph node recurrence,  
distant metastasis or unspecified progression

1-KM 1 minus Kaplan-Meier estimator  
 (“inverse” Kaplan-Meier estimator)

CI Cumulative incidence  
Death as competing risk (according to Kalbfleisch und Prentice)

PPS Post-progression survival Survival since first progression / relapse (Kaplan-Meier estimate)  
Date of entry (progression / relapse): first local-, lymph node  
recurrence, distant metastasis or unspecified progression  
Event: death from any cause

## Recommended Citation

Munich Cancer Registry. Survival ICD-10 C73: Thyroid cancer [Internet]. 2018 [updated 2018 Aug 22; cited 2018 Oct 1]. Available from: [https://www.tumorregister-muenchen.de/en/facts/surv/sC73\\_\\_E-ICD-10-C73-Thyroid-cancer-survival.pdf](https://www.tumorregister-muenchen.de/en/facts/surv/sC73__E-ICD-10-C73-Thyroid-cancer-survival.pdf)

## Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

## Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.