


Munich Cancer Registry



- ▶ Incidence and Mortality
- ▶ Selection Matrix
- ▶ Homepage
- ▶ *Deutsch*

BNET: Pulm. neuroend. tumor

Survival

Year of diagnosis	1998-2020
Patients	1,264
Diseases	1,266
Cases evaluated	957
Creation date	04/27/2022
Database export	12/20/2021
Population	4.92 m



Munich Cancer Registry
Cancer Registry Bavaria - Upper Bavaria Regional Center
at Klinikum Grosshadern/IBE
Marchioninistr. 15
Munich, 81377
Germany

<https://www.tumorregister-muenchen.de/en>

<https://www.tumorregister-muenchen.de/en/facts/surv/shBNETE-BNET-Pulm.-neuroend.-tumor-survival.pdf>

Index of figures and tables

Fig./Tbl.		Page
1a	Relative survival by period of diagnosis (chart)	3
1b	Survival by period of diagnosis (table)	3
2a	Survival by sex (chart)	4
2b	Survival by sex (table)	4
2c	Conditional survival by sex (chart)	5
2d	Conditional survival by sex (table)	5
3a	Relative survival by age category (chart)	6
3b	Survival by age category (table)	6
4a	Relative survival by extent of disease (chart)	7
4b	Survival by extent of disease (table)	7
5a	Time to first progression (chart)	8
5b	Time to first progression (table)	8
5c	Observed post-progression survival (chart)	9
5d	Observed post-progression survival (table)	9
5e	Observed post-progression survival by period of progression (chart)	10
5f	Observed post-progression survival by period of progression (table)	10




Figure 1a. Relative survival of patients with pulm. neuroend. tumor by period of diagnosis. Included in the evaluation are 953 cases diagnosed between 1998 and 2020.

The survival results of the SEER program (Surveillance, Epidemiology, and End Results) of the American National Cancer Institute (NCI) are summarized as the period of diagnosis from 2000 to 2018, and are represented by colored diamonds in order to facilitate comparisons between MCR and SEER.

The presented survival curves are derived from clinical records with valid follow-up informations, which means that death certificate cases (DCO) cases are omitted from the analysis. With this one restriction, the MCR has provided population-based statistics since 1998, collecting data on all cancer cases in the region of southern Bavaria. Historical data of previous time periods can be heavily selected, therefore, univariate survival comparisons of the presented time periods must be carefully considered. Nonetheless, all calculable survival curves are depicted to facilitate the comparison of long time follow-up analyses of relative survival between particular cancers.

Years	Period of diagnosis					
	1998-2006 n=179		2007-2015 n=461		2016+ n=313	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0
1	82.9	83.4	70.1	71.1	75.0	76.1
2	77.6	79.7	60.9	62.9	67.0	69.3
3	76.4	78.9	56.9	59.7	61.8	64.9
4	72.8	76.2	54.9	58.5	59.9	63.5
5	70.3	75.1	53.1	57.8		
Median	15.2		6.9			

Table 1b. Observed (obs.) and relative (rel.) survival of patients with pulm. neuroend. tumor by period of diagnosis for period 1998-2020 (N=953).




Figure 2a. Survival of patients with pulm. neuroend. tumor by sex. Included in the evaluation are 953 cases diagnosed between 1998 and 2020.

Years	Sex			
	Male n=481		Female n=472	
	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0
1	63.6	64.7	85.0	86.1
2	53.5	55.7	79.1	81.1
3	51.0	53.9	74.0	76.9
4	48.5	52.2	71.8	75.6
5	46.2	51.0	70.3	75.0
6	45.9	51.6	67.8	73.5
7	43.8	50.3	65.5	71.9
8	43.4	50.5	65.1	72.5
9	41.2	49.3	62.8	71.0
10	39.7	48.4	60.3	69.7
11	37.7	46.9	58.5	68.7
12	37.0	47.1	57.8	68.6
13	36.3	46.8	57.0	68.6
14	33.2	44.4	55.0	67.4
15	31.3	42.9	53.7	67.3
Median	3.7			

Table 2b. Observed (obs.) and relative (rel.) survival of patients with pulm. neuroend. tumor by sex for period 1998-2020 (N=953).




Figure 2c. Conditional relative 2 / 5-year survival of patients with pulm. neuroend. tumor by sex. For 953 of 953 cases diagnosed between 1998 and 2020 valid data could be obtained for this item.

Years	Sex					
	Male			Female		
	n	Cond. surv. % 2 yrs	5 yrs	n	Cond. surv. % 2 yrs	5 yrs
0	481	55.7	51.0	472	81.1	75.0
1	297	83.1	79.6	370	89.3	85.5
2	237	93.9	90.6	323	93.3	88.9
3	219	94.6	93.9	279	97.7	94.6
4	186	98.8	94.9	246	97.4	94.4
5	156	99.0	95.8	212	96.0	93.4
6	141	98.2	91.9	183	98.8	94.1
7	115	98.7	95.1	163	98.7	96.0
8	103	95.7	93.5	149	96.2	95.4
9	89	95.7	91.7	130	96.9	95.8
10	78	97.8	90.2	116	99.1	98.1

Table 2d. Conditional relative 2 / 5-year survival of patients with pulm. neuroend. tumor by sex for period 1998-2020 (N=953).

Conditional relative survival rates refer to the relative survival probability, in this case for 2 and 5 years after cancer diagnosis, compared to the age- and sex-matched population (=100 %) under the condition of being alive for a certain time period (x-axis in Figure 2a). The results illustrate to what extent the cancer induced mortality of particular subgroups declines in the subsequent years after detection of the malignancy. For instance, according to the presented survival statistics, patients in the subgroup sex="Male", who are alive at least 3 years after cancer diagnosis, the conditional relative 2-year survival rate is 94.6% (n=219).




Figure 3a. Relative survival of patients with pulm. neuroend. tumor by age category. Included in the evaluation are 953 cases diagnosed between 1998 and 2020.

Years	Age category							
	0-49 yrs n=154		50-59 yrs n=205		60-69 yrs n=300		70+ yrs n=294	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	94.5	94.2	75.2	75.4	71.8	72.3	65.3	67.6
2	91.6	91.6	66.7	67.5	61.8	63.3	56.9	61.2
3	90.1	90.4	63.5	64.2	58.7	60.6	50.9	57.1
4	88.3	88.7	61.2	62.5	56.2	58.9	48.3	56.4
5	87.3	87.7	60.5	62.3	54.5	57.8	44.7	55.2
Median			15.0		8.7		3.6	

Table 3b. Observed (obs.) and relative (rel.) survival of patients with pulm. neuroend. tumor by age category for period 1998-2020 (N=953).




Figure 4a. Relative survival of patients with pulm. neuroend. tumor by extent of disease. For 853 of 953 cases diagnosed between 1998 and 2020 valid data could be obtained for this item. For a total of 845 cases an evaluable classification was established. The grey line represents the subgroup of 108 patients with missing values regarding extent of disease (11.3 % of 953 patients, the percent values of all other categories are related to n=845).

Years	Extent of disease							
	Local n=419		Regional n=151		Distant n=275		NA/NOS n=108	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	96.3	97.2	70.8	71.3	39.7	40.2	82.6	83.5
2	92.8	95.2	57.7	59.8	26.8	27.8	76.6	79.5
3	90.9	94.4	50.1	52.6	21.8	22.6	74.5	78.1
4	89.9	94.6	44.3	47.0	19.7	21.2	70.4	75.4
5	88.2	94.3	41.2	45.4	19.1	20.8	66.2	72.9
Median			3.2		0.7		13.7	

Table 4b. Observed (obs.) and relative (rel.) survival of patients with pulm. neuroend. tumor by extent of disease for period 1998-2020 (N=845).




Figure 5a. Time to first progression of 674 patients with pulm. neuroend. tumor diagnosed between 1998 and 2020 (in solid cancers M0 only) estimated by cumulative incidence function (CI, solid line) accounting for death as competing risk and by inverse Kaplan-Meier estimate (1-KM, dashed line). The frequency of events may be underestimated due to underreporting.

	Type of progression	
	Any type (CI)	Any type (1-KM)
N	674	674
Events	172	172
compet.	40	
Years	%	%
0	0.0	0.0
1	17.5	17.7
2	21.8	22.2
3	24.2	24.6
4	26.4	27.0
5	27.2	27.8

Table 5b. Time to first progression of patients with pulm. neuroend. tumor for period 1998-2020 (N=674), also showing the total of progression events (Events) and of deaths as competing risk (compet.).




Figure 5c. Observed post-progression survival of 418 patients with pulm. neuroend. tumor diagnosed between 1998 and 2020. These 418 patients with documented progression events during their course of disease represent 44.2 % of the totally 946 evaluated cases (incl. M1, n=272, 28.8 %). Patients with cancer relapse documented via death certificates only were excluded (n=49, 5.2 %).

Medical record documentation often lacks the linguistic severity to distinguish between local relapse, regional lymph node metastasis and distant spread in solid cancers. Frequently, the statement “not specified” is the only information in registries regarding relapse of the disease. The category “Any type” denotes all cases who suffered from at least one relapse during the course of disease (incl. primary M1-status). Although, the real number of relapsed patients is likely to be much higher. The accumulated percentage of patients with local relapse or distant metastasis exceeds the 100 % value because patients are potentially considered in more than one subgroup.

Type of progression	
Any type n=418	
Years	%
0	100.0
1	46.5
2	33.4
3	27.5
4	24.4
5	23.2
6	20.2
7	19.6
8	18.9
9	18.9

Table 5d. Observed post-progression survival of patients with pulm. neuroend. tumor for period 1998-2020 (N=418).




Figure 5e. Observed post-progression (distant metastasis) survival of 368 patients with pulm. neuroend. tumor diagnosed between 1998 and 2020 by period of progression.

Years	Period of progression		
	1998-2006 n=36 %	2007-2015 n=190 %	2016+ n=142 %
0	100.0	100.0	100.0
1		35.8	51.1
2		23.2	36.0
3		17.9	26.9
4		14.1	
5		13.0	
6		11.9	

Table 5f. Observed post-progression (distant metastasis) survival of patients with pulm. neuroend. tumor for period 1998-2020 by period of progression (N=368).

Shortcuts

MCR Munich Cancer Registry, Germany

NCI National Cancer Institute, USA

SEER Surveillance, Epidemiology, and End Results, USA

UICC Union for International Cancer Control, Geneva

DCO Death certificate only Death certificate provides the only notification to the registry.

NA Not available

NOS Not otherwise specified

OS Overall/Observed survival Overall/Observed survival (Kaplan-Meier estimate)

Date of entry: diagnosis
Event: death from any cause

RS Relative survival Survival compared to “general population”, ratio of observed to expected survival (Ederer II method), reflecting cancer specific survival

AS Assembled survival Assembled chart of observed, expected, relative survival

CS Conditional survival Survival probability under the condition of surviving a given period of time

TTP Time to progression Time to first progression / relapse
Date of entry: diagnosis
Event: (progression / relapse): first local-, lymph node recurrence, distant metastasis or unspecified progression

1-KM 1 minus Kaplan-Meier estimator (“inverse” Kaplan-Meier estimator)

CI Cumulative incidence
Death as competing risk (according to Kalbfleisch und Prentice)

PPS Post-progression survival Survival since first progression / relapse (Kaplan-Meier estimate)
Date of entry (progression / relapse): first local-, lymph node recurrence, distant metastasis or unspecified progression
Event: death from any cause

Recommended Citation

Munich Cancer Registry. Survival BNET: Pulm. neuroend. tumor [Internet]. 2022 [updated 2022 Apr 27; cited 2022 Jun 1]. Available from: <https://www.tumorregister-muenchen.de/en/facts/surv/shBNETE-BNET-Pulm.-neuroend.-tumor-survival.pdf>

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.