

Munich Cancer Registry



- ▶ [Baseline statistics](#)
- ▶ [Selection Matrix](#)
- ▶ [Homepage](#)

Munich Cancer Registry at Munich Cancer Center
Marchioninstr. 15
Munich, 81377
Germany

<http://www.tumorregister-muenchen.de/en>

Cancer statistics: Survival

CLL: Chronic lymph. leukaemia (morph.)

Year of diagnosis	1978-1987	1988-1997	1998-2011
Patients	233	476	2,452
Diseases	233	476	2,455
Creation date	04/01/2013		
Export date	01/03/2013		
Population	4.5 m		



http://www.tumorregister-muenchen.de/en/facts/surv/surv_hCLL_E.pdf




Figure 1a. Relative survival of patients with chronic lymph. leukaemia (morph.) by period of diagnosis. Included in the evaluation are 1,572 cases diagnosed between 1978 and 2011.

The presented survival curves are derived from clinical records with valid follow-up informations, which means that death certificate cases (DCO) cases are omitted from the analysis. With this one restriction, the MCR has provided population-based statistics since 1998, collecting data on all cancer cases in the region of southern Bavaria. Historical data of previous time periods can be heavily selected, therefore, univariate survival comparisons of the presented time periods must be carefully considered. Nonetheless, all calculable survival curves are depicted to facilitate the comparison of long time follow-up analyses of relative survival between particular cancers.

Years	Period of diagnosis					
	1978-1987 n=211		1988-1997 n=405		1998+ n=956	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0
1	98.1	99.5	96.0	98.7	92.1	94.8
2	93.8	98.4	92.3	97.5	85.9	91.0
3	87.1	93.8	87.9	95.1	78.3	85.2
4	80.4	89.4	82.2	91.6	72.7	81.1
5	70.3	80.1	72.5	83.3	65.6	74.9
6	63.6	74.7	66.8	78.9	60.4	70.6
7	54.3	65.3	62.8	76.2	56.0	67.1
8	49.5	61.5	58.8	73.2	50.6	61.9
9	44.2	56.4	51.6	66.0	46.8	58.3
10	39.8	52.2	44.4	58.5	42.1	54.2

Table 1b. Observed (obs.) and relative (rel.) survival of patients with chronic lymph. leukaemia (morph.) by period of diagnosis for period 1978-2011 (N=1,572).




Figure 2a. Survival of patients with chronic lymph. leukaemia (morph.) by gender. Included in the evaluation are 1,361 cases diagnosed between 1988 and 2011.

Years	Gender			
	Male n=837		Female n=524	
	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0
1	93.9	96.8	92.2	94.7
2	88.8	94.4	86.3	90.7
3	82.5	90.1	79.7	85.9
4	75.5	84.9	76.3	83.8
5	67.9	78.8	67.2	75.4
6	62.4	74.5	62.2	71.4
7	57.9	71.0	58.7	68.8
8	53.5	67.4	54.2	64.8
9	47.8	61.9	48.3	59.1
10	41.2	54.9	42.7	53.8

Table 2b. Observed (obs.) and relative (rel.) survival of patients with chronic lymph. leukaemia (morph.) by gender for period 1988-2011 (N=1,361).




Figure 2c. Conditional relative 2 / 5-year survival of patients with chronic lymph. leukaemia (morph.) by gender. For 1,361 of 1,361 cases diagnosed between 1988 and 2011 valid data could be obtained for this item.

Years	Gender					
	n	Male		Female		
		Cond. surv. %	2 yrs	5 yrs	n	Cond. surv. %
0	837	94.4	78.8	524	90.7	75.4
1	755	93.1	77.2	462	90.8	75.6
2	672	90.3	75.8	410	92.5	76.1
3	576	87.7	75.5	347	88.2	76.0
4	478	88.1	73.8	301	85.4	71.2
5	380	90.9	70.8	245	91.6	72.3
6	306	91.3	65.7	205	91.3	69.2
7	262	88.0	59.7	177	86.6	68.0
8	222	82.1	58.2	146	83.5	63.7
9	185	78.3	61.1	116	82.7	65.9
10	143	76.3	66.0	90	86.5	73.3

Table 2d. Conditional relative 2 / 5-year survival of patients with chronic lymph. leukaemia (morph.) by gender for period 1988-2011 (N=1,361).

Conditional relative survival rates refer to the relative survival probability, in this case for 2 and 5 years after cancer diagnosis, compared to the age- and sex-matched population (=100%) under the condition of being alive for a certain time period (x-axis in Figure 2a). The results illustrate to what extent the cancer induced mortality of particular subgroups declines in the subsequent years after detection of the malignancy. For instance, according to the presented survival statistics, patients in the subgroup gender="Male", who are alive at least 3 years after cancer diagnosis, the conditional relative 2-year survival rate is 87.7% (n=576).




Figure 3a. Relative survival of patients with chronic lymph. leukaemia (morph.) by age category. Included in the evaluation are 1,361 cases diagnosed between 1988 and 2011.

Years	Age category					
	0-59 yrs n=374		60-74 yrs n=658		75+ yrs n=329	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0
1	97.0	97.4	96.0	97.9	83.3	89.8
2	95.1	96.1	92.1	96.0	70.1	81.8
3	92.4	94.0	85.6	91.1	58.9	74.4
4	87.6	89.7	79.9	87.2	52.2	70.8
5	83.6	86.4	70.1	78.4	41.5	62.5
6	81.7	84.9	63.9	73.4	32.6	54.1
7	78.5	82.5	59.5	70.4	27.5	50.0
8	75.1	79.3	54.9	67.1	21.2	42.8
9	68.1	72.8	50.2	63.4	14.4	32.3
10	61.7	66.9	44.5	58.2		

Table 3b. Observed (obs.) and relative (rel.) survival of patients with chronic lymph. leukaemia (morph.) by age category for period 1988-2011 (N=1,361).




Figure 5a. Observed post-progression survival of 646 patients with chronic lymph. leukaemia (morph.) diagnosed between 1978 and 2011 (incl. M1). These 646 patients with documented progression events during their course of disease represent 41.1% of the totally 1,572 evaluated cases. Patients with cancer relapse documented via death certificates only were excluded (n=111, 7.1%).

Medical record documentation often lacks the linguistic severity to distinguish between local relapse, regional lymph node metastasis and distant spread in solid cancers. Frequently, the statement “not specified” is the only information in registries regarding relapse of the disease. The category “All types” denotes all cases who suffered from at least one relapse during the course of disease (incl. primary M1-status). Although, the real number of relapsed patients is likely to be much higher. The accumulated percentage of patients with local relapse or distant metastasis exceeds the 100% value because patients are potentially considered in more than one subgroup.

Type of progression	
All types n=646	
Months	%
0	100.0
12	80.0
24	67.9
36	58.0
48	50.2
60	40.5

Table 5b. Observed post-progression survival of patients with chronic lymph. leukaemia (morph.) for period 1978-2011 (N=646).

Shortcuts

AS	Assembled survival chart (observed, expected, relative)
CS	Conditional survival
DCO	Death certificate only. The death certificate provides the only notification to the registry.
MCR	Munich Cancer Registry, Germany (Tumorregister München)
NA	Not available
NCI	National Cancer Institute, USA
NOS	Not otherwise specified
OS	Observed/overall survival (Kaplan-Meier estimate)
PPS	Post-progression survival
RS	Relative Survival. Ratio of observed and expected survival (derived from the normal population)
SEER	Surveillance, Epidemiology, and End Results, USA
TTP	Time to progression Date of entry: Date of diagnosis Event (Progression): First local recurrence, lymph node relapse or distant metastasis, unspecified recurrence First all-cause recurrence is illustrated by survival curves (Kaplan-Meier estimate). First local recurrence, lymph node relapse or distant metastasis are depicted cumulatively, where applicable ("reverse" Kaplan-Meier estimate).
UICC	Union for International Cancer Control, Geneva

Recommended Citation

Munich Cancer Registry. Survival CLL: Chronic lymph. leukaemia (morph.) [Internet]. 2013 [updated 2013 Apr 1; cited 2013 May 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/surv/surv_hCLL_E.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

Fig./Tbl.		Page
1a	Relative survival by period of diagnosis (chart)	2
1b	Survival by period of diagnosis (table)	2
2a	Survival by gender (chart)	3
2b	Survival by gender (table)	3
2c	Conditional survival by gender (chart)	4
2d	Conditional survival by gender (table)	4
3a	Relative survival by age category (chart)	5
3b	Survival by age category (table)	5
5a	Observed post-progression survival (chart)	6
5b	Observed post-progression survival (table)	6