Munich Cancer Registry



- ▶ Incidence and Mortality
- ▶ Selection Matrix
- ▶ Homepage
- ▶ Deutsch

ICD-10 C37: Thymus cancer

Survival

Year of diagnosis	1988-1997	1998-2014
Patients	22	205
Diseases	22	206
Cases evaluated	20	169
Creation date	03/02/2016	
Export date	12/23/2015	
Population	4.64 m	



Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

http://www.tumorregister-muenchen.de/en/facts/surv/sC37__E-ICD-10-C37-Thymus-cancer-survival.pdf




Figure 1a. Relative survival of the total cohort with thymus cancer. Included in the evaluation are 187 cases diagnosed between 1988 and 2014.

Group				
	Total cohort			
	n=187			
Years	obs. % rel. %			
0	100.0	100.0		
1	91.0	92.1		
2	79.7	81.8		
3	73.8	76.7		
4	70.2	73.8		
5	64.0	68.0		
6	58.6	63.4		
7	54.1	59.8		
8	52.5	58.4		
9	50.8	56.9		
10	45.1	52.5		

Table 1b. Observed (obs.) and relative (rel.) survival of the total cohort with thymus cancer for period 1988-2014 (N=187).




Figure 2a. Survival of patients with thymus cancer by gender. Included in the evaluation are 187 cases diagnosed between 1988 and 2014.

Gender						
Ma	ale	Fen	nale			
n=98		n=89				
obs. %	rel. %	obs. %	rel. %			
100.0	100.0	100.0	100.0			
91.5	92.2	90.6	91.5			
78.7	80.6	80.7	82.5			
73.8	76.5	74.0	75.9			
68.1	71.5	72.5	74.7			
63.4	67.9	64.6	67.8			
54.3	58.7	62.9	66.0			
49.7	55.8	58.5	62.8			
49.7	55.8	55.4	58.8			
49.7	55.8	52.0	55.4			
	Man= obs. % 100.0 91.5 78.7 73.8 68.1 63.4 54.3 49.7	Male n=98 obs. % rel. % 100.0 100.0 91.5 92.2 78.7 80.6 73.8 76.5 68.1 71.5 63.4 67.9 54.3 58.7 49.7 55.8	Male n=98 Fem n=			

Table 2b. Observed (obs.) and relative (rel.) survival of patients with thymus cancer by gender for period 1988-2014 (N=187).




Figure 5a. Time to first progression of 132 patients with thymus cancer diagnosed between 1998 and 2014 (M0 only in solid cancers) estimated by cumulative incidence function (CI, solid line) accounting for death as competing risk and by inverse Kaplan-Meier estimate (1-KM, dashed line). The frequency of events may be underestimated due to underreporting.

Type of progression						
	Any type (CI)	Any type (1-KM)				
	n=132	n=132				
Years	%	%				
0	0.0	0.0				
1	11.3	11.5				
2	21.6	22.3				
3	27.4	28.5				
4	29.4	30.7				
5	31.8	33.4				
6	34.9	37.3				
7	34.9	37.3				

Table 5b. Time to first progression of patients with thymus cancer for period 1998-2014 (N=132).






Figure 5c. Observed post-progression survival of 59 patients with thymus cancer diagnosed between 1998 and 2014. These 59 patients with documented progression events during their course of disease represent 35.5% of the totally 166 evaluated cases (incl. M1, n=34, 20.5%). Patients with cancer relapse documented via death certificates only were excluded (n=14, 8.4%). Multiple progression types on different sites are included in the evaluation even when not occuring synchronously. The NOS (not otherwise specified) class is included under the condition, that it is the one and only progression type during the course of disease. Subgroups with sample size <15 are dropped from the chart.

Medical record documentation often lacks the linguistic severity to distinguish between local relapse, regional lymph node metastasis and distant spread in solid cancers. Frequently, the statement "not specified" is the only information in registries regarding relapse of the disease. The category "Any type" denotes all cases who suffered from at least one relapse during the course of disease (incl. primary M1-status). Although, the real number of relapsed patients is likely to be much higher. The accumulated percentage of patients with local relapse or distant metastasis exceeds the 100% value because patients are potientially considered in more than one subgroup.

Type of progression				
	Any type	Distant metastasis	Locoregional relapse	NOS
	n=59	n=50	n=13	n=5
Years	%	%	%	%
0	100.0	100.0	100.0	100.0
1	84.3	83.6		
2	58.3	53.1		
3	48.2	45.7		
4	39.0			

Table 5d. Observed post-progression survival of patients with thymus cancer for period 1998-2014 (N=59).

Shortcuts

MCR	Munich Cancer Registry, Germany				
NCI	National Cancer Institute, USA				
SEER	Surveillance, Epidemiology, and End Results, USA				
UICC	Union for International Cancer Control, Geneva				
DCO	Death certificate only	Death certificate provides the only notification to the registry.			
NA	Not available				
NOS	Not otherwise specified				
os	Overall/Observed survival	Overall/Observed survival (Kaplan-Meier estimate) Date of entry: diagnosis Event: death from any cause			
RS	Relative survival	Survival compared to "general population", ratio of observed to expected survival (Ederer II method), reflecting cancer specific survival			
AS	Assembled survival	Assembled chart of observed, expected, relative survival			
CS	Conditional survival	Survival probability under the condition of surviving a given period of time			
TTP	Time to progression	Time to first progression / relapse Date of entry: diagnosis Event: (progression / relapse): first local-, lymph node recurrence, distant metastasis or unspecified progression			
	1-KM	1 minus Kaplan-Meier estimator ("inverse" Kaplan-Meier estimator)			
	CI	Cumulative incidence Death as competing risk (according to Kalbfleisch und Prentice)			
PPS	Post-progression survival	Survival since first progression / relapse (Kaplan-Meier estimate) Date of entry (progression / relapse): first local-, lymph node recurrence, distant metastasis or unspecified progression Event: death from any cause			

Recommended Citation

Munich Cancer Registry. Survival ICD-10 C37: Thymus cancer [Internet]. 2016 [updated 2016 Mar 2; cited 2016 May 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/surv/sC37__E-ICD-10-C37-Thymus-cancer-survival.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

F	ig./Tb	l.	Page
	1a	Relatives survival of total cohort (chart)	2
	1b	Survival of total cohort (table)	2
	2a	Survival by gender (chart)	3
	2b	Survival by gender (table)	3
	5a	Time to first progression (chart)	4
	5b	Time to first progression (table)	4
	5c	Observed post-progression survival (chart)	5
	5d	Observed post-progression survival (table)	5

