

# Munich Cancer Registry



- Incidence and Mortality
- Selection Matrix
- Homepage
- *Deutsch*

## ICD-10 C32, C10.1: Larynx cancer

### Survival


Year of diagnosis	1988-1997	1998-2014
Patients	827	2,295
Diseases	827	2,298
Cases evaluated	757	1,746
Creation date	04/11/2016	
Export date	12/23/2015	
Population	4.64 m	



Munich Cancer Registry at Munich Cancer Center  
Marchioninstr. 15  
Munich, 81377  
Germany

<http://www.tumorregister-muenchen.de/en>

[http://www.tumorregister-muenchen.de/en/facts/surv/sC32\\_\\_E-ICD-10-C32-C10.1-Larynx-cancer-survival.pdf](http://www.tumorregister-muenchen.de/en/facts/surv/sC32__E-ICD-10-C32-C10.1-Larynx-cancer-survival.pdf)




**Figure 1a.** Relative survival of patients with larynx cancer by period of diagnosis. Included in the evaluation are 2,492 cases diagnosed between 1988 and 2014.

The survival results of the SEER program (Surveillance, Epidemiology, and End Results) of the American National Cancer Institute (NCI) are summarized as the period of diagnosis from 1998 to 2011, and are represented by gray diamonds in order to facilitate comparisons between MCR and SEER.

The presented survival curves are derived from clinical records with valid follow-up informations, which means that death certificate cases (DCO) cases are omitted from the analysis. With this one restriction, the MCR has provided population-based statistics since 1998, collecting data on all cancer cases in the region of southern Bavaria. Historical data of previous time periods can be heavily selected, therefore, univariate survival comparisons of the presented time periods must be carefully considered. Nonetheless, all calculable survival curves are depicted to facilitate the comparison of long time follow-up analyses of relative survival between particular cancers.

Years	Period of diagnosis					
	1988-1997 n=754		1998-2006 n=905		2007+ n=833	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0
1	88.3	90.4	87.5	89.2	85.3	87.1
2	77.4	81.2	77.8	81.0	74.5	77.6
3	69.9	75.0	69.8	74.1	67.0	71.5
4	63.5	69.8	65.4	71.0	62.9	68.6
5	58.4	65.7	60.8	67.5	58.4	65.1
6	52.9	61.2	56.1	63.7	55.6	63.9
7	49.5	58.7	51.4	59.7	51.1	59.4
8	46.4	56.2	48.4	57.8		
9	43.2	53.5	45.3	55.2		
10	41.3	52.6	42.9	53.7		
11	38.3	50.0	40.0	51.4		
12	35.9	48.0	37.1	48.7		
13	32.6	44.7	35.2	47.5		
14	29.9	42.1	33.9	47.0		
15	27.9	40.4	29.3	41.6		
16	24.9	37.1				
17	22.5	34.5				
18	20.4	32.2				
19	18.9	30.8				
20	18.1	30.1				


**Table 1b.** Observed (obs.) and relative (rel.) survival of patients with larynx cancer by period of diagnosis for period 1988-2014 (N=2,492).



**Figure 2a.** Survival of patients with larynx cancer by gender. Included in the evaluation are 2,492 cases diagnosed between 1988 and 2014.

Years	Gender			
	Male n=2,177		Female n=315	
	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0
1	87.0	88.9	87.2	88.5
2	76.5	80.0	76.9	79.3
3	69.0	73.8	68.6	72.0
4	63.7	69.8	65.5	69.7
5	59.0	66.2	60.2	65.1
6	54.5	62.8	55.1	60.8
7	50.5	59.6	50.3	56.0
8	47.3	57.3	48.8	55.1
9	44.3	54.8	44.6	51.6
10	42.0	53.5	44.0	51.4
11	39.0	51.1	40.5	48.6
12	36.5	48.9	37.3	45.7
13	33.5	46.3	34.4	43.3
14	31.0	43.9	34.4	43.3
15	28.2	41.2	33.3	42.7
16	25.0	37.6	30.6	40.4
17	22.7	35.2	27.2	37.3
18	20.3	32.6	27.2	36.9
19	18.7	30.9	27.2	36.5
20	17.8	30.0	27.2	36.1


**Table 2b.** Observed (obs.) and relative (rel.) survival of patients with larynx cancer by gender for period 1988-2014 (N=2,492).



**Figure 3a.** Relative survival of patients with larynx cancer by age category. Included in the evaluation are 2,492 cases diagnosed between 1988 and 2014.

Years	Age category							
	0-49 yrs n=290		50-59 yrs n=737		60-69 yrs n=861		70+ yrs n=604	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	93.0	93.0	89.7	90.3	86.9	88.4	81.0	85.6
2	84.5	85.0	78.6	79.7	76.8	79.6	69.9	78.2
3	78.7	79.2	71.4	73.2	69.0	72.9	60.9	72.6
4	74.4	75.4	68.5	70.8	63.5	68.4	53.6	68.2
5	70.0	71.0	64.4	67.1	58.1	64.0	48.7	65.7
6	67.0	68.4	60.3	63.5	53.4	60.4	42.5	62.7
7	64.7	66.3	56.8	60.5	49.4	57.2	36.5	58.3
8	62.7	64.3	53.6	57.9	47.1	56.1	31.8	54.8
9	62.2	64.3	51.3	56.1	42.7	52.4	27.7	50.9
10	61.6	64.0	49.9	55.4	40.4	51.4	23.5	47.5
11	58.6	61.7	47.2	53.1	37.2	49.3	20.0	44.3
12	57.3	60.6	44.5	50.8	34.5	47.5	16.6	40.1
13	55.3	58.4	41.4	48.2	31.5	45.1	13.1	35.9
14	51.7	55.1	39.3	46.5	29.1	43.9	11.4	33.8
15	50.0	54.1	35.4	42.7	26.5	41.9	10.1	32.4
16	48.1	52.5	31.2	38.8	22.4	37.4		
17	48.1	52.4	27.9	35.1	18.7	33.5		
18	47.1	51.8	25.6	32.9	15.4	29.5		
19	44.9	50.1	25.0	32.8	13.0	26.3		
20	43.8	48.9	25.0	32.8	11.2	24.1		


**Table 3b.** Observed (obs.) and relative (rel.) survival of patients with larynx cancer by age category for period 1988-2014 (N=2,492).



**Figure 4a.** Relative survival of patients with larynx cancer by TNM staging. For 2,292 of 2,492 cases diagnosed between 1988 and 2014 valid data could be obtained for this item. For a total of 2,283 cases an evaluable classification was established. The accumulated percentage exceeds the 100% value because patients are potentially considered in more than one subgroup. The grey line represents the subgroup of 209 patients with missing values regarding TNM staging (8.4 % of 2,492 patients, the percent values of all other categories are related to n=2,283).

TNM staging														
	T1 n=894		T2 n=490		T3+4 n=876		N0 n=1,485		N+ n=601		NX n=104		NA/NOS n=209	
Years	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	96.8	99.0	93.1	94.9	76.6	78.1	92.7	94.8	77.3	78.5	82.4	84.3	74.5	76.1
2	91.6	96.0	82.9	86.2	59.2	61.5	85.4	89.4	55.6	57.5	75.4	78.6	69.5	73.3
3	86.9	93.3	73.2	77.8	49.5	52.4	79.8	85.5	41.8	44.0	70.8	75.3	61.7	66.9
4	82.5	90.9	68.7	74.5	43.8	47.2	74.9	82.0	36.4	38.8	67.3	72.6	55.3	61.3
5	78.3	88.5	62.7	69.2	39.3	43.0	70.4	78.9	31.9	34.4	63.5	70.3	49.7	56.9
6	72.5	84.3	57.2	64.5	36.2	40.5	65.2	75.1	28.4	31.3	62.1	70.0	45.0	52.7
7	68.5	81.7	52.0	59.9	32.9	37.3	60.7	71.5	24.9	27.9	58.2	67.9	41.6	50.9
8	65.4	80.1	47.2	55.6	30.3	35.1	56.6	68.3	23.7	26.8	58.2	67.9	40.8	50.5
9	61.8	77.3	44.3	53.4	27.4	32.4	53.3	65.7	20.6	23.8	52.1	62.8	39.2	50.4
10	59.2	76.4	42.1	52.0	25.4	30.6	50.8	64.5	19.5	22.8			38.4	50.4


**Table 4b.** Observed (obs.) and relative (rel.) survival of patients with larynx cancer by TNM staging for period 1988-2014 (N=2,283).



**Figure 4c.** Relative survival of patients with larynx cancer by TNM staging. For 1,596 of 1,738 cases diagnosed between 1998 and 2014 valid data could be obtained for this item. For a total of 1,588 cases an evaluable classification was established. The accumulated percentage exceeds the 100% value because patients are potentially considered in more than one subgroup. The grey line represents the subgroup of 150 patients with missing values regarding TNM staging (8.6 % of 1,738 patients, the percent values of all other categories are related to n=1,588).

TNM staging														
Years	T1 n=636		T2 n=349		T3+4 n=587		N0 n=998		N+ n=419		NX n=93		NA/NOS n=150	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	97.0	98.8	93.8	95.5	75.1	76.5	93.0	94.9	77.1	78.3	81.4	83.1	69.1	70.7
2	92.2	96.4	83.7	86.3	57.2	59.3	86.0	89.7	55.8	57.5	74.6	77.8	64.2	67.4
3	87.3	93.4	73.3	77.6	47.7	50.4	80.2	85.6	41.3	43.4	72.0	76.5	56.7	61.1
4	83.5	91.6	69.2	74.6	42.9	46.1	75.3	82.2	38.0	40.4	69.3	74.8	51.2	56.3
5	79.1	89.1	64.5	70.6	38.3	41.8	71.3	79.6	32.6	35.1	65.0	72.1	46.3	52.4
6	74.1	85.9	58.1	64.8	36.3	40.4	66.4	75.8	30.0	33.2	63.4	71.5	42.7	49.5
7	70.2	83.3	51.7	58.7	32.0	36.0	61.3	71.7	25.3	28.3	58.8	68.8	40.5	49.3
8	66.6	81.5	48.3	56.2	29.7	34.2	57.7	69.3	23.1	26.4	58.8	68.6	39.0	49.0
9	62.3	77.9	44.9	53.4	27.3	31.9	54.4	66.7	20.0	22.9			39.0	49.0
10	59.2	76.4	43.1	51.9	25.1	29.6	52.0	65.4	19.3	21.8			37.4	48.5

**Table 4d.** Observed (obs.) and relative (rel.) survival of patients with larynx cancer by TNM staging for period 1998-2014 (N=1,588).




**Figure 4e.** Conditional relative 2-year survival of patients with larynx cancer by extent of disease. For 2,292 of 2,492 cases diagnosed between 1988 and 2014 valid data could be obtained for this item. For a total of 2,282 cases an evaluable classification was established. The grey line represents the subgroup of 210 patients with missing values regarding extent of disease (8.4% of 2,492 patients, the percent values of all other categories are related to n=2,282).

Years	Extent of disease							
	Local		Regional		Distant		NA/NOS	
	n	Cond. surv. % 2 yrs	n	Cond. surv. % 2 yrs	n	Cond. surv. % 2 yrs	n	Cond. surv. % 2 yrs
0	1,661	89.0	566	60.3	55		210	75.5
1	1,494	90.7	443	56.9			157	87.4
2	1,321	92.6	304	68.2			142	84.7
3	1,173	92.9	212	78.2			122	86.4
4	1,042	91.9	173	80.4			107	86.9
5	921	91.5	140	80.5			92	90.4

**Table 4f.** Conditional relative 2-year survival of patients with larynx cancer by extent of disease for period 1988-2014 (N=2,282).

Conditional relative survival rates refer to the relative survival probability, in this case for 2 years after cancer diagnosis, compared to the age- and sex-matched population (=100%) under the condition of being alive for a certain time period (x-axis in Figure 4c). The results illustrate to what extent the cancer induced mortality of particular subgroups declines in the subsequent years after detection of the malignancy. For instance, according to the presented survival statistics, patients in the subgroup extent of disease="Local", who are alive at least 3 years after cancer diagnosis, the conditional relative 2-year survival rate is 92.9% (n=1,173).




**Figure 5a.** Time to first progression of 1,685 patients with larynx cancer diagnosed between 1998 and 2014 (M0 only in solid cancers) estimated by cumulative incidence function (CI, solid line) accounting for death as competing risk and by inverse Kaplan-Meier estimate (1-KM, dashed line). The frequency of events may be underestimated due to underreporting.

Years	Type of progression						
	Any type (CI)	Any type (1-KM)	Local relapse (CI)	Local relapse (1-KM)	Distant metastasis (CI)	Distant metast. (1-KM)	Locoreg. relapse (CI)
	n=1,685 %	n=1,685 %	n=1,685 %	n=1,685 %	n=1,685 %	n=1,685 %	n=1,685 %
0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1	18.3	18.9	6.8	7.5	3.7	4.1	7.7
2	28.0	29.5	10.8	12.3	8.3	9.5	12.5
3	31.6	33.7	12.3	14.4	10.0	11.8	14.0
4	33.2	35.7	13.7	16.5	10.7	12.8	15.4
5	34.7	37.6	14.1	17.2	11.4	13.8	15.9
6	36.2	39.7	14.9	18.6	11.9	14.7	16.8
7	37.5	41.7	15.7	20.0	12.2	15.2	17.9
8	38.6	43.2	15.9	20.4	12.8	16.2	18.1
9	39.3	44.5	16.4	21.6	12.9	16.5	18.7
10	40.1	45.8	17.1	23.0	12.9	16.5	19.3
11	40.4	46.4	17.2	23.3	13.1	16.8	19.5
12	40.4	46.4	17.2	23.3	13.3	17.3	19.5
13	40.7	47.0	17.5	24.2	13.3	17.3	19.8
14	40.7	47.0	17.5	24.2	13.3	17.3	19.8
15	41.2	48.2	17.5	24.2	13.3	17.3	19.8

<i>cont'd</i> Years	Type of progression		
	Locoreg. rel. (1-KM)	NOS (CI)	NOS (1-KM)
	n=1,685 %	n=1,685 %	n=1,685 %
0	0.0	0.0	0.0
1	8.4	3.3	3.6
2	14.2	4.2	4.8
3	16.3	4.9	5.9
4	18.4	5.1	6.3
5	19.3	5.3	6.6
6	20.8	5.5	6.9
7	22.7	5.6	7.1
8	23.1	5.7	7.4
9	24.5	5.7	7.4
10	25.8	5.7	7.4
11	26.2	5.7	7.4
12	26.2	5.7	7.4
13	27.0	5.7	7.4
14	27.0	5.7	7.4
15	27.0	5.7	7.4

**Table 5b.** Time to first progression of patients with larynx cancer for period 1998-2014 (N=1,685).




**Figure 5c.** Observed post-progression survival of 462 patients with larynx cancer diagnosed between 1998 and 2014. These 462 patients with documented progression events during their course of disease represent 26.8 % of the totally 1,727 evaluated cases (incl. M1, n=42, 2.4 %). Patients with cancer relapse documented via death certificates only were excluded (n=191, 11.1 %). Multiple progression types on different sites are included in the evaluation even when not occurring synchronously. The NOS (not otherwise specified) class is included under the condition, that it is the one and only progression type during the course of disease.

Medical record documentation often lacks the linguistic severity to distinguish between local relapse, regional lymph node metastasis and distant spread in solid cancers. Frequently, the statement “not specified” is the only information in registries regarding relapse of the disease. The category “Any type” denotes all cases who suffered from at least one relapse during the course of disease (incl. primary M1-status). Although, the real number of relapsed patients is likely to be much higher. The accumulated percentage of patients with local relapse or distant metastasis exceeds the 100% value because patients are potentially considered in more than one subgroup.

Years	Type of progression				
	Any type n=462 %	Local relapse n=253 %	Distant metastasis n=161 %	Locoregional relapse n=289 %	NOS n=89 %
0	100.0	100.0	100.0	100.0	100.0
1	62.1	72.6	34.7	71.0	63.9
2	49.7	60.1	21.9	57.6	53.6
3	43.4	55.2	15.5	52.4	46.1
4	39.2	48.0	14.6	46.2	44.7
5	35.6	44.9		42.9	38.7
6	32.1	41.3		39.2	34.8
7	30.6	39.5		36.8	34.8
8	27.2	33.3		31.3	34.8
9	25.3	31.0		29.3	
10	23.9	28.5		27.0	

**Table 5d.** Observed post-progression survival of patients with larynx cancer for period 1998-2014 (N=462).



**Figure 5e.** Observed post-progression (distant metastasis) survival of 227 patients with larynx cancer diagnosed between 1988 and 2014 by period of progression.

Years	Period of progression		
	1988-1997 n=47	1998-2006 n=77	2007+ n=103
0	100.0	100.0	100.0
1		28.9	39.6
2			26.8

**Table 5f.** Observed post-progression (distant metastasis) survival of patients with larynx cancer for period 1988-2014 by period of progression (N=227).

## Shortcuts

MCR Munich Cancer Registry, Germany

NCI National Cancer Institute, USA

SEER Surveillance, Epidemiology, and End Results, USA

UICC Union for International Cancer Control, Geneva

DCO Death certificate only Death certificate provides the only notification to the registry.

NA Not available

NOS Not otherwise specified

OS Overall/Observed survival Overall/Observed survival (Kaplan-Meier estimate)

Date of entry: diagnosis  
Event: death from any cause

RS Relative survival Survival compared to “general population”, ratio of observed to expected survival (Ederer II method), reflecting cancer specific survival

AS Assembled survival Assembled chart of observed, expected, relative survival

CS Conditional survival Survival probability under the condition of surviving a given period of time

TTP Time to progression Time to first progression / relapse  
Date of entry: diagnosis  
Event: (progression / relapse): first local-, lymph node recurrence, distant metastasis or unspecified progression

1-KM 1 minus Kaplan-Meier estimator  
 (“inverse” Kaplan-Meier estimator)

CI Cumulative incidence  
Death as competing risk (according to Kalbfleisch und Prentice)

PPS Post-progression survival Survival since first progression / relapse (Kaplan-Meier estimate)  
Date of entry (progression / relapse): first local-, lymph node recurrence, distant metastasis or unspecified progression  
Event: death from any cause

## Recommended Citation

Munich Cancer Registry. Survival ICD-10 C32, C10.1: Larynx cancer [Internet]. 2016 [updated 2016 Apr 11; cited 2016 Jun 1]. Available from: [http://www.tumorregister-muenchen.de/en/facts/surv/sC32\\_\\_E-ICD-10-C32-C10.1-Larynx-cancer-survival.pdf](http://www.tumorregister-muenchen.de/en/facts/surv/sC32__E-ICD-10-C32-C10.1-Larynx-cancer-survival.pdf)

## Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

## Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

## Index of figures and tables

Fig./Tbl.		Page
1a	Relative survival by period of diagnosis (chart)	2
1b	Survival by period of diagnosis (table)	3
2a	Survival by gender (chart)	4
2b	Survival by gender (table)	4
3a	Relative survival by age category (chart)	5
3b	Survival by age category (table)	5
4a	Relative survival by TNM staging 1988+ (chart)	6
4b	Survival by TNM staging 1988+ (table)	6
4c	Relative survival by TNM staging 1998+ (chart)	7
4d	Survival by TNM staging 1998+ (table)	7
4e	Conditional survival by extent of disease (chart)	8
4f	Conditional survival by extent of disease (table)	8
5a	Time to first progression (chart)	9
5b	Time to first progression (table)	9
5c	Observed post-progression survival (chart)	11
5d	Observed post-progression survival (table)	11
5e	Observed post-progression survival by period of progression (chart)	12
5f	Observed post-progression survival by period of progression (table)	12