

Munich Cancer Registry



- ▶ Incidence and Mortality
- ▶ Selection Matrix
- ▶ Homepage
- ▶ *Deutsch*

ICD-10 C62: Testicular cancer

Survival

Year of diagnosis	1988-1997	1998-2014
Patients	935	3,262
Diseases	950	3,330
Cases evaluated	905	3,073
Creation date	04/11/2016	
Export date	12/23/2015	
Population (males)	2.28 m	



Munich Cancer Registry at Munich Cancer Center
Marchioninstr. 15
Munich, 81377
Germany

<http://www.tumorregister-muenchen.de/en>

http://www.tumorregister-muenchen.de/en/facts/surv/sC62__E-ICD-10-C62-Testicular-cancer-survival.pdf




Figure 1a. Relative survival of patients with testicular cancer by period of diagnosis. Included in the evaluation are 3,958 cases diagnosed between 1988 and 2014.

The survival results of the SEER program (Surveillance, Epidemiology, and End Results) of the American National Cancer Institute (NCI) are summarized as the period of diagnosis from 1998 to 2011, and are represented by gray diamonds in order to facilitate comparisons between MCR and SEER.

The presented survival curves are derived from clinical records with valid follow-up informations, which means that death certificate cases (DCO) cases are omitted from the analysis. With this one restriction, the MCR has provided population-based statistics since 1998, collecting data on all cancer cases in the region of southern Bavaria. Historical data of previous time periods can be heavily selected, therefore, univariate survival comparisons of the presented time periods must be carefully considered. Nonetheless, all calculable survival curves are depicted to facilitate the comparison of long time follow-up analyses of relative survival between particular cancers.

Years	Period of diagnosis					
	1988-1997 n=905		1998-2006 n=1,461		2007+ n=1,592	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0
1	98.4	98.7	98.1	98.3	98.6	98.8
2	97.4	97.9	97.3	97.8	97.8	98.2
3	96.8	97.6	96.7	97.4	96.7	97.4
4	96.2	97.3	96.3	97.3	96.4	97.2
5	95.7	97.0	95.9	97.2	96.2	97.2
6	95.6	97.0	95.7	97.2	95.4	96.7
7	94.9	97.0	95.3	97.2	94.9	96.2
8	94.5	96.9	94.8	97.1		
9	94.4	96.9	94.5	97.1		
10	93.6	96.7	94.1	97.0		
11	93.5	96.7	93.7	97.0		
12	93.4	96.7	93.7	97.0		
13	93.0	96.7	93.3	97.0		
14	92.6	96.7	92.7	97.0		
15	92.3	96.7	91.9	96.9		
16	91.6	96.7	91.9	96.6		
17	91.3	96.7				
18	90.5	96.7				
19	89.3	96.5				
20	88.9	96.4				

Table 1b. Observed (obs.) and relative (rel.) survival of patients with testicular cancer by period of diagnosis for period 1988-2014 (N=3,958).




Figure 2a. Observed, expected and relative survival of the total cohort with testicular cancer. Included in the evaluation are 3,958 cases diagnosed between 1988 and 2014.

Years	Group	
	Total cohort n=3,958	
	obs. %	rel. %
0	100.0	100.0
1	98.4	98.6
2	97.5	98.0
3	96.7	97.5
4	96.3	97.3
5	96.0	97.2
6	95.6	97.2
7	95.2	97.1
8	94.7	96.9
9	94.5	96.9
10	93.9	96.8
11	93.7	96.8
12	93.6	96.8
13	93.2	96.8
14	92.7	96.8
15	92.3	96.8
16	91.7	96.8
17	91.3	96.8
18	90.5	96.8
19	89.3	96.3
20	88.9	96.3

Table 2b. Observed (obs.) and relative (rel.) survival of the total cohort with testicular cancer for period 1988-2014 (N=3,958).




Figure 3a. Relative survival of patients with testicular cancer by age category. Included in the evaluation are 3,958 cases diagnosed between 1988 and 2014.

Years	Age category					
	0-29 yrs n=1,115		30-49 yrs n=2,415		50+ yrs n=428	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0
1	99.2	99.2	98.6	98.8	94.6	95.7
2	98.3	98.4	98.1	98.4	91.9	94.2
3	98.0	98.2	97.5	98.0	89.3	92.9
4	97.6	97.9	97.2	97.8	88.0	92.5
5	97.2	97.5	96.9	97.7	87.3	92.5
6	96.8	97.2	96.6	97.7	86.9	92.5
7	96.8	97.2	96.3	97.7	84.0	92.5
8	96.3	96.9	96.1	97.6	82.1	92.4
9	96.3	96.8	96.1	97.6	80.0	91.6
10	96.1	96.8	95.5	97.6	78.3	91.0
11	96.1	96.8	95.2	97.6	77.6	91.0
12	96.1	96.8	95.1	97.6	77.6	91.0
13	96.1	96.8	94.8	97.6	74.6	91.0
14	96.1	96.8	94.5	97.6	71.4	90.2
15	95.8	96.8	94.2	97.6	70.3	90.1
16	95.8	96.8	93.4	97.6	68.9	90.1
17	95.8	96.8	92.7	97.6	68.9	90.1
18	95.8	96.8	91.7	97.4	67.1	90.1
19	94.7	96.3	91.0	97.4	61.1	89.1
20	94.2	96.2	90.7	97.4	61.1	89.1

Table 3b. Observed (obs.) and relative (rel.) survival of patients with testicular cancer by age category for period 1988-2014 (N=3,958).




Figure 4a. Relative survival of patients with testicular cancer by NM staging. For 3,833 of 3,958 cases diagnosed between 1988 and 2014 valid data could be obtained for this item. For a total of 3,832 cases an evaluable classification was established. The grey line represents the subgroup of 126 patients with missing values regarding NM staging (3.2% of 3,958 patients, the percent values of all other categories are related to n=3,832).

Years	NM staging							
	NOM0 n=3,025		N+M0 n=482		M1 n=325		NA/NOS n=126	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	99.5	99.7	99.1	99.3	89.2	89.3	92.5	93.0
2	99.1	99.6	97.8	98.0	85.2	85.5	89.8	90.8
3	98.6	99.4	97.1	97.5	83.0	83.6	87.0	89.4
4	98.2	99.2	96.8	97.4	82.6	83.2	86.1	88.6
5	97.9	99.1	96.0	96.8	82.2	82.9	86.1	88.6
6	97.7	99.1	95.4	96.4	81.8	82.7	85.0	88.6
7	97.1	99.0	95.1	96.4	81.8	82.6	85.0	88.6
8	96.7	98.9	94.8	96.3	80.7	82.1	83.7	88.6
9	96.5	98.9	94.8	96.3	80.7	81.8	82.4	88.0
10	96.0	98.9	94.4	96.3	79.2	80.7	81.1	87.8
11	95.7	98.9	94.4	96.3	78.5	80.5	81.1	87.8
12	95.6	98.9	94.4	96.3	78.5	80.4	81.1	87.8
13	95.1	98.9	94.4	96.3	78.5	80.2	81.1	87.8
14	94.7	98.9	94.4	96.3	77.3	79.7	79.2	87.8
15	94.4	98.9	94.4	96.3	74.8	77.7	79.2	87.8
16	94.1	98.9	92.9	96.3	74.8	77.6	77.0	85.9
17	93.7	98.9	92.9	96.3	74.8	77.5	74.8	84.4
18	93.2	98.9	90.9	95.5	74.8	77.4	72.3	82.4
19	92.2	98.9	88.7	93.4	74.8	77.2	69.0	80.4
20	92.2	98.9	87.6	93.1	72.6	75.2	69.0	78.6

Table 4b. Observed (obs.) and relative (rel.) survival of patients with testicular cancer by NM staging for period 1988-2014 (N=3,832).




Figure 4c. Relative survival of patients with testicular cancer by NM staging. For 2,972 of 3,053 cases diagnosed between 1998 and 2014 valid data could be obtained for this item. The grey line represents the subgroup of 81 patients with missing values regarding NM staging (2.7% of 3,053 patients, the percent values of all other categories are related to n=2,972).

Years	NM staging							
	NOMO n=2,382		N+M0 n=342		M1 n=248		NA/NOS n=81	
	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1	99.6	99.8	99.1	99.2	88.3	88.4	89.4	89.5
2	99.4	99.7	98.1	98.4	83.9	84.1	85.1	85.8
3	98.7	99.5	97.4	97.8	81.4	82.0	83.5	84.6
4	98.3	99.3	97.0	97.5	81.4	81.9	83.5	84.1
5	98.1	99.3	96.2	96.9	81.4	81.9	83.5	83.5

Table 4d. Observed (obs.) and relative (rel.) survival of patients with testicular cancer by NM staging for period 1998-2014 (N=2,972).




Figure 4e. Conditional relative 5-year survival of patients with testicular cancer by extent of disease. For 3,833 of 3,958 cases diagnosed between 1988 and 2014 valid data could be obtained for this item. For a total of 3,829 cases an evaluable classification was established. The grey line represents the subgroup of 129 patients with missing values regarding extent of disease (3.3% of 3,958 patients, the percent values of all other categories are related to n=3,829).

Extent of disease								
Years	Local		Regional		Distant		NA/NOS	
	n	Cond. surv. % 5 yrs	n	Cond. surv. % 5 yrs	n	Cond. surv. % 5 yrs	n	Cond. surv. % 5 yrs
0	3,022	99.2	482	96.8	325	82.9	129	89.5
1	2,756	99.6	450	97.1	275	92.5	110	96.3
2	2,542	99.4	419	98.2	242	96.5	103	98.7
3	2,332	99.6	394	98.7	223	98.3	95	101.2
4	2,137	99.9	379	98.9	207	98.1	88	100.5
5	1,951	99.9	339	99.7	191	97.2	83	100.0

Table 4f. Conditional relative 5-year survival of patients with testicular cancer by extent of disease for period 1988-2014 (N=3,829).

Conditional relative survival rates refer to the relative survival probability, in this case for 5 years after cancer diagnosis, compared to the age- and sex-matched population (=100%) under the condition of being alive for a certain time period (x-axis in Figure 4c). The results illustrate to what extent the cancer induced mortality of particular subgroups declines in the subsequent years after detection of the malignancy. For instance, according to the presented survival statistics, patients in the subgroup extent of disease="Local", who are alive at least 3 years after cancer diagnosis, the conditional relative 5-year survival rate is 99.6% (n=2,332).




Figure 5a. Time to first progression of 2,765 patients with testicular cancer diagnosed between 1998 and 2014 (M0 only in solid cancers) estimated by cumulative incidence function (CI, solid line) accounting for death as competing risk and by inverse Kaplan-Meier estimate (1-KM, dashed line). The frequency of events may be underestimated due to underreporting.

Years	Type of progression			
	Any type (CI)	Any type (1-KM)	Distant metastasis (CI)	Distant metastasis (1-KM)
	n=2,765 %	n=2,765 %	n=2,765 %	n=2,765 %
0	0.0	0.0	0.0	0.0
1	4.3	4.4	1.9	1.9
2	5.8	5.8	2.5	2.5
3	6.7	6.7	3.1	3.1
4	7.1	7.2	3.5	3.5
5	7.6	7.7	3.8	3.8
6	7.9	8.0	4.0	4.1
7	8.1	8.2	4.1	4.2
8	8.1	8.2	4.1	4.2
9	8.1	8.2	4.1	4.2
10	8.1	8.2	4.1	4.2
11	8.4	8.5	4.1	4.2
12	8.4	8.5	4.1	4.2
13	8.6	8.7	4.1	4.2
14	8.6	8.7	4.1	4.2
15	9.1	9.2	4.6	4.7

Table 5b. Time to first progression of patients with testicular cancer for period 1998-2014 (N=2,765).




Figure 5c. Observed post-progression survival of 431 patients with testicular cancer diagnosed between 1998 and 2014. These 431 patients with documented progression events during their course of disease represent 14.3 % of the totally 3,007 evaluated cases (incl. M1, n=242, 8.0 %). Patients with cancer relapse documented via death certificates only were excluded (n=13, 0.4 %). Multiple progression types on different sites are included in the evaluation even when not occurring synchronously.

Medical record documentation often lacks the linguistic severity to distinguish between local relapse, regional lymph node metastasis and distant spread in solid cancers. Frequently, the statement “not specified” is the only information in registries regarding relapse of the disease. The category “Any type” denotes all cases who suffered from at least one relapse during the course of disease (incl. primary M1-status). Although, the real number of relapsed patients is likely to be much higher. The accumulated percentage of patients with local relapse or distant metastasis exceeds the 100% value because patients are potentially considered in more than one subgroup.

Years	Type of progression	
	Any type n=431 %	Distant metastasis n=330 %
0	100.0	100.0
1	90.3	88.4
2	87.2	84.1
3	84.2	81.1
4	83.9	81.1
5	83.9	81.1
6	83.5	80.6
7	83.0	80.0
8	81.4	78.7
9	81.4	78.7
10	81.4	78.7

Table 5d. Observed post-progression survival of patients with testicular cancer for period 1998-2014 (N=431).




Figure 5e. Observed post-progression (distant metastasis) survival of 458 patients with testicular cancer diagnosed between 1988 and 2014 by period of progression.

Years	Period of progression		
	1988-1997 n=111	1998-2006 n=159	2007+ n=188
	%	%	%
0	100.0	100.0	100.0
1	91.8	87.3	89.9
2	87.1	82.1	85.0
3	85.3	80.8	80.4
4	84.3	80.8	80.4
5	83.2	80.8	80.4
6	83.2	80.8	78.9
7	83.2	80.1	76.8
8	83.2	78.0	76.8
9	83.2	78.0	
10	81.2	78.0	

Table 5f. Observed post-progression (distant metastasis) survival of patients with testicular cancer for period 1988-2014 by period of progression (N=458).

Shortcuts

MCR Munich Cancer Registry, Germany

NCI National Cancer Institute, USA

SEER Surveillance, Epidemiology, and End Results, USA

UICC Union for International Cancer Control, Geneva

DCO Death certificate only Death certificate provides the only notification to the registry.

NA Not available

NOS Not otherwise specified

OS Overall/Observed survival Overall/Observed survival (Kaplan-Meier estimate)

Date of entry: diagnosis
Event: death from any cause

RS Relative survival Survival compared to “general population”, ratio of observed to expected survival (Ederer II method), reflecting cancer specific survival

AS Assembled survival Assembled chart of observed, expected, relative survival

CS Conditional survival Survival probability under the condition of surviving a given period of time

TTP Time to progression Time to first progression / relapse
Date of entry: diagnosis
Event: (progression / relapse): first local-, lymph node recurrence, distant metastasis or unspecified progression

1-KM 1 minus Kaplan-Meier estimator
 (“inverse” Kaplan-Meier estimator)

CI Cumulative incidence
Death as competing risk (according to Kalbfleisch und Prentice)

PPS Post-progression survival Survival since first progression / relapse (Kaplan-Meier estimate)
Date of entry (progression / relapse): first local-, lymph node recurrence, distant metastasis or unspecified progression
Event: death from any cause

Recommended Citation

Munich Cancer Registry. Survival ICD-10 C62: Testicular cancer [Internet]. 2016 [updated 2016 Apr 11; cited 2016 Jun 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/surv/sC62__E-ICD-10-C62-Testicular-cancer-survival.pdf

Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

Index of figures and tables

Fig./Tbl.		Page
1a	Relative survival by period of diagnosis (chart)	2
1b	Survival by period of diagnosis (table)	3
2a	Survival of total cohort (chart)	4
2b	Survival of total cohort (table)	4
3a	Relative survival by age category (chart)	5
3b	Survival by age category (table)	5
4a	Relative survival by NM staging 1988+ (chart)	6
4b	Survival by NM staging 1988+ (table)	6
4c	Relative survival by NM staging 1998+ (chart)	7
4d	Survival by NM staging 1998+ (table)	7
4e	Conditional survival by extent of disease (chart)	8
4f	Conditional survival by extent of disease (table)	8
5a	Time to first progression (chart)	9
5b	Time to first progression (table)	9
5c	Observed post-progression survival (chart)	10
5d	Observed post-progression survival (table)	10
5e	Observed post-progression survival by period of progression (chart)	11
5f	Observed post-progression survival by period of progression (table)	11