# **Munich Cancer Registry**



- Incidence and Mortality
- Selection Matrix
- Homepage
- Deutsch

### ICD-10 C72: Spinal cord cancer

## Survival


Year of diagnosis	1998-2014
Patients	120
Diseases	120
Cases evaluated	108
Creation date	04/11/2016
Export date	12/23/2015
Population	4.64 m



Munich Cancer Registry at Munich Cancer Center Marchioninistr. 15 Munich, 81377 Germany

http://www.tumorregister-muenchen.de/en

 $http://www.tumorregister-muenchen.de/en/facts/surv/sC72\_E-ICD-10-C72-Spinal-cord-cancer-survival.pdf$ 




**Figure 1a.** Relative survival of the total cohort with spinal cord cancer. Included in the evaluation are 107 cases diagnosed between 1998 and 2013.

Group					
	Total cohort n=107				
Years	obs. %	rel. %			
0	100.0	100.0			
1	87.6	87.7			
2	80.3	81.0			
3	77.0	77.8			
4	73.2	74.8			
5	73.2	74.3			
6	73.2	73.9			
7	68.2	70.2			
8	64.2	67.2			

**Table 1b.** Observed (obs.) and relative (rel.) survival of the total cohort with spinal cord cancer for period 1998-2013 (N=107).





**Figure 1c.** Relative survival of patients with spinal cord cancer by year of diagnosis. Included in the evaluation are 107 cases diagnosed between 1998 and 2013.

The survival results of the SEER program (Surveillance, Epidemiology, and End Results) of the American National Cancer Institute (NCI) are summarized as the period of diagnosis from to, and are represented by gray diamonds in order to facilitate comparisons between MCR and SEER.

Year of diagnosis														
	19	98	1999 2000		20	01	2002		2003		2004			
	n=	=3	n=	=2	n=3		n=4 n=5		n=9		n=4			
Years	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1			100.0	69.6	100.0	92.3	100.0	90.0	80.0	77.9	77.8	75.3	100.0	89.8
2					100.0	84.6	100.0	79.9	80.0	73.3	77.8	70.4	100.0	79.7
3					100.0	76.9	75.0	73.9	80.0	68.7	77.8	65.4	75.0	73.5
4					100.0	69.2	75.0	70.4	80.0	64.1			75.0	68.1
5					100.0	61.5	75.0	66.8	80.0	59.5			75.0	62.8
6					100.0	53.9	75.0	63.3	80.0	54.9			75.0	57.4
7							75.0	59.7	80.0	50.3				
8							75.0	56.1						

Year of diagnosis														
cont'd	d 2005 2006		2007		2008 200		09 2010		10	2011				
	n=	n=2 n=2		n=9 n=9		n=13		n=7		n=13				
Years	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %	obs. %	rel. %
0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
1					100.0	93.0	88.9	83.5	92.3	90.3			92.3	88.2
2									92.3	88.1				
3									92.3	85.8				

Year of diagnosis								
cont'd	20	12	20	13				
	n=	12	n=	10				
Years	obs. %	rel. %	obs. %	rel. %				
0	100.0	100.0	100.0	100.0				
1	75.0	72.7						
2	66.7	63.2						

**Table 1d.** Observed (obs.) and relative (rel.) survival of patients with spinal cord cancer by year of diagnosis for period 1998-2013 (N=107).

#### Shortcuts

MCR	Munich Cancer Registry, Germany								
NCI	National Cancer Institute, USA								
SEER	Surveillance, Epidemiology, and End Results, USA								
UICC	Union for International Can	cer Control, Geneva							
DCO	Death certificate only	Death certificate provides the only notification to the registry.							
NA	Not available								
NOS	Not otherwise specified								
OS	Overall/Observed survival	Overall/Observed survival (Kaplan-Meier estimate) Date of entry: diagnosis Event: death from any cause							
RS	Relative survival	Survival compared to "general population", ratio of observed to expected survival (Ederer II method), reflecting cancer specific survival							
AS	Assembled survival	Assembled chart of observed, expected, relative survival							
CS	Conditional survival	Survival probability under the condition of surviving a given period of time							
TTP	Time to progression	Time to first progression / relapse Date of entry: diagnosis Event: (progression / relapse): first local-, lymph node recurrence, distant metastasis or unspecified progression							
	1-КМ	1 minus Kaplan-Meier estimator ("inverse" Kaplan-Meier estimator)							
	CI	Cumulative incidence Death as competing risk (according to Kalbfleisch und Prentice)							
PPS	Post-progression survival	Survival since first progression / relapse (Kaplan-Meier estimate) Date of entry (progression / relapse): first local-, lymph node recurrence, distant metastasis or unspecified progression Event: death from any cause							

#### **Recommended Citation**

Munich Cancer Registry. Survival ICD-10 C72: Spinal cord cancer [Internet]. 2016 [updated 2016 Apr 11; cited 2016 Jun 1]. Available from: http://www.tumorregister-muenchen.de/en/facts/surv/sC72\_E-ICD-10-C72-Spinal-cord-cancer-survival.pdf

#### Copyright

The content of the public web site provided by the Munich Cancer Registry is available worldwide and free of charge. All documents are free to download, utilize, copy, print-out and distribute, providing that the MCR is referenced.

#### Disclaimer

The Munich Cancer Registry reserves the right to not be responsible for the topicality, correctness, completeness or quality of the information provided. Liability claims regarding damage caused by the use of any information provided, including any kind of information which is incomplete or incorrect, will therefore be rejected.

### Index of figures and tables

Fig./Tb	ol.	Page
1a	Relatives survival of total cohort (chart)	2
1b	Survival of total cohort (table)	2
1c	Relative survival by year of diagnosis (chart)	3
1d	Survival by year of diagnosis (table)	3

MCR